



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The Kepler-19 System: A Thick-envelope Super-Earth with Two Neptune-mass Companions Characterized Using Radial Velocities and Transit Timing Variations

Citation for published version:

Malavolta, L, Borsato, L, Granata, V, Piotto, G, Lopez, E, Vanderburg, A, Figueira, P, Mortier, A, Nascimbeni, V, Affer, L, Bonomo, AS, Bouchy, F, Buchhave, LA, Charbonneau, D, Collier Cameron, A, Cosentino, R, Dressing, CD, Dumusque, X, Fiorenzano, AFM, Harutyunyan, A, Haywood, RD, Johnson, JA, Latham, DW, Lopez-Morales, M, Lovis, C, Mayor, M, Micela, G, Molinari, E, Motalebi, F, Pepe, F, Phillips, DF, Pollacco, D, Queloz, D, Rice, K, Sasselov, D, Ségransan, D, Sozzetti, A, Udry, S & Watson, C 2017, 'The Kepler-19 System: A Thick-envelope Super-Earth with Two Neptune-mass Companions Characterized Using Radial Velocities and Transit Timing Variations', *Astronomical Journal*, vol. 153, no. 5, pp. 224. <https://doi.org/10.3847/1538-3881/aa6897>

Digital Object Identifier (DOI):

[10.3847/1538-3881/aa6897](https://doi.org/10.3847/1538-3881/aa6897)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Astronomical Journal

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



THE KEPLER-19 SYSTEM: A THICK-ENVELOPE SUPER-EARTH WITH TWO NEPTUNE-MASS COMPANIONS CHARACTERIZED USING RADIAL VELOCITIES AND TRANSIT TIMING VARIATIONS

LUCA MALAVOLTA^{1,2}, LUCA BORSATO^{1,2}, VALENTINA GRANATA^{1,2}, GIAMPAOLO PIOTTO^{1,2}, ERIC LOPEZ³, ANDREW VANDERBURG⁴, PEDRO FIGUEIRA⁵, ANNELES MORTIER⁶, VALERIO NASCIMBENI^{1,2}, LAURA AFFER⁷, ALDO S. BONOMO⁸, FRANCOIS BOUCHY⁹, LARS A. BUCHHAVE¹⁰, DAVID CHARBONNEAU⁴, ANDREW COLLIER CAMERON⁶, ROSARIO COSENTINO¹¹, COURTNEY D. DRESSING^{12,13}, XAVIER DUMUSQUE⁹, ALDO F. M. FIOREZZANO¹¹, AVET HARUTYUNYAN¹¹, RAPHAËLLE D. HAYWOOD⁴, JOHN ASHER JOHNSON⁴, DAVID W. LATHAM⁴, MERCEDES LOPEZ-MORALES⁴, CHRISTOPHE LOVIS⁹, MICHEL MAYOR⁹, GIUSI MICELA⁷, EMILIO MOLINARI^{11,14}, FATEMEH MOTALEBI⁹, FRANCESCO PEPE⁹, DAVID F. PHILLIPS⁴, DON POLLACCO¹⁵, DIDIER QUELOZ^{9,16}, KEN RICE³, DIMITAR SASSELOV⁴, DAMIEN SÉGRANSAN⁹, ALESSANDRO SOZZETTI⁸, STÉPHANE UDRY⁹, CHRIS WATSON¹⁷

¹Dipartimento di Fisica e Astronomia “Galileo Galilei”, Università di Padova, Vicolo dell’Osservatorio 3, 35122 Padova, Italy; luca.malavolta@unipd.it

²INAF - Osservatorio Astronomico di Padova, Vicolo dell’Osservatorio 5, 35122 Padova, Italy

³SUPA, Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh, EH93HJ, UK

⁴Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, Massachusetts 02138, USA

⁵Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal

⁶Centre for Exoplanet Science, SUPA, School of Physics and Astronomy, University of St. Andrews, St. Andrews KY16 9SS, UK

⁷INAF - Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90124 Palermo, Italy

⁸INAF - Osservatorio Astrofisico di Torino, via Osservatorio 20, 10025 Pino Torinese, Italy

⁹Observatoire Astronomique de l’Université de Genève, 51 ch. des Maillettes, 1290 Versoix, Switzerland

¹⁰Centre for Star and Planet Formation, Natural History Museum of Denmark & Niels Bohr Institute, University of Copenhagen, Øster Voldgade 5-7, DK-1350 Copenhagen K, Denmark

¹¹INAF - Fundación Galileo Galilei, Rambla José Ana Fernandez Pérez 7, 38712 Breña Baja, Spain

¹²Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125, USA

¹³NASA Sagan Fellow

¹⁴INAF - IASF Milano, via Bassini 15, 20133, Milano, Italy

¹⁵Department of Physics, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK

¹⁶Cavendish Laboratory, J J Thomson Avenue, Cambridge CB3 0HE, UK

¹⁷Astrophysics Research Centre, School of Mathematics and Physics, Queens University Belfast, Belfast BT7 1NN, UK

ABSTRACT

We report a detailed characterization of the *Kepler*-19 system. This star was previously known to host a transiting planet with a period of 9.29 days, a radius of $2.2 R_{\oplus}$ and an upper limit on the mass of $20 M_{\oplus}$. The presence of a second, non-transiting planet was inferred from the transit time variations (TTVs) of *Kepler*-19b over 8 quarters of *Kepler* photometry, although neither mass nor period could be determined. By combining new TTVs measurements from all the *Kepler* quarters and 91 high-precision radial velocities obtained with the HARPS-N spectrograph, we measured through dynamical simulations a mass of $8.4 \pm 1.6 M_{\oplus}$ for *Kepler*-19b. From the same data, assuming system coplanarity, we determined an orbital period of 28.7 days and a mass of $13.1 \pm 2.7 M_{\oplus}$ for *Kepler*-19c and discovered a Neptune-like planet with a mass of $20.3 \pm 3.4 M_{\oplus}$ on a 63 days orbit. By comparing dynamical simulations with non-interacting Keplerian orbits, we concluded that neglecting interactions between planets may lead to systematic errors that could hamper the precision in the orbital parameters when the dataset spans several years.

With a density of $4.32 \pm 0.87 \text{ g cm}^{-3}$ ($0.78 \pm 0.16 \rho_{\oplus}$) *Kepler*-19b belongs to the group of planets with a rocky core and a significant fraction of volatiles, in opposition to low-density planets characterized by transit-time variations only and the increasing number of rocky planets with Earth-like density. *Kepler*-19 joins the small number of systems that reconcile transit timing variation and radial velocity measurements.

1. INTRODUCTION

After the discovery of thousands of planets with radii smaller than $2.7 R_{\oplus}$ from the NASA *Kepler* mission (Borucki et al. 2011; Coughlin et al. 2016), a consistent effort has been

devoted to understanding the formation scenario and chemical composition of such planets (e. g., Weiss & Marcy 2014, Dressing et al. 2015 and Wolfgang & Lopez 2015). To distinguish between a rocky composition and the presence of a thick envelope of water or volatile elements, the radius

derived from the transit depth must be coupled with a precise mass determination (better than 20%), either from radial velocity (RV) measurements or transit timing variations (TTVs). Planets that have been characterized with such level of precision appear to fall into two populations, a first one following an Earth-like composition and a second one with planets larger than 2 Earth radii requiring a significant fraction of volatiles (e. g., [Rogers 2015](#), [Gettel et al. 2016](#) and [López-Morales et al. 2016](#)). Recently, improved mass and radius determinations of known planets have uncovered the existence of super-Earths which fall between these two populations, such as 55 Cancri e ([Demory et al. 2016](#)) and *Kepler*-20b ([Buchhave et al. 2016](#)).

We need more planets in that mass regime with precise mass and radius measurements to understand the undergoing physics. For this reason we carried out a RV follow-up of *Kepler*-19b (hereafter *K19b*), a planet with period of 9.287 days and radius $2.209 \pm 0.048 R_{\oplus}$, orbiting a relatively bright ($V = 12.1$, $K = 11.9$) solar-type star ($T_{\text{eff}} = 5541 \pm 60$ K, $\log g = 4.59 \pm 0.10$, $[\text{Fe}/\text{H}] = -0.13 \pm 0.06$). The planet was detected by [Borucki et al. \(2011\)](#) and subsequently validated by [Ballard et al. \(2011\)](#), hereafter B11 using adaptive optics and speckle imaging to exclude a secondary source in the *Kepler* light curve, *Spitzer* observations to verify the achromaticity of the transit and Keck-HIRES high-resolution spectroscopy to rule out the presence of massive, non-planetary perturbors. From BLENDER analysis ([Torres et al. 2011](#)) the probability of a false-positive scenario was constrained to less than 1.5×10^{-4} . RVs measured on high-resolution spectra were consistent with a mass of $1.5 M_{\oplus}$ (0.5 m s^{-1}) and activity-induced RV jitter of 4 m s^{-1} , but ultimately they lacked the required precision for a robust determination of the planetary mass, and only an upper limit of $20.3 M_{\oplus}$ was set. The existence of an additional planet, *Kepler*-19c (hereafter *K19c*), with period $\lesssim 160$ days and mass $\lesssim 6 M_{\text{Jup}}$, and a further confirmation of the planetary nature of *K19b* were inferred by B11 from the presence of TTVs on 8 quarters of *K19b* light curve.

In this paper we couple high-precision RV measurements obtained with HARPS-N with updated measurements of transit times (T_0) encompassing all 17 quarters of *Kepler* data to determine the orbital parameters of *K19b*, *K19c* and a third, previously unknown planet in the system, *Kepler*-19d (hereafter *K19d*). TTV and RV datasets are analyzed independently, to understand which constraints they can provide to the characterization of the system. Subsequently, a simultaneous TTV and RV fit is performed using dynamical simulation to take into account gravitational interactions between planets. We perform this analysis under the assumption of coplanarity between planets, and then investigate the effect of different mutual inclinations on the goodness of the fit. We confirm that only the inner planet is seen transiting the host star. A comparison between RV obtained from dynamical simulations and when assuming non-interacting planets

is performed. We conclude describing the role of *K19b* in understanding the bulk densities of small planets.

2. RADIAL VELOCITIES

We collected 101 spectra using HARPS-N at the Telescopio Nazionale Galileo, in La Palma. Observations spanned over two years, from June 2012 to November 2014, overlapping the *Kepler* observations during the first year. Every observation consisted of a 30 minute exposure, with a median signal-to-noise ratio of 37 at 550 nm, corresponding to a RV nominal error of 2.8 m s^{-1} . Given the faintness of the target, observations were gathered with the *objAB* setting, i. e., the second fiber (fiber B) was observing the sky instead of acquiring a simultaneous thorium-argon (ThAr) lamp spectrum. Several observations demonstrated that the stability of the instrument over 24 hours is within 1 m s^{-1} (e. g., [Cosentino et al. 2014](#)), thus the precision of the measurements was dictated largely by photon noise. Data were reduced using the standard Data Reduction Software (DRS) using a G8 flux template (the closest available one to the spectral type of the target) to correct for variations in the flux distribution as a function of the wavelength, and a G2 binary mask was used to perform the cross-correlation ([Baranne et al. 1996](#); [Pepe et al. 2002](#)). The resulting RV data with their formal $1-\sigma$ uncertainties, the Full Width Half Maximum (FWHM) of the cross-correlation function (CCF) and its contrast (i. e., the depth normalized to the continuum), the bisector inverse span (BIS), and the $\log R'_{\text{HK}}$ activity index are listed in Table 1.

2.1. Effect of Moon Illumination

A simple procedure was adopted to check the influence of the moon illumination on the science fiber (labeled as fiber A). First, the cross-correlation function of the sky spectrum acquired with fiber B, CCF_B , was recomputed¹ using the same flux correction coefficients as for the target (CCF_A) for that specific acquisition. Then, CCF_B was subtracted to the corresponding CCF_A and radial velocities were computed again using the script from [Figueira et al. \(2013\)](#), which uses the same algorithm implemented in the DRS. For 10 observations the difference between the sky-corrected RV and the DRS RV was greater than twice the photon noise, so we rejected those observations, and used the remaining 91 RVs from the DRS in the following analysis. A flag has been included in Table 1 to identify the rejected observations.

While the rejected observations have in common a fraction of the illuminated Moon greater than 0.9 and a barycentric RV correction within 15 km s^{-1} from the absolute RV of the target star, not all the observations that met this criterium were affected by the sky contamination, suggesting that other

¹ When using the *objAB* setting, CCF_B is computed by the DRS without flux correction by default.

Table 1. HARPS-N radial velocities and ancillary measurements of *Kepler*-19. Epoch of the observations, RV with associated noise, FWHM and contrast of the CCF, inverse bisector span, $\log R'_{\text{HK}}$ activity index with the associated error, and a flag indicating if the data has been contaminated by the Moon (1) or not (0), see Section 2.1.

| BJD _{UTC} | RV | σ_{RV} | FWHM | Contrast | BIS | $\log R'_{\text{HK}}$ | $\sigma_{\log R'_{\text{HK}}}$ | Moon flag |
|--------------------|----------------------|----------------------|-----------------------|----------|----------------------|-----------------------|--------------------------------|-----------|
| [d] | [m s ⁻¹] | [m s ⁻¹] | [km s ⁻¹] | | [m s ⁻¹] | [dex] | [dex] | |
| 2456100.608 | -10608.31 | 2.31 | 6.745 | 48.83 | -41.90 | -4.975 | 0.039 | 0 |
| 2456100.629 | -10610.99 | 2.09 | 6.735 | 48.91 | -40.73 | -5.039 | 0.031 | 0 |
| 2456101.606 | -10613.07 | 3.56 | 6.732 | 48.70 | -55.78 | -4.961 | 0.060 | 0 |
| ... | ... | ... | ... | ... | ... | ... | ... | . |

NOTE—Table 1 is published in its entirety in the machine-readable format. A portion is shown here for guidance regarding its form and content.

unidentified factors can determine whether or not contamination is negligible. An in-depth analysis of the outcome of the observations is then advised when measuring RVs for faint stars.

3. KEPLER PHOTOMETRY

Kepler-19 was initially observed in long-cadence (LC) mode during the first 0-2 quarters, and then in short-cadence (SC) mode from quarter 3 until the end of the mission in 2013 (at quarter 17). At the time of publishing, B11 had at their disposal only the first 8 quarters. In an effort to make use of any additional information coming from *Kepler* photometry, we redetermined the transit times for all the quarters now available, in both LC and SC light curves. Quarters already analyzed by B11 were examined as well, to validate our T_0 determination and to provide a homogeneous set of measurements.

Transit identification was performed by propagating the linear ephemeris of B11, with the inclusion of 3 hours of pre-ingress and post-egress around the expected transit time.

For each transit time, we firstly detrended the transit light curve with a polynomial between 1st and 10th degree, with the best-fit degree chosen according to the Bayesian information criterion (BIC). Then, we determined a new T_0 guess with an automatic selection among different search algorithms² fitting the Mandel & Agol (2002) transit model, implemented in PyTransit³ (Parviainen 2015), fixing all other parameters to the literature value. Finally, the T_0 s were refined using JKTEBOP program (Southworth et al. 2004) and the associated errors were determined with a classical bootstrap approach.

Transit times from LC and SC light curves were matched together, keeping the SC measurements when available. Transit times are reported in Table 2. A comparison with B11 measurements of the observed minus predicted time of

transit ($O - C$), using their linear ephemeris for both datasets, is shown in Figure 1. The scatter of the residuals, well within the error bars, shows that the methodologies are perfectly consistent, i.e., that we are limited by photon noise, data sampling and/or unknown systematics rather than the exact procedure followed to measure the transit times.




Figure 1. The difference between the observed and the predicted (from the linear ephemeris) times of transit for *K19b*. In red, the measurements from Ballard et al. (2011), in black our new measurements for all the *Kepler* quarters. In the lower plot the difference between the two measurements is shown for the data points in common, with the error bars obtained by summing in quadrature the errors from the two estimates. The small scatter of the residuals with respect to the size of the error bars demonstrate that we are not influenced by the exact methodology used to measure the T_0 s.

Due to an error in *Kepler* archiving system, at the time of B11 publication the time stamps of all the *Kepler* light curves were reported in Coordinated Universal Time system (UTC) instead of the Barycentric Dynamical Time system (TDB)⁴.

² Levenberg-Marquardt (Moré et al. 1980), Nelder-Mead (Nelder & Mead 1965; Wright 1996), COBYLA (Powell 1994) as implemented in `scipy.optimize`, and the Affine Invariant Markov Chain Monte Carlo (MCMC) Ensemble sampler implemented in `emcee` (Foreman-Mackey et al. 2013) available at <https://github.com/dmcm/cmcee>

³ Available at <https://github.com/hpparvi/PyTransit>

⁴ http://archive.stsci.edu/kepler/timing_error.html

Table 2. Transit times of *Kepler*-19 from Q0-Q17.

| Transit Number | T_0 [BJD _{UTC}] | σ_{T_0} [d] |
|----------------|-----------------------------|--------------------|
| 0 | 2454959.7074 | 0.0014 |
| 1 | 2454968.9935 | 0.0023 |
| 2 | 2454978.2801 | 0.0020 |
| ... | ... | ... |

NOTE—Table 2 is published in its entirety in the machine-readable format. A portion is shown here for guidance regarding its form and content.

While this error was not affecting the internal consistency of B11 analysis, it must be taken into account when comparing time-series with timing accuracy better than a few minutes. We corrected for this error before comparing B11 data with our new T_0 measurements.

4. PHYSICAL PARAMETERS

Atmospheric stellar parameters of *Kepler*-19 were determined in B11 using Spectroscopy Made Easy (SME, [Valenti & Fischer 2005](#)). Since this method may suffer of correlation between derived parameters ([Torres et al. 2012](#)), and having at our disposal several high-resolution spectra from HARPS-N, we decided to carry out an independent determination with an alternative approach, i. e., equivalent width measurements of individual spectral lines instead of fitting of the whole spectrum. We used all the spectra free from sky contamination to obtain a coadded spectrum with an average S/N of 350.

Stellar atmospheric parameters were determined using the classical line-of-growth approach. For this purpose we used the 2014 version of the line analysis and synthetic code MOOG⁵ ([Sneden 1973](#)), which works in the assumption of local thermodynamic equilibrium (LTE), and the ATLAS9 grid of stellar model atmosphere from [Castelli & Kurucz \(2004\)](#) with the new opacity distribution functions and no convective overshooting. Equivalent Width measurements were carried out with the code ARESv2⁶ ([Sousa et al. 2015](#)) coupled with the updated linelist of [Malavolta et al. \(2016\)](#), where the oscillator strength of the atomic lines have been modified to correctly take into account the chemical abundances from [Asplund et al. \(2009\)](#).

Temperature and microturbulent velocity were determined by minimizing the trend of iron abundances from individual lines with respect to excitation potential and reduced equivalent width respectively, while the gravity $\log g$ was adjusted by imposing the same average abundance from neutral and ionized iron lines. For a detailed description of the procedure for the atmospheric parameters and associated errors we refer the reader to [Dumusque et al. \(2014\)](#). The derived at-

Table 3. Astrophysical parameters of the star.

| Parameter | B11 | This work |
|-------------------------------|-------------------|------------------|
| T_{eff} [K] | 5541 ± 60 | 5544 ± 20 |
| $\log g$ | 4.59 ± 0.10 | 4.51 ± 0.03 |
| ξ_t [km s ⁻¹] | - | 0.88 ± 0.05 |
| [Fe/H] | -0.13 ± 0.06 | -0.08 ± 0.02 |
| M_\star [M_\odot] | 0.936 ± 0.040 | - |
| R_\star [R_\odot] | 0.859 ± 0.018 | - |
| Age (Gyr) | 1.9 ± 1.7 | - |
| $\log R'_{\text{HK}}$ | -4.95 ± 0.05 | -5.00 ± 0.04 |

mospheric parameters are summarized in Table 3.

Our stellar atmospheric parameters agree within the uncertainties with the ones determined by B11, including the surface gravity which is usually the parameter most difficult to derive, and there is only a difference of 3 K in T_{eff} despite the use of two complementary approaches and independent datasets. For this reason we adopted their determination for mass and radius of the star and physical radius of K19b based on light curve analysis.

5. STELLAR ACTIVITY

Recently a considerable effort has been devoted to analyze the effect of stellar activity on RV measurements as well as T_0 determination ([Mazeh et al. 2015](#); [Ioannidis et al. 2016](#)). The HARPS-N DRS automatically delivers several diagnostics for activity such as the Full Width Half Maximum of the CCF, the bisector inverse span and the $\log R'_{\text{HK}}$ index, while several other chromospheric indexes such as the $H\alpha$ index ([Gomes da Silva et al. 2011](#); [Robertson et al. 2013](#)) can be determined from the spectra themselves⁷. In Figure 2 the analysis of BIS and $\log R'_{\text{HK}}$ are reported as representative of all the indexes. For each index we checked the presence of any correlation with time, either by visual inspection (upper panels of Figure 2) and with the Generalized Lomb-Scargle (GLS) periodogram ([Zechmeister & Kürster 2009](#)), where the 1% and 0.1% false alarm probability (FAP) have been computed with a bootstrap approach (middle panels of Figure 2). Finally, the presence of any correlation with RVs was verified by calculating the Spearmans rank correlation coefficient ρ , the slope of the linear fit m with its error and the p-value using the weighted least-square regression⁸ (lower panels of Figure 2). We omitted the FWHM from the analysis since a few changes of the spectrograph focus during the first year of observations modified the instrumental profile, hence the measured FWHM, without however affecting the measured RVs.

⁵ Available at <http://www.as.utexas.edu/~chris/moog.html>

⁶ Available at <http://www.astro.up.pt/~sousasag/ares/>

⁷ The code to retrieve the activity indexes is available at <https://github.com/LucaMalavolta/>

⁸ StatsModels available at <http://statsmodels.sourceforge.net/>




Figure 2. The bisector inverse span (panels on the left side) and the $\log R'_{\text{HK}}$ index (panels on the right side) are shown as example of the analysis conducted on CCF asymmetry and activity indexes. Upper panels: indexes as a function of time, the seasonal medians with the first and third quartiles indicated in red. Middle panels: GLS periodograms of the indexes, the rotational period of the star is indicated with a red vertical line. The 1% and 0.1% FAP levels are displayed as dashed and dotted horizontal lines, respectively. Lower panels: indicators as a function of RV. The best fit is represented by the dashed red line.

The absence of significant peaks in the periodogram of the indexes under analysis around the expected rotational period of 32 days (from B11 following Noyes et al. 1984), 34 ± 6 days and 36 ± 3 days (from HARPS-N $\log R'_{\text{HK}}$, following Noyes et al. 1984 and Mamajek & Hillenbrand 2008 respectively), and the lack of statistically significant correlations between the indexes and RVs, confirmed the low activity level of the star already deduced by B11 and consistent with $\log R'_{\text{HK}} = -5.00 \pm 0.04^9$.

We also searched for stellar variability on the most recent Kepler Pre-search Data Conditioning (PDC) light curve, which presents several improvements in correcting instrumental trends (and thus is better suited to search for activity modulation) with respect to the light curve available at the time of B11 publication. For most of the time the star is photometrically quiet, while in some parts of the light curve a clear signal, likely due to stellar activity, is detected. We applied the autocorrelation function technique over these portions of light curve and we estimated a rotational period of 30 days. This signal is characterized by a short time scale of decay (a few rotational periods) and a rapid loss in coherence.

We comment on the impact on RV in section 7.

6. TTV ANALYSIS

Dynamical analysis of the system was performed with TRADES¹⁰ (TRANSITS and Dynamics of Exoplanetary Systems, Borsato et al. 2014), a N-body integrator with the capability of fitting RVs and T_0 s simultaneously to determine the orbital parameters of the system through χ^2 minimization.

Since only the innermost planet is transiting, it is extremely difficult to constrain the orbital parameters of all the planets in the system by TTV alone. In fact, attempts to fit the T_0 s with a two-planet model resulted in a strong degeneracy between the mass and the period of the second planet, i. e., O-C diagrams with similar shape could be produced by jointly increasing the mass and the period of *K19c*. The amplitude and shape of the T_0 s however can still give us upper limits on the mass and period of the non-transiting planet if we compare the outcome of the dynamical simulations with the maximum mass compatible with the observed semi-amplitude of the RVs.

We proceeded as follows. We used TRADES to perform a fit of the T_0 s by assuming a two-planet model and fixing the mass of *K19b* to a grid of values between 2.5 and $20 M_{\oplus}$ and a spacing of $2.5 M_{\oplus}$, with its period already measured from the *Kepler* observations. To each point of this grid we assigned several values for the mass of *K19c* randomly selected between 5 and $250 M_{\oplus}$. For each combination of *K19b* and *K19c* masses, the other orbital parameters of the system were left free to vary and their best-fitting values were determined by χ^2 minimization through the Levenberg-Marquardt algorithm. We discarded those solutions with at least one planet having eccentricity greater than 0.3 , assuming that interacting planets meeting this condition are likely to be unstable.

In Figure 3 we show the results obtained for the mass of *K19c* as a function of its period. The expected RV semi-amplitude of *K19c* as a function of mass and period are superimposed. We note that many of the orbital configurations reported in the plot could be unstable, since dynamical stability was not checked yet at this stage (stability analysis is introduced in Section 8). We can attempt to estimate a lower limit to the periods and masses of the non-transiting planets according to the observed semi-amplitude of the RVs, by taking advantage of the correlation between the mass and period of *K19c*. Our RVs have a peak-to-peak variation of 23 m s^{-1} , so if we keep into account the additional signal of *K19b* (a few m s^{-1} in the case of a Neptune-like density), *K19c* amplitude should necessarily lie below the $K = 10 \text{ m s}^{-1}$ line. This fact suggests that a short period ($\lesssim 50$ days) for *K19c* should be expected, while nothing can be said regarding *K19b* from TTV alone. We remind the reader that this analysis cannot be

⁹ Note however that this value could be affected by interstellar medium absorption and be higher than measured (Fossati et al. 2017).

¹⁰ TRADES is available at <https://github.com/lucaborsato/trades>




Figure 3. The mass of K19c versus its period, determined by fitting K19b T_0 s for a grid of masses of the two planets. The three lines represent the expected RV semi-amplitude of K19c only as a function of its mass and period. A linear fit to the data is marked with a dash-dot blue line.

considered conclusive due to the reduced number of points and their scatter around the best linear fit.

7. RV ANALYSIS

While only one planet is transiting *Kepler*-19, the presence of at least one additional planet was inferred by the presence of TTVs. To reveal such planets, we first performed a frequentist analysis on the RV dataset by computing the GLS periodogram and iterating over the residuals until no significant periodicity was present. This analysis revealed two signals at 28.6 days and 62.3 days with false alarm probability lower than 1%, while the signal at 9.3 days corresponding to K19b was barely detected (Figure 4).

The signal at 28 days is very close to the second order 3:1 mean motion resonance (MMR), which is among the possible configurations listed by B11 as the cause of the TTVs of K19b. While the false alarm probability of this signal is below the traditional threshold for RV planet detection claims ($\approx 10^{-3}$), we can compare it with the probability of observing K19c near another MMR resonance. In order to do so, we scrambled once again the RV observations and calculated the fraction of periodograms which had a stronger peak with respect to the untouched dataset, in a frequency range of 5% around the interior and exterior 1:2, 2:3, 3:4 first order resonances, the 1:3 and 3:5 second order resonances, and the 1:4 and 1:5 higher order resonances. The FAP of the signal at 28 days computed in this way decreases to 8×10^{-4} , i.e., it is unlikely that the signal at 28 days is a spurious signal caused by a planet in another MMR configuration.




Figure 4. Upper panel: the RV dataset. The vertical line divides data taken before and after the substitution of HARPS-N CCD. Lower panels: GLS periodograms computed starting from the initial dataset and iterating over the residuals. The 1% and 0.1% false alarm probabilities were estimated following a bootstrap approach for each periodogram. Red lines mark the best-fit period at 28.6 days, 9.29 days and 62.3 days.

The signal at ≈ 28.6 days is consistent with the rotational period of the star obtained from the active regions on *Kepler* light curve, which however have the characteristic of rapidly decaying and reappearing later at different phase and intensity. We checked if the RV signal has the same properties by performing a jackknife analysis by splitting the RV dataset in two parts (at BJD = 2456700d) and determining the phase and amplitude of the signal on each dataset¹¹. We obtained $K = 2.6 \pm 1.1 \text{ m s}^{-1}$, $\phi = 3.0 \pm 0.3 \text{ rad}$ for the first half of the dataset, and $K = 3.0 \pm 0.6 \text{ m s}^{-1}$, $\phi = 2.8 \pm 0.2 \text{ rad}$ for the second half, i.e., the signal is stable over several years, and therefore unlikely to be due to stellar activity. As an additional check, we computed the stacked Bayesian Generalized Lomb-Scargle (BGLS) periodogram (Mortier & Collier Cameron 2017). The S/N is increasing with the square of the number of observations, as expected for a coherent (planetary) signal, although the plot is heavily affected by the poor sampling and the signal of K19b near the 3:1 MMR, as we verified by performing the same analysis on synthetic datasets with the same temporal sampling. The absence of RV modulation due to stellar activity is supported by the analysis performed in Section 5, where no periodicity or correlation with RVs is seen for any of the activity indexes.

The semi-amplitudes of these signals however cannot be

¹¹ We imposed as a prior the period $P = 28.57 \pm 0.02$ day to compensate for the scarcity and poor sampling of the split datasets

Table 4. Orbital parameters for a 3-planet model for the *Kepler*-19 system obtained from RV only, except for the period of K19c which is constrained by the *Kepler* light curve. The reference time for the orbital elements is $T_{\text{ref}} = 2456624.82263024$ days.

| | K19b | K19c | K19d |
|------------------------|-----------------------|------------------|-------------------------|
| Circular orbits | | | |
| Period [d] | 9.28699 ± 10^{-5} | 28.61 ± 0.24 | 63.0 ± 0.3 |
| K [m s ⁻¹] | 2.3 ± 0.5 | 1.7 ± 0.8 | 3.8 ± 0.6 |
| ϕ [deg] | 194.4 ± 0.3 | 185 ± 43 | 174 ± 6 |
| Mass [M_{\oplus}] | 7.4 ± 1.7 | 7.8 ± 3.6 | 22.5 ± 3.8 |
| Keplerian orbits | | | |
| Period [d] | 9.28699 ± 10^{-5} | 28.54 ± 0.27 | 62.9 ± 0.3 |
| K [m s ⁻¹] | 2.6 ± 0.6 | 1.95 ± 0.8 | 4.0 ± 0.7 |
| ϕ [deg] | 198 ± 8 | 192 ± 29 | 172 ± 8 |
| $\sqrt{e} \cos \omega$ | 0.0 ± 0.2 | 0.0 ± 0.4 | $0.31^{+0.20}_{-0.31}$ |
| $\sqrt{e} \sin \omega$ | 0.0 ± 0.3 | 0.0 ± 0.3 | $-0.25^{+0.35}_{-0.22}$ |
| Mass [M_{\oplus}] | 8.0 ± 1.8 | 8.6 ± 3.5 | 23.1 ± 3.8 |
| e | ≤ 0.13 | ≤ 0.30 | ≤ 0.32 |
| ω [deg] | unconstrained | unconstrained | -36^{+65}_{-35} |

inferred by the frequentist analysis alone, since an offset in RV between the data taken before and after September 2012 may exist due to the failure of the first CCD of HARPS-N (see Bonomo et al. 2014). The value of this offset cannot be determined a priori, even when observations of RV standard stars are available, since it may depend primarily on the spectral type of the star (i.e., the two CCD may have a different efficiency as a function of wavelength). Introducing an RV offset as a free parameter is a common procedure for non-overlapping datasets, e.g., see Benatti et al. (2016).

We then performed a tentative fit using the Markov-Chain Monte Carlo (MCMC) code PyORBIT¹² (Malavolta et al. 2016), allowing for two different systemic velocity γ of the star and using a 3-planet model to fit the data. We attempted two different fits with both circular and Keplerian orbits. For planet *c* and *d* we set uniform priors in the range [10, 50] days and [50, 90] days respectively, while the other parameters were left to vary within their physically meaningful range (e.g., positive-definite RV semi-amplitude). The results are shown in Table 4.

Eccentricity and argument of pericenter of each of the three planets are poorly constrained by the RVs alone, thus affecting the precision on mass of the presumed planets. A combined analysis of RVs and TTV is then required to unambiguously detect and characterize the planets in the system.

8. COMBINED RV AND TTV ANALYSIS

The amplitude of dynamical perturbations between planets is very sensitive to the eccentricity and angular parameters

(i.e., argument of pericenter ω and mean anomaly at reference time M_0) of the planetary orbits, which however can be only poorly constrained by the RVs, especially for planets in nearly circular orbits. For this reason we expect an overall improvement on the precision of the orbital parameters by simultaneously fitting RVs and TTVs.

Dynamical simulations are extremely time-consuming, and we have to use all the information at our disposal to reduce the extension of the parameter space. We used the results from the RV fit in Section 7 to put a constraint to the range of period, mass and orbital phase of each planet.

We followed an iterative approach to avoid being trapped in a local minimum of the χ^2 . We started 10 separate runs of TRADES with the Particle Swarm Optimization algorithm and loose priors on the periods (± 10 days around the expected period of each planet from the RV analysis), masses (between 0 and 40 M_{\oplus}) and eccentricities ($e \leq 0.5$), taking into account the limits imposed by photometry and RVs. We checked the stability of the outcome of each run and then we ran TRADES again on a range of parameters which was half the size than in the previous run and centered of the outcome of the previous run with the lowest χ^2 among those that satisfied the stability requirement. Convergence was considered achieved when all the runs resulted in similar parameters (within 5% from the mean) and similar χ^2 (10% from the mean).

Following Gladman (1993), during the numerical integration we checked the stability criterion for each pair of planets: $\Delta = 2\sqrt{3}R_H(i, j)$, where $\Delta = a_j - a_i$ is the semi-major axis difference between the j_{th} and i_{th} planet, and $R_H(i, j)$ is the mutual Hill radius between planet *i* and *j*. At the end of each TRADES fit we performed a N-body integration with SymBA (Duncan et al. 1998) and checked the stability of the result with the Frequency Map Analysis tool (FMA, Laskar et al. 1992; Laskar 1993a,b) with the prescriptions of Marzari et al. (2002). A system is considered stable if the coefficient of diffusion is lower than 10^{-5} , in an unstable or chaotic state otherwise.

We used the value measured by B11 for the inclination *i* of K19b. Several attempts to fit the mutual inclinations of the non-transiting planets along with the other parameters resulted always in unstable solutions with high mutual inclinations. We took advantage of the low mutual inclinations of planetary orbital planes inferred from *Kepler* multi-planet systems (Fabrycky et al. 2014) and the additional information coming from systems characterized with high-precision RVs (Figueira et al. 2012) to impose coplanarity with K19b for the other planets, while the longitude of the ascending node Ω was fixed to zero for all the planets. This assumption allowed us to drastically reduce computational time. The orbital period and mass of the planet, the eccentricity *e*, the argument of periastron ω and the mean anomaly at the reference epoch M were left as free parameters.

Differently from (Borsato et al. 2014), we fitted $\sqrt{e} \cos \omega$ and $\sqrt{e} \sin \omega$ instead of $e \cos \omega$ and $e \sin \omega$, and the mean lon-

¹² Available at <https://github.com/LucaMalavolta/PyORBIT>

gitude at the reference epoch $\lambda = \omega + \mathcal{M} + \Omega$ (where Ω is the longitude of the ascending node, fixed to zero for being unconstrained by the data) instead of \mathcal{M} . We used scaled stellar and planetary masses as commonly done in TTV analysis (e.g., Nesvorný et al. 2012). A RV offset between the data taken before and after September 2012 was included as a free parameter to take into account the change of the CCD (see Section 7).

We used the solution obtained with the global exploration of the parameter space as a starting point for the Bayesian analysis. For this purpose we expanded TRADES functionalities with the `emcee` package (Foreman-Mackey et al. 2013), an affine invariant Markov chain Monte Carlo (MCMC) ensemble sampler, and made it available to the community in the TRADES repository. Following Feigelson & Babu (2012), we calculated the log-likelihood $\ln \mathcal{L}$ from the χ^2 using Equation 1, where *dof* is the degree of the freedom of the problem.

$$\ln \mathcal{L} = -\ln(2\pi) \frac{\text{dof}}{2} - \frac{\sum \ln \sigma^2}{2} - \frac{\chi^2}{2} \quad (1)$$

We tested three different scenarios. The first model assumed that the signal at ≈ 28 days (the first one being detected in the RV periodogram) is the only planet in the system other than K19b. The second model is still a two-planet model, but here we assumed that the signal at ≈ 28 days is due to activity (without any effects on the T_0 s) and that the system consists of two planets at ≈ 9.2 and ≈ 62 days. The third model assumed that all the RV signals have planetary origin. The results are presented in the following subsections. We note that a planet in a strong MMR with K19b could still produce the TTVs while having a RV semi-amplitude below our detection sensitivity. Following B11, the TTVs of K19b could be explained by a planet in [2:1, 3:2, 4:3] MMRs with masses [4, 2, 1] M_\oplus respectively. The signal of the perturbing planet would have an RV semi-amplitude on the order of [1, 0.6, 0.3] m s^{-1} , i.e., beyond the reach of modern velocimeters given the magnitude of our target. This scenario however requires an activity origin for the 28 days signals, which is not supported by the analysis of Section 5 and the coherent nature over years of the RV signal in opposition to the short time-scale decay of the spots observed in *Kepler* photometry, as shown in Section 7. For these reasons we can regard this scenario as unlikely.

8.1. Two-planet model

We tested the two-planet model following the steps described in Section 8, with the only additional constraint of K19c having a period lower than 40 days. We ran the MCMC sampler for 50000 steps, with a number of chains twice the dimensionality of the problem. We checked the convergence of the chains using the Gelman-Rubin statistic ($\hat{R} < 1.03$, Gelman & Rubin 1992; Ford 2006), and we built the posterior distributions with the last 20000 steps and a thinning factor of 200.

Results are listed in Table 5. Rather than using the median or the mode, we summarize the outcome of the analysis by selecting from the chains a sample with the nearest χ^2 to the median of its distribution and with each parameter within its confidence interval. Confidence intervals are computed by taking the 15.87th and 84.14th percentiles of the distributions, and are reported in the table with respect to the selected sample.

8.2. Two-planet and stellar activity model

The presence of the activity signal can potentially affect the analysis, but TRADES is not equipped to deal with non-planetary signals in RV datasets. For this reason we decided to remove such signal from the RV time series before starting the global exploration of the parameter space, using as dataset the residuals of the first iteration of the GLS analysis in Section 7. The global solution was used as starting point for a MCMC analysis with PyORBIT, this time using the original RV dataset and the activity signal modeled with a Keplerian curve (as similarly done in Pepe et al. 2013). RVs and T_0 s calculations regarding the two interacting planets were performed by calling the dynamical integrator of TRADES through a FORTRAN90 wrapper. As additional test-cases, we run TRADES using the two-planet model and no activity modeling on both the original RV dataset (imposing a lower limit of 40 days for the period of K19c, to avoid falling in the case analyzed in Section 8.1) and the *GLS-cleaned* dataset. In all cases we repeated the analysis without imposing any constraint on eccentricity, since the eccentricities of both planets were moving towards the upper boundary imposed in Section 8. We followed the same methodology described in Section 8.1 to run the MCMC and extract the results reported in Table.

The eccentricity of K19b is extremely high in all the cases we considered. The most likely explanation is that high eccentricities for both planets are required to produce the same T_0 s while keeping their masses within the boundaries set by the RVs. Following Burke (2008) we computed the ratio τ of the transit duration of an eccentric orbit with respect to a circular orbit, and we obtained $\tau = 0.64 \pm 0.04$ for the original RV dataset, $\tau = 0.58 \pm 0.06$ for the *GLS-cleaned* case, and 0.54 ± 0.10 for the planets+activity case. All these values are well below the value of $\tau = 0.7$ that B11 considered as the minimum reasonable value for the transit duration ratio due to the eccentricity of the planet from the analysis of the *Kepler* light curve. We note that the period-mass combinations we obtain for the outer planet fall in an empty region of Figure 3, which however was obtained by selecting those solutions with $e < 0.3$ for at least one of the planets.

8.3. Three-planet model

Bayesian analysis for the three-planet model were performed using TRADES combined with `emcee` (see Section 8). From a preliminary analysis we noticed that the chains were

affected by poor mixing, with a Gelman-Rubin statistic $\hat{R} \simeq 1.3$ (Gelman & Rubin 1992; Ford 2006). We decided to run 100 chains for an extensive number of steps (250000) to overcome the poor mixing and perform a proper exploration of the parameter space, knowing that we already reside near the global minimum of the χ^2 . After the first 150000 iterations we did not see any variation of the posterior distributions while increasing the length of the chains, making us confident of the robustness of our result. We finally built the posterior probability by drawing 40000 independent samples from the chains, after removing the burn-in part and applying a thinning factor equal to their auto-correlation time ($\simeq 100$ steps).

Our results are listed in Table 5. The confidence intervals of the posteriors are computed by taking the 15.87th and 84.14th percentiles of the distributions, and they are reported as error bars around a sample solution selected as in 8.2. From now on, we will use the reported values as a representative *solution* of the orbital parameters of the planets. In Figure 6 we show the solution overplotted on T_0 s and RVs, with their respective residuals.

In Table 6 we compare the outcome of the different models under examination. The three-planet solution is the favorite one according to the BIC (having a $\Delta\text{BIC} > 10$ with respect to all the other models under analysis, Kass & Raftery 1995), and the considerations at the end of Section 8.2 further strengthen this result. Our three-planet solution has a total reduced $\chi_r^2 = 1.25$ ($\chi^2 = 272.7$, BIC = 365.5), where the contribution from RV and TTV is respectively 0.37 and 0.88¹³. The standard deviation of the residual RVs is 2.9 m s^{-1} , consistent with the average error of the measurements, and while several peaks can be found in the periodogram, none of them reaches the 1% false-alarm probability threshold (Figure 7). Similarly, the standard deviation of the TTV of 2.2 minutes is consistent with the average error of 2.1 minutes.

For the three planets we determined their masses with a precision of $\sigma_{M_b} = 1.6 \text{ M}_\oplus$ (error of 19% on planetary mass) for K19b, $\sigma_{M_c} = 2.7 \text{ M}_\oplus$ (error of 21%) for K19c and $\sigma_{M_d} = 3.4 \text{ M}_\oplus$ (error of 17%) for K19d. We included in the computation the uncertainty on the stellar mass and the effect of orbital inclinations, assuming as representative distributions for the latter a normal distribution with $i_c = i_d = 89.94^\circ$ and dispersion $\sigma_i = 5^\circ$ for K19c and $\sigma_i = 15.0^\circ$ for K19d, after Section 9. Knowing that K19c and K19d should have a transit depth greater than 0.5 mmag (Winn 2010), a visual inspection of the *Kepler* light-curve confirmed that the planets are not transiting.

Our solution agrees with the solution in Table 4, except for the mass of K19c, with a $2\text{-}\sigma$ difference between the RV-only

and RV+TTV determinations. The origin of this discrepancy is likely due to the uncertainty in the orbital phase and eccentricity from the RV-only analysis, with one planet absorbing the signal of the other, possibly coupled with the effect described in Section 10.

The RV offset between the *old* and *new* CCD is $\simeq 2 \text{ m s}^{-1}$, i. e., within the RV noise. We do not expect any influence of the RV offset on the outcome of the analysis, since the three planets have periods fully contained within the time span of both *old* and *new* CCD datasets.

9. MUTUAL INCLINATIONS

In Section 8 we assumed co-planarity with K19b (the only planet with known inclination) to derive the orbital parameters of the additional planets in the system. To check if this assumption was still valid after determining the orbital period of the additional planets, we determined the upper limit on inclination for which a transit is visible for a given planet, by inverting Equation 7 of Winn (2010) with the assumption for the impact parameter $b_{\text{tra}} = 1$. To properly take into account variation with time of e and ω induced by dynamical interactions, we selected 1000 (e, ω) pairs for each planet by randomly sampling in time the integration our best solution over the *Kepler* observational time span. Error in the stellar radius was included by generating random samples from a normal distribution with mean R_\star and standard deviation σ_{R_\star} , as reported in Table 3. We obtained $i_{\text{min}} = 88.72 \pm 0.03^\circ$ for K19c and $i_{\text{min}} = 89.24 \pm 0.02^\circ$ for K19d. Since we assumed $i_b = 89.94$ (from B11), then the coplanarity assumption cannot hold for K19c. We assessed the influence of orbital inclinations on the validity of our solution by running dynamical simulations for a grid of i_c and i_d (from 60 to 120 degrees, with step of 0.25° for i_c and 0.5° for i_d), with the remaining orbital parameters fixed to our best solution, and determining the reduced χ^2 as in Section 8. As shown in Figure 8, the reduced χ^2 increases rapidly with K19c going farther from coplanarity with K19b, reaching a value of ≈ 1.6 for $|i_b - i_c| \simeq 5^\circ$, while K19d can span a larger interval in inclinations without affecting the outcome (although increasing $|i_b - i_d|$ would affect negatively the stability of the system). Assuming $i_c = 88.72$ (grazing scenario) the outcome of the fit is nearly the same $\chi_{\text{red}}^2 = 1.26$, which is very close to the value obtained when assuming coplanarity ($\chi^2 = 1.25$, see Section 8). It is likely that the system is very close to orbital alignment, as observed for the majority of transiting multi-planet systems, e. g., Figueira et al. 2012; Fabrycky et al. 2014; Ballard & Johnson 2016; Becker & Adams 2016.

10. DYNAMICAL VERSUS NON-INTERACTING ORBITS

Dynamical simulations include by definition the effects of gravitational interactions between planets. This is different from the usual approach followed in the exoplanet literature, where a series of non-interacting Keplerian orbits are used to

¹³ These two values do not correspond to the individual χ_{red}^2 of RVs and TTVs, since the sum of residuals of each dataset is divided by the total number of data points.

Table 5. Orbital parameters of the planets in the *Kepler*-19 system obtained from TTV+RV MCMC analysis using different assumptions for the number of planets and the stellar activity. The reference time for the orbital elements is $T_{\text{ref}} = 2456624.82263024$ d.

| Planet | Period [d] | Mass [M_{\oplus}] | $\sqrt{e} \cos \omega$ | $\sqrt{e} \sin \omega$ | λ [deg] | e | ω [deg] | \mathcal{M} [deg] |
|---|----------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|---------------------------|-------------------------|-------------------------|
| Two-planet model, $P_c < 40$ days | | | | | | | | |
| K19b | $9.287108^{+0.00006}_{-0.00003}$ | $8.2^{+1.6}_{-1.4}$ | $0.23^{+0.04}_{-0.02}$ | $0.21^{+0.03}_{-0.05}$ | $188.9^{+0.9}_{-1.5}$ | $0.10^{+0.01}_{-0.01}$ | $42.5^{+5.7}_{-11.6}$ | $146.4^{+10.5}_{-5.1}$ |
| K19c | $28.723^{+0.0005}_{-0.007}$ | $17.0^{+2.4}_{-2.4}$ | $0.48^{+0.01}_{-0.04}$ | $0.15^{+0.02}_{-0.11}$ | $175.5^{+3.0}_{-7.0}$ | $0.25^{+0.01}_{-0.04}$ | $17.3^{+2.9}_{-12.7}$ | $158.1^{+6.8}_{-1.0}$ |
| Two-planet model, $P_c > 40$ days | | | | | | | | |
| K19b | $9.286970^{+0.00009}_{-0.00005}$ | $8.6^{+1.9}_{-1.7}$ | $-0.09^{+0.11}_{-0.06}$ | $0.64^{+0.04}_{-0.05}$ | $201.6^{+3.5}_{-6.3}$ | $0.42^{+0.06}_{-0.05}$ | $97.9^{+5.5}_{-9.9}$ | $103.7^{+3.6}_{-2.2}$ |
| K19c | $63.128^{+0.010}_{-0.007}$ | $21.7^{+1.8}_{-3.7}$ | $-0.11^{+0.25}_{-0.02}$ | $-0.59^{+0.02}_{-0.02}$ | $167.7^{+14.4}_{-0.1}$ | $0.36^{+0.011}_{-0.02}$ | $-100.4^{+23.9}_{-1.2}$ | $268.1^{+0.2}_{-10.1}$ |
| Two-planet model, <i>GLS-cleaned</i> RV dataset | | | | | | | | |
| K19b | $9.28696^{+0.00006}_{-0.00006}$ | $8.4^{+1.3}_{-1.6}$ | $-0.08^{+0.02}_{-0.17}$ | $0.74^{+0.02}_{-0.11}$ | $201.1^{+10.2}_{-1.5}$ | $0.551^{+0.03}_{-0.11}$ | $95.9^{+14.8}_{-1.7}$ | $105.2^{+0.3}_{-4.8}$ |
| K19c | $63.128^{+0.011}_{-0.008}$ | $17.4^{+2.9}_{-2.9}$ | $-0.07^{+0.11}_{-0.08}$ | $-0.57^{+0.01}_{-0.03}$ | $169.3^{+10.2}_{-3.3}$ | $0.332^{+0.035}_{-0.006}$ | $-97.4^{+10.9}_{-7.4}$ | $266.7^{+6.26}_{-3.4}$ |
| Two-planet and activity model | | | | | | | | |
| K19b | $9.28696^{+0.00008}_{-0.00004}$ | $7.8^{+2.1}_{-1.9}$ | $-0.02^{+0.19}_{-0.11}$ | $0.71^{+0.09}_{-0.06}$ | $197.9^{+6.4}_{-11.3}$ | $0.50^{+0.17}_{-0.07}$ | $91.7^{+9.2}_{-14.8}$ | $106.2^{+2.6}_{-1.8}$ |
| Act | $28.57^{+0.08}_{-0.11}$ | $(2.3^{+1.1}_{-0.3})^a$ | $0.22^{+0.31}_{-0.68}$ | $0.34^{+0.31}_{-0.23}$ | $178.4^{+2.0}_{-23.6}$ | $0.16^{+0.45}_{-0.01}$ | $56.4^{+84.4}_{-36.1}$ | $122.0^{+31.4}_{-86.7}$ |
| K19c | $63.139^{+0.003}_{-0.019}$ | $16.2^{+2.4}_{-2.7}$ | $0.11^{+0.02}_{-0.24}$ | $-0.58^{+0.04}_{-0.02}$ | $178.3^{+1.16}_{-15.6}$ | $0.35^{+0.02}_{-0.04}$ | $-79.0^{+1.3}_{-23.6}$ | $257.4^{+9.0}_{-1.7}$ |
| Three-planet model | | | | | | | | |
| K19b | $9.28716^{+0.00004}_{-0.00006}$ | $8.4^{+1.6}_{-1.5}$ | $0.17^{+0.05}_{-0.03}$ | $0.29^{+0.04}_{-0.06}$ | $190.3^{+1.0}_{-1.9}$ | $0.12^{+0.02}_{-0.02}$ | $59.1^{+68.9}_{-11.7}$ | $131.2^{+10.4}_{-6.1}$ |
| K19c | $28.731^{+0.012}_{-0.005}$ | $13.1^{+2.7}_{-2.7}$ | $0.42^{+0.04}_{-0.03}$ | $0.19^{+0.10}_{-0.05}$ | $181.5^{+6.0}_{-4.5}$ | $0.21^{+0.05}_{-0.07}$ | $23.8^{+11.1}_{-6.8}$ | $157.6^{+3.1}_{-6.6}$ |
| K19d | $62.95^{+0.04}_{-0.30}$ | $22.5^{+1.2}_{-5.6}$ | $0.13^{+0.17}_{-0.38}$ | $0.18^{+0.21}_{-0.19}$ | $170.1^{+12.3}_{-0.3}$ | $0.05^{+0.16}_{-0.01}$ | $55.2^{+76.8}_{-37.4}$ | $114.9^{+63.7}_{-70.5}$ |

^a Semi-amplitude of the signal in m s^{-1} **Table 6.** Statistical indexes for the different models under exams. The number of parameters in the fit, the degree of freedom (*dof*), the χ^2 and its reduced value, the log-likelihood $\ln \mathcal{L}$ and the Bayesian Information Criterion (BIC) are reported.

| Model | N. parameters | <i>dof</i> | χ^2 | χ^2_{red} | $\ln \mathcal{L}$ | BIC |
|-----------------------------------|---------------|------------|------------------------|------------------------|-----------------------|-----------------------|
| 2 planets $P_c < 40$ days | 12 | 223 | $323.8^{+5.5}_{-7.4}$ | $1.45^{+0.01}_{-0.02}$ | $498.6^{+1.0}_{-3.7}$ | $389.4^{+5.5}_{-1.9}$ |
| 2 planets, $P_c > 40$ days | 12 | 223 | $324.0^{+4.4}_{-3.4}$ | $1.45^{+0.02}_{-0.01}$ | $499.5^{+1.6}_{-2.2}$ | $389.5^{+4.4}_{-3.4}$ |
| 2 planets, <i>GLS-cleaned</i> RVs | 12 | 223 | $318.8^{+4.4}_{-3.5}$ | $1.43^{+0.02}_{-0.02}$ | $502.1^{+1.8}_{-2.2}$ | $384.3^{+4.4}_{-3.5}$ |
| 2 planets and activity | 17 | 218 | $319.5^{+8.8}_{-6.1}$ | $1.47^{+0.04}_{-0.03}$ | $506.3^{+3.0}_{-4.4}$ | $412.3^{+8.8}_{-6.1}$ |
| 3 planets | 17 | 218 | $276.7^{+6.9}_{-5.27}$ | $1.27^{+0.03}_{-0.02}$ | $527.7^{+2.6}_{-3.4}$ | $369.5^{+6.9}_{-5.3}$ |

derive the planet parameters in multiple system. While the assumption of negligible interactions between planets may hold in most of the cases, in the presence of TTVs we know that such interactions are happening. It is then worthy to analyze the differences in the RVs between the two approaches, i. e., dynamical vs. non-interacting Keplerian orbits. In order to do so, we have simulated the expected RVs of the *Kepler*-19 system at the observational epochs of our dataset, using the planetary parameters in Table 5 and assuming non-interacting orbits, and then we have subtracted the outcome to the dynamically-derived RVs from the same orbital parameters. The results are shown in Fig. 9. The $\text{RV}_{\text{TTV}} - \text{RV}_{\text{Kep}}$ residuals show a peak-to-peak variation of 0.30 m s^{-1} with a prominent periodicity at 29.3 days, i. e., very close to the orbital period of K19c, although these values depend strongly on the assumed orbital parameters. It is worthy to note that the difference between dynamical and Keplerian RVs diverges while moving further from the reference time of the orbital parameters T_{ref} . The reason resides in the small varia-

tions of the orbital parameters with time caused by planet interactions, which are implicitly taken into account in dynamical integration but ignored in the Keplerian approach. For the *Kepler*-19 system, after two years of observations from T_{ref} the peak-to-peak difference has already reached $\approx 0.6 \text{ m s}^{-1}$, i. e., a value that can negatively affect the mass determination of the planets. Assuming non-interacting planets when modeling the RVs can thus mislead the determination of the orbital parameters, e. g. in the case of dataset spanning several years.

11. DISCUSSION

The transiting planet *Kepler*-19b was previously validated by Ballard et al. (2011). A period of 9.23 days and an upper limit of $20 M_{\oplus}$ were determined from the analysis of the light curve and high-resolution spectroscopy. From the presence of TTV in 8 *Kepler* quarters they deduced the presence of a second, non-transiting planet with period ≤ 160 days and mass $\leq 6 M_{\text{Jup}}$. In this paper we presented the first precise




Figure 5. Posterior distributions of the fitted parameters for the three-planet model for the *Kepler*-19 system. Dashed blue lines identify the reference solution listed in Table 5.

mass measurement for K19b ($8.4 \pm 1.6 M_\oplus$) and the characterization of two non-transiting Neptune-mass companions, K19c ($P = 28.73 \pm 0.01$ days, $M = 13.1 \pm 2.7 M_\oplus$) and K19d ($P = 62.9 \pm 0.2$ days, $M = 20.3 \pm 3.4 M_\oplus$), obtained by simultaneously modeling TTVs and RVs through dynamical integration. We excluded stellar activity as a possible origin for the RV signal at $P \approx 28$ days using all the activity diagnostics at our disposal, including the latest reduction of *Kepler* light curve. Nevertheless we performed standard model selection between the three-planet model and two-planet models, ei-

ther with the hypothesis of the outer planet causing the TTVs and different assumptions for the activity signal, and with the hypothesis of the perturber having period ≈ 28 days and no outer planets. In all cases the three-planet model resulted the favorite one with high degree of confidence. A planet in a strong MMR with K19b could still produce the TTVs while having a RV semi-amplitude below our detection sensitivity, but only at the condition of assuming that the 28 signal is due to stellar activity, which is not supported by our data.

With a period ratio $P_c/P_b = 3.09$, the system is very close




Figure 6. Top panels: Observed-Calculated (O-C) transit times and residuals for K19b. Lower panels: RVs and residuals of the *Kepler*-19 system. Red empty circles represent the predicted values from the solution obtained with TRADES. The radial velocity curve of the *Kepler*-19 system obtained from the dynamical integration is shown in gray.

to a 3:1 mean motion resonance. The sinusoidal shape of the TTV is then induced by the 3:1 MMR of the inner planets, with a modulation caused by the outer planet. We performed a comparison between dynamical RVs with those calculated assuming non-interacting planets, using the orbital parameters of the *Kepler*-19 system, and showed that, for this specific system, the difference for a dataset spanning several years is at the limit of detection with the state-of-the-art instruments used for planet search and characterization.

Our new determination of K19b mass is in disagreement with the most-likely value of $1.6 M_{\oplus}$ (semi-amplitude of 0.5 m s^{-1}) obtained by B11 while attempting a fit with only 8 Keck-HIRES RVs and assuming only one planet in the system. As a consequence of this assumption, they derived a likely RV jitter of $\approx 4 \text{ m s}^{-1}$, which is not confirmed by our analysis. In general, RV analysis performed on a small number of measurements should always be handled with extreme care, since the results could be affected by additional non-transiting planets with little influence on the TTV of the transiting planets but with significant RV semi-amplitude, such as K19d.

K19b falls in the region of super-Earths with rocky cores and a significant fraction of volatiles or H/He gas, in opposition to the low-density planets characterized by TTV only (Weiss & Marcy 2014; Jontof-Hutter et al. 2016), and well separated from the group of rocky planets with radii smaller than $2 R_{\oplus}$, as can be seen in Figure 10. If we assume that the planet composition is a mixture of H/He enve-

lope with solar composition atop a rocky core with Earth-like rock/iron abundances, we can estimate the internal structure of K19b. By employing theoretical models from Lopez & Fortney (2014), and assuming an age of 2 Gyr, an envelope of $0.4 \pm 0.3\%$ of the total mass is required to explain the observed mass and radius of K19b. Despite the fairly high level of irradiation, its atmosphere is only moderately vulnerable to photo-evaporation due to the relatively large mass of the planet (Lopez et al. 2012). Employing the same models, we found that the mass of the primordial envelope was approximately twice ($\approx 1\%$) the current envelope mass.

Although the radii of the Neptune-mass planets are unknown, we can still speculate on their possible internal composition. By using the gas accretion scaling relations from Lee & Chiang (2015), we expect for K19c and K19d to have accreted significantly larger envelopes than K19b, likely unaffected by photo-evaporation due to the larger masses of these two planets. Using Equation 22 from Lee & Chiang (2015), we can get a rough estimate for the size of these envelopes and find that K19c should have an H/He envelope of $\approx 4.5\%$ of its total mass and a radius of $\approx 3.2 R_{\oplus}$, while K19d should have a fraction of volatiles around 10-20% and with a radius of $4 - 5 R_{\oplus}$.

Our results confirm that TTV and RV techniques can converge to planetary densities similar to the ones obtained with RV dataset only, when enough data is involved from both sides. This result support the analysis of Steffen (2016), where the discrepancy between planetary density obtained from TTV and RV noted by Weiss & Marcy (2014) is explained as a selection effect rather than an intrinsic problem of one of the two techniques. Previous notable examples of agreement between RV and TTV masses are represented by the WASP-47 (Becker et al. 2015; Weiss et al. 2016) and the *Kepler*-18 systems (Cochran et al. 2011), and the independent RV confirmation by Dai et al. (2016) of the TTV-derived masses of the planets in the K2-19 system (Barros et al. 2015). It should be noted however that Nespral et al. (2016) derived an higher RV mass for K2-19b, while the mass obtained by combining the two datasets resides halfway the two extreme values. In the case of the KOI-94 system, RV (Weiss et al. 2013) and TTV (Masuda et al. 2013) planetary masses agree except for planet *d*, where the TTV mass measurement is half the mass obtained by high-precision RVs. Up to now, most of the targets characterized with TTVs are too faint for a precise, independent mass characterization with RVs. Future space-borne missions such as TESS (Transiting Exoplanet Survey Satellite, Ricker et al. 2014) and PLATO (PLANetary Transits and stellar Oscillations, Rauer et al. 2014) will finally shed light on this problem by providing a large number of targets bright enough for mass measurement with both techniques independently.

The HARPS-N project was funded by the Prodex Pro-




Figure 7. Orbital solution and RV residuals for *K19b* (upper-left panels), *K19c* (lower-left panels) and *K19d* (upper-right panels), phased on the period of the corresponding planet after removing the RV contribution from the other planets. These plots have been obtained using non-interacting Keplerian orbits, and are intended for illustrative purpose only. In the lower-right panels, both the RV residuals after subtracting the dynamical solution from TRADES and their periodogram show no evidence for additional signals that are statistically significant.




Figure 8. Distribution of χ^2_{red} as a function of *K19c* and *K19d* inclinations, with the other orbital parameters fixed to our solution (Table 5). Contour lines for several values of the χ^2_{red} are shown for reference. Values of inclination for which one of the planets would transit are shadowed in gray.

gram of the Swiss Space Office (SSO), the Harvard- University Origin of Life Initiative (HUOLI), the Scottish Universities Physics Alliance (SUPA), the University of Geneva, the Smithsonian Astrophysical Observatory (SAO), and the Italian National Astrophysical Institute (INAF), University of St. Andrews, Queen’s University Belfast and University of Edinburgh.




Figure 9. Analysis of the RV difference (ΔRV) obtained by subtracting to the RV from dynamical simulation (RV_{dyn}) the RV from non-interacting Keplerian orbits (RV_{kep}), assuming planets with the same orbital parameters in both cases. In the upper panel, ΔRV at the HARPS-N epochs and for a 5-year time span are shown as black dots and a black line respectively. The reference time of the orbital parameters T_0 is marked with a dashed red line, to emphasize the increase in ΔRV when moving further from T_0 . In the middle panel, the periodogram of such difference reveals the presence of a periodicity at 29.3 d (red line). As comparison, the periods of the three planets in the system are highlighted in blue. The ΔRV phased at such period is shown in the lower panel, with the sinusoidal model marked with a red line.




Figure 10. Mass-radius for transiting exoplanets with measured masses^b. Planets are color-coded according to the incident flux on the planet F_p , relative to the solar constant F_\oplus . Line thickness reflects the precision on density measurements. Blue error bars identify planets with TTV detection. The dashed lines are theoretical mass-radius curves for different internal compositions, while Earth-like composition and a mixture of Earth-like core plus H/He envelope of 1% in mass are represented by solid red and blue lines respectively (Lopez et al. 2012). The shaded gray region marks the maximum value of iron content predicted by collisional stripping (Marcus et al. 2010). K19b is highlighted with a black dot.

^aAutomatically retrieved in February 2017 from the Nasa Exoplanet Archive, <http://exoplanetarchive.ipac.caltech.edu>

^bAutomatically retrieved in February 2017 from the Nasa Exoplanet Archive, <http://exoplanetarchive.ipac.caltech.edu>

The research leading to these results received funding from the European Union Seventh Framework Programme (FP7/2007- 2013) under grant agreement number 313014 (ETA-EARTH).

This work was performed in part under contract with the California Institute of Technology/Jet Propulsion Laboratory funded by NASA through the Sagan Fellowship Program executed by the NASA Exoplanet Science Institute.

AV is supported by the NSF Graduate Research Fellowship, Grant No. DGE 1144152.

This publication was made possible by a grant from the John Templeton Foundation. The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. This material is based upon work supported by NASA under grant No. NNX15AC90G issued through the Exoplanets Research Program.

PF acknowledges support by Fundação para a Ciência e a Tecnologia (FCT) through Investigador FCT contract of reference IF/01037/2013 and POPH/FSE (EC) by FEDER funding through the program “Programa Operacional de Factores de Competitividade - COMPETE”, and further support in the form of an exploratory project of reference IF/01037/2013CP1191/CT0001.

XD is grateful to the Society in Science–Branco Weiss Fellowship for its financial support.

REFERENCES

- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
- Ballard, S., & Johnson, J. A. 2016, *ApJ*, 816, 66
- Ballard, S., Fabrycky, D., Fressin, F., et al. 2011, *ApJ*, 743, 200
- Baranne, A., Queloz, D., Mayor, M., et al. 1996, *A&AS*, 119, 373
- Barros, S. C. C., Almenara, J. M., Demangeon, O., et al. 2015, *MNRAS*, 454, 4267
- Becker, J. C., & Adams, F. C. 2016, *MNRAS*, 455, 2980
- Becker, J. C., Vanderburg, A., Adams, F. C., Rappaport, S. A., & Schwengeler, H. M. 2015, *ApJL*, 812, L18
- Benatti, S., Desidera, S., Damasso, M., et al. 2016, *ArXiv e-prints*, arXiv:1611.09873
- Bonomo, A. S., Sozzetti, A., Lovis, C., et al. 2014, *A&A*, 572, A2
- Borsato, L., Marzari, F., Nascimbeni, V., et al. 2014, *A&A*, 571, A38
- Borucki, W. J., Koch, D. G., Basri, G., et al. 2011, *ApJ*, 736, 19
- Buchhave, L. A., Dressing, C. D., Dumusque, X., et al. 2016, *AJ*, 152, 160
- Burke, C. J. 2008, *ApJ*, 679, 1566
- Castelli, F., & Kurucz, R. L. 2004, *arXiv, astro-ph*
- Cochran, W. D., Fabrycky, D. C., Torres, G., et al. 2011, *ApJS*, 197, 7
- Cosentino, R., Lovis, C., Pepe, F., et al. 2014, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 9147, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 8
- Coughlin, J. L., Mullally, F., Thompson, S. E., et al. 2016, *ApJS*, 224, 12
- Dai, F., Winn, J. N., Albrecht, S., et al. 2016, *ApJ*, 823, 115
- Demory, B.-O., Gillon, M., de Wit, J., et al. 2016, *Nature*, 532, 207
- Dressing, C. D., Charbonneau, D., Dumusque, X., et al. 2015, *ApJ*, 800, 135
- Dumusque, X., Bonomo, A. S., Haywood, R. D., et al. 2014, *ApJ*, 789, 154
- Duncan, M. J., Levison, H. F., & Lee, M. H. 1998, *AJ*, 116, 2067
- Fabrycky, D. C., Lissauer, J. J., Ragozzine, D., et al. 2014, *ApJ*, 790, 146
- Feigelson, E. D., & Babu, G. J. 2012, *Modern Statistical Methods for Astronomy*
- Figueira, P., Santos, N. C., Pepe, F., Lovis, C., & Nardetto, N. 2013, *A&A*, 557, A93
- Figueira, P., Marmier, M., Boué, G., et al. 2012, *A&A*, 541, A139
- Ford, E. B. 2006, *ApJ*, 642, 505
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
- Fossati, L., Marcellja, S. E., Staab, D., et al. 2017, *ArXiv e-prints*, arXiv:1702.02883
- Gelman, A., & Rubin, D. B. 1992, *Statistical Science*, 7, 457
- Gettel, S., Charbonneau, D., Dressing, C. D., et al. 2016, *ApJ*, 816, 95
- Gladman, B. 1993, *Icarus*, 106, 247
- Gomes da Silva, J., Santos, N. C., Bonfils, X., et al. 2011, *A&A*, 534, A30
- Ioannidis, P., Huber, K. F., & Schmitt, J. H. M. M. 2016, *A&A*, 585, A72
- Jontof-Hutter, D., Ford, E. B., Rowe, J. F., et al. 2016, *ApJ*, 820, 39

- Kass, R. E., & Raftery, A. E. 1995, *Journal of the American Statistical Association*, 90, 773
- Laskar, J. 1993a, *Physica D: Nonlinear Phenomena*, 67, 257
- . 1993b, *Celestial Mechanics and Dynamical Astronomy*, 56, 191
- Laskar, J., Froeschl, C., & Celletti, A. 1992, *Physica D: Nonlinear Phenomena*, 56, 253
- Lee, E. J., & Chiang, E. 2015, *ApJ*, 811, 41
- Lopez, E. D., & Fortney, J. J. 2014, *ApJ*, 792, 1
- Lopez, E. D., Fortney, J. J., & Miller, N. 2012, *ApJ*, 761, 59
- López-Morales, M., Haywood, R. D., Coughlin, J. L., et al. 2016, *AJ*, 152, 204
- Malavolta, L., Nascimbeni, V., Piotto, G., et al. 2016, *A&A*, 588, A118
- Mamajek, E. E., & Hillenbrand, L. A. 2008, *ApJ*, 687, 1264
- Mandel, K., & Agol, E. 2002, *ApJL*, 580, L171
- Marcus, R. A., Sasselov, D., Hernquist, L., & Stewart, S. T. 2010, *ApJL*, 712, L73
- Marzari, F., Tricarico, P., & Scholl, H. 2002, *ApJ*, 579, 905
- Masuda, K., Hirano, T., Taruya, A., Nagasawa, M., & Suto, Y. 2013, *ApJ*, 778, 185
- Mazeh, T., Holczer, T., & Shporer, A. 2015, *ApJ*, 800, 142
- Moré, J. J., Garbow, B. S., & Hillstom, K. E. 1980, *User Guide for MINPACK-1*, Report ANL-80-74
- Mortier, A., & Collier Cameron, A. 2017, *ArXiv e-prints*, arXiv:1702.03885
- Nelder, J. A., & Mead, R. 1965, *The Computer Journal*, 7, 308
- Nespral, D., Gandolfi, D., Deeg, H. J., et al. 2016, *ArXiv e-prints*, arXiv:1604.01265
- Nesvorný, D., Kipping, D. M., Buchhave, L. A., et al. 2012, *Science*, 336, 1133
- Noyes, R. W., Weiss, N. O., & Vaughan, A. H. 1984, *ApJ*, 287, 769
- Parviainen, H. 2015, *MNRAS*, 450, 3233
- Pepe, F., Mayor, M., Galland, F., et al. 2002, *A&A*, 388, 632
- Pepe, F., Cameron, A. C., Latham, D. W., et al. 2013, *Nature*, 503, 377
- Powell, M. J. D. 1994, *A Direct Search Optimization Method That Models the Objective and Constraint Functions by Linear Interpolation*, ed. S. Gomez & J.-P. Hennart (Dordrecht: Springer Netherlands), 51–67
- Rauer, H., Catala, C., Aerts, C., et al. 2014, *Experimental Astronomy*, 38, 249
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2014, in *Proc. SPIE*, Vol. 9143, *Space Telescopes and Instrumentation 2014: Optical, Infrared, and Millimeter Wave*, 914320
- Robertson, P., Endl, M., Cochran, W. D., & Dodson-Robinson, S. E. 2013, *ApJ*, 764, 3
- Rogers, L. A. 2015, *ApJ*, 801, 41
- Snedden, C. 1973, *ApJ*, 184, 839
- Sousa, S. G., Santos, N. C., Adibekyan, V., Delgado-Mena, E., & Israelian, G. 2015, *A&A*, 577, A67
- Southworth, J., Maxted, P. F. L., & Smalley, B. 2004, *MNRAS*, 351, 1277
- Steffen, J. H. 2016, *MNRAS*, 457, 4384
- Torres, G., Fischer, D. A., Sozzetti, A., et al. 2012, *ApJ*, 757, 161
- Torres, G., Fressin, F., Batalha, N. M., et al. 2011, *ApJ*, 727, 24
- Valenti, J. A., & Fischer, D. A. 2005, *ApJS*, 159, 141
- Weiss, L. M., & Marcy, G. W. 2014, *ApJL*, 783, L6
- Weiss, L. M., Marcy, G. W., Rowe, J. F., et al. 2013, *ApJ*, 768, 14
- Weiss, L. M., Deck, K., Sinukoff, E., et al. 2016, *ArXiv e-prints*, arXiv:1612.04856
- Winn, J. N. 2010, *ArXiv e-prints*, arXiv:1001.2010
- Wolfgang, A., & Lopez, E. 2015, *ApJ*, 806, 183
- Wright, M. H. 1996, in *Pitman Research Notes in Mathematics*, Vol. 344, *Numerical Analysis 1995 (Proceedings of the 1995 Dundee Biennial Conference in Numerical Analysis)*, ed. D. F. Griffiths & G. A. Watson (CRC Press), 191–208
- Zechmeister, M., & Kürster, M. 2009, *A&A*, 496, 577