



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Pain related-visual imagery is associated with distress in chronic pain sufferers

Citation for published version:

Gillanders, D, Potter, L & Morris, PG 2012, 'Pain related-visual imagery is associated with distress in chronic pain sufferers', *Behavioural and Cognitive Psychotherapy*, vol. 40, no. 5, pp. 577-589.
<https://doi.org/10.1017/S1352465812000045>

Digital Object Identifier (DOI):

[10.1017/S1352465812000045](https://doi.org/10.1017/S1352465812000045)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Behavioural and Cognitive Psychotherapy

Publisher Rights Statement:

hosted here with consent of copyright holder, Cambridge University Press

Publisher's Version/PDF: subject to Restrictions below, author can archive publisher's version/PDF
General Conditions: Publishers version/PDF may be used in an institutional repository or PubMed Central after 12 month embargo

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Pain Related-Visual Imagery is Associated with Distress in Chronic Pain Sufferers

David Gillanders

University of Edinburgh and Lothian Chronic Pain Service, UK

Louise Potter

NHS Grampian, Elgin, UK

Paul Graham Morris

University of Edinburgh, UK

Background: Chronic pain patients often describe their pain in ways that suggest vivid mental images, with some reporting images relating to their pain. Despite these clinical observations, there are few studies describing the nature and consequences of these images. This study examined whether mental imagery of pain is associated with levels of reported distress, cognitions, disability or pain severity. **Method:** In a postal survey, 83 adult chronic pain patients indicated whether or not they experienced mental images of their pain. They also completed standardized measures of depression, anxiety, cognitions, disability and pain severity. Those reporting images were compared to those not reporting images on their responses to the other variables. **Results:** People with pain imagery reported significantly higher levels of anxiety, depression and catastrophizing than patients who did not report such images. No differences were found on measures of physical disability. **Discussion:** Mental images of pain appear to be associated with higher levels of anxiety, depression and catastrophizing. It is possible that these images play a role in maintaining such difficulties. For these patients, imagery may provide a route via which clinicians can work with patients to help them reinterpret or respond more flexibly to their pain.

Keywords: Chronic pain, mental imagery, cognitive behavioural therapy, catastrophizing, anxiety, depression.

Introduction

The role of cognitive factors in chronic pain has been well documented, with cognitions shown to influence reported levels of pain, levels of disability and treatment outcomes (e.g. Flor and

Turk, 1988; Jensen, Turner, Romano and Karoly, 1991; Turk and Rudy, 1992). Meta-analyses of randomized control trials conclude that cognitive and behavioural therapies are effective for chronic pain (Morley, Eccleston and Williams, 1999; Eccleston, Williams and Morley, 2009). Cognitive behavioural interventions help patients understand how maladaptive beliefs about pain affect emotions and behavioural responses and how these in turn influence the experience of pain.

Many tools have been developed to identify and measure maladaptive beliefs, both in chronic pain samples and other clinical and community samples. Most studies and tools have conceptualized maladaptive beliefs as predominantly verbal in nature. However, cognitive therapy manuals usually encourage clinicians to also enquire about non-verbal dimensions of thinking, such as visual images (e.g. White, 2001; Turk, Meichenbaum and Genest, 1983; Beck, 1976).

Studies of visual imagery have contributed towards our understanding of several psychological disorders, including social phobia (Hackmann, Surawy and Clark, 1998), obsessive-compulsive disorder (de Silva, 1986; Rachman, 2007) and agoraphobia (Day, Holmes and Hackmann, 2004). Characteristic mental images have also been noted in disorders such as depression (Patel et al., 2007), body dysmorphic disorder (Osman, Cooper, Hackmann and Veale, 2004) and bulimia nervosa (Somerville, Cooper and Hackmann, 2007). Research in this area is starting to integrate these image-based cognitions within models of the development and maintenance of disorders, which can then be used to inform imagery-based psychological interventions (Holmes, Arntz and Smucker, 2007).

Most current research on imagery in chronic pain has focused upon generating imagery to facilitate relaxation, rather than on working with spontaneously occurring mental images. In acute pain settings, guided imagery has been reported to be effective in reducing perioperative symptoms (Tusek, Church, Strong, Grass and Fazio, 1997), cancer pain (Sloman, 1995) and chemotherapy-related nausea (Troesch, Rodehaver, Delaney and Yanes, 1993). There has also been some investigation of imagery as a coping response or distracter to experimentally induced pain in non-clinical samples (Fernandez and Turk, 1989). There is currently little empirical evidence about the prevalence of spontaneous pain related mental images in chronic pain populations, the characteristics and functions of such images, or the potential for such images to be targets for psychological therapies. Jamani and Clyde (2008) present a conceptual paper and case study that begins to explore some of these questions, showing how imagery-based cognition in pain may be incorporated into cognitive behavioural models and developed as a potential treatment target.

In addition, Phillips (2011) reports an interview-based study in which chronic pain patients report vivid visual images in association with pain. In Phillips' study, chronic and acute pain patients reported images related to memories, feared future catastrophe and anatomical substrates of pain. These images are described as producing an intense affective response for participants, including tearfulness, flushing and averting gaze. Participants gave ratings of emotions (anxiety, sadness, anger, happiness, calm) as well as pain, using numerical rating scales before and after voluntary image exposure. Participants' ratings on these distress related emotions changed significantly following the exposure procedure; they also reported greater awareness of pain intensity. In addition, participants completed self-report measures of trauma symptoms, depression and anxiety. Whilst around one-third of the images reported in this study were related to a pain causing traumatic event (e.g. a road traffic accident), two-thirds were not trauma related. Level of trauma symptomatology was relatively low in this

sample. Surprisingly, mean scores for the standardized depression and anxiety measures are not reported. It is clear, however that mental images related to the self in pain are associated with distress in this sample. Phillips describes the implications of these images as targets for cognitive behavioural treatment.

Carruthers et al. (2009) describe a study of 109 patients with Irritable Bowel Syndrome (IBS) undergoing hypnotherapy. In this study participants were asked if they had a mental image of their IBS, and they worked with a medical illustrator to produce these pictures. In this study 49% of the participants could report an IBS image, and these were also significantly associated with greater distress. Interestingly, those with mental imagery were more likely to respond to hypnotherapy intervention.

Finally, Berna et al. (2011) report an interview-based study of 10 women with longstanding pelvic pain and significant distress. In their study, all 10 participants reported mental images. The content of these images suggests themes of threat and vulnerability, the relationship the person has with their pain, the sensory qualities of pain, the meaning of pain and the implications it has for the person. All these images are negatively valenced and strongly affective for participants. In addition, 8 of the 10 participants also described comforting or coping images, which they used purposively in response to pain and pain imagery. This study also showed that pain related verbal thoughts are strongly linked to the themes expressed non-verbally in the pain related mental images.

The current paper evaluates the presence of spontaneously occurring pain related mental imagery in persons with chronic pain and uses standardized self-report measures to determine whether those who report experiencing such images differ in terms of emotions, cognitions, physical disability and pain levels from those who report not experiencing such images. The study predicted that those who reported mental images of pain would have higher levels of anxiety, depression, maladaptive cognitions and physical disability than those who reported not experiencing such images. This study examines mental imagery in a broad sample of chronic pain patients and also employs a novel self-report method to determine presence or absence of mental imagery, using a postal questionnaire.

Method

Participants

Participants were 83 attenders at an outpatient based interdisciplinary chronic pain clinic. The service sees patients from the age of 16 years upwards. The current study represents a heterogeneous sample of chronic pain patients, with pain arising from a number of different causes (e.g. musculoskeletal pain, chronic daily headache, neuropathic pain, arthritis) and in different locations of the body. Demographic characteristics of the sample, including, age, gender, duration of pain and self-reported history of mental health difficulties can be seen in Table 1.

Inclusion and exclusion criteria

All patients with a diagnosis of chronic pain who attended chronic pain service clinics were invited to take part, unless they met the exclusion criteria. Patients were excluded if they had pain from a malignant condition such as cancer or if they were not fluent in English. Those

Table 1. Demographic Information

		Whole sample <i>n</i> = 83	Imagers <i>n</i> = 19	Non-imagers <i>n</i> = 64	Test statistic (<i>df</i> 1, 81)	<i>p</i>
Gender	Female	30	7	23	$\chi^2 = .024$.99 <i>ns</i>
	Male	30	7	23		
	Missing	23*	5	18		
Age (years)	Mean (<i>SD</i>)	55.06 (15.3)	46.4 (15)	58.1 (14)	<i>t</i> = 3.349	.001 ^a
Duration of pain (months)		94.7 (91.3)	60.9 (69.3)	91.2 (83.6)	<i>t</i> = 1.78	.08 <i>ns</i>
History of mental health problems	Yes	19 (23%)	7 (36.8%)	12 (18.8%)	$\chi^2 = 3.2$.07 <i>ns</i>
	No	64 (77%)	12 (63.2%)	52 (81.2%)		

*an administrative problem led to a proportion of gender data not being recorded. ^aAlthough the imagery group were significantly younger, age is not correlated with any of the dependant variables. Analyses were performed with and without age as a covariate and the pattern of results was not affected by age.

with a diagnosis of psychotic illness were also excluded, as these individuals may have a different perception of the mental images discussed in this study.

Ethical suitability

Ethical permission was granted by the local NHS Research Ethics Committee (Project reference number: 06/s1104/47) A University Ethics panel and the local Health Board Research and Development Department also approved the project.

Procedure

The questionnaire packs were piloted on five patients attending the clinical psychology service, from which it was estimated that it would take an average of 45 minutes to complete the questionnaire pack. Three hundred and fifty questionnaire packs were then sent to patients who attended the clinic. These packs contained information about the study and a stamped addressed envelope for their return. Eighty-three responses were received (a response rate of approximately 24%). The questionnaire packs contained the following measures:

Hospital Anxiety and Depression Scale (HADS; Zigmond and Snaith, 1983). This measure is widely used in physical health settings as it avoids somatic items that may reflect illness symptoms rather than genuine depressive or anxious symptoms (for example tiredness and changes in appetite). There are 14 questions, each scored from 0 to 3 depending on severity of symptomatology. A review by Bjelland and colleagues reports a cut off of ≥ 8 provides the best balance between sensitivity and specificity, giving both scales scores of approximately .80 for each of these parameters. The same review found Cronbach's alpha for the Anxiety subscale to vary from 0.68 to 0.93 (mean 0.83) and for the Depression subscale from 0.67 to 0.90 (mean 0.82) (Bjelland, Dahl, Haug and Neckelmann, 2002).

Roland and Morris Sickness Impact Profile (R+M SIP; Roland and Morris, 1983). This is a 24-item measure of physical disability in which respondents tick a sentence if it describes

their activity, (e.g. “Because of my pain I lie down to rest more often”). Higher scores indicate greater physical disability. This scale was adapted from the original 136-item Sickness Impact Profile (Bergner, Bobbitt, Carter and Gilson, 1981), which was intended for those with back pain. The R+M SIP compares favourably to the original SIP and has been reported to be more sensitive to change (Deyo and Centor, 1986; Deyo, 1986). Jensen, Strom, Turner and Romano (1992) provide evidence that the R+M SIP scale correlates well with the original SIP ($r = .78$) and is as temporally stable as the original SIP in chronic pain patients with pain in sites other than the lower back (1 month test – retest: $r = .71$).

McGill Pain Questionnaire (Short Form) (McGill-SF; Melzack, 1987). The Short Form of the McGill Pain Questionnaire correlates very highly ($r = .70-.93$) with the standard McGill Pain Questionnaire and has been adopted for clinical trials (Melzack, 1987; Sloman, 1995). The short form gives 15 words that may describe pain (e.g. throbbing, shooting, stabbing) and for each word respondents are asked to indicate whether this type of pain is non-existent, mild, moderate or severe. The words can be used to derive scores for the sensory and affective dimensions of pain. There is also a Likert type rating scale of current pain intensity from 1 – 5. Summing ratings for all words, plus the current intensity score, yields a total score, with higher numbers representing more intense and distressing pain.

Pain Related Control Scale (PRCS; Flor, Behle and Birbaumer, 1993). This scale measures an individual’s beliefs about his/her ability to cope and divides into two factors: Resourcefulness (beliefs about the manageability and predictability of pain, perceived control and ability to cope) and Helplessness (locus of pain control and hopelessness). The two subscales were found to be valid and sensitive to change whilst being closely related to pain intensity and interference from pain experience (Flor et al., 1993). The scale has 15 questions, which are rated from 0 “no, not at all” to 5 “very much”. The scales have adequate reliability (Cronbach’s $\alpha = .77-.83$) and adequate test – retest reliability over a one-week period ($r = .86-.88$).

Pain Related Self Statement Scale (PRSS; Flor et al., 1993). This measures an individual’s thoughts about pain and divides into two factors: Catastrophizing (anxious thoughts associated with loss of control, hopelessness and external locus of control) and Coping (thoughts of coping, predictability and internal control). Both of these subscales were demonstrated to be valid and sensitive to change, and closely related to pain intensity and interference from pain experiences (Flor et al., 1993). There are 18 questions that are rated from 0 “almost never” to 5 “almost always”. The scales have excellent reliability (Cronbach’s $\alpha = .88-.92$) and good test – retest reliability over a 1-week period ($r = .87-.86$).

Pain Anxiety Symptoms Scale (PASS; Roelofs et al., 2004). The PASS is a 20-item scale measuring cognitive, emotional and behavioural responses to chronic pain. It consists of 4 subscales: Pain Related Fear; Cognitive Anxiety; Escape/Avoidance Behaviours; and Physiological Arousal. Respondents answer on a 6-point scale anchored from “never” to “always”; higher scores are indicative of higher levels of pain related anxiety. All subscales have satisfactory reliability (Cronbach’s $\alpha = .73-.84$; total scale = .92); in addition, the PASS has demonstrated good validity via correlations with other relevant measures.

Chronic pain imagery

The Vividness of Visual Imagery Questionnaire (Marks, 1973) is the only standardized measure of visual imagery. However, it focuses upon vividness of intentionally generated

images suggested by the interviewer, rather than examining the presence or content of spontaneously arising mental imagery. Similarly, there are other measures that measure the individual's propensity to use visual imagery such as the Spontaneous Use of Imagery Scale (SUIS: Reisberg, Pearson and Kosslyn, 2003) and the Questionnaire upon Mental Imagery (Sheehan, 1967). None of these measures explore the content of pain related mental imagery. Other studies that have explored mental imagery in either other psychological disorders or in pain have tended to use interview based methods to assess imagery. There is no currently standardized measure of mental imagery in pain.

Consequently, in the absence of a suitable standardized measure of spontaneously arising mental imagery, participants were simply asked to respond to the written question: "Some people report having mental images and/or pictures of their pain, do you have these?" Participants then circled either "yes" or "no". Those participants who reported experiencing mental images of their pain were asked to describe these images in a free text box.

Results

Nineteen respondents (22.9%) reported having mental images of their pain; 12 of these did not have a history of mental health difficulties, whilst 7 of them reported such a history ($\chi^2 = 3.2$, $df = 1$, $p = .07$). It is clear that those who report pain related mental imagery are not simply those who have a history of mental health problems, a potential confounding variable in this novel area of study.

Participants' descriptions of their mental images are outlined in Table 2. As can be seen, the types of images experienced seem very individual and idiosyncratic and do not easily form discrete categories.

Kolmogorov-Smirnov tests and exploratory data analysis indicated that the standardized measures data were sufficiently normally distributed to justify using parametric tests. Effect sizes were calculated using the Dunst guidelines for calculating Cohen's d (Dunst, Hamby and Trivette, 2004), and using the formula provided by Field for calculating r for General Linear Models (Field, 2005, p. 385).

Table 3 outlines differences between those who reported pain related images (imagers) and those who reported no such images (non-imagers) on each of the measures of functional outcome. Whilst, technically, there is no statistically significant difference in self-reported pain severity between imagers and non-imagers, the value of p is close to the accepted cut-off of .05 (McGill-SF total for imagers was 19.4, relative to 14.8 for non-imagers, $F = 3.87$, $df = 1,81$, $p = 0.053$, $r = .15$). Intensity of pain is potentially a confounding variable in the current study, given that the pain imagery group do differ from the non-imagery group on this variable. In order to take a relatively conservative stance the analysis was conducted using ANCOVA to control for the effect of pain intensity on other functional outcomes.

Anxiety and depression

The mean HADS anxiety score for those who reported mental images of their pain (12.89, $SD = 5.39$) was significantly higher than for those who reported that they did not experience such mental images (8.02, $SD = 4.29$). Even when controlling for pain intensity, this difference is statistically significant and is of a moderate effect size ($F = 12.73$, $df = 1,81$,

Table 2. Respondents' description of mental images of their pain

Someone gripping the muscles in my hand
 Fire or a volcano and lava flowing out from my scar
 The displaced bone pressing on the nerve and recalling the anatomical distribution of the nerve and how accurate it is represented by the pain I experience
 A dark cloud
 Someone stabbing and sawing my leg
 Things under my rib cage that they must be able to remove
 A lightning bolt
 Imagining I have a balloon inside me
 Imagine my pain as a solid mass that I push out of my leg
 A sharp kitchen knife right through my knee
 Like someone has set of a thousand fireworks at once and they won't stop
 Being burned alive
 Being poked by knitting needles
 I see my leg as a plastic bottle with one hand clasped tightly around the middle and the other hand twisting the bottle round
 Hot metal wires jiggling up and down my leg
 Flashbulb images related to my accident
 An image of being tormented by the devil and his three pronged fork of pain
 Electric shocks
 Pain as nerve related and amputation would not help

Table 3. Mean differences between imagers and non-imagers

	Mean imagers	<i>SD</i>	Mean non-imagers	<i>SD</i>	<i>F</i> (<i>df</i> _{1, 81})	<i>p</i>	<i>r</i> **
McGill Total*	19.42	9.75	14.81	8.73	3.87	.053	.15
HADS Anxiety	12.89	5.39	8.02	4.29	12.73	.001	.37
HADS Depression	11	6.06	7.56	3.92	6.12	.016	.27
R+M SIP	13.42	6.97	12.19	5.53	.003	.953 <i>ns</i>	–
PRSS: Catastrophizing	27	12.31	19.9	8.72	4.45	.038	.23
PRSS: Coping	20.68	9.70	23.95	7.90	2.5	.12 <i>ns</i>	–
PRCS: Resourcefulness	23.94	6.57	25.03	7.84	1.0	.32 <i>ns</i>	–
PRCS: Helplessness	15.39	8.40	13.35	6.27	.25	.62 <i>ns</i>	–
PASS Cognitive	16.63	7.5	11.98	6.66	3.73	.057 <i>ns</i>	.21
PASS Fear	11.0	8.67	8.03	6.84	.69	.41 <i>ns</i>	–
PASS Escape	13.21	7.59	12.31	6.4	.16	.69 <i>ns</i>	–
PASS Physiology	10.0	7.4	7.13	6.88	.68	.41 <i>ns</i>	–

(HADS = Hospital Anxiety and Depression Scale, R+M SIP = Roland Morris Sickness Impact Profile, PRSS = Pain Related Self Statements, PRCS = Pain Related Control Scale, PASS = Pain Anxiety Symptoms Scale). *All statistical tests are ANCOVA, with McGill Total as a covariate, apart from McGill Total itself. ***r* = effect size estimate: small = .10, medium = .30, large = .50 (Field, 2005, p.33).

p = .001, *r* = .37), suggesting that those who report mental images of their pain also report significantly higher levels of anxiety than those who do not report having images of their pain.

Additionally, those who reported mental images were significantly more likely to score within the range considered to indicate possible clinical anxiety (a score of 8 or above) than those who reported not having such images (84% vs 53%, $\chi^2 = 5.91$, $df = 1$, $p = .015$).

The mean HADS Depression score for those who reported mental images of their pain was 11.0 ($SD = 6.06$) compared to 7.56 ($SD = 3.92$) for those who indicated that they did not experience such images. This result was statistically significant ($F = 6.12$, $df = 1, 81$, $p = .016$) and represents a small to medium effect size ($r = .27$), suggesting that those who report mental images of their pain report higher levels of depression than those who do not report mental images of their pain.

Those who reported mental images were more likely to score within the range considered to indicate possible clinical depression (a score of 8 or above) than those who reported not having such images (73% vs 44%, $\chi^2 = 5.25$, $df = 1$, $p = .022$).

Disability

Those who reported mental images had a slightly higher mean score on the Roland and Morris Sickness Impact profile (mean: 13.42, $SD = 6.97$) than those who did not experience such images (mean: 12.19, $SD = 5.53$), although this was not statistically significant ($F = .003$, $df = 1, 81$, $p = .95$).

Adaptive and maladaptive cognitions

Those reporting mental images of their pain had significantly higher mean scores on the PRSS Catastrophizing scale than those who did not report such images (mean imagers: 27.0, $SD = 12.3$ vs non imagers: 19.9, $SD = 8.72$; $F = 4.45$, $df = 1, 81$, $p = 0.038$, $r = .23$). As can be seen in Table 3, there were no differences between the two groups on the scales of Coping, Resourcefulness or Helplessness.

Pain related anxiety

None of the subscales of the Pain Anxiety Symptoms Scale (PASS) showed any significant differences between the imagery and non-imagery groups. The subscale of "Cognitive Anxiety" did approach significance ($F = 3.73$, $df = 1, 81$, $p = .057$, $r = .21$), which is coherent with the finding related to Catastrophizing.

Discussion

Those who reported mental images of their pain had significantly higher scores on measures of anxiety, depression and catastrophizing than those who reported not experiencing such images. These differences were present despite similar levels of physical disability between the two groups. The group differences were also present after statistically controlling for marginally higher pain intensity being associated with the presence of mental imagery.

There are a number of possible explanations for these findings. It is possible that people who are more distressed are more likely to generate images of their pain. Depression is known to cause people to spend longer ruminating on their predicament (Just and Alloy, 1997) and spending time thinking about their pain may have made the imagers more likely

to generate images of that pain. Rumination is also a key feature of pain catastrophizing and it is noteworthy that imagers were higher in catastrophizing than non-imagers. Alternatively, it is known that some individuals have more of a propensity toward experiencing visually based cognition over more verbal modalities (see for example Reisberg et al., 2003). It may be that when people with this propensity for visual imagery develop chronic pain they are simply more inclined to generate mental images of this pain. This study suggests that the presence of these images may then dispose them towards greater distress. A third possibility is that both the increased distress and the disposition towards experiencing mental images are caused by an unmeasured third factor, possibly a personality variable. Of course, the current study cannot address these issues of causal direction and future research will be required to fully explore this novel clinical phenomenon. This study has shown, however, that chronic pain intensity, pain-related imagery, catastrophizing and distress are related in a substantial proportion of people with chronic pain.

Future studies might be able to determine whether the generation of mental images of pain precedes increased distress, or whether increased distress precedes the generation of mental images. It is likely to be difficult to determine this, as chronic pain patients would by their nature have experienced any such images and distress for some years and any responses as to which occurred first would be subject to retrospective bias.

However, given the possibility that the mental images may play a role in the maintenance of distress amongst some patients, it would seem reasonable to explore whether helping patients to become aware of, understand and perhaps manipulate their pain related mental images may have a role in reducing their distress. Thus, in addition to imagery based relaxation work, which serves to distract individuals from their pain (Main and Spanswick, 2000), there may be a role for interventions that focus upon pain related mental images. This may involve working to change image-based maladaptive cognitions in a similar manner to work undertaken to change verbally based maladaptive cognitions.

It is also likely (though not investigated by the present study) that these images may play a role in cueing avoidance behaviours. Greater awareness of images and their impact upon behavioural responses could help chronic pain patients to reduce unhelpful safety behaviours, as suggested by Jamani and Clyde (2008). Although these authors write from a cognitive modification perspective, it is also possible that mindfulness and acceptance based interventions in relation to chronic pain imagery could help chronic pain patients to develop the psychological flexibility necessary to fully engage in behavioural exposure to feared movements.

Future research would benefit from developing reliable and valid methods of measuring chronic pain imagery, based on a fuller description of the phenomenology of chronic pain imagery. Our research group has now completed two projects further exploring the phenomenology of chronic pain imagery (Gosden, 2008; Lonsdale, 2010). Manuscripts describing these promising results are currently in preparation. In addition, qualitative methods may be useful as a further step in this endeavour. Ultimately, research is needed to investigate the degree to which chronic pain imagery is a suitable target for cognitive behavioural treatment of chronic pain and pain related distress. Such research should attempt to address questions such as whether imagery modification is a useful addition to verbal cognitive modification strategies. As indicated, the function of pain related imagery in inhibiting behavioural flexibility is a further important dimension. Such questions are also shaped by the theoretical or conceptual frameworks adopted by researchers and clinicians and

future research in this area would benefit from developing and articulating a workable and empirically based conceptual framework within which to understand chronic pain imagery, cognitive processes, distress and behavioural flexibility.

Limitations of the study

The questionnaire packs sent to participants took approximately 45 minutes to complete, which may have deterred some respondents. Whilst the 23% response rate is consistent with other surveys within health care settings (e.g. Harrison and Cock, 2004), it is possible that those who participated differed in some respects from those who did not take part. The study was unable to control for the use of medication by the participants, although most pain clinic patients take analgesic medication and some take psychiatric medication, primarily anti-depressants. The use of these pharmaceuticals may have influenced reported severity of pain and reported levels of anxiety and depression. All measures used within the study were self-report measures and thus open to biases. However, the main measures are widely used research tools with good evidence for their validity.

The measurement of chronic pain imagery in this study is at a preliminary stage of development. It is possible that many potential participants decided not to take part because the suggestion of pain related mental imagery may be misinterpreted as suggesting that their pain is “all in the mind”. This may reflect the relatively low response rate in the current study, suggesting that care must be taken in how far these results may be generalized to the larger population of chronic pain patients.

Pain patients may also find it difficult to readily distinguish between thoughts and images and so it is possible that the self-report postal questionnaire method is not the most reliable method of measurement. Furthermore, some of the descriptions in Table 2 appear more imagery-like than others. There remains a possibility that some of these imagery descriptions are simply metaphorical ways of describing pain rather than an actual intrusive image. Using a postal survey method makes it hard to fully disambiguate this. Future studies may need to continue to use interview-based methods whilst we refine our methods of assessing mental imagery.

Interestingly, in our subsequent studies (still using self-report postal survey methods) we have provided a slightly more detailed description of what is meant by pain-related visual imagery, as follows:

“We are particularly interested in finding out if you have a picture or a mental image of what your pain is like. A mental image is like having a picture in your head, which may include things you can imagine seeing, hearing or feeling. Do you ever have a mental image like this of your pain?”

In these studies we find around 40% of chronic pain patients report mental imagery (Gosden, 2008; Lonsdale, 2010). Finally, the link between visual imagery and emotions has been well established, and so the findings here may be interpreted as being relevant to visual images per se, rather than specifically being related to mental imagery of pain. As the current study did not measure other types of imagery, this cannot be ruled out, although our more focused exploration of pain related mental imagery suggests this as a promising line of further enquiry, consistent with the small literature that is developing in this area.

Strengths of the study

Although the assessment of chronic pain imagery was a non-validated single item question, all dependant variables were measured using well-standardized measures that have previously been validated within the chronic pain population. Finally, participants were recruited using very broad inclusion criteria, with few exclusions. This allows the results reported here to be possibly widely generalizable to heterogeneous populations of chronic pain patients.

Conclusions

The present study indicates that a substantial proportion of chronic pain patients experience pain-related mental images and that these individuals have significantly higher levels of anxiety and distress than other chronic pain patients. Whilst there could be several explanations for these findings, it is possible that interventions aimed at manipulating such images may help to reduce distress and increase behavioural flexibility amongst some individuals suffering from chronic pain.

Acknowledgements

We appreciate support from the Lothian Chronic Pain Service and statistical advice provided by Professor Dave Peck and Dr Emily Newman.

References

- Beck, A. T.** (1976). *Cognitive Therapy and the Emotional Disorders*. New York: American Library.
- Bergner, M., Bobbitt, R. A., Carter, W. B. and Gilson, B. S.** (1981) The Sickness Impact Profile: development and final revision of a health status measure, *Medical Care*, 19, 787–805.
- Berna, C., Vincent, K., Moore, J., Tracey, I., Goodwin, G. M. and Holmes, E. A.** (2011). Presence of mental imagery associated with chronic pelvic pain: a pilot study. *Pain Medicine*, 12, 1086–1093. doi: 10.1111/j.1526-4637.2011.01152.x
- Bjelland, I., Dahl, A., Haug, T. T. and Neckelmann, D.** (2002). The validity of the Hospital Anxiety and Depression Scale: an updated literature review. *Journal of Psychosomatic Research*, 52, 69–77.
- Carruthers, H. R., Miller, V., Morris, J., Evans, R., Tarrier, N. and Whorwell, P. J.** (2009). Using art to help understand the imagery of irritable bowel syndrome and its response to hypnotherapy. *The International Journal of Clinical and Experimental Hypnosis*, 57, 162–173. doi: 10.1080/00207140802665401
- Day, S. J., Holmes, E. A. and Hackmann, A.** (2004). Occurrence of imagery and its link with early memories in agoraphobia. *Memory*, 12, 416–427.
- de Silva, P.** (1986). Obsessional-compulsive imagery. *Behaviour Research and Therapy*, 24, 333–350.
- Deyo, R. A.** (1986). Comparative validity of Sickness Impact Profile and shorter scales for functional assessment in low back pain. *Spine*, 11, 951–954.
- Deyo, R. A. and Centor, R. M.** (1986). Assessing the responsiveness of functional scales to clinical change: analogy to diagnostic test performance. *Journal of Chronic Diseases*, 39, 897–906.
- Dunst, C. J., Hamby, D. W. and Trivette, C. M.** (2004). Guidelines for calculating effect sizes for practice based research syntheses, *Centerscope*, 3, 1–10.

- Eccleston, C., Williams, A.C.D.C. and Morley, S.** (2009). Psychological therapies for the management of chronic pain (excluding headache) in adults (Review). *Cochrane Database of Systematic Reviews*, (2), 2–4. doi: 10.1002/14651858.CD007407.pub2
- Fernandez, E. and Turk, D.C.** (1989). The utility of cognitive coping strategies for altering pain perception: a meta-analysis. *Pain*, 38, 123–135.
- Field, A.** (2005). *Discovering Statistics using SPSS* (2nd edn.). London: Sage.
- Flor, H., Behle, D.J. and Birbaumer, N.** (1993). Assessment of pain-related cognitions in chronic pain patients. *Behaviour Research and Therapy*, 31, 63–73.
- Flor, H. and Turk, D. C.** (1988). Chronic back pain and rheumatoid arthritis: predicting pain and disability from cognitive variables. *Journal of Behavioural Medicine*, 11, 251–265.
- Gosden, T.** (2008). *Mental Imagery in Chronic Pain: prevalence and characteristics*. Unpublished Doctoral Thesis, University of Edinburgh.
- Hackmann, A., Surawy, C. and Clark, D. M.** (1998). Seeing yourself through others' eyes: a study of spontaneously occurring images in social phobia. *Behavioural and Cognitive Psychotherapy*, 26, 3–12.
- Harrison, R. and Cock, D.** (2004). Increasing response to a postal survey of sedentary patients: a randomised controlled trial. *BMC Health Services Research*, 4, 31.
- Holmes, E. A., Arntz, A. and Smucker, M. R.** (2007). Imagery rescripting in cognitive behaviour therapy: images, treatment techniques and outcomes. *Journal of Behavior Therapy and Experimental Psychiatry*, 38, 297–305.
- Jamani, N. and Clyde, Z.** (2008). Treatment of pain related fear in chronic (persistent) pain: the role of safety seeking behaviour and imagery. *The Cognitive Behaviour Therapist*, 1, 3–15.
- Jensen, M. P., Strom, S. E., Turner, J. A. and Romano, J. M.** (1992). Validity of the Sickness Impact Profile Roland Scale as a measure of dysfunction in chronic pain patients. *Pain*, 50, 157–162.
- Jensen, M. P., Turner, J. A., Romano, J. M. and Karoly, P.** (1991). Coping with chronic pain: a critical review of the literature. *Pain*, 47, 249–283.
- Just, N. and Alloy, L. B.** (1997). The response styles theory of depression: tests and an extension of the theory. *Journal of Abnormal Psychology*, 106, 221–229.
- Lonsdale, J.** (2010). *Imagery and Emotion in Chronic Pain*. Unpublished Doctoral Thesis, University of Edinburgh.
- Main, C. and Spanswick, C.** (2000). *Pain Management: an interdisciplinary approach*. New York: Harcourt Publishers, Ltd.
- Marks, D. F.** (1973). Visual imagery differences in the recall of pictures. *British Journal of Psychology*, 64, 17–24.
- Melzack, R.** (1987). The short form McGill Pain questionnaire. *Pain*, 30, 191–197.
- Morley, S., Eccleston, C. and Williams, A.** (1999). Systematic review and meta-analysis of randomised controlled trials of cognitive behaviour therapy for chronic pain in adults, excluding headache. *Pain*, 80, 1–13.
- Osman, S., Cooper, M., Hackmann, A. and Veale, D.** (2004). Spontaneously occurring images and early memories in people with body dysmorphic disorder. *Memory*, 12, 428–436.
- Patel, T., Brewin, C., Wheatley, J., Wells, A., Fisher, P. and Myers, S.** (2007). Intrusive images and memories in major depression. *Behaviour Research and Therapy*, 45, 2573–2580.
- Philips, H. C.** (2011). Imagery and pain: the prevalence, characteristics, and potency of imagery associated with pain. *Behavioural and Cognitive Psychotherapy*, 39, 1–18. doi: 10.1017/S1352465811000282
- Rachman, S.** (2007). Unwanted intrusive images in obsessive compulsive disorders. *Journal of Behavior Therapy and Experimental Psychiatry*, 38, 402–410.
- Reisberg, D., Pearson, D. G. and Kosslyn, S. M.** (2003). Intuitions and introspections about imagery: the role of imagery experience in shaping an investigator's theoretical views. *Applied Cognitive Psychology*, 17, 147–160.

- Roelofs, J., McCracken, L., Peters, M. L., Crombez, G., Breukelen, G. van and Vlaeyen, J. W.** (2004). Psychometric evaluation of the Pain Anxiety Symptoms Scale (PASS) in chronic pain patients. *Journal of Behavioral Medicine*, 27, 167–183.
- Roland, M. and Morris, R.** (1983). Study of natural history of low back pain. Part II: development of guidelines for trials of treatment in primary care. *Spine*, 8, 145–150.
- Sheehan, P. W.** (1967). A shortened form of Betts' questionnaire upon mental imagery. *Journal of Clinical Psychology*, 23, 386–389.
- Sloman, R.** (1995). Relaxation and the relief of cancer pain. *Nursing Clinics of North America*, 30, 697–709.
- Somerville, K., Cooper, M. and Hackmann, A.** (2007). Spontaneous imagery in women with bulimia nervosa: an investigation into content, characteristics and links to childhood memories. *Journal of Behavior Therapy and Experimental Psychiatry*, 38, 435–446.
- Troesch, L. M., Rodehaver, C. B., Delaney, E. A. and Yanes, B.** (1993). The influence of guided imagery on chemotherapy-related nausea and vomiting. *Oncology Nursing Forum*, 20, 1179–1185.
- Turk, D. C., Meichenbaum, D. and Genest, M.** (1983). *Pain and Behavioral Medicine: a cognitive-behavioral perspective*. New York: Guilford Press.
- Turk, D. C. and Rudy, T.E.** (1992). Neglected topics in the treatment of chronic pain patients: relapse, non-compliance and adherence enhancement. *Pain*, 44, 5–28.
- Tusek, D. L., Church, J. M., Strong, S. A, Grass, J. A. and Fazio, V. W.** (1997). Guided imagery: a significant advance in the care of patients undergoing elective colorectal surgery. *Diseases of the Colon and Rectum*, 40, 172–178.
- White, C.** (2001). *Cognitive Behaviour Therapy for Chronic Medical Problems: a guide to assessment and treatment in practice*. Chichester: John Wiley and Sons.
- Zigmond, A. S. and Snaith, R. P.** (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67, 361–370.