THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Correlations of particle number concentrations and metals with nitrogen oxides and other traffic-related air pollutants in Glasgow and London

Citation for published version:

Sánchez Jiménez, A, Heal, MR & Beverland, IJ 2012, 'Correlations of particle number concentrations and metals with nitrogen oxides and other traffic-related air pollutants in Glasgow and London', *Atmospheric Environment*, vol. 54, pp. 667-678. <https://doi.org/10.1016/j.atmosenv.2012.01.047>

Digital Object Identifier (DOI):

[10.1016/j.atmosenv.2012.01.047](https://doi.org/10.1016/j.atmosenv.2012.01.047)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Atmospheric Environment

Publisher Rights Statement:

Author's Post-print: author can archive post-print (ie final draft post-refereeing)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Post-print of peer-reviewed article published by Elsevier.

Published article available at: <http://dx.doi.org/10.1016/j.atmosenv.2012.01.047>

Cite as:

Sánchez Jiménez, A., Heal, M. R., Beverland, I. J., 2012. Correlations of particle number concentrations and metals with nitrogen oxides and other traffic-related air pollutants in Glasgow and London. *Atmospheric Environment* 54, 667-678.

Correlations of particle number concentrations and metals with nitrogen oxides and other traffic-related air pollutants in Glasgow and London

Authors:

Araceli Sánchez Jiménez^{1,2 *}

Mathew R Heal³

Iain J Beverland¹

¹Environmental Health Unit, Department of Civil Engineering, Strathclyde University, Glasgow G4 0NG, UK

²Currently at Institute of Occupational Medicine (IOM) Edinburgh

³ School of Chemistry, University of Edinburgh, West Mains Road, Edinburgh EH9 3JJ, UK

* Correspondence to: araceli.sanchez@iom-world.org

Research Highlights

- Particle number concentration (PNC) and transition metals in particulate matter have been associated with health outcomes.
- PNC and metals are difficult and expensive to measure, so they are not routinely monitored.
- NO₂ and NO_x can be used as surrogates for PNC and water-soluble metal at background sites but not at street canyons.
- Weekly NO₂ measurements derived from passive diffusion tubes could be used as metrics for long-term exposure to PNC.

Abstract

Particle number concentration (PNC) and transition metal content are implicated in the health effects of airborne particulate matter (PM) but they are difficult to measure so consequently their temporal and spatial variations are not well characterized. Daily concentrations of PNC and particle-bound water-soluble metals (V, Cr, Mn, Fe, Ni, Cu, As, Cd and Pb) were measured at background and kerbside sites in Glasgow and London to examine if other metrics of air pollution such as optical darkness (absorbance) of collected filter samples of PM, gravimetric PM, and NO, NO₂ and CO gas concentrations, can be used as surrogates for the temporal and spatial variations of the former. NO₂ and NO_x exhibited a high degree of within-site correlation and with PNC and water-soluble metals (Fe, Cu, As, Cd, Pb) at background sites in both cities. There is therefore potential to use NO₂ and NO_x as surrogates for PNC and water-soluble metal at background sites. However, correlation was weaker in complex street canyon environments where pollutant concentrations are strongly affected by local sources and the small-scale variations in pollutant dispersion induced by the wind regimes within street canyons. The corollary of the high correlation between NO₂ and PNC and water-soluble metals at the background sites is that the latter pollutants may act as confounders for health effects attributed to NO₂ from such sites. Concentrations of CO cannot be used as a surrogate for PNC. Increments in daily NO_x and NO₂ concentrations between trafficked and background sites were shown to be a simple and novel surrogate for daily spatial variation of PNC; for example, increments in NO_x explained 78-79% of the variance in PNC at the paired sites in both Glasgow and London, but relationships were city specific. The increments in NO_x also explained 70 % of the spatial variation in Cu and Ni in Glasgow but not in London. Weekly NO₂ measurements derived from passive diffusion tubes were also shown to correlate well with increments in PNC. A high temporal correlation between PNC and 1,3-butadiene and benzene (which can also be measured by passive sampler) implies that passive sampler measurements may be a straightforward tool for deriving long-term spatial patterns in PNC.

Keywords: particle number concentration, nitrogen oxides, transition metals, air pollution, Glasgow, London

1 Introduction

Epidemiological studies have shown consistent associations between health outcomes and exposure to airborne particulate matter (PM) (Hoek et al., 2002; Pope et al., 2004). Toxicological studies suggest that specific components within particles and the number of particles, more than particle mass alone, are responsible for these health effects. Particle-bound transition metals (especially Fe, Cu, Ni, V, and Zn) and ultrafine particles have been associated with inflammatory responses in human cells (Jiménez et al., 2000; Wilson et al., 2002, Donaldson et al., 2005; and Monteiller et al., 2007). Ultrafine particles are better characterized by the particle number concentration (PNC) rather than by particle mass concentration, as they dominate the former but contribute negligibly to the latter.

PNC and particle-bound metals are expensive and time-consuming to measure and, except for a few specific metals, their ambient levels are not currently subject to regulation. They are therefore not routinely monitored in air quality networks. Consequently there is limited epidemiology on their health effects. Sulphates of Fe, Ni and Zn in PM_{2.5} were associated with mortality in a time-series study of 8 cities in Canada from 1986 to 1996 (Burnett et al., 2000). The effect of the metals was greater than for the mass alone. In contrast, a study in Edinburgh (UK) did not show stronger associations between metals and adverse acute cardiopulmonary health outcomes after adjustment for PM₁₀ mass (Heal et al., 2009). A growing number of studies have reported associations between ultrafine particles and cardiovascular disease (von Klot et al., 2005, Lanki et al. 2006, Nicklas et al. 2009, Hoek et al., 2010) and mortality (Hoek et al., 2010).

It would be useful, for epidemiological studies in particular, if other easier and less expensive to measure pollutant metrics (e.g. NO, NO₂ or PM optical absorption) could be used as quantitative relative measures of the concentrations of transition metals and PNC. Total metal content in PM₁₀ has been found to correlate with NO_x, PNC and PM light absorption (Harrison et al., 2003; Götschi et al., 2005). Less attention has been paid to the water-extractable fraction, despite its likely greater bioavailability in the human respiratory system (Costa et al., 1997; Adamson et al., 2000). Heal et al. (2005) reported significant correlations

between the water-extractable fraction of metals in PM_{10} and $PM_{2.5}$ and black smoke (equivalent to filter darkness) at an urban background site in Edinburgh.

The aim of this study was to examine the spatio-temporal associations of transition metals and PNC with traffic-related air pollutants at paired kerbside and urban background sites in two cities with a view (1) to investigate a 'simple tracer' for transition metals and PNC, (2) to determine the extent of between and within site inter-pollutant correlations to facilitate better interpretation of epidemiological studies of the health effects of traffic-related air pollution.

2 Experimental methods

2.1 Sampling sites

Associations between traffic-related air pollutants were evaluated for central Glasgow and central London. Both are large, congested cities in which road traffic is the major contributor of local PM_{10} , NO_2 , CO, benzene and 1,3-butadiene, and which have nearby kerbside and urban background monitoring stations. This enables comparison of the extent of consistency in within-site inter-pollutant relationships across locations with differing levels of air pollution and of the inter-pollutant relationships in the roadside increments of the pollutants (concentrations at the kerbside site with the concentrations at the background site subtracted).

All monitoring sites are part of the UK national Automatic Urban & Rural Network (AURN). The Glasgow sites were located at:

- Hope St. (kerbside, 55° 51' 30" N, 4° 15' 11" W): Hope St. is a busy road with two carriageways each way and average traffic flow of around 25,000 vehicles per day comprising mostly buses and taxis. The street is orientated NS and forms a 30 m high (20 m wide) symmetric canyon with the monitoring site located on the E side of the street. A junction ~10 m S of the monitoring station is subject to frequent congestion during peak traffic periods. The monitor inlets are ~1 m from the kerb and 3 m above the ground. The monitoring station is located on the downwind side of the street.
- St. Enoch Sq. (now designated urban background but formerly designated urban centre, 55° 51' 84" N, 4° 15' 45" W): St. Enoch Sq. is a pedestrian area surrounded by commercial premises. The nearest road is ~20 m N from the site and is orientated WE with typical traffic

flows up to 20,000 vehicles per day. The monitor inlets are ~3 m above the ground. The site is only ~300 m E of the kerbside site in Hope St. and is therefore a good background site for evaluating the contribution of traffic to pollutant concentrations at the latter.

- Montrose St. (urban background, 55° 51' 22" N, 4° 14' 44" W): monitor inlets at this site are located on the second floor of Glasgow City Chambers ~12 m above ground level facing south. Montrose St. is a symmetric street canyon of depth ~30 m, but width ~10 m, with typical traffic flows in the range 8,000-15,000 vehicles per day.

The London sites were located at:

- Marylebone Rd. (kerbside, 51° 31' 21" N, 0° 9' 17" W): Marylebone Rd. is a very congested road with three carriageways in both directions and carrying around 80,000 vehicles per day. The carriageway adjacent to the monitoring station is bus and taxi only. The surrounding area forms an asymmetrical street canyon orientated in a NE-SW direction with the monitoring station on the E side of the street. The canyon width is ~40 m and the average building height is ~25 m, comprising educational and commercial buildings plus housing. The monitor inlets are ~1 m from the kerb and ~3 m above the ground and located on the upwind side of the street.

- North Kensington (urban background, 51° 31' 15" N, 0° 12' 49" W): this site is in a residential area in the grounds of a school, 4 km S of Marylebone Rd. The inlets are 3 m above the ground.

Further information on the monitoring sites can be found at <http://uk-air.defra.gov.uk/>.

2.2 Sampling instrumentation

Instrumentation and data sources at each site are summarised in Table 1. Sampler operation and gravimetric determination of filter masses at the London sites was undertaken as part of the London Air Quality Network. Sampler operation in Glasgow was carried out as part of this project with the exception of the butanol-based condensation particle counter (CPC), which was operated by National Physical Laboratories as part of the UK network. Filters collected in London were shipped to Glasgow for the optical and chemical analyses. Samples for metal analysis were collected using a Partisol 2025 (Rupprecht & Patashnick Co. Inc.) with a PM₁₀ size-selective inlet at Glasgow (on to Zefluor PTFE membrane filters) and a

PM_{2.5} inlet at London (on to quartz filters). Filters were changed automatically at midnight and total daily volume of air sampled was 24 m³. In Glasgow, the Partisol sampler was operated in accordance with guidance specified by the USEPA (1998); pre- and post-exposed filters were conditioned for 24 h in a temperature and humidity controlled room before being weighed three times on an analytical balance of 1 µg precision (Sartorius MC5) equipped with anti-static ionising blower. A set of 6 blank filters were weighed together with the pre- and post-exposed filters and their mean mass change used to correct the mass change of the sample filters.

The optical absorbance of the sample filters (PM₁₀ and PM_{2.5}) was determined by measuring sample reflectance, R , with an EEL Model 43D reflectometer (Diffusion Systems Limited, London, U.K), and converting to an absorption coefficient, α (m⁻¹), with the relationship $\alpha = \frac{A}{2V} \ln\left(\frac{R_0}{R}\right)$, where R_0 is the reflectance of an unloaded filter (100%), A the exposed area of the filter (m²) and V the volume of air sampled (m³).

PNC at Hope St. and Montrose St. in Glasgow were measured using a water-based, CPC (TSI Model 3785, minimum detectable D_{50} particle diameter: 5 nm) operated according to manufacturer instructions. A butanol-based CPC (TSI Model 3022A; minimum detectable D_{50} particle diameter: 7 nm) was used at St. Enoch Sq. and at the London sites. PNC data were recorded at 1 min and 15 min resolution for the water- and butanol-CPCs, respectively, and averaged as required.

Hourly concentrations of nitrogen oxides (NO, NO₂ and NO_x), CO, PM₁₀ as measured by TEOM, 1,3-butadiene and benzene, and daily black smoke (BS), were obtained from the UK Air Quality Archive (<http://uk-air.defra.gov.uk/>). The gases 1,3-butadiene and benzene are not measured at St Enoch Sq. and North Kensington and BS is only measured at Marylebone Rd.

One-week averaged NO₂ was also measured at St. Enoch Sq. using Palmes-type diffusion tubes (PDTs). Tubes were prepared and analysed colorimetrically, as described by Heal et al. (1999). Tubes were exposed in duplicate together with a field blank.

Hourly meteorological data were obtained from the British Atmospheric Data Centre (www.badc.nerc.ac.uk) for the Bishopton site ~18 km W of Glasgow city centre and for Heathrow Airport ~16 km and 20 km W of the London North Kensington and Marylebone Rd. sites, respectively.

In Glasgow a total of 16/17 days of metal data were collected between 8 June and 31 August 2006. Individual pollutants had longer time series measurements. NO₂ PDTs data was collected over 12 weeks. In London 26 days in 2006 were chosen on the basis of providing the maximum pairs of pollutant concentrations at both sites simultaneously. These days were: 13, 15, 16 May; 23, 24, 25, 26 June; 24, 25, 26, 31 October; 3, 8, 11, 12, 13, 17, 23, 27 November; 4, 6, 8, 10, 12, 14, 15 December.

2.3 Chemical analysis

The Glasgow PM₁₀ filters were sequentially extracted for polycyclic aromatic hydrocarbons (PAHs) and water-soluble metals following a method reported by Piñeiro-Iglesias et al. (2003). Each filter was first digested in 2.5 mL of 1:1 v/v hexane:acetone in an ultrasonic bath for 30 min to extract the PAHs. D-10 acenaphthene and D-12-indenol (1,2,3-c,d) were added as internal standards to yield a final concentration of 10 ng mL⁻¹. Extracts were concentrated to 0.1 µL with a flow of N₂. Filters were then digested in 7 mL of ultrapure water (18.4 MΩ) in an ultrasonic bath for 1 h following a protocol described by Heal et al. (2005). The first extracts were analysed for (benzo(b,k)fluoranthene, benzo(a)pyrene, dibenzo(a,h)anthracene, benzo(g,h,i)perylene and indenol(1,2,3-cd)pyrene) by gas-chromatography mass spectrometry (GC-MS) at the Natural Environment Research Council (NERC) facility at the University of Bristol. Concentrations for all PAHs were below the limit of detection (LOD) and are therefore not discussed further here. The second extracts were analyzed for V, Cr, Mn, Fe, Ni, Cu, As, Cd and Pb by inductively coupled plasma mass spectroscopy (ICP-MS, Thermo PlasmaQuad instrument) at the NERC facility at Kingston University. The PM_{2.5} filters from the London sites were subject only to the extraction step for metal determination, but otherwise the procedure was as described above.

3 Quality Control

3.1 Limits of detection

The LOD for PM₁₀ was calculated as 3 times the mean standard deviation (SD) of day-to-day mass change of blank filters $\times \sqrt{\frac{n+1}{n}}$, divided by the nominal sampling volume, as defined by Vaughan et al. (1989), and was 0.1 $\mu\text{g m}^{-3}$.

The LODs for each metal were defined as three times the standard deviation of the concentrations in the field blanks (Table 2). The median concentration of the field blanks was subtracted from the concentration measured in the exposed filters.

The LOD for NO₂ PDT measurement, defined as the mean of the field blank tubes plus three times the SD of the blanks, was calculated to be equivalent to 6.4 $\mu\text{g m}^{-3}$. NO₂ measurements were also field blank corrected with the corresponding blank for each week and site.

All PM₁₀ and NO₂ measurements were above their LODs. There were only three instances of metal concentration below its LOD, one each for Ni and Cd at Hope St. and one for Ni at St. Enoch Sq., and these were retained in the dataset as recommended by Brown et al. (2008).

3.2 Precision

The precision of the PDTs expressed as the coefficient of variation (CV) from duplicate measurements was 8.8%. The temporal correlations of the PDTs and the co-located chemiluminescence analyser was high, with R values of 0.79 ($n=12$).

An inter-comparison study at St. Enoch Sq. between the water-CPCs showed a CV of 7% and $R = 0.98$ ($n = 624$, 15-mins average). The comparison of the two water-CPC and butanol-CPCs showed CVs of 25% and 28%. The water-CPC read lower than the butanol-CPC for PNC $>30,000$ particles cm^{-3} (as measured by the butanol-CPC), possibly because of reduced condensation efficiency in the particle growth tube in the water-CPC. Results from the inter-comparison study are shown elsewhere (Sánchez Jiménez, 2010).

3.3 Metal losses during PAH extraction

To examine potential losses of particle-bound metal during the PAH extraction step of the PM₁₀ filters, parallel determinations were undertaken on filters undergoing both extractions or the metal extraction step only. Six filters (F) were cut in half (h), (F₁h₁, F₁h₂, F₂h₁, F₂h₂...) and two halves of different filters underwent extraction together so that the metal concentration in F₁h₁&F₂h₁ was the same as the metal concentration in F₁h₂&F₂h₂ (assuming the metals were distributed homogeneously in the filters). Although loss of metal during the PAH extraction step could be substantial (5-60%), it was systematic as indicated by the large correlation ($R = 0.97$) between the metal concentration in the pairs of filters undergoing either one or two extraction steps. Therefore, although metal concentrations in the Glasgow data have been underestimated this does not affect the strength of the correlations in the relationships between metals and co-pollutants.

4 Statistical analysis

Pollutant measurements were averaged to produce 24 h concentrations, as this is the averaging period used in time-series epidemiology with these metrics. Daily and hourly data were not normally distributed so non-parametric Spearman correlation coefficients were used to describe the strength of the correlation. Hourly and daily correlations were examined to determine which averaging period predicted better the PNC. In addition, the inter-pollutant correlations between daily averages was examined to determine whether PNC and metals can act as confounders or collinear pollutants on the associations between health outcomes and nitrogen oxides and PM. Analyses were conducted using SPSS 16.0. Correlation coefficients between 0.50-0.69 were referred to as moderate, above 0.69 as high and below 0.50 as low.

The inter-pollutant relationships, including those between roadside increments, were examined using reduced major axes regression which allows for statistical uncertainty in both variables (Ayers, 2001). Coefficients of determination (R^2) were used to describe the strength of the association.

5 Results

5.1 Pollutant concentrations

A summary of the pollutant concentrations at each location is given in Table 3. Direct comparison between the pollutant concentrations at London and Glasgow is compromised because the sampling periods and sample sizes were different for both cities. The data from London were selected to maximise the range of observed pollutant concentrations. The datasets were also of insufficient duration to represent intra-annual and inter-annual variations. Nevertheless, concentrations of PM and gaseous pollutants were much higher at the kerbside sites (Hope St. and Marylebone Rd.) than at the background sites (North Kensington, St. Enoch Sq. and Montrose St.) because of the major impact of nearby traffic at the former sites. The variability of pollutant concentrations, expressed as the inter quartile range (IQR), was higher at the London sites than in Glasgow, likely because the data for London were from 26 days spread through winter and summer whereas in Glasgow the data were from a consecutive period in summer only.

5.2 Correlations of the same pollutant measured at different sites

Individual pollutants showed low to moderate correlations between kerbside and background sites in each city (Table 4). Correlations between Hope St. and St. Enoch Sq. in Glasgow were greater for PM₁₀ ($R = 0.67$ and 0.73 , for hourly and daily averaged data, as measured by TEOM) than for PNC ($R = 0.47, 0.43$) and NO_x ($R = 0.52, 0.33$). The contrast in absolute pollution concentrations between the kerbside and background sites was greater in London than in Glasgow but between-site correlation coefficients for the particle metrics PM₁₀ ($R = 0.50, 0.23$) and PNC ($R = 0.58, 0.46$) were similar. The correlation of NO_x between the two London sites was lower, particularly for daily averages ($R = 0.43, 0.06$) which appears to be driven by the much greater local NO_x source strength at the highly polluted Marylebone Rd. site. Concentrations of NO at North Kensington (background) were very low ($<5 \mu\text{g m}^{-3}$) except on the 3 November where concentrations reached $200 \mu\text{g m}^{-3}$ (data points deviating from the general trend in the scatter plot in Table 4). On this day, PM₁₀ concentrations in North Kensington were also higher (scattered points in Table 4). This was probably the result of the bonfires that took place to celebrate Guy Fawkes' Night, which fell on the following Sunday (5 November). When data for this day were removed from the dataset, correlation coefficients did not change substantively.

PM₁₀, PNC and constituent NO_x species correlated more strongly between the two background sites in Glasgow (Montrose St. and St. Enoch Sq.), particularly NO_x, NO₂ and NO (R values ≈ 0.8) (Table 4), than between kerbside and background sites, as expected given the absence of strong kerbside source influences at the background sites.

CO concentrations were more weakly correlated between sites, if at all, with the exception of between Hope St. and Montrose St. but the low variability in CO concentrations at St. Enoch Sq. (Table 4) reduces the numerical ‘discrimination’ in statistical calculations.

The averaging period did not significantly influence the correlation coefficients between the Glasgow sites. However, for the London sites, hourly averaging yielded greater correlation coefficients than daily averaging, especially for NO_x, NO₂ and NO.

For particle-bound metals, correlations between the kerbside and background sites in London were highly significant ($P < 0.01$) for each metal measured (Table 5). In Glasgow, between kerbside and background site correlations were high for V and As ($R > 0.7$), moderate for Mn ($R = 0.5$) but low for the remaining metals quantified (Table 5). The lower between-site correlations between metals in Glasgow may be due to greater inaccuracies in the metal concentration because of losses during the sequential extraction.

5.3 Correlations of different pollutants measured at the same site

At all sites in this study, PNC and NO_x were the pollutant pair with the greatest and most consistent inter-pollutant correlations ($R = 0.60$ - 0.90 , Tables 6-8). Generally, there were no substantive differences between correlations derived from daily and hourly averaged data (hourly data not shown), with the exception of Hope St. At this site daily averaged data had greater R values than their hourly counterparts, presumably due to the underestimation at high PNCs by the water-CPC.

Correlations between PNC and PM₁₀ were high only at Hope St. ($R = 0.74$). At the other sites the correlation coefficients were <0.5 .

In Glasgow, correlations between PNC and optical absorbance of PM₁₀ were low ($R = 0.47$ and 0.35 at St. Enoch Sq. and Hope St., respectively) (Table 6). In contrast, in London, PNC was more strongly correlated with PM_{2.5} optical absorbance at the more polluted site (Marylebone Rd., $R = 0.87$) than at the background site (North Kensington), $R = 0.34$ (Table 8).

Correlations between PNC and 1,3-butadiene or benzene were moderate to high ($R = 0.71$ and 0.55 , respectively) at Hope St. (Table 6) and high ($R = 0.81$ and 0.78 , respectively) at Marylebone Rd. (Table 8).

Correlations between PNC and CO varied substantially across the sites ($R = 0.27$ - 0.70 , Tables 6-8).

In Glasgow, the strength of the correlations of metals with PNC were generally greater than with PM₁₀ or with PM₁₀ optical absorbance. Correlations between metal concentrations and NO_x or NO₂ were high at St. Enoch Sq. but not at Hope St. At St. Enoch Sq., Mn, Fe, Cu, As, Cd and Pb were significantly correlated with NO₂ and NO_x ($R = 0.6$ - 0.7), PNC ($R = 0.7$ - 0.8), PM₁₀ ($R = 0.5$ - 0.8) and PM₁₀ optical absorbance ($R = 0.5$ - 0.8) (Table 6). Cr was also significantly correlated with PNC and NO_x and NO₂ ($R \approx 0.7$) but not with PM₁₀ and PM₁₀ optical absorbance ($R \approx 0.4$), and Ni showed moderate correlation with PNC and PM₁₀ ($R \approx 0.6$) but not with nitrogen oxides and PM₁₀ optical absorbance ($R = 0.3$ - 0.4). CO did not show significant correlation with any of the metals.

A different pattern of inter-pollutant correlations was present at Hope St. For example, unlike at St. Enoch Sq., Fe, As, Cd, and Pb were not significantly correlated with PNC, NO_x or NO₂ (Fe was significantly but negatively correlated with NO_x and NO₂). However, V, Cr, Mn, Ni and Cu were significantly correlated with PNC ($R = 0.7$ - 0.9), Ni and Cu were significantly correlated with PM₁₀ ($R = 0.58$ and 0.70 , respectively) and Ni was also highly correlated with

NO₂ ($R = 0.75$) (Table 8). None of the metals were correlated with PM₁₀ optical absorbance except Cd ($R = 0.50$). Concentrations of 1,3-butadiene and benzene (available only at Hope St.) were moderately correlated with Cu ($R = 0.58$ and 0.52 , respectively), but not with the other metals (Table 8).

At Marylebone Rd. (Table 8) there was no significant correlation between any of the particle-bound metals and NO_x, NO₂, PNC and optical absorbance of PM_{2.5}, except for V and Ni with PM₁₀ ($R = 0.48$ and 0.44 , respectively) and for Mn and Fe with PM_{2.5} ($R = 0.54$ and 0.48 , respectively). Curiously, Cu concentrations were significantly negatively correlated with NO_x, NO₂, PNC, BS and optical absorbance of PM_{2.5}.

At North Kensington (Table 8), Fe, Cu, Zn, As, Cd and Pb were moderately to highly correlated with NO_x and NO₂ ($R \approx 0.5$ - 0.8). These metals (except for Cu) were also moderately to highly correlated with PM_{2.5} ($R \approx 0.6$ - 0.8) and moderately correlated with PM₁₀ ($R \approx 0.5$). Metal concentrations did not correlate with PNC. A few metals, Cu, As, Cd and Pb, had low positive correlations with CO ($R \approx 0.5$).

5.4 Effects of wind on pollutant correlations

In both cities the prevailing wind direction was W-SW as is usual in the UK. At Marylebone Rd. the monitoring station was located at the upwind side of the street so southerly winds (perpendicular to the street) may induce a recirculation vortex blowing pollutants against the station. This might explain the observed increase in the increments in pollutant concentrations (i.e. the on road emissions at Marylebone Rd.). In contrast, Hope St. was located on the downwind side of the street, relative to westerly winds. Therefore, wind blowing in this direction dilutes pollutant concentration resulting in a decrease of the increments and absolute concentrations.

At the background sites, St. Enoch Sq. and North Kensington, wind speed was negatively correlated with all pollutants. At the street canyon sites pollutant concentrations varied in their response to wind speed. PNC was positively correlated with wind speed at Marylebone Rd. ($R = 0.40$) but negatively correlated at Hope St. ($R = -0.52$). PM₁₀ was not correlated with wind

speed at Marylebone Rd. ($R = -0.02$), and showed a moderate but not significant negative association at Hope St. ($R = -0.40$). Metal concentrations decreased with increasing wind speed at both sites. NO_x and NO_2 showed a small positive association with increasing wind speed at Marylebone Rd. ($R = 0.36$ and 0.41), and a moderate negative correlation at Hope St. ($R = -0.47$ and -0.51 , respectively).

5.5 Regressions of roadside increments of pollutants

Table 9 summarises the relationships between the increments in pollutant concentrations at the kerbside sites (Hope St. and Marylebone Rd.) and their respective background sites (St. Enoch Sq. and North Kensington). The increments in NO_x explained 79% and 78% of the variance in the increments in PNC in Glasgow and London, respectively. At St. Enoch Sq., NO_2 as measured by the PDTs explained 70% ($n = 12$) of the variance in the weekly averaged PNC.

The increments in PM_{10} explained 70% of the variance in the increments in PNC in Glasgow and 55% in London ($p < 0.05$ in Glasgow and $p < 0.001$ in London). The increments in CO explained less than 50% of the variance in both cities indicating CO cannot be used as a surrogate for PNC.

The increments in NO_x and NO_2 were in general not strongly associated with the increments in metal concentrations (Table 10). Only in Glasgow the increments in NO_2 explained 47% of the variance in Ni and 67% of the variance in Cu. The increments in NO_x explained 65% of the variance in Cu. The increments in PNC showed a strong correlation with the increments in Cu and Ni in Glasgow, explaining 66% and 76% of the variance, respectively. In London explanatory power between the remaining metals and co-pollutants was below 40% and are not shown.

6 Discussion

6.1 *Spatial variability of air pollutants*

Direct comparisons between the concentrations at London and Glasgow have to be made with caution because sampling periods and sample sizes were different in the two cities.

A metal enrichment at the kerbside compared to background site St. Enoch Sq. was observed in Glasgow, reflecting the contribution of traffic to metal concentrations. That this trend was not observed at the London sites, where concentrations of V, Pb, As and Ni at North Kensington were higher than at Marylebone Rd. could be the result of a local contribution of metals in water-soluble form at North Kensington. These emissions could originate from a Ni refinery located 3.2 km NW from North Kensington (data retrieved from the UK Environment Agency). It could be that on the timescale in which the air reaches Marylebone Rd. these metals are oxidised and therefore insoluble in water.

6.2 *Correlations between air pollutants*

Nitrogen oxides showed the most consistent associations with PNC across all the studied sites in Glasgow ($R = 0.7-0.8$) and London ($R = 0.6-0.9$) for hourly and daily averaging periods and therefore would be the best surrogate for PNC. The averaging period, hourly and daily did not significantly influence the strength of the correlation coefficients, suggesting that correlations are mostly driven by the diurnal pattern of traffic flows. If the correlations between pollutants were mainly driven by meteorology and long-transport, then correlations between daily averages would be higher than their hourly counterparts, since the homogenising effects of meteorology and long-transport are more noticeable over an entire day rather than on an hourly basis. The slightly higher correlation coefficients at the London sites for hourly data is attributed to the larger contrast in the pollutant concentrations, compared to Glasgow, which result in a higher statistical power. Daily averages are principally reported here as this is the relevant exposure period for time-series epidemiological studies.

Pollutants were highly correlated at locations close to traffic source but considerably less so a few tens of metres away from the traffic source, which supports the supposition that correlations are driven by traffic sources.

6.2.1 Correlations of the same pollutant measured at different sites

The correlations of PNC between sites observed in this study are lower than those reported elsewhere in Europe. For example, Aalto et al. (2005) reported correlation coefficients for daily PNC between urban background and traffic sites of 0.67, 0.69 and 0.84 in Rome ($n = 293$), Stockholm ($n = 534$) and Barcelona ($n = 89$), respectively, and Puustinen et al. (2007) reported correlation coefficients in the range of 0.67-0.76 between background sites and high and low traffic sites in Amsterdam, Athens, Birmingham and Helsinki for the years 2002 to 2004. The correlation coefficients for hourly averaged data were lower (0.56-0.66). However, different studies have different characteristics such as distance between monitoring sites, sample sizes, and, in particular, meteorological influences. Higher correlation between background and roadside sites likely reflects the dominance of the homogenising effect of meteorology, whereas lower correlation indicates that local, traffic-related sources of the pollutant are important at the roadside site. The moderate to low correlations of PNC between paired sites in this study suggest the latter. The between-site correlations for PM_{10} and NO_2 in this study show site dependence: for example, correlation coefficients for NO_2 ranged from 0.37 between the two sites in London to 0.5-0.8 between pairs of sites in Glasgow. These results indicate that measurements of these pollutant metrics at one site may not well represent daily variations at other locations, so assigning PNC, PM_{10} and NO_2 data from a single monitoring site as the exposure for a large urban population in epidemiological studies may lead to exposure misclassification.

PNC did not show higher correlations between the background sites compared to background vs. traffic site in Glasgow, as observed for nitrogen oxides. This indicates that nitrogen oxides are more homogeneously distributed than PNC, suggesting meteorological factors affect in a different way the formation and dispersion of both pollutants. In addition PNC do not only depend on traffic emissions but also on particle formation processes, nucleation of particles by condensation of pre-existing vapours and condensation of vapours into existing particles (Kulmala et al., 2004). These can be different at each site, resulting in a lower concentration between sites compared to NO_x and NO_2 which mostly originate from traffic and follow the same pattern across sites.

6.2.2 Correlations between PNC and nitrogen oxides measured at the same site

The within-site correlation coefficients observed in this study between PNC and NO_x or NO₂ were higher at the roadside sites, where concentrations are driven by traffic emissions, than at the urban background sites, where concentrations can be influenced by local sources resulting in weaker correlations.

Harrison and Jones (2005) in a study that examined the correlations between hourly PNC vs. NO_x at five UK sites including St. Enoch Sq., North Kensington and Marylebone Rd. reported similar correlations to those found in this study at North Kensington and Marylebone Rd. (0.51, 0.76, respectively) and slightly lower at St. Enoch Sq. ($r=0.42$).

The higher correlations found at the roadside compared to the background sites suggest that the use of daily NO_x and NO₂ concentrations as surrogates for daily PNC is appropriate at trafficked sites but not at background sites.

6.2.3 Correlations between water-soluble metals and other traffic-related air pollutants

The large correlation coefficients between paired sites for metals reveal that the day-to-day variation in metal concentration is similar at both sites. The low correlation coefficients of Fe and Cu between the London sites, compared with those for V and As, and the lack of correlation of Cr, Ni and Pb between the Glasgow sites could be the result of the higher uncertainty on the calculation of the concentrations of these metals, as their concentrations were close to their respective LODs. The high correlation between metals but not PM_{2.5} concentrations between the London sites suggests different dispersion processes for both pollutants. This is partly supported by the fact that metals were negatively correlated with wind speed at both sites, whereas PM_{2.5} was negatively correlated with wind speed at North Kensington but not at Marylebone Rd.

The different pattern of correlations between metals and PNC between background sites and street canyon, and between both cities, is attributed to the different dispersion in open spaces compared with street canyons. Dispersion of on-road emissions within street canyons is a function of the orientation of the canyon relative to wind direction. The wind regimes at both

sites influence the dispersion of pollutant in different ways, as indicated by the different correlations between wind speed and pollutant concentrations at each site. Therefore, the position of the monitoring station in the canyon becomes very important when measuring the concentration of pollutants. Results from this study showed that for open spaces but not for street canyons NO_2 appear to represent well the temporal variations of water-soluble metals. However, in the street canyons the relationships found in this study may not translate to other urban locations. Harrison et al. (2005) also found low correlations for PNC vs. NO_x at street canyon sites. Noble et al. (2003) found better correlation between PNC vs. CO, PM_{10} and nitrogen oxides at wind speeds below 3 m s^{-1} . These earlier findings together with those from this study suggest that correlation between pollutants seem to be affected by small-scale variations in pollutant dispersion induced by the wind regimes created within street canyons and therefore results found at one location cannot be generalised.

The high correlations between NO_2 and metals (Fe, Cu, As Cd, Pb) at St. Enoch Sq. and North Kensington, which are sites representative of population exposure, means that metals are a potential confounding factor for the health effects associated with NO_2 in time-series epidemiological studies based on data from a single exposure site. Further studies are required to investigate whether personal exposures of metals and NO_2 are also correlated.

The lack of correlation between PNC and metals at North Kensington is attributed to the fact that metal concentrations were possibly influenced by long range transport of particles, as opposed to traffic sources, as all metals except Cu correlated with PM_{10} , which at background sites is largely influenced by background air masses. This is confirmed by the low correlation between PM_{10} and NO_x ($r=0.38$). In contrast, at St. Enoch Sq. PM_{10} , was highly correlated to both NO_x and metals, suggesting a common traffic source.

6.3 Potential surrogates for PNC

The increments in daily NO_x and NO_2 concentrations between trafficked and background sites have shown to be a simple and novel surrogate for daily spatial variation of PNC. The increments in NO_x explained 78-79% of the variance in PNC at the paired sites in both Glasgow and London. Likewise, weekly NO_2 PDT values have been shown to be a good relative measure of weekly variability in PNC, explaining 70% of the variability of the latter.

The different relationship between PNC and nitrogen oxides is partly attributed to the fact that PNC do not depend only on traffic emissions but also on particle formation processes, as discussed above. Therefore, particles can still be formed in the atmosphere in the absence of traffic emissions leading to high background concentrations and the different relationships with nitrogen oxides, as observed in this study. This reduces the value of nitrogen oxides as a surrogate for PNC as a different relationship has to be derived for each location.

Benzene and 1,3-butadiene were also highly correlated with PNC and are therefore also potentially useful surrogates for PNC. Further examination of the relationships between the increments in benzene and 1,3-butadiene vs. increments in PNC are required. In addition, benzene and 1,3-butadiene can be measured by passive samplers, which therefore suggests a relatively non-technical and inexpensive surrogate for determining long time-averaged spatial patterns of PNC.

The particle size fraction dominating the PNC metric contributes comparatively little to the particle mass measured by PM_{10} . Therefore unless PM_{10} at a site is overwhelmingly dominated by the ultrafine fraction, correlation between PNC and PM_{10} is not expected to be high, and this is what is observed in this study. The low correlation between PNC and PM_{10} at Marylebone Rd. has been noted previously and is attributed to the large contribution of resuspended particles at this site (Charron and Harrison, 2003). The use of PM_{10} as a surrogate for PNC is therefore not appropriate.

The PM_{10} and $PM_{2.5}$ optical absorbance is a measure of black carbon (BC) (Cyrus et al. 2003), and has been found to correlate strongly with PNC in some previous studies (Zhu et al. 2002, Ketzel et al. 2003, Noble et al. 2003). In this study, PM optical absorbance was strongly correlated with PNC at Marylebone Rd. ($R = 0.87$) but not at the other sites ($R < 0.5$). Marylebone Rd. has substantial heavy goods vehicle and bus traffic, considerably more even than Glasgow Hope St. The optical 'blackness' depends not only on the BC content but also on the size of the particles (decreasing for smaller sizes) and the degree of dilution of optically absorbing particles with non-absorbing/transparent particles: the absorption of BC increases considerably when it is included in a large transparent particle, due to the lower density (Horvath, 1986). Particle size distributions shift to larger sizes and become

compositionally mixed as they disperse (Ketzel et al., 2003; Gidhagen et al., 2005), changing their optical properties. Therefore, it is not surprising that at background sites the optical properties of the mix of particles collected over a day is not correlated with average PNC. The lack of correlation at Hope St., where BC concentrations may be expected to be high remains unclear.

6.4 Limitations

This study clearly had some major limitations. The relationships are derived from a relatively small data set and may not represent the annual trend. Only one pair of sites was studied in each city. Measurements at paired sites in a given urban area over longer periods are needed to confirm the relationships found in this study. However, the different slopes and intercepts between NO_x , NO_2 and PNC suggest that for each location a different relationship has to be derived. Despite this limitation NO_x and NO_2 are a simple and non expensive method to represent PNC.

7 Conclusions

One of the mayor challenges in epidemiological studies of air pollution and health is the lack of data on spatial variability on PNC. This study has shown the potential for using NO_2 and NO_x increments between background and street sites to represent daily variations in PNC and water-soluble metals: increments in NO_x explained 78-79% of the variance in PNC at the paired sites in both Glasgow and London. The increments in NO_x also explained in 70 % of the spatial variation in Cu and Ni at Glasgow but not at London. Weekly NO_2 measurements derived from passive diffusion tubes were also shown to correlate well with increments in weekly PNC

The corollary of the high correlation between NO_2 and PNC and water-soluble metals at the background sites is that the latter pollutants may act as cofounders for health effects attributed to NO_2 data from such sites. Concentrations of CO cannot be used as a surrogate for PNC.

Further validation on the geographical extension of these relationships is necessary. This could be achieved for example by examining the spatio-temporal relationships of aggregated

data on the increments of a reference site and different additional sites (with different influence of traffic sources). A high correlation would mean that air pollution maps derived from NO₂ could be used to represent spatial and temporal variations in PNC and water-soluble metals. There is a need of regulating PNC. The relationships derived from these validated studies could be use to estimate the PNC which correspond to the EU limit value for NO₂ as a guide to establish a threshold value for PNC.

Acknowledgements

ASJ acknowledges PhD scholarship funding from the University of Strathclyde. Analysis of metals was funded by the National Environmental Research Council (NERC) Inductively Coupled Plasma Spectrometry Facility in Kingston University (Grant no. OSS/315/0506) and analysis of PAHs by the NERC Life Sciences Mass Spectrometry Facility in Bristol (Grant no. LSMSBRIS013 10/06). The authors also acknowledge Glasgow City Council for allowing siting of the monitoring equipment in the air quality monitoring stations, Defra for providing the filters collected at the London sites, and Gerry Carr, Gavin Gibson and Francis McGilligan for technical assistance during the project.

References

- Aalto, P., Hameri, K., Paatero, P., Kulmala, M., Bellander, T., Berglind, N., Bouso, L., Castano-Vinyals, G., Sunyer, J., Cattani, G., Marconi, A., Cyrys, J., Von Klot, S., Peters, A., Zetzsche, K., Lanki, T., Pekkanen, J., Nyberg, F., Sjoval, B. and Forastiere, F., 2005. Aerosol particle number concentration measurements in five European cities using TSI-3022 condensation particle counter over a three-year period during health effects pollution on susceptible subpopulations. *Journal of the Air & Waste Management Association* 55, 1064-1076.
- Adamson I.Y.R., Prieditis H., Hedgecock C. and Vincent R., 2000. Zinc is the toxic factor in the lung response to an atmospheric particulate sample. *Toxicology and Applied Pharmacology* 166, 111-119
- Ayers, G. P., 2001. Comment on regression analysis of air quality data. *Atmospheric Environment* 35, 2423-2425.
- Nicklas, B., Bellander, T., Forastiere, F., von Klot, S., Aalto, P., Elosua, R., Kulmala, M., Lanki, T., Löwel, H., Hannelored; P., Peters, A., Picciotto, S., Veikkoi; S., Salomaa, V., Stafoggia, M., Sunyer, J., Nyberg, F., 2009. Ambient air pollution and daily mortality among survivors of myocardial infarction. *Epidemiology* 20, 110-118
- Brown, R. J. C., 2008. The use and abuse of limits of detection in environmental analytical chemistry. *The Scientific World Journal* 8, 796-801.
- Burnett, R. T., Brook, J., Dann, T., Delocla, C., Philips, O., Cakmak, S., Vincent, R., Goldberg, M.S, and Krewski, D., 2000. Association between particulate- and gas-phase components of urban air pollution and daily mortality in eight Canadian cities. *Inhalation Toxicology* 12, 15-39.
- Charron, A. and Harrison, R. M., 2003. Primary particle formation from vehicle emissions during exhaust dilution in the roadside atmosphere. *Atmospheric Environment* 37, 4109-4119.

- Costa, D. L. and Dreher, K.L., 1997. Bioavailable transition metals in particulate matter mediate cardiopulmonary injury in healthy and compromised animal models. *Environmental Health Perspectives* 105, 1053-1060.
- Cyrys, J., Heinrich, J., Hoek, G., Meliefste, K., Lewne, M., Gehring, U., Bellander, T., Fischer, P., Van Vliet, P., Brauer, M., Wichmann, H. E. and Brunekreef, B., 2003. Comparison between different traffic-related particle indicators: Elemental carbon EC, PM_{2.5} mass, and absorbance. *Journal of Exposure Science and Environmental Epidemiology* 13, 134-143.
- Donaldson, K., Tran, L., Jimenes, L.A., Duffin, R., Newby, E.E., Mills, N., 2005. Combustion-derived nanoparticles: a review of their toxicology following inhalation exposure. *Particle & Fibre Toxicology* 2, 10, doi:10.1186/1743-8977-2-10.
- Gidhagen, L., Johansson, C., Langner, J., Foltescu, V. L., 2005. Urban scale modelling of particle number concentration in Stockholm. *Atmospheric Environment* 39, 1711-1725.
- Götschi, T., Hazenkamp-von Arx, M.E., Heinrich, J., Bono, R., Burney, P., Forsberg, B., Jarvis, D., Maldonado, J., Norbäck, D., Stern, W.B., Sunyer, J., Torén, D., Verlato, G., Villani, S. and Künzli, N., 2005. Elemental composition and reflectance of ambient fine particles at 21 European cities. *Atmospheric Environment* 39, 5947-5958.
- Harrison R.M., Tilling R., Romero Callen M.S., Harrad S. and Jarvis K. 2003. A study of trace metals and PAHs in the roadside environment. *Atmospheric Environment* 37, 2391-2402.
- Harrison, R. M. and Jones, A. M. 2005.. Multisite study of particle number concentrations in urbanair. *Environmental Science and Technology*. 39, 6063-6070.
- Heal, M. R., O'Donoghue, M. A. and Cape, J. N., 1999. Overestimation of urban nitrogen dioxide by passive diffusion tubes: a comparative exposure and model study. *Atmospheric Environment* 33, 513-524.

- Heal, M. R., Hibbs, L. R., Agius, R. M. and Beverland, I. J., 2005. Total and water-soluble trace metal content of urban background PM₁₀, PM_{2.5} and Black Smoke in Edinburgh, UK, *Atmospheric Environment* 39, 1417-1430.
- Heal, M. R., Elton, R.A., Hibbs L. R., Agius, R. M. and Beverland, I. J., 2009. A time-series study of the health effects of water-soluble and total-extractable metal content of airborne particulate matter. *Occupational and Environmental Medicine* 66, 636-638.
- Hoek, G., Brunekreef, B., Goldbohm, S., Fisher, P., van den Brandt, A., 2002. Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *The Lancet* 360, 1203-1209.
- Hoek, G., Boogaard, H., Knol, A., de Hartog, J., Slottje, P., Ayres, J. G., Borm, P., Brunekreef, B., Donaldson, K., Forastiere, F., Holgate, S., Kreyling, W. G., Nemery, B., Pekkanen, J., Stone, V., Wichmann W.G., van der Sluijs, J., 2010. Concentration response functions for ultrafine particles and all-cause mortality and hospital admissions: results of a European expert panel elicitation. *Environmental Science & Technology* 44, 476–482
- Horvath, H., Jager, J., Kreiner, I., Norek, C., 1986. Determination of the size-dependent light absorption coefficient of aerosols. *Journal of Aerosol Science* 17, 258-260.
- Jiménez, L.A., Thompson, J., Brown, D.A., Rahman, I., Antonicelli, F., Duffin, R., Drost, E. M., Hay, R. T., Donaldson, K. & Macnee, W., 2000. Activation of NF-κB by PM₁₀ occurs via an iron-mediated mechanism in the absence of IκB degradation. *Toxicology and Applied Pharmacology*, 166, 101-110.
- Ketzel, M., Wåhlin, P., Berkowicz, R. and Palmgren, F., 2003. Particle and trace gas emission factors under urban driving conditions in Copenhagen based on street and roof-level observations. *Atmospheric Environment* 37, 2735-2749.

- Kulmala, M., Vehkamäki, H., Petaja, T., Dal Maso, M., Lauri, A., Kerminen, V. M., Birmili, W. & McMurry, P. H. (2004) Formation and growth rates of ultrafine atmospheric particles, a review of observations. *Journal of Aerosol Science* 35, 143-176
- Lanki, T., Pekkanen, J., Aalto, P., Elosua, R., Berglind, N., D'ippoliti, D., Kulmala, M., Nyberg, F., Peters, A., Picciotto, S., Salomaa, V., Sunyer, J., Tiittanen, P., Von Klot, S., Forastiere, F. & For the, H. S. G., 2006. Associations of traffic related air pollutants with hospitalisation for first acute myocardial infarction: the HEAPSS study *Occupational and Environmental Medicine* 63, 844-851.
- Monteiller, C., Tran, L., Macnee, W., Faux, S., Jones, A., Miller, B. & Donaldson, K. (2007) The pro-inflammatory effects of low-toxicity low-solubility particles, nanoparticles and fine particles, on epithelial cells in vitro: the role of surface area.
- Noble, C., Mukerjee, S., Gonzales, M., Rodes, C.E., Lawless, P.A., Natarajan, S., Myers, E.A., Norris, G.A., Smith, L., Ā-Zkaynak, H. and Neas, L.M., 2003. Continuous measurement of fine and ultrafine particulate matter, criteria pollutants and meteorological conditions in urban El Paso, Texas. *Atmospheric Environment* 37, 827-840.
- Piñeiro-Iglesias, M., Lopex-Mahia, P., Muniategui-Lorenzo, S., Prada-Rodriguez, D., Querol, X. and Alastuey, A., 2003. A new method for the simultaneous determination of PAH and metals in samples of atmospheric particulate matter. *Atmospheric Environment* 37, 4171-7175.
- Pope, C. A., Burnett, R. T., Thurston, G. D., Thun, M. J., Calle, E. E., Krewski, D. and Godleski, J. J., 2004. Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease. *Circulation* 109, 71-77.
- Puustinen, A., Hämeri, K., Pekkanen, J., Kulmala, M., De Hartog, J., Meliefste, K., Ten Brink, H., Kos, G., Katsouyanni, K., Karakatsani, A., Kotronarou, A., Kavouras, I., Meddings, C., Thomas, S., Harrison, R., Ayres, J. G., Van Der Zee, S., Hoek, G., 2007. Spatial variation of particle number and mass over four European cities. *Atmospheric Environment* 41, 6622-6636.

Sánchez Jiménez, A., 2010. Air pollution and health: identification of chemical and physical exposure metrics for epidemiological research. PhD Thesis, University of Strathclyde, Glasgow, UK.

USEPA (1998) Quality Assurance Guidance Document 2.12 Monitoring PM_{2.5} in Ambient Air Using Designated Reference or Class I Equivalent Methods.

Vaughan, N. P., Milligan, B. D. and Ogden, T. L., 1989. Filter weighing reproducibility and the gravimetric detection limit. *Annals of Occupational Hygiene* 33, 331-337.

von Klot, S., Peters, A., Aalto, P., Bellander, T., Berglind, N., D'Ippoliti, D., Elosua, R., Allmut Hörmann, M. D., Kulmala, M., Lanki, T., Löwel, H., Pekkanen, J., Picciotto, S., Sunyer, J., Forastiere, F., 2005. Ambient air pollution is associated with increased risk of hospital cardiac readmissions of myocardial infarction survivors in five European cities. *Circulation* 112, 3073-3079.

Wilson M. R., Lightbody, J. H., Donaldson, D., Sales, J., Stone, V., 2002. Interactions between ultrafine particles and transition metals in vivo and in vitro. *Toxicology and Applied Pharmacology* 184, 172–179.

Zhu, Y., Hinks, W. C., Kim, S., Shen, S., Sioutas, C., 2002. Study of ultrafine particles near a major highway with heavy-duty diesel traffic. *Atmospheric Environment* 36, 4323-4335.

Table captions

Table 1: Sampling equipment used at each monitoring site and data source.

Table 2: Limits of detection (ng m^{-3}) for metal concentrations derived from 24 h Partisol samples of PM_{10} on PTFE filters and of $\text{PM}_{2.5}$ on quartz filters.

Table 3: Descriptive statistics of traffic-related air pollutants at Glasgow (Hope St: Kerbside/street canyon, St:Enoch Sq., urban centre and Montrose St: background/street canyon) and London (Marylebone Rd: street canyon, North Kensington: urban background).

Table 4: Spearman correlation coefficients (p-values and sample sizes) between concentrations of the same pollutant measured at paired sites.

Table 4: Spearman correlation coefficients (*p*-values and sample sizes) of water-soluble metal concentrations in PM_{10} between Hope St. and St. Enoch Sq. in Glasgow and in $\text{PM}_{2.5}$ between Marylebone Rd. and North Kensington in London.

Table 6: Spearman correlation coefficients (and sample sizes) between daily-averaged air pollutant measurements at Hope St. (upper table) and St. Enoch Sq. (lower table).

Table 7: Spearman correlation coefficients (and sample sizes) between daily-averaged air pollutant measurements at Montrose St. (urban background).

Table 8: Spearman correlation coefficients (and sample sizes) between daily-averaged traffic-related air pollutants at Marylebone Rd. (upper table) and North Kensington (lower table).

Table 9: Slope, intercept and R^2 values for Reduced Major Axis regressions of the increments in particle number concentrations (PNC) vs. increments in other pollutant measures for the paired sites Hope St vs. St. Enoch Sq. in Glasgow and Marylebone Rd vs. North Kensington in London. Particle optical absorbance (absPM) is for PM₁₀ in Glasgow and PM_{2.5} in London.

Table 10: Slope, intercept and R^2 values for Reduced Major Axis regressions of the increments in Cu and Ni vs. the increments in particle number concentrations (PNC) and nitrogen oxides at Hope St. vs. St. Enoch Sq. in Glasgow.

Table 1 Sampling equipment used at each monitoring site and data source.

Pollutant	Equipment	Monitoring sites				
		Glasgow		London		
		Hope St.	St. Enoch Sq.	Montrose St.	Marylebone Rd.	North Kensington
PM ₁₀	PARTISOL	√	√		NAQA	NAQA
PM _{2.5}	PARTISOL				NAQA	NAQA
BS	BS monitor				NAQA	
PNC	water based-CPC	√	√	√		
	butanol based-CPC		NAQA		NAQA	NAQA
Absorbance	measured on PM ₁₀	√	√		√	√
PM	PARTISOL filters					
Metals	extracted from PM ₁₀	√	√		√	√
	PARTISOL filters					
NO ₂	PDTs	√	√			
	Chemilumiscence	NAQA	NAQA	NAQA	NAQA	NAQA
	Analyser					
CO	Infra Red Analyser	NAQA	NAQA	NAQA	NAQA	NAQA
Benzene & 1,3-butadiene	Ultra Violet Analyser	NAQA			NAQA	

√: measured as part of this project; NAQA: provided by the National Air Quality Archive ;BS:Black Smoke; PNC: particle number concentration;CPC: condensation particle counter; PDTs: passive diffusion tubes.

Table 2 Limits of detection (ng m^{-3}) for metal concentrations derived from 24 h Partisol samples of PM_{10} on PTFE filters and of $\text{PM}_{2.5}$ on quartz filters.

Metal	V	Cr	Mn	Fe	Ni	Cu	As	Cd	Pb
Quartz filters									
(London sites) n=6	0.017	1.52	0.106	3.02	0.123	0.630	0.006	0.005	0.053
PTFE filters									
(Glasgow sites) n=2	0.013	0.113	0.128	2.67	0.325	0.538	0.004	0.013	0.232

Table 3 Descriptive statistics of traffic-related air pollutants at Glasgow (Hope St: kerbside/street canyon, St:Enoch Sq., urban centre and Montrose St: background/street canyon) and London (Marylebone Rd: street canyon, North Kensington: urban background).

	Hope St.			St. Enoch Sq.			Montrose St.			Marylebone Rd.			North Kensington		
	n	Median	IQR	n	Median	IQR	n	Median	IQR	n	Median	IQR	n	Median	IQR
PNC (particles cm ⁻³)	35	23,564	10,475	79	12,851	3,128	63	11,095	2,940	26	109,953	35,393	26	23,407	9,784
PM ₁₀ PARTISOL (µg m ⁻³)	31	32.9	12.0	24	21.6	12.5	-	-	-	26	45.0	9.00	26	25.5	10.8
PM _{2.5} PARTISOL (µg m ⁻³)	-	-	-	-	-	-	-	-	-	26	30.0	7.00	26	15.5	7.00
BS (µg m ⁻³)	-	-	-	-	-	-	-	-	-	26	57.0	28.2	-	-	-
AbsPM (10 ⁻⁵ m ⁻¹)	31	2.65	2.39	24	1.47	1.05	-	-	-	26	12.0	2.00	26	6.96	0.61
NO _x (µg m ⁻³)	63	157	132	81	35.7	15.2	83	66.0	27.4	26	389	157	26	34.3	15.2
NO ₂ (µg m ⁻³)	63	50.1	31.9	81	22.4	9.65	83	39.2	13.4	26	133	37.4	23	28.8	13.7
NO (µg m ⁻³)	63	71.0	60.7	81	8.04	4.91	83	17.3	9.72	26	167	79.8	26	3.51	3.11
CO (mg m ⁻³)	75	0.296	0.133	54	0.191	0.061	83	0.248	0.124	26	0.974	0.34	26	0.240	0.083
1,3- Butadiene (µg m ⁻³)	31	0.077	0.082	-	-	-	-	-	-	19	0.463	0.157	-	-	-
Benzene (µg m ⁻³)	33	0.563	0.313	-	-	-	-	-	-	21	2.35	1.32	-	-	-
V (ng m ⁻³)	17	0.202	0.354	17	0.183	0.565	-	-	-	26	1.20	2.00	26	1.49	2.06
Cr (ng m ⁻³)	17	0.190	0.187	17	0.141	0.328	-	-	-	-	-	-	-	-	-
Mn (ng m ⁻³)	17	0.428	1.23	17	1.57	1.46	-	-	-	26	0.723	0.802	26	0.352	0.488
Fe (ng m ⁻³)	17	24.4	15.9	17	26.0	16.5	-	-	-	26	6.91	11.1	26	5.04	5.91
Ni (ng m ⁻³)	17	0.392	0.675	17	0.473	0.593	-	-	-	26	0.429	0.473	26	0.803	0.97
Cu (ng m ⁻³)	17	9.54	4.85	17	3.06	3.401	-	-	-	26	2.60	1.81	26	1.56	1.32
Zn (ng m ⁻³)	-	-	-	-	-	8.468	-	-	-	26	11.2	7.66	26	7.73	11.3
As (ng m ⁻³)	17	0.070	0.094	17	0.040	0.135	-	-	-	26	0.440	0.300	26	0.524	0.519
Cd (ng m ⁻³)	17	0.057	0.033	17	0.054	0.036	-	-	-	26	0.058	0.042	26	0.061	0.078
Pb (ng m ⁻³)	17	1.32	0.894	17	0.714	1.51	-	-	-	26	1.96	2.22	26	2.44	3.45
Wind speed	84	5.54	3.43	84	11.7	4.69	84	5.54	3.43	25	11.7	3.25	25	11.7	4.69

PNC: particle number concentration; BS: black smoke; Wind speed data in Glasgow sites was measured at Bishopton (circa 18Km from the sites) and in London at Heathrow Airport (circa 16 km from North Kensington and 20 Km from Marylebone Rd).

Table 4 Spearman correlation coefficients (*p*-values and sample sizes) between concentrations of the same pollutant measured at paired sites.

		Hope St. vs St. Enoch Sq.	Marylebone Rd. vs. North Kensington	Hope St. vs Montrose St.	Montrose St. vs St. Enoch Sq.
PNC	Hourly	0.47** 0.000 717	0.58** 0.000 609	0.56** 0.000 828	0.57** 0.000 1349
	Daily	0.43* 0.017 30	0.46* 0.018 26	0.55** 0.002 35	0.41** 0.002 57
PM ₁₀ TEOM	Hourly	0.67** 0.000 1386	0.50** 0.000 608	NA	NA
	Daily	0.73** 0.000 56	0.23 0.267 25	NA	NA
PM ₁₀ PARTISOL	Daily	0.34 0.158 17	0.34 0.094 25	NA	NA
PM _{2.5} PARTISOL	Daily	NA	0.35 0.080 26	NA	NA
AbsPM*	Daily	0.43 0.082 17	-0.03 0.900 26	NA	NA
NO _x	Hourly	0.52** 0.000 1552	0.43** 0.000 611	0.63** 0.000 1551	0.78** 0.000 1967
	Daily	0.33** 0.000 62	0.06 0.766 26	0.55** 0.001 63	0.72** 0.000 81
NO ₂	Hourly	0.50** 0.000 1552	0.37** 0.000 611	0.62** 0.000 1551	0.80** 0.000 1967
	Daily	0.49** 0.000 62	0.05 0.819 26	0.61** 0.000 63	0.80** 0.000 81
NO	Hourly	0.47** 0.000 1552	0.45** 0.000 611	0.61** 0.000 1551	0.74** 0.000 1967
	Daily	0.33** 0.009 62	0.036 0.860 26	0.47** 0.000 63	0.61** 0.000 81
CO	Hourly	0.26** 0.000 1168	0.33** 0.000 605	0.58** 0.000 1817	0.27** 0.000 1326
	Daily	0.17 0.257 47	0.06 0.766 26	0.64** 0.000 75	0.17 0.227 54

PNC: particle number concentration; * AbsPM₁₀ at Glasgow sites and AbsPM_{2.5} at London sites; NA: not available;

**Correlation is significant at the 0.01 level.

Table 5 Spearman correlation coefficients (*p*-values and sample sizes) of water-soluble metal concentrations in PM₁₀ between Hope St. and St. Enoch Sq. in Glasgow and in PM_{2.5} between Marylebone Rd. and North Kensington in London.

		Hope St. vs St. Enoch Sq.	Marylebone Rd. vs. North Kensington
V	Correlation Coefficient Sig. (2-tailed) N	0.89** 0.000 17	0.94** 0.000 26
Cr	Correlation Coefficient Sig. (2-tailed) N	0.126 0.629 17	NA
Mn	Correlation Coefficient Sig. (2-tailed) N	0.53* 0.035 17	0.79** 0.000 26
Fe	Correlation Coefficient Sig. (2-tailed) N	0.25 0.350 17	0.49* 0.010 26
Ni	Correlation Coefficient Sig. (2-tailed) N	-0.01 0.955 17	0.77** 0.000 26
Zn	Correlation Coefficient Sig. (2-tailed) N	NA	0.67** 0.000 26
Cu	Correlation Coefficient Sig. (2-tailed) N	0.42 0.092 17	0.45** 0.010 26
As	Correlation Coefficient Sig. (2-tailed) N	0.85** 0.000 17	0.85** 0.000 26
Cd	Correlation Coefficient Sig. (2-tailed) N	0.17 0.552 17	0.57** 0.002 26
Pb	Correlation Coefficient Sig. (2-tailed) N	0.36 0.172 17	0.63** 0.001 26

** Correlation is significant at 0.01 level (2 tailed);* Correlation is significant at 0.05

Table 6 Spearman correlation coefficients (and sample sizes) between daily-averaged air pollutant measurements at Hope St. (upper table) and St. Enoch Sq. (lower table).

Hope St. (Kerbside)	V	Cr	Mn	Fe	Ni	Cu	As	Cd	Pb	PNC	PM ₁₀ P	Abs PM ₁₀	NO _x	NO ₂	CO
PNC	0.76**	0.71**	0.93**	0.661	0.92**	0.94**	0.54	0.22	0.35	1					
	11	11	10	10	11	11	10	10	11	35					
PM ₁₀ P	0.29	0.21	0.46	0.07	0.58*	0.70**	0.33	0.08	0.02	0.74**	1				
	17	17	16	16	17	17	16	16	17	25	25				
Abs PM ₁₀	0.06	0.13	0.23	-0.09	0.20	0.29	0.47	0.50*	0.13	0.35	0.28	1			
	17	17	16	16	17	17	16	16	17	25	25	25			
NO _x	-0.17	0.05	-0.27	-0.70*	0.32	0.55	-0.32	-0.29	-0.57	0.87**	0.85**	0.53*	1		
	9	9	9	9	9	9	9	8	9	19	17	17	63		
NO ₂	0.12	0.29	-0.23	-0.59*	0.75*	0.48	-0.09	0.07	-0.29	0.85**	0.83**	0.45*	0.90**	1	
	9	9	9	9	9	9	9	8	9	19	17	17	63	63	
CO	0.20	0.22	0.25	-0.26	0.55*	0.66*	0.20	0.22	0.04	0.64**	0.76**	0.43*	0.78**	0.60**	1
	14	14	13	13	14	14	14	13	14	25	22	22	63	63	75
1,3-Butadiene	0.20	0.00	0.29	-0.39	0.41	0.58*	0.31	0.38	-0.09	0.71**	0.71**	0.54**	0.95**	0.91**	0.89**
	15	15	14	14	15	15	15	14	15	33	23	23	19	19	25
Benzene	0.15	0.05	0.25	-0.23	0.41	0.52*	0.06	0.02	-0.04	0.55**	0.76**	0.32	0.93**	0.88**	0.93**
	15	15	14	14	15	15	15	14	15	33	23	23	19	19	25
Wind speed	-0.55*	-0.45	-0.81**	-0.49	-0.48	-0.84**	-0.68**	0.09	-0.28	-0.52**	-0.40	-0.62**	-0.47*	-0.51*	-0.51*
	16	16	16	16	15	16	15	14	16	34	24	24	19	19	24
St. Enoch Sq. (Urban background)	V	Cr	Mn	Fe	Ni	Cu	As	Cd	Pb	PNC	PM ₁₀ P	Abs PM ₁₀	NO _x	NO ₂	CO
PNC	0.52	0.62*	0.81**	0.73**	0.57*	0.83**	0.81**	0.70**	0.72**	1					
	17	17	17	17	17	17	17	17	17	17					
PM ₁₀ P	0.13	0.44	0.77**	0.56*	0.55*	0.60*	0.52*	0.53*	0.53*	0.29	1				
	17	17	17	17	17	17	17	17	17	23	24				
AbsPM ₁₀	0.69**	0.41	0.49*	0.61**	0.23	0.69**	0.76**	0.55*	0.52*	0.47*	0.23	1			
	17	17	17	17	17	17	17	17	17	23	24				
NO _x	0.52	0.71**	0.66*	0.75**	0.42	0.74**	0.76**	0.59*	0.72**	0.65**	0.64**	0.415	1		
	15	15	15	15	15	15	15	15	15	75	20	20			
NO ₂	0.51	0.67*	0.62*	0.71**	0.39	0.74**	0.76**	0.53	0.65*	0.52**	0.69**	0.49*	0.91**	1	
	15	15	15	15	15	15	15	15	15	75	20	20	81		
CO	0.31	0.12	0.19	0.00	-0.10	0.05	0.12	-0.31	-0.24	0.34*	0.26	-0.28	0.49**	0.40**	1
	10	10	10	10	10	10	10	10	10	49	11	11	54	54	54
Wind	-0.57*	-0.63**	-0.49	-0.51*	-0.38	-0.61*	-0.64*	-0.43	-0.59*	-0.30**	-0.51*	-0.35	-0.47**	-0.57**	-0.38**
	17	17	17	17	17	17	17	17	17	78	23	23	80	80	53

Significant correlations are in bold; **Correlation is significant at the 0.01 level (2-tailed), *Correlation is significant at the 0.05 level (2-tailed).

PNC: particle number concentration; PM10 P: PM₁₀ concentrations measured with a Partisol sampler

Table 7 Spearman correlation coefficients (and sample sizes) between daily-averaged air pollutant measurements at Montrose St. (urban background).

	PNC	NO _x	NO ₂	CO
NO _x	0.62** 61	1 83		
NO ₂	0.66** 61	0.93** 83	1 83	
CO	0.67** 61	0.65** 83	0.74** 83	1 83
Wind speed	-0.45** 62	-0.30** 82	-0.42** 82	-0.44** 82

**Correlation is significant at the 0.01 level (2-tailed);

PNC: particle number concentration.

Table 8 Spearman correlation coefficients (and sample sizes) between daily-averaged traffic-related air pollutants at Marylebone Rd. (upper table) and North Kensington (lower table).

Marylebone Rd. (Kerbside)	V	Mn	Fe	Ni	Cu	Zn	As	Cd	Pb	PNC	PM ₁₀ P	PM _{2.5} P	BS	Abs PM _{2.5}	NO _x	NO ₂	CO	1,3- butadiene	Benzene	Wind speed
PNC	-0.12	-0.24	-0.21	-0.24	-0.60**	-0.26	-0.26	-0.36	-0.08	1.00										
	26	26	26	26	26	26	26	26	26	26										
PM ₁₀ P	0.48*	0.31	0.12	0.44*	-0.18	0.25	0.02	-0.03	0.10	0.40*	1.00									
	25	25	25	25	25	25	25	25	25	25	25									
PM _{2.5} P	0.38	0.54**	0.48*	0.34	0.03	0.36	0.24	0.19	0.19	0.47*	0.71**	1.00								
	26	26	26	26	26	26	26	26	26	26	25	26								
BS	-0.03	-0.26	-0.28	-0.19	-0.49*	-0.25	-0.14	-0.33	-0.14	0.71**	0.35	0.28	1.00							
	26	26	26	26	26	26	26	26	26	26	25	26	26							
Abs PM _{2.5}	0.07	-0.13	-0.23	-0.06	-0.54**	-0.14	-0.27	-0.39*	-0.07	0.87**	0.60**	0.47*	0.75**	1.00						
	26	26	26	26	26	26	26	26	26	26	25	26	26	26						
NO _x	-0.08	-0.28	-0.35	-0.21	-0.67**	-0.20	-0.27	-0.39*	-0.03	0.90**	0.57**	0.40*	0.73**	0.93**	1.00					
	26	26	26	26	26	26	26	26	26	26	25	26	26	26	26					
NO ₂	0.03	-0.22	-0.29	-0.10	-0.63**	-0.17	-0.29	-0.34	-0.07	0.87**	0.67**	0.49*	0.68**	0.92**	0.97**	1.00				
	26	26	26	26	26	26	26	26	26	26	25	26	26	26	26	26				
CO	0.44*	0.20	0.14	0.39*	-0.18	0.11	0.29	0.25	-0.07	0.27	0.49*	0.56**	0.24	0.23	0.26	0.30	1.00			
	26	26	26	26	26	26	26	26	26	26	25	26	26	26	26	26	26			
1,3-butadiene	-0.02	-0.27	-0.27	-0.17	-0.61	-0.21	-0.24	-0.29	0.05	0.81**	0.67**	0.45**	0.69**	0.74**	0.87**	0.83**	0.50**	1.00		
	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19		
Benzene	-0.04	-0.19	-0.13	-0.19	-0.55	-0.18	-0.25	-0.31	0.16	0.78**	0.62**	0.47**	0.74**	0.72**	0.83**	0.77**	0.45	0.98**	1.00	
	21	21	21	21	21	21	21	21	21	21	20	21	21	21	21	21	21	19	21	
Wind speed	-0.26	-0.53**	-0.46*	-0.27	-0.57**	-0.52**	-0.86**	-0.48*	-0.40*	0.40*	-0.02	-0.20	0.12	0.32	0.36	0.41*	-0.26	0.18	0.17	1.00
	25	25	25	25	25	25	25	25	25	25	24	25	25	25	25	25	25	18	20	25

North Kensington (Urban background)	V	Mn	Fe	Ni	Cu	Zn	As	Cd	Pb	PNC	PM ₁₀ P	PM _{2.5} P	Abs PM _{2.5}	NO _x	NO ₂	NO	CO	Wind speed		
PNC	-0.32	-0.25	-0.12	-0.53**	0.24	0.19	0.25	0.20	0.32	1.00										
	26	26	26	26	26	26	26	26	26	26										
PM ₁₀ P	0.44*	0.58**	0.59**	0.54**	0.21	0.47*	0.55**	0.63**	0.49*	0.03	1.00									
	26	26	26	26	26	26	26	26	26	26	26									
PM _{2.5} P	0.48*	0.72**	0.77**	0.63**	0.34	0.64**	0.69**	0.81**	0.64**	-0.12	0.82**	1.00								
	26	26	26	26	26	26	26	26	26	26	26	26								
Abs PM _{2.5}	0.31	0.60**	0.74**	0.25	0.58**	0.71**	0.64**	0.64**	0.74**	0.34	0.55**	0.65**	1.00							
	26	26	26	26	26	26	26	26	26	26	26	26	26							
NO _x	0.04	0.35	0.56**	-0.04	0.64**	0.68**	0.61**	0.57**	0.71**	0.60**	0.38	0.44*	0.89**	1.00						
	26	26	26	26	26	26	26	26	26	26	26	26	26	26						
NO ₂	0.06	0.34	0.54**	-0.04	0.62**	0.69**	0.63**	0.56**	0.74**	0.59**	0.37	0.42*	0.89**	0.98**	1.00					
	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26					
CO	-0.20	0.08	0.08	-0.36	0.46*	0.34	0.45*	0.47*	0.54**	0.70**	0.09	0.16	0.48*	0.59**	0.61**	0.54**	1.00			
	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26	26			
Wind	-0.17	-0.64**	-0.75**	-0.35	-0.63**	-0.75**	-0.66**	-0.69**	-0.64**	0.06	-0.52**	-0.69**	-0.70**	-0.62**	-0.65**	-0.55**	-0.23	1.00		
	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	25	

Significant correlations are in bold; * Correlation is significant at the 0.05 level (2-tailed); ** Correlation is significant at the 0.01 level (2-tailed). Wind data was measured at Heathrow Airport (circa 20 Km from the monitoring site). Data were provided by the British Atmospheric Data Centre (BADC).

Table 9 Slope, intercept and R^2 values for Reduced Major Axis regressions of the increments in particle number concentrations (PNC) vs. increments in other pollutant measures for the paired sites Hope St vs. St. Enoch Sq. in Glasgow and Marylebone Rd vs. North Kensington in London. Particle optical absorbance (absPM) is for PM₁₀ in Glasgow and PM_{2.5} in London.

Relationship	Hope St. - St. Enoch Sq.					Marylebone Rd. - North Kensington				
	N	intercept	slope	R ²		N	intercept	slope	R ²	
$\Delta\text{PNC vs. } \Delta\text{PM}_{10}$	10	-15,408	1,273	0.70		25	26,073	3,078	0.55
$\Delta\text{PNC vs. } \Delta\text{PM}_{2.5}$	NA	NA	NA			26	37,570	3,520	0.43
$\Delta\text{PNC vs. } \Delta\text{AbsPM}^*$	10	-10,980	8,657	0.03		26	-22,769	21,533	0.71
$\Delta\text{PNC vs. } \Delta\text{NO}_x$	14	7,408	42.3	0.79		26	732	233	0.78
$\Delta\text{PNC vs. } \Delta\text{NO}_2$	14	6,880	155	0.59		26	-12,534	933	0.74
$\Delta\text{PNC vs. } \Delta\text{CO}$	19	1,881	75,842	0.30		26	652	119,440	0.08

Table 10 Slope, intercept and R^2 values for Reduced Major Axis regressions of the increments in Cu and Ni vs. the increments in particle number concentrations (PNC) and nitrogen oxides at Hope St. vs. St. Enoch Sq. in Glasgow.

Relationship	N	intercept	slope	R^2
ΔCu vs. ΔPNC	11	3.63	4.0E-04	0.66
ΔCu vs. ΔNO_x	8	0.499	0.037	0.65
ΔCu vs. ΔNO_2	8	0.739	0.151	0.67
ΔCu vs. ΔNO	8	0.696	0.072	0.59
ΔNi vs. ΔPNC	9	0.018	5.3E-05	0.76
ΔNi vs. ΔNO_x	8	-2.33	0.014	0.19
ΔNi vs. ΔNO_2	8	-2.24	0.055	0.47
$R^2 > 0.5$ are in bold				