



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Carbonate delta drift: a new sediment drift type

Citation for published version:

Lüdmann, T, Betzler, C, Eberli, GP, Reolid, J, Reijmer, JJG, Sloss, CR, Bialik, OM, Alvarez-zarikian, CA, Alonso-garcía, M, Blättler, CL, Guo, JA, Haffen, S, Horozal, S, Inoue, M, Jovane, L, Kroon, D, Lanci, L, Laya, JC, Mee, ALH, Nakakuni, M, Nath, BN, Niino, K, Petruny, LM, Pratiwi, SD, Slagle, AL, Su, X, Swart, PK, Wright, JD, Yao, Z & Young, JR 2018, 'Carbonate delta drift: a new sediment drift type', *Marine geology*. <https://doi.org/10.1016/j.margeo.2018.04.011>

Digital Object Identifier (DOI):

[10.1016/j.margeo.2018.04.011](https://doi.org/10.1016/j.margeo.2018.04.011)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Marine geology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 Carbonate delta drift: a new sediment drift type

2 Thomas Lüdmann¹, Christian Betzler¹, Gregor P. Eberli², Jesús Reolid¹, John J.G. Reijmer³, Craig
 3 R. Sloss⁴, Or M. Bialik⁵, Carlos A. Alvarez-Zarikian⁶, Montserrat Alonso-García^{7,8}, Clara L.
 4 Blättler⁹, Junhua Adam Guo¹⁰, Sébastien Haffen¹¹, Senay Horozal¹², Mayuri Inoue¹³, Luigi
 5 Jovane¹⁴, Dick Kroon¹⁵, Luca Lanci¹⁶, Juan Carlos Laya¹⁷, Anna Ling Hui Mee², Masatoshi
 6 Nakakuni¹⁸, B. Nagender Nath¹⁹, Kaoru Niino²⁰, Loren M. Petruny²¹, Santi D. Pratiwi²², Angela L.
 7 Slagle²³, Xiang Su²⁴, Peter K. Swart², James D. Wright²⁵, Zhengquan Yao^{26,27}, Jeremy R. Young²⁸

8 ¹*Institute of Geology, CEN, University of Hamburg, Bundesstrasse 55, Hamburg 20146, Germany.*

9 ²*Department of Marine Geosciences, Rosenstiel School of Marine & Atmospheric Science, University of Miami, Miami*
 10 *FL 33149, USA.*

11 ³*College of Petroleum Engineering and Geosciences, Fahd University of Petroleum and Minerals, Dhahran 31261,*
 12 *Saudi Arabia.*

13 ⁴*Earth and Environmental Sciences, University of Technology Queensland, R-Block 317, 2 George Street, Brisbane*
 14 *Queensland 4001, Australia.*

15 ⁵*Dr. Moses Strauss Department of Marine Geosciences, The Leon H. Charney School of Marine Sciences, University of*
 16 *Haifa, Carmel 31905, Israel.*

17 ⁶*International Ocean Discovery Program, Texas A&M University, Discovery Drive, College Station TX 77845, USA.*

18 ⁷*Instituto Portugues do Mar e da Atmosfera (IPMA), Divisão de Geologia e Georecursos Marinhos, Rua Alfredo*
 19 *Magalhães Ramalho, 6, 1495-006, Portugal.*

20 ⁸*Centro de Ciencias do Mar (CCMAR), Universidade do Algarve, Faro, Portugal.*

21 ⁹*Department of Geosciences, Princeton University, Guyot Hall, Princeton NJ 08544, USA.*

22 ¹⁰*Department of Geological Sciences, California State University Bakersfield, 9001 Stockdale Highway, Bakersfield,*
 23 *CA 93311, USA.*

24 ¹¹*Physical Properties Specialist, Ecole Nationale Supérieure de Géologie, Université de Lorraine, 2 rue du Doyen*
 25 *Marcel Roubault, Vandœuvre-les-Nancy 54501, France.*

26 ¹²*Petroleum and Marine Research Division, Korea Institute of Geoscience & Mineral Resources (KIGAM), Gwahang-*
 27 *no 124, Yuseong-gu, Daejeon 305-350, Korea.*

28 ¹³*Graduate School of Natural Science and Technology, Okayama University, 3-1-1 Tsushima-naka 700-8530, Japan.*

¹⁴Instituto Oceanográfico da Universidade de São Paulo, Praça do Oceanográfico, 191, São Paulo, SP 05508-120, Brazil.

¹⁵School of GeoSciences, University of Edinburgh, Grant Institute, The King's Buildings, James Hutton, Edinburgh EH9 3FE, United Kingdom.

¹⁶Istituto di Scienze della Terra, Università di Urbino, Via S. Chiara 27, Urbino 61029, Italy.

¹⁷Department of Geology and Geophysics, Texas A&M University, Mail Stop 3115, College Station TX 77843-3115, USA.

¹⁸Department of Environmental Engineering for Symbiosis, Soka University, 1-236 Tangi-cyo, Hachioji-shi Tokyo 192-0003, Japan.

¹⁹Geological Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula Goa 403004, India.

²⁰Graduate School of Science and Engineering, Yamagata University, 1-4-12 Kojirakawa-machi, Yamagata City 990-8560, Japan.

²¹Environmental Science and Policy Department, David King Hall Rm 3005, MSN 5F2, George Mason University, University Drive, Fairfax, VA 22030-4444, USA.

Universitas Padjadjaran, Geological Engineering, Jl. Raya Bandung, Sumedang KM 21, Jatinangor 45363, Indonesia.

²³Lamont-Doherty Earth Observatory, Columbia University, Borehole Bldg. 61 Route 9W, Palisades NY 10964, USA.

²⁴Key Laboratory of Marginal Sea Geology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, West Xingang Road, Guangzhou 510301, P.R. China.

²⁵Department of Geological Sciences, Rutgers, The State University of New Jersey, 610 Taylor Road, Piscataway NJ 08854-8066, USA.

²⁶Department of Marine Geology, First Institute of Oceanography (FIO) State Oceanic Administration (SOA), #6 Xian Xia Ling Road, Qingdao Shandong Province 266061, P.R. China.

²⁷Laboratory for Marine Geology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, P.R. China.

²⁸Department of Earth Sciences, University College London, Gower Street, London WC1E 6BT, United Kingdom.

ABSTRACT

Based on high-resolution reflection seismic and core data from IODP Expedition 359 we present a new channel-related drift type attached to a carbonate platform slope, which we termed delta drift. Like a river delta, it is comprised of several stacked lobes and connected to

a point source. The delta drifts were deposited at the exit of two gateways that connect the Inner Sea of the Maldives carbonate platform with the open ocean. The channels served as conduits focusing and accelerating the water flow; Entrained material was deposited at their mouth where the flows relaxed. The lobe-shaped calcareous sediment drifts must have formed under persistent water through flow. Sediment supply was relatively high and continuous, resulting in an average sedimentation rate of 17 cm ka^{-1} . The two delta drifts occupy 342 and 384 km^2 , respectively; with a depositional relief of approximately 500 m . They have a sigmoidal clinoform reflection pattern with a particular convex upward bending of the foresets. In the Maldives the drift onset marks the transition from a sea-level controlled to a progressively current dominated depositional regime. This major event occurred in the Serravallian about 13 Ma ago, leading to the partial drowning of the carbonate platform and the creation of shallow seaways. The initial bank-enclosed topography resembles an “empty bucket” geometry which is rapidly filled by the drift sediments that aggrade and prograde into the basin. Thereby the depositional environment of the delta drifts changes from deep water (>500) to shallow-water conditions at their topsets, indicated by the overall coarsening upward trend in grain size and the presence of shallow water large benthic foraminifers at their top.

Keywords: delta drift, carbonate platform, drift sedimentation, bottom current, clinoform, Maldives

INTRODUCTION

Current-controlled carbonate deposits have so far not been systematically investigated in contrast to their intensively studied siliciclastic counterparts (Faugères et al., 1999; Stow et al., 2002b; Viana and Rebesco, 2007; Rebesco and Camerlenghi, 2008; Rebesco et al., 2014). Several studies, however, document their importance, especially in tropical carbonate platforms (Anselmetti et al., 2000; Betzler et al., 2009, 2013, 2014; Isern et al., 2004; Eberli et al., 2010; Lüdmann et al., 2012, 2013). For the Maldives, Lüdmann et al. (2013) demonstrated

that since the Middle Miocene carbonate sedimentation in the Inner Sea was dominated by ocean currents entering the archipelago interior via gateways between the atolls. This situation resulted in the deposition of 10 mega-drift sequences. This is in contrast to the standard sequence stratigraphic model that describes carbonate platform geometry and depositional setting as a response to relative sea-level changes (Schlager, 2005 and references therein). Recent studies show that sea-level-controlled highstand shedding plays an essential role in sediment supply; However, currents could be the main agent transporting carbonate debris from the platform top and distributing it to the surrounding margins (Betzler et al., 2013, 2014, 2015; Lüdmann et al., 2013).

The Maldives, a large N-S elongated isolated carbonate platform southwest of the southern tip of India, are situated on an approximately 900 km long and 100 to 125 km wide submarine ridge consisting of a double row of atolls enclosing a deep basin, the Inner Sea (Fig. 1). It can be considered as a type locality for calcareous drift deposits. Here, in the deep water realm of the Inner Sea giant elongated drift bodies formed with geometric and seismic characteristics comparable to their siliciclastic counter parts with a typical mounded geometry and an associated moat (Lüdmann et al., 2013). Based on geometries depicted in reflection seismic profiles, we identified a new calcareous drift type at the mouth of the gateways. These drifts have a lobe-shaped external geometry with a clinoformal, prograding internal reflection configuration. We named the new sediment drifts delta drifts because they have much in common with river or tidal deltas. In 2015, during IODP Expedition 359 two platform-to-basin transects were drilled north and south of Goidhoo Atoll as well as in the Inner Sea (Betzler et al., 2016a; 2017a, b). The cores and well logs through the delta drift deposits provide the sedimentological and stratigraphic data for the comprehensive analysis of the new drift type that is presented here (Fig. 1). This research presents new diagnostic criteria that allow the classification of carbonate sediment drifts and provide the base for further detailed studies of its sedimentological characteristics and facies associations. Results from this research will also

potentially provide depositional models that could lead to a re-evaluation of carbonate deposits elsewhere and in the geological record that meet the new diagnostic criteria.

GEOLOGICAL BACKGROUND

The Maldives carbonate platform rests on a 55-57 Myrs old volcanic ridge. The Inner Sea basin, which is 300 to 350 m deep on average, is underlain by a fault-controlled en-echelon graben system (Purdy and Bertram, 1993; Aubert and Droxler, 1996) (Fig. 1). Reconstruction of the long term evolution of the Maldives was based on seismic data and industrial wells NMA-1 and ARI-1 from Elf Aquitaine and Shell as well as scientific drillholes of ODP leg 115 (Backman et al. 1988; Aubert and Droxler, 1992, 1996; Purdy and Bertram, 1993; Belopolsky and Droxler, 2003, 2004), and on the M74/4 cruise data (Betzler et al., 2009, 2014; Lüdmann et al., 2013). Subsidence was generally low averaging about 0.15 mm yr⁻¹ to 0.045 mm yr⁻¹ based on Pleistocene grainstones of Rashdoo Atoll (Gischler et al., 2008) and calculated from basalt depth of ODP Site 715 (Backman et al., 1988) located at the eastern margin of the Maldives (Fig. 1).

The Maldives are an isolated platform system far from any siliciclastic source assembling an almost complete Cenozoic sedimentary succession of exclusively calcareous material (Aubert and Droxler, 1992; Purdy and Bertram, 1993; Backman et al., 1988), with minor amount of aeolian dust (Betzler et al., 2016b). Sedimentation started with lacustrine deposits filling Eocene grabens accumulating sedimentary rocks in an anoxic environment (Aubert and Droxler, 1996). Eocene relative sea-level rise provoked neritic carbonate bank growth on the shoulders of the graben structures. A rimmed platform with a protected lagoon developed during the Early to Late Oligocene transition. In the Early Miocene (21.5 Ma) banks aggraded and prograded and a large central basin (the palaeo-Inner Sea) developed, surrounded by a narrow peripheral reef complex that faces the Indian Ocean (Aubert and Droxler, 1992, 1996; Betzler et al., 2009, 2012, 2016b). Seismic data show that the palaeo-Inner Sea was connected

via the NE-Kardiva Channel (Fig. 1) with the Indian Ocean since the Middle Miocene (Aubert
 and Droxler 1996; Lüdmann et al., 2013). Bottom currents could enter the semi-enclosed basin
 from the NE and flow along its western flank (Lüdmann et al., 2013). Aggradation of the
 platform margin continued into the Middle Miocene forming an “empty bucket” geometry of
 the palaeo-Maldives. During the late Early Miocene (18.15 Ma), the flat-topped carbonate
 bank margins started outbuilding towards the central part, thus beginning to narrow the size
 of the “empty bucket” (Betzler et al., 2016b). At the end of the Middle Miocene, the sea-level
 controlled depositional regime abruptly changed to a predominately current dominated
 system. This significant transition occurred about 13.0 Ma ago and is attributed to the onset
 and/or intensification of the Indian monsoon (Betzler et al., 2009; 2012; 2016b). This change to
 a current dominated system resulted in partial bank drowning of the platform margin, leading
 to the opening of passages that connect the Inner Sea with the Indian Ocean. Since then, 10
 mega-drift sequences were deposited in the Inner Sea (Lüdmann et al., 2013; Betzler et al.
 2017a). Contemporaneously, the remaining atolls switched from a prograding to an aggrading
 mode (Betzler et al., 2009, 2016c). Drift sedimentation initiated in the northern part of the
 Inner Sea with the deposition of lobe-shaped drift bodies at the mouth of the gateways
 adjacent to the Goidhoo Atoll (Fig. 1). The northern gateway has a present width of ca. 12 km
 and a length of ca. 17 km as well as a swell depth of 510 m. The dimension of the southern one
 is almost the same but with 420 m it is shallower. At 5.8 Myr ago, when the depocenter
 migrated eastward, large elongated mounded drifts developed with associated moats along
 the eastern basin flank (Lüdmann et al., 2013). The latter are related to a northward flowing
 bottom current that entered the Maldives from the south. Recent studies demonstrate that
 the shallow water inner atoll environment likewise is current dominated (Betzler et al., 2015).
 Present information on the oceanographic setting of the Maldives is rare (Knox, 1976). Figure 1
 shows the general bottom current pattern based on our cruises M74 (winter monsoon) and
 SO236 (summer monsoon) in the northern part of the archipelago (Lüdmann et al., 2013).

During the summer monsoon, prevailing wind direction is towards east. Bottom currents (below 150 m water depth) enter the Inner Sea from the western gateways and exiting the Maldives to the east. In the central Inner Sea a southward flow dominates. The situation turns back during the winter monsoon, then bottom water masses entering the Inner Sea from NE and leaving it through the western gateways. The central Inner Sea is marked by a northward flow. This hydrodynamic pattern is overprinted by tidal currents that act especially in the narrow gateways and shallow channels of the atolls (Kench et al., 2009).

DATA SET AND METHODS

The seismic data based on industrial and two multidisciplinary scientific cruises in 2007 and 2014, respectively. During these cruises a 144-channel digital streamer system with an active length of 600 m was used. Details about data acquisition and processing can be found in the initial IODP report Expedition 359 (Betzler et al., 2016a) and an earlier work by Lüdmann et al. (2013). IODP Expedition 359 was aimed to reconstruct the paleoceanographic evolution of the Maldives over the past 23 Myr. To achieve the scientific goals, eight sites were drilled (U1465-U1472), aligned along two transects covering shallow to deep-water deposits (Fig. 1). The standard coring systems, the advanced piston corer (APC), extended core barrel (XCB), and rotary core barrel (RCB) were used. The APC was utilized in the upper portion of each site to obtain higher quality cores, with the exception of the platform top Site U1469 where only the RCB system was applied. Total penetration for the entire expedition was about 8,725 m with the deepest drilled single hole reaching 1,003.7 m below seafloor (Hole U1471E). However, due to lithification of the carbonates in the deeper sections at each site, total recovery was less than 50 % (3,096 m). Downhole wireline logging was successfully performed at 4 sites (U1466 to U1468 and U1471). For the characterization of the sediment drift the lithostratigraphy, biostratigraphy, downhole logs as well as the physical properties of the IODP Leg 359 cores were used. A detailed description of methods can be found in the IODP Expedition 359 reports (Betzler et al., 2016a, 2017a).

For time/depth conversion of the mapped sequence boundaries previous velocity models (Lüdmann et al., 2013) were fine-tuned by using the VSI velocity data acquired during IODP Expedition 359. Additionally, core lithology was correlated with the seismic facies and major lithological boundaries were tied to the mapped seismic unconformities (Betzler et al., 2017b). By the use of post-cruise data some of the drift sequences ages were slightly corrected. Seismic-core-log correlation was carried out with the interpretation software package Petrel (Schlumberger). The 3D sequence surfaces are calculated in Petrel from the picked horizons using the convergent interpolation algorithm and a 100 x 100 m grid size.

SEISMIC ARCHITECTURE AND FACIES

Seismic line M74-65 that runs through the western Kardiva Channel into the Inner Sea basin delineating a high-resolution cross-sectional view of the platform margin displays an apparent change in the geometry of prograding clinoforms (Fig. 2). The older clinoforms (marked as carbonate platform in Fig. 2) have horizontal topsets and steep concave foresets that are part of a prograding shallow-water platform. These concave clinoforms are onlapped by a large prograding sediment body with convex upward clinoforms, the delta drift (delineated as delta drift in Fig. 2). This abrupt change of geometry marks the transition from sea-level controlled platform progradation to current-controlled drift deposition at ca. 13 Ma ago (Betzler et al., 2016b). The delta drift in front of the channel north of the Goidhoo Atoll has a slightly mounded geometry in which the apex of the mound (located at the position of Site U1468, Fig. 2) is higher than the top of the former prograding platform despite being partially eroded (Site U1466; Fig. 2). Seismic line SO236-21 that is perpendicular to the dip of the delta drift displays channels carved into the prograding clinoform bodies (Fig. 3). Additionally, seismic line SO-236-21 shows the typical bi-directional downlap pattern of a lobe and its mounded across-strike geometry. A second prograding delta drift body, with similar geometries and seismic

212 facies occurs in front of the channel south of Goidhoo Atoll (Fig. 4). The stacked lobes can be
 213 subdivided into three seismic mega-drift sequences (DS1-DS3) separated by major angular
 214 unconformities, characterized by onlap and downlap terminations (Figs. 2-4, red arrows).
 215 Basinward, the unconformities pass over into correlative conformities. North of Goidhoo Atoll
 216 mega-sequence DS1 can be further divided into two larger subsequences DS1a and DS1b. The
 217 seismic sections perpendicular to the depositional strike of the lobes (Figs. 2 and 4) display the
 218 nearly sigmoidal external shape of the sequences (DS1-DS3). They are characterized by gently
 219 dipping upper and lower segments and a thicker, more steeply inclined middle segment of
 220 prograding foresets. The upper segment shows a divergent reflection pattern with dip angles
 221 of 1°-3°. The thicker middle segment has foresets dipping basinward with angles of 3°-5°. The
 222 transition zone between the upper and middle segments is indicated by a continuous
 223 steepening of the slope forming a convex break-in-slope morphology (Figs. 2B and 4B, black
 224 curved lines). The lower segment is characterized by basinward thinning bottomsets that
 225 exhibit real or apparent downlap termination when their thickness falls below seismic
 226 resolution. Commonly the reflections indicate that strata are parallel and concordant with the
 227 sequence boundaries. Seismic line M74-62 south of Goidhoo Atoll runs southward of the
 228 deepest channel bed (Fig. 4). Here, the sequences DS1 to DS3 apparently pass over into the
 229 reef at the platform edge which was drowned at a later stage in the Tortonian compared to
 230 the northern channel. However, according to margin collapse the reflector transition from the
 231 edge to the slope strata is obscured and cannot be accurately determined (CS; Fig. 4). Most of
 232 the sediment was probably supplied from the immediately adjacent channel to the north that
 233 has cut into the platform top.

234 North of Goidhoo Atoll, the delta drift is marked in cross section by a depression like incision at
 235 the contact to the drowned Middle Miocene platform edge (Fig. 2). Here, parts of DS1 to DS3
 236 are eroded and the depression is unconformably filled with younger sediments. At Site U1466,
 237 the hiatus spans ca. 6 Mill. yrs. (Middle Miocene to Pliocene time) (Fig. 5). The channel axial

profiles (Fig. 1) allow its spatial reconstruction which forms a local elliptical incision elongated along the drowned platform edge at the mouth of the northern channel (E, Fig. 5). Its major and semi axis is about 6.5 and 1 km, respectively. The maximum incision depth of 60 m is reached at profile M74-65 (Fig. 2).

Another specific feature seen in downdip view of the delta drifts is horizontal to down-cutting reflections that interrupt the uniform clinoform geometry as demonstrated in profile M74-65 (dashed black box; Fig. 2). In cross sectional view, this zone is the rim of a dip-parallel channel cut deeply into the foreset strata (dashed black box; Fig. 3). The southward dipping channel layers that are cut along strike create side echos and generate pseudo-horizontal down-stepping reflections. These dip-parallel channels in the drift sequences DS1 and DS2 cut at their base into underlying deposits, which is of particular importance. The channels are concentrated in DS1b, reaching varied widths of 500 m to 1 km and depths of 10s of meters to 150 m (Fig. 3). The channels depict linear conduits that are restricted to the steeper middle segment of the drift clinoforms and fading out basinward. There are no deep-sea fans attached to the mouth of these channels. After initial incision, the channels exhibit a divergent, aggradational infill pattern while the channel axis remains in the center. Laterally, the channel infill passes over into the layering of the drift sequences that form the delta drift, while the channel represents a local depocenter. In places, the channel can be traced as a smooth depression in the overlying foresets strata (depression; Fig. 3). In contrast to the aforementioned channels the platform sequences show a series of gullies (Fig. 3). They are considerably smaller in dimension reaching widths of only 100-300 m and depths of 10-15 m. Line SO236-21, oriented strike-parallel to the platform sequences displays a successive downward transition from proximal platform strata below DS1 to more distal strata. Here, the gullies are restricted and aligned along the most proximal strata representing the middle slope

part of the platform clinoforms (Figs. 2 and 3). In general, our profiles depict that the gullies occur at the steeper upper to middle slope part.

At the base of DS1 wavy bottomsets appear near the toe-of-slope (Figs. 2 and 4). The waves have wavelengths of ca. 500 m and heights of 10-15 m. They have a distinct asymmetric shape and increasing wavelength in down-dip and in up-dip direction. A 3D image of their distribution demonstrates a slope-parallel N-S trend with the steep and shorter flank facing basinward (Fig. 6). Their stacking pattern indicates an upslope-migration (arrows; Fig. 6A).

The formation and early evolution of the delta drifts is depicted in Figure 7. The computed surfaces of DS01 to DS04 illustrate their development of in front of the gateways north and south of the Goidhoo Atoll. Surface DS01 represents the antecedent depositional topography with a continuous platform to the west of the paleo-Inner Sea, which was an approximately 500 m deep “empty bucket” surrounded by steep platform flanks (Fig. 7A). The onset of currents at about 13.0 Ma gradually carved channels into the carbonate platform. In the course of time these channels became deeper and wider (Figs. 7B-D). Compared to the southern channel, the northern one is much wider leading to the assumption that the effect of the bottom current was more pronounced in the north. Surfaces DS02 to DS04 (Figs. 7B-D) document the coeval progradation of the delta drifts into the Inner Sea and their successive deflection to the south. 3D geometry and cross sectional view of the delta drifts reveal their overall lobe-shaped morphology (Figs. 3 and 7). The lobes attained a maximal width of 16-17 km and a length of ca. 25 km, resulting in an area of 342 and 384 km², respectively. Their stratigraphic thickness reached up to 535 m and thereby filling the empty bucket in front of the channels.

The seismic facies of the delta drifts generally consists of continuous parallel reflections of low- to medium-amplitudes but in places, packages of strong reflections are observed (Figs. 2 and 4). These are particularly distinctive for the foresets of the southern delta drift clinoforms.

Here, in the interval between 1,000 and 1,700 m very strong amplitude reflections occur in DS1-DS3 at the steepest part of their slope (Fig. 4). In contrast, the northern delta drift shows intervals of alternating medium to strong amplitude reflectors at the bottomsets of DS1 and the flat topped apex of DS2, between 26,000 and 30,800 m and 25,200 to 28,400 m, respectively (Fig. 2).

LITHOLOGY AND AGE CONTROL

Sedimentation rates

The base of drift sequence DS1 at Sites U1466 and U1468 is dated to ca. 13.0 Ma, 11.7 Ma for DS2, to 10.55 Ma for DS3 and to 8.8 Ma for DS4 (Betzler et al., 2016b, 2017b). In the thickest portions of the drift sequences, sedimentation rates lie around 17.0 cm ka⁻¹ (DS1), 21.0 cm ka⁻¹ (DS2) and 13.5 cm ka⁻¹ (DS3). The entire delta drift was accumulated over 4.2 Myr with an average rate of 17 cm ka⁻¹. For the underlying platform sequences the sedimentation rate is significantly lower with about 4.4 cm ka⁻¹ at the proximal location of Site U1466 and 3.2 cm ka⁻¹ at the more distal location of Site U1468 (Betzler et al., 2016a).

Lithological content

Cores from Expedition 359 Sites U1466, U1468, U1471 and U1472 were drilled in order to characterize the facies of the drift sequences (Figs. 5 and 8A). Unfortunately, because of the deep incision at the former platform edge, large parts of the proximal deposits of DS1 and DS2 are missing in the northern transect, which makes the study of the downslope trends more difficult (Fig. 5). The delta drift deposits overlie the slope and basinal deposits of the drowned Middle Miocene platform. The platform sequences exhibit alternations of light (highstand) and dark, organic-rich (lowstand) layers (Fig. 9A). The lighter layers are thicker and formed by highstand shedding when the platform is flooded and most sediment is produced and exported, in contrast, the thinner darker layers are deposited during lowstand phases when the platform is exposed and sediment supply is reduced as is the case for similar deposits

elsewhere (Droxler and Schlager, 1985; Schlager et al., 1994; Eberli et al., 1997). Bioturbation is common in the platform sequences and individual trace fossils can be discriminated. In contrast, the overlying drift strata are nearly uniform in color and a distinct lamination is absent. Bioturbation is detectable; however, the bioturbation degree is too high to identify individual burrows.

The lower part of the northern delta drift deposits (DS1a), consists of lithified, medium- to coarse-grained grainstone and packstone with abundant planktonic and benthic foraminifers, including minor large benthic foraminifers such as *Miogypsinoidea* sp., *Lepidocyclina* sp., *Amphistegina* sp., and *Borelis* sp., together with other fragmented bioclasts in the proximal part, Site U1466. At the distal part, Site U1468, the lowermost facies of the delta drift consists exclusively of planktonic and small benthic foraminifers (Fig. 5). Bioturbation is very intense throughout, destroying the original texture. The vertical and lateral grain-size distribution for the northern delta drift shows coarser grained materials for the foresets of DS1a at Site U1466 (Fig. 10A) associated with a thin interval of finer bottomsets at Site U1468 (Figs. 2 and 5). With continuous progradation of the delta drift lobes, the deposits become finer at the proximal Site U1466 (Fig. 5).

At Site U1466, delta drift sequence DS1b is composed of medium- to coarse-grained, unlithified to partially lithified wackestone that gradually changes to packstone and grainstone towards the top of the sequence. The main components are small-sized benthic foraminifers, locally with some large specimen, planktonic foraminifers and minor bioclasts including red algae, bryozoans and *Halimeda* plates. Towards the top of the sequence, the facies changes into packstone and grainstone and large benthic foraminifera become the dominant component. This trend abruptly ends at the erosional unconformity with a sharp increase in planktonic foraminifers in the facies. In the more distal Site U1468, the sequence comprises packstone (Fig. 10A) that gradually changes upcore into wackestone (Fig. 10B). The main components are planktonic and small benthic foraminifers with rare sponge spicules. For DS1b,

339 Site U1466 displays a coarsening upward trend; however, its top is eroded. Comparing the
 340 time equivalent strata of DS1b at Site U1468, the aforementioned trend of the topsets at the
 341 proximal Site is associated with no significant grain size variations of the bottomsets at the
 342 distal site. The facies of DS2 (missing at the proximal Site U1466) is composed of medium- to
 343 fine-grained packstones starting from the base of the sequence up to 178.5 mbsf at Site
 344 U1468. From this depth up to 150 mbsf, there is an overall coarsening-upward trend to a
 345 rudstone at the top of the sequence (Fig. 5). Locally, shorter intervals of coarsening-upward
 346 and fining-upward occur. The coarser intervals are rich in large benthic foraminifers
 347 (*Amphistegina* sp., *Lepidocyclina* sp., *Miogypsinoidea* sp., *Heterostegina* sp., *Operculina* sp.,
 348 and *Sphaerogypsina globulus*) and locally in echinoid spines, red algae, mollusk fragments,
 349 branching and encrusting bryozoans, *Halimeda* plates. Aggregated grains are also present (Fig
 350 10C and 10E). Planktonic foraminifers are absent in this sequence and the abundance of large
 351 benthic foraminifers increases upcore reaching a maximum at the top of the sequence. The
 352 facies of DS3 is similar to those from the top of DS2 with large benthic foraminifer-rich
 353 rudstones (Fig. 5). Bioturbation in sequences DS2 and DS3 is intense and single burrows are
 354 not identified. In the southern transect there was an active carbonate platform by the time of
 355 the initial delta drift deposition (Fig. 4). The platform sediments at Site U1470 (Fig. 8A) are
 356 coeval to the drift sequences DS1 to DS3 and consist of shallow water carbonates with
 357 abundant corals and coralline algae. Site U1471 (Fig. 8A), in a more distal position, displays a
 358 distinct facies for the delta drift sequences. Here, the facies in DS1 consists of alternating fine-
 359 grained planktonic foraminifer-rich packstone and grainstone. Planktonic foraminifers,
 360 calcareous bioclasts, and calcareous nannofossils are abundant, and benthic foraminifers are a
 361 minor component. Bioturbation is abundant to common. Grainstone intervals often have a
 362 sharp basal contact with the underlying packstone. The overlying sequence DS2 consists of
 363 very fine to fine-grained wackestone to packstone with abundant planktonic foraminifers, and
 364 common to present sponge spicules, radiolarians, and calcareous nannofossils. Delta drift

sequence DS3 in its proximal position (Site U1472; Fig. 8A), comprises medium- to coarse-grained partially lithified planktonic foraminifer-rich grainstone and packstone. The main components are abundant planktonic foraminifers, benthic foraminifers, and aggregate grains/intraclasts. Among the large benthic foraminifers there are *Amphistegina*, *Lepidocyclina*, and *Miogypsina*. Mollusk fragments, gastropods, echinoid fragments, coral remains, as well as *Halimeda* and red algae fragments are present to rare. Basinwards, at Site U1471, DS3 facies changes into fully lithified very fine to coarse-grained planktonic foraminifer-rich wackestone to packstone that gradually evolve to packstone and grainstone, at the top of the sequence. Towards the top of the sequence benthic foraminifers and mollusk fragments are common.

Logs and physical properties

Figures 5 and 8A show the core (NGR) and high-resolution downhole (HSGR) natural gamma radiation profiles. The total gamma radiation signal is dominated by the contribution of uranium, with only minor contribution from thorium or potassium (Betzler et al., 2016a). Uranium variations most likely relate to the amount of organic matter. There is a clear discrepancy between NGR and HSGR at Site U1468 (Fig. 7) which may be explained by a closed caliper during downhole logging, leading to an inadequate borehole size correction of the HSGR. Here, the NGR measurements are more reliable. In general, the sequences DS1 to DS3 of the delta drift show a gamma radiation with a smoother overall trend but with a clear difference in the distal part of the northern and southern transect, respectively (U1468 and U1471; Figs. 7 and 8A). In the north the values are lower and show less variability. However, at both sites the sequence boundaries DS1-DS3 are marked by changes in radiation, especially at Site U1471. As a distinctive feature at Site U1468 there is a pronounced increase in NGR at the top of sequence DS2. Both NGR und HSGR data from Site U1468, as well as NGR data from Site

U1466 (below the depth where logging data could be recorded), show a pronounced interval of increased organic matter content within the bottom sets of the platform sequences with a high variability of the gamma radiation

Physical properties at the proximal Site U1466 exhibit a continuous increase in bulk density with depth to about 300 mbsf while porosity decreases (Fig. 9B). Below that depth density and porosity remain nearly constant. This trend is also expressed by the seismic facies, the less dense drift sequences exhibit much lower reflection amplitudes compared to the platform strata (Fig. 5). At distal Site U1468, there is an abrupt change at about 45 mbsf to nearly constant values below. This depth coincides with the unconformable contact between delta drift topset and overlying younger sheeted drift strata. The latter are characterized by low bulk density and high porosity, whereas, generally, the delta drift sequences DS1 to DS3 show a normal density and porosity trend with increasing burial depth.

DISCUSSION

We classify the herein described lobe-shaped sedimentary bodies as drift deposits that mainly accumulated through the action of persistent bottom currents. We denominate these bodies delta drifts, they represent different features compared with contourite fans. The latter, originally described a siliciclastic fine grained fan-shaped body from the Vema Channel in the Brazilian basin, downstream of a deep-water gateway which was assigned it to abyssal plain contourites (Mézerai et al., 1993; Faugères et al., 2002; Hernández-Molina et al., 2008). . Carter and McCave (1994) used the term fan drift attributed to a turbidite fan that extended into a drift. Locker and Laine (1992) introduced a companion system comprising a submarine fan/drift interaction. Such deposits, however, are different from the delta drifts we observed in the Maldives. *Sensu stricto* a contourite drift is defined as a sediment drift, principally formed by deep-water bottom currents (Stow et al. 2002a), however, the delta drifts observed

in the Maldives are not related to deep water bottom currents. Additionally, the term contourite emphasizes current orientation with respect to the bathymetric contours. The newly discovered delta drift is related to a relatively shallow water gateway and predominantly formed by currents that expand at the exit of the gateway and preferably run downslope as an underflow perpendicular to the isobaths. It can be best described as delta-like current-controlled deposits. The northern delta drift consist of four lobes and the southern one of three lobes resembling sequences DS1a, DS1b, DS2-DS3 and DS1-DS3, respectively.

Delta drift characteristics

There are several lines of evidence that indicate the current-controlled nature of the delta drifts. A compilation of their characteristics is found in Figure 11. First a comparison of the clinoform architecture of the platform and the drift sequences (DS1-DS3) clearly underpin their differences, because they exhibit an obvious contrast in slope curvature: the sea-level driven platform clinoforms have a distinct concave shape created by the foresets whereas the drift sequences exhibit a convex upward geometry (Figs. 2, 4 and 11). The 3D image clearly documents that the lobe-shaped sedimentary bodies are connected to the basinward channel exits and so do not represent a sea-level controlled regression of the platform margin (Fig. 7). Furthermore, typical gravitational foreslope deposits like slumps or debris flows have not been identified in either core samples or the seismic data. In addition, gullies as we found in the platform sequences (Fig. 3) do not exist in the delta drift. The only indication for downslope gravity-induced mass transport is observed in the distal part of the drift at Site U1471. Here, convolute bedding and soft-sediment deformation occurs at the base of the wackestone interval in drift sequence DS1 (Fig. 8A). They might be products of sediment remobilization (Phillips, 2006). However, taken into account that no clear grading was identified in these layers and that some contacts were gradational at both, the base and top of the grainstone

440 layers, it is more likely that these intervals represent high-current events related to the
 441 migration of bottom current.

442 Several observations argue for the action of long-lasting bottom currents and meet the criteria
 443 established by Stow and Faugères (2008) for the recognition of drift deposits. (1) The absent of
 444 bedding and lamination structures. In contrast to the gravity-controlled deposits of the
 445 platform sequences typical black and white layers or distinct laminations are lacking in the
 446 delta drift strata which are uniform in color. (2) Grain size from sand to clay. The grain size
 447 varies from coarse sand (e.g. large benthic foraminifers) to fine carbonate debris. (3) The
 448 degree of bioturbation. The latter is intensive and continuous (see results section). (4)
 449 Variable, low to moderate sedimentation rates. Cycles of normal and inverse grading point to a
 450 persistent bottom current flow with variation in mean current velocity or sediment supply. The
 451 latter was relatively high and the average sedimentation rate of 17 cm ka^{-1} of the delta drifts
 452 notably differs from the platform sequences with 4.4 to 3.2 cm ka^{-1} . It is comparable to rates
 453 on mounded drifts, assigned to $5\text{-}30 \text{ cm ka}^{-1}$ by Stow et al. (2008). When accommodation
 454 space was filled and the top of the delta drifts reached the swell depth of the channels, current
 455 speed increased as indicated by the dominance of grain- and rudstones (Fig. 5, U1466 and
 456 U1468). This point to intensification in transport energy of the current leading to the re-
 457 deposition of larger components and the winnowing of the finer fraction. The filled
 458 accommodation space resulted in a shallowing of the environment as evidenced by the
 459 dominance and diversity of large benthic foraminifers and the occurrence of coralline algae
 460 and *Halimeda* plates.

461 In general, the delta drifts show a proximal to distal fining trend. Basinward, very fine to fine
 462 sand-sized carbonate grains dominate and grainstones are rare. The occurrence of grainstones
 463 could be related to winnowing of the fine fraction by the postulated slope-parallel current that
 464 slightly deflected the depositional center of the delta drifts to the south (red arrow; Figs. 7B to
 465 7C). However, this current played only a minor rule in the formation of the delta drifts.

The wavy bottomsets at the base of the delta drift are herein interpreted as cyclic steps, as described for similar up-slope migrating down-slope asymmetrical sedimentary structures (Cartigny et al., 2011). The cyclic steps near the toe-of-slope are typical for the early stage of delta drift development, when the slope profile is steeper (Figs. 2 and 4). In their up-section they fade out, most likely related to a change in hydraulic regime associated with successive reduction in slope curvature. Unfortunately, the recovery in DS1a at Site U1468 is very low and details of the sedimentary facies of the cyclic steps cannot be determined. Principally, they represent a lithified packstone interval with no significant differences to the overlying strata.

Another specific feature of the delta drift are channel-like features that cut into the lobe surface down-current from the lobe apex (Fig. 3). They might be an indication for a multicore bottom current with individual flows (Fig. 11). The current remained stationary along these incised tracks while sedimentation continued over the entire lobe. Contemporaneously, the current channels were filled above the incision level and prevailed as depressions where the current stayed active (Fig. 3). The depositional elements of the mapped delta drift slope channels are not comparable to siliciclastic deep water channels or slope valleys as described for example by Deptuck et al. (2007), McHargue et al. (2011), Janocko et al. (2013). These include high-amplitude reflection patches, lateral accretion packages, onlap fill patterns and channel-levee-overbank complexes.

Giant excavation structures in the topsets of the delta drift lobes like the oval-shaped depression at the contact to the northern channel mouth represent another distinctive feature. Here, the loosely packed, non-cohesive delta drift strata onlap the former cemented platform edge (Fig. 2) and a bottom current intensification may have been responsible for the excavation. When the confined and accelerated bottom current that exits the channel reaches a threshold velocity it may generate a turbulent flow at the lithological contact, eroding into the unconsolidated delta drift beds by sweeping away the sand-sized carbonates. When this process lasted for longer periods it created the observed large oval-shaped depression in front

of the platform edge that was later refilled by younger sediments (Figs. 2 and 7b-7D).
According to the bio-stratigraphy of Site U1466 the excavations formed after the deposition of
delta drift sequences DS1 to DS3 (Fig. 5).

Depositional environment

Prerequisites for a delta drift development are: a persistent sediment supply from a point
source and sufficient available accommodation space. Suitable point sources in carbonate
platform setting are inter-atoll gateways that connect the open ocean with the platform
interior shedding huge volumes of material from different source areas into a receiving basin.
The water masses flowing through the channels catch up fallout from the water column of
dominantly planktonic microfossil assemblages from the Indian Ocean side as well as
calcareous debris driven from the adjacent atolls and produced by the organisms living in the
gateways (Fig. 7B, green arrows). An additional source are benthic organisms including large
foraminifers colonized the flattened top of the delta drift and planktic organisms living in the
water column above the drift. Erosion of the former drowned platform top played only a minor
role in the sediment budget. We calculate the total volume of the northern delta drift to be
 142 km^3 and the redeposited material eroded from the channel base accounts for only 0.05%.
On the western side of the archipelago, the west-east flowing current driven by summer
monsoon winds is stronger and the dominant driver pushing Indian Ocean water masses into
the Inner Sea (Purdy and Bertram 1993; Storz and Gischler, 2011). This dominant current
direction explains the distinct lobe-shaped morphology at the exit of the two gateways.
Changes in the flow regime possibly triggered by major oceanographic or climatic alternations
have result in the formation of major unconformities separating the delta drift sequences DS1-
DS3.

In the case of the Maldives, these gateways formed by partial drowning of the surrounding platform top in the late Middle Miocene ca. 13 Ma ago and were initiated by a modification in water mass circulation induced by the beginning of monsoon intensification (Betzler et al., 2009, 2016b). Smaller inlets are still common for the modern Maldives atolls implying that the reef rim facing the ocean is generally not a closed feature. The mechanism behind the transformation of these inlets into large gateways remains speculative and needs further research. Compared to the small inlets, the gateways allow the exchange of large volume of water between the surrounding ocean.

Accommodation changes related to relative sea-level may have played only a minor role in the formation the delta drifts. Figure 8B documents that the drift sequences DS1-DS3 were deposited during a phase of minor sea-level fluctuations as shown by benthic foraminiferal $\delta^{18}\text{O}$ values for the Atlantic and Pacific (Cramer et al., 2011). Coevally, there was no substantial influence on sea-level based on ocean basin dynamics (Müller et al., 2008). A rough estimation of the post-Middle Miocene subsidence rate provides a low value of 0.044 mm yr^{-1} that fits very well with the rate deduced from ODP Site 715 (see Geological Background). For the calculation we used the top of the carbonate platform at the base of DS1 (13.0 Ma) at Site U1465 as a past sea-level indicator (Fig. 2). It lies at the present depth of ca. 570 mbsl. Because of its thin sedimentary cover consisting of loosely packed carbonate sand (Fig. 5; Site U1465) compaction is negligible. Paleo-sea level was almost the same as today (Kominz et al., 2008). Coral reefs growing at rates of 10 mm yr^{-1} (Schlager, 2005) could easily catch up with the low subsidence rate.

Accommodation during the growth of the delta drifts was mainly controlled by the preexisting depositional topography of the Inner Sea and the magnitude of sediment supply. At the ocean facing margin of a carbonate platform a delta drifts cannot develop because accommodation space is generally too large; the slope profile generally rapidly declines to abyssal depth and sediment export outside the platform through the gateways is negligible. The delta drifts

developed in a shallower basin with few hundred meters of water depth. The Maldives palaeo-
 Inner Sea with an approximate 500 m water depth fits in this category (Fig. 7A). The gateway
 delivering the sediments is not changing its base level, in fact the delta drift attached to the
 gateway exit aggrades up to this level and at the same time outbuilds into the interior basin
 filling the Inner Sea and eventually has a crest that is higher than the spill height of the channel
 (Fig. 2). The bottom current in the gateway provides a steady sediment supply and distributes
 the sediments after leaving the gateway along the slope profile, generating a Gaussian
 curvature and keeping the angle-of-repose below a threshold for triggering significant slope
 failure. Distinct changes in current speed occurred as documented by the waning and waxing
 structures of the sediments. They are possibly related to variations in monsoon wind strength
 as well as widening and/or deepening of the gateway induced by sea-level fluctuations. These
 variations in current speed could have also influenced the organic matter proportion in the
 delta drift sequences as indicated by the natural gamma radiation. Wetzel et al. (2008)
 proposed that higher current speeds generally winnow out organic matter. There is a slight
 trend in increasing gamma ray values at the top of each sequence, which in turn is related to a
 decrease in current speed. However, variations in nutrient supply is another important factor.

CONCLUSIONS

Reflection seismic data together with cores and logs from IODP Expedition 359 document a
 new sediment drift type located in front of two gateways. We term this new type a delta drift
 and classify it as a channel-related drift that is emplaced under a long-term, current-driven
 sediment flux regime. Our data do not show indication of significant gravity-controlled
 sedimentation in the delta drift. The drift bodies form stacks of individual lobes in front of two
 gateways. Both delta drifts are nearly identical and are deposited in front of the edge of a

566 drowned carbonate platform. The main characteristic of the delta drifts discovered on
 567 Maldivian carbonate platform are:

- 568 • The delta drifts are situated in the Inner Sea basin at the downstream exit of a shallow
 569 and over time deepening gateway. In contrast to a fluvial delta system, the delta drifts
 570 accumulated considerably below base level;
- 571 • Individual lobes may built up stacks that resemble a delta drift;
- 572 • Current flow was wide-ranging on the drift body perpendicular to the isobaths. Smaller
 573 channels running downdip are developed on top of the delta drift lobes indicating that
 574 they must have been fed by a multicore current;
- 575 • The delta drifts exhibit progradation with pronounced sigmoidal clinoform geometry.
 576 Their spatial extent is about 342-384 km² and the depositional relief reaches up to 500
 577 m;
- 578 • The delta drifts show a distinct coarsening upward trend in grain size, related to a
 579 shallowing of the depositional setting when accommodation space is filled up to the
 580 level of the feeder channel bed;
- 581 • During the early phase of deposition, when the depositional profile of the slope was
 582 still concave from the underlying distal slope clinoforms, cyclic steps developed at the
 583 toe-of-slope;
- 584 • Large excavations with depths of up to 80 m may occur where the bottom current
 585 overflows beds of consolidated and unconsolidated material and turns into a turbulent
 586 flow. They reach dimension of 10 km² and preferably develop at the onlap termination
 587 of the former cemented platform edge and the loosely packed coarse delta drift
 588 topsets.

589

590 Acknowledgements

Our special thanks go to the IODP Expedition 359 drilling crew, ship's crew and scientific staff of the Drillship JOIDES Resolution. The expedition was funded by the US National Science Foundation (NSF); the European Consortium for Ocean Research Drilling (ECORD); the Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT); the Ministry of Science and Technology (People's Republic of China); the Korea Institute of Geoscience and Mineral Resources; the Australian Research Council and the New Zealand Institute for Geological and Nuclear Sciences; and the Ministry of Earth Sciences (India). The German Federal Ministry of Education and Research is thanked for funding the seismic surveys (03S0405, 03G0236A). Additional acknowledgements go to the companies HALLIBURTON-LANDMARK, SCHLUMBERGER and CEGAL Geosciences for providing university grants for the seismic processing software PROMAX and seismic interpretation software PETREL as well as its BLUEBACK plugin. We thank M. Rebesco, as the editor of this manuscript, and T. Mulder as well as D.A.V. Stow, as reviewers, for their very helpful comments that considerably improved the manuscript.

References

- Anselmetti, F.S., Eberli, G.F., Ding, Z.-D., 2000. From the Great Bahama Bank into the Straits of Florida: A margin architecture controlled by sea-level fluctuations and ocean currents. Geological Society of America Bulletin 112, 829–844.
- Aubert, O., Droxler, A.W., 1992. General Cenozoic evolution of the Maldives carbonate system (equatorial Indian Ocean). Bulletin Centres Recherche Exploration Production Elf Aquitaine 16, 113-136.
- Aubert, O., Droxler, A.W., 1996. Seismic stratigraphy and depositional signatures of the Maldivian Carbonate System (Indian Ocean). Marine and Petroleum Geology 13, 503-536.

- 615 Backman, J., Duncan, R.A., et al., 1988. Mascarene Plateau - Sites 705-716. Proceedings of the
616 Ocean Drilling Program, Initial Reports 115, College Station, TX.
- 617 Belopolsky, A., Droxler, A., 2003. Imaging Tertiary Carbonate System - the Maldives, Indian
618 Ocean: Insights into Carbonate Sequence Interpretation. *The Leading Edge*, 22, 646-652.
- 619 Belopolsky, A.V., Droxler, A.W., 2004. Seismic expressions of prograding carbonate bank
620 margins: middle Miocene, Maldives, Indian Ocean, in: Eberli, G.P., Masferro, J.L., Sarg, J.F.
621 (Eds.), *Seismic Imaging of Carbonate Reservoirs and Systems*, American Association Petroleum
622 Geologists Memoir 81, Tulsa, pp. 267–290..
- 623 Betzler, C., Eberli, G.P., Alvarez Zarikian, C.A., and the Expedition 359 Scientists, 2017a.
624 Maldives Monsoon and Sea Level. Proceedings of the International Ocean Discovery Program,
625 359: College Station, TX (International Ocean Discovery Program).
626 <http://dx.doi.org/10.14379/iodp.proc.359.2017>.
- 627 Betzler, C., Eberli, G.P., Alvarez Zarikian, C.A., and the Expedition 359 Scientists, 2016a.
628 Expedition 359 Preliminary Report: Maldives Monsoon and Sea Level. International Ocean
629 Discovery Program. <http://dx.doi.org/10.14379/iodp.pr.359.2016>.
- 630 Betzler, C., Eberli, G.P., Kroon, D., Wright, J.D., Swart, P.K., Nath, B.N., Alvarez-Zarikian, C.A.,
631 Alonso-Garcia, M., Bialik, O.M., Blatter, C.L., Guo, J.A., Haffen, S., Horozal, S., Inoue, M.,
632 Jovane, L., Lanci, L., Laya, J.C., Mee, A.L., Lüdmann, T., Nakakuni, M., Niino, K., Petruny, L.M.,
633 Pratiwi, S.D., Reijmer, J.J., Reolid, J., Slagle, A.L., Sloss, C.R., Su, X., Yao, Z., Young, J.R., 2016b.
634 The Abrupt Onset of the Modern South Asian Monsoon Winds. *Science Report*, **6**, 29838.
- 635 Betzler, C., Eberli, G.P., Lüdmann, T., Reolid, J., Kroon, D., Reijmer, J.J.G., Swart, P.K., Wright, J.,
636 Young, J.R., Alvarez-Zarikian, C., Alonso-García, M., Bialik, O.M., Blättler, C.L., Guo, J.A., Haffen,
637 S., Horozal, S., Inoue, M., Jovane, L., Lanci, L., Laya, J.C., Hui Mee, A.L., Nakakuni, M., Nath,
638 B.N., Niino, K., Petruny, L.M., Pratiwi, S.D., Slagle, A.L., Sloss, C.R., Su, X., Yao, Z., 2017b.

- 639 Refinement of Miocene sea-level and monsoon events from the sedimentary record of the
 640 Maldives (Indian Ocean). *Progress in Earth and Planetary Science*, in press.
- 641 Betzler, C., Fürstenau, J., Lüdmann, T., Hübscher, C., Lindhorst, S., Paul, A., Reimer, J., Droxler,
 642 A.W., 2012. Sea-level and ocean-current control on carbonate platform growth, Maldives,
 643 Indian Ocean. *Basin Research* 24, 1-15.
- 644 Betzler, C., Hübscher, C., Lindhorst, S., Lüdmann, T., Reijmer, J.J.G., Braga, J.-C., 2016c.
 645 Lowstand wedges in carbonate platform slopes (Quaternary, Maldives, Indian Ocean). *The*
 646 *Depositional Record* 2, 196-207.
- 647 Betzler, C., Hübscher, C., Lindhorst, S., Reijmer, J.J.G., Römer, M., Droxler, A., Fürstenau, J.,
 648 Lüdmann, T., 2009. Monsoon-Induced Partial Carbonate Platform Drowning (Maldives, Indian
 649 Ocean). *Geology* 37, 867-870.
- 650 Betzler, C., Lindhorst, S., Eberli, G.P., Lüdmann, T., Möbius, J., Ludwig, J., Schütter, I., Wunsch,
 651 M., Reijmer, J.J.G., Hübscher, C., 2014. Periplatform drift: the combined result of contour
 652 current and off-bank transport along carbonate platforms. *Geology* 42, 871-874.
- 653 Betzler, C., Lindhorst, S., Lüdmann, T., Weiss, B., Wunsch, M., Braga, J.C., 2015. The Leaking
 654 Bucket of a Maldives Atoll: Implications for the Understanding of Carbonate Platform
 655 Drowning. *Marine Geology* 366, 16-33.
- 656 Betzler, C., Lüdmann, T., Hübscher, C., Fürstenau, J., 2013. Current and Sea-Level Signals in
 657 Periplatform Ooze (Neogene, Maldives, Indian Ocean). *Sedimentary Geology* 290, 126-137.
- 658 Carter, L., McCave, I.N., 1994. Development of drift sediments approaching an active plate
 659 margin und the SW Pacific Deep Western Boundary Current. *Paleoceanography* 9(6), 1061-
 660 1085.

- 661 Cartigny, M.J.B., Postma, G., van den Berg, J.H., Mastbergen, D.R., 2011. A comparative study
 662 of sediment waves and cyclic steps based on geometries, internal structures and numerical
 663 modeling. *Marine Geology* 280(1-4), 40-56.
- 664 Cramer, B.S., Miller, K.G., Barrett, P.J., Wright, J.D., 2011. Late Cretaceous–Neogene trends in
 665 deep ocean temperature and continental ice volume: Reconciling records of benthic
 666 foraminiferal geochemistry ($\delta^{18}\text{O}$ and Mg/Ca) with sea level history Ocean overturning since
 667 the Late Cretaceous: Inferences from a new benthic foraminiferal isotope compilation. *Journal*
 668 *Geophysical Research* 116. Doi:10.1029/2011JC007255.
- 669 Deptuck, M.E., Sylvester, Z., Pirmez, C., O’Byrne, C., 2007. Migration–Aggradation History and
 670 3-D Seismic Geomorphology of Submarine Channels in the Pleistocene Benin-Major Canyon,
 671 Western Niger Delta Slope. *Marine and Petroleum Geology* 24, 406-433.
- 672 Droxler, A. W., Schlager, W., 1985. Glacial versus interglacial sedimentation rates and turbidite
 673 frequency in the Bahamas. *Geology* 13, 799-802.
- 674 Eberli, G.P., Anselmetti, F.S., Isern, A.R., Delius, H., 2010. Timing of changes in sea-level and
 675 currents along the Miocene platforms of the Marion Plateau, Australia, in: Morgan, W.A.,
 676 George, A.D., Harris, P.M., Kupecz, J.A., Sarg J.E. (Eds.), *Cenozoic Carbonate Systems of*
 677 *Australasia* by), Society for Sedimentary Geology Special Publication 95, pp. 219-242.
- 678 Eberli, G.P., Swart, P.K., Malone, M.J., et al., 1997. *Proceedings ODP, Initial Reports* 166,
 679 College Station, TX (Ocean Drilling Program).
- 680 Faugères, J.-C.S., D.A.V., Imbert, P., Viana, A., 1999. Seismic features diagnostic of contourite
 681 drifts. *Marine Geology* 162, 1-38.
- 682 Faugères, J.-C., Zaragosi, S., Mézerais, M.L., Massé, L., 2002. The Vema contourite fan in the
 683 South Brazilian basin, in: Stow, D.A.V., Pudsey, C.J., Howe, J.A., Faugères, J.-C., Viana, A.R.

- 684 (Eds.), *Deep-Water Contourite Systems: Modern Drifts and Ancient Series*, Seismic and
 685 *Sedimentary Characteristics*, Geological Society London Memoir 22, pp. 209-238.
- 686 Gischler, E., Hudson, H.J., Pisera, A., 2008. Late Quaternary reef growth and sea level in the
 687 Maldives (Indian Ocean). *Marine Geology* 250, 104-113.
- 688 Hernández-Molina, F.J., Maldonado, A., Stow, D.A.V., 2008. Abyssal plain contourites, in:
 689 Rebesco, M., Camerlenghi, A. (Eds.), *Contourites, Developments in Sedimentology* 60, Elsevier,
 690 pp. 347-378.
- 691 Isern, A.R., Anselmetti, F.S., Blum, P., 2004. A neogene carbonate platform, slope, and shelf
 692 edifice shaped by sea-level and ocean currents, Marion Plateau (Northeast Australia), in:
 693 Eberli, G.P., Masferato, J.L., Sarg, J.F.R. (Eds.), *Seismic Imaging of Carbonate Reservoirs and*
 694 *Systems*, American Association Petroleum Geologists Memoir 81, 291-307.
- 695 Janocko, M., Nemec, W., Henriksen, S., Warchol, M., 2013. The diversity of deep-water sinuous
 696 channel belts and slope valley-fill complexes. *Marine and Petroleum Geology* 41, 7-34.
- 697 Kench, P.S., Parnell, K.E., Brander, R.W., 2009. Monsoonally influenced circulation around coral
 698 reef islands and seasonal dynamics. *Marine Geology* 266, 91-108.
- 699 Knox, R.A., 1976. On a long series of measurements of Indian Ocean equatorial currents near
 700 Addu Atoll. *Deep-Sea Research* 23, 211-221.
- 701 Kominz, M.A., Browning, J.V., Miller, K.G., Sugarman, P.J., Mizintseva, S., Scotese, C.R., 2008.
 702 Late Cretaceous to Miocene sea-level estimates from the New Jersey and Delaware coastal
 703 plain coreholes: an error analysis. *Basin Research* 20, 211–226.
- 704 Locker, S.D., Laine, E.P., 1992. Paleogene-Neogene depositional history of the middle U.S.
 705 Atlantic continental rise: mixed turbidite and contourite depositional systems. *Marine Geology*
 706 103, 137-164.

- 707 Lüdmann, T., Kalvelage, C., Betzler, C., Fürstenau, J., Hübscher, C., 2013. The Maldives, a Giant
 708 Isolated Carbonate Platform Dominated by Bottom Currents. *Marine and Petroleum Geology*
 709 43, 326-340.
- 710 Lüdmann, T., Wiggershaus, S., Betzler, C., Hübscher, C., 2012. Southwest Mallorca Island: A
 711 cool-water carbonate margin dominated by drift deposition associated with giant mass
 712 wasting. *Marine Geology* 307-310, 73-87.
- 713 McHargue, T., Pyrcz, M.J., Sullivan, M.D., Clark, J.D., Fildani, A., Romans, B.W., Covault, J.A.,
 714 Levy, M., Posamentier, H.W., Drinkwater, N.J., 2011. Architecture of Turbidite Channel Systems
 715 on the Continental Slope: Patterns and Predictions. *Marine and Petroleum Geology* 28, 728-
 716 743.
- 717 Mézerais, M.L., Faugères, J.-C., Figueiredo, A., Massé, L., 1993. Contour current accumulation
 718 off Vema Channel mouth, Southern Brazil basin. *Sedimentary Geology* 82(1-4), 173-188.
- 719 Müller, D.R., Sdrolias, M., Gaina, C., Steinberger, B., Heine, C., 2008. Long-term sea-level
 720 fluctuations driven by ocean basin dynamics. *Science* 319, 1357-1362.
- 721 Phillips, E., 2006. Micromorphology of a debris flow deposit: evidence of basal shearing, hydro-
 722 fracturing, liquefaction and rotational deformation during emplacement. *Quaternary Science*
 723 *Reviews* 25(7-8), 720-738.
- 724 Purdy, E.G., Bertram, G.T., 1993. Carbonate Concepts from the Maldives, Indian Ocean.
 725 *American Association Petroleum Geologists Studies in Geology* 34, pp. 56.
- 726 Rebesco, M., Camerlenghi, A., 2008. Contourites. *Developments in Sedimentology* 60, Elsevier,
 727 pp. 663.

- 728 Rebesco, M., Hernández-Molina, F.J., Van Rooij, D., Wåhlin, A., 2014. Contourites and
 729 associated sediments controlled by deep-water circulation processes: State-of-the-art and
 730 future considerations. *Marine Geology* 352, 111-154.
- 731 Schlager, W., 2005. Carbonate Sedimentology and Sequence Stratigraphy. Concepts in
 732 Sedimentology and Paleontology 8, Society for Sedimentary Geology, pp. 200.
- 733 Schlager, W., Reijmer, J.J.G., Droxler, A., 1994. Highstand shedding of carbonate platforms.
 734 *Journal of Sedimentary Research; Section B – Stratigraphy and Global Studies*, 64(3), 270-281.
- 735 Storz, D., Gischler, E., 2011. Coral extension rates in the NW Indian Ocean I: reconstruction of
 736 20th century SST variability and monsoon current strength. *Geo-Marine-Letters* 31, 141-154.
- 737 Stow, D.A.V., Faugères, J.-C., 2008. Contourite facies and the facies model, in: Rebesco, M.,
 738 Camerlenghi A. (Eds.), *Contourites, Developments in Sedimentology* 60, Elsevier, 223-256.
- 739 Stow, D.A.V., Faugères, J.-C., Howe, J.A., Pudsey, C.J., Viana, R., 2002a. Bottom currents,
 740 contourites and deep-sea sediment drifts: current state-of-the-art, in: Stow, D.A.V., Pudsey,
 741 C.J., Howe, J.A., Faugères, J.-C., and Viana, A.R. (Eds.), *Deep-Water Contourite Systems:*
 742 *Modern Drifts and Ancient Series, Seismic and Sedimentary Characteristics*, Geological Society
 743 London Memoir 22, 7-20.
- 744 Stow, D.A.V., Hunter, S., Wilkinson, D, Hernández-Molina, F.J., 2008. The Nature of Contourite
 745 Deposition, in: Rebesco, M., Camerlenghi A. (Eds.), *Contourites, Developments in*
 746 *Sedimentology* 60, Elsevier, 143-155.
- 747 Stow, D.A.V., Pudsey, C.J., Howe, J.A., Faugères, J.-C., Viana, A.R., 2002b. Deep-Water
 748 Contourite Systems: Modern Drifts and Ancient Series, Seismic and Sedimentary
 749 Characteristics. Geological Society London Memoir 22, pp. 464.

750 Viana, A.R., Rebesco, M., 2007. Economic and Palaeoceanographic Significance of Contourite
 751 Deposits. Geological Society of London, Special Publications 276, pp. 343.

752 Wetzel, A., Werner, F., Stow, D.A.V., 2008. Bioturbation and biogenic sedimentary structures in
 753 contourites, in: Rebesco, M., Camerlenghi, A. (Eds.), Contourites, Developments in
 754 Sedimentology 60, Elsevier, 183-202.

755

756 Fig. 1: A) Map of the study area (red box) in the northern part of the Maldives. B) Distribution
 757 of seismic lines (black lines) and the location of IODP Expedition 359 Sites as well as ODP Site
 758 716 and 715. Indicated is the general present bottom current pattern below a water depth of
 759 about 200 m (compiled after Lüdmann et al., 2013). Example seismic profiles are marked in
 760 orange. Present reefs are color-coded (yellow: island; blue: reef; green: 10-100 fathoms). WM:
 761 winter monsoon; SW: summer monsoon.

762 Fig. 2: A) Part of reflection W-E seismic line M74-65 that crosses the top of the Middle
 763 Miocene carbonate platform and delta drift north of Goidhoo Atoll in dip direction (for
 764 location see Fig. 1). B) Interpreted version showing mapped boundaries of the drift mega-
 765 sequences. Drift sequence DS1 can be subdivided into subsequence a and b. Indicated are the
 766 positions of the IODP Expedition 359 drillsites. Indicated in grey is the delta drift which rests on
 767 the exponential clinoforms of the former platform margin. Red wavy line marks a local
 768 erosional unconformity. Dashed lines exhibit possible fault traces. Red arrows show reflector
 769 termination. Black boxes reveal to an area of side echoes (see also cross line in Fig. 3) and an
 770 excavation zone (see text for further explanation). CS: collapse structure. Dip indicator:
 771 average velocity applied 1600 m/s.

772 Fig. 3: A) Part of reflection N-S seismic line SO-236-21 that crosses the former platform talus
 773 and delta drift north of Goidhoo Atoll in strike direction (for location see Fig. 1). B) Interpreted
 774 version showing mapped boundaries of the drift mega-sequences. Drift sequence DS1 can be

775 subdivided into subsequence a and b. Displayed in grey are the mounted delta drift lobe strata
 776 in strike view, underlain by the former platform foresets. Dotted black line indicates base of
 777 incised channel that are crossed perpendicular to their downdip trend. Red wavy line marks a
 778 local erosional unconformity. Black triangles point to the location of gullies that occur
 779 exclusively in the platform sequences and are absent in the drift sequences. Red arrows show
 780 reflector termination. Dashed black lines exhibit possible fault traces. Black boxes indicate an
 781 area of side echoes (see also cross line in Fig. 2 and text for further explanation).

782 Fig. 4: A) Part of reflection W-E seismic line M74-62 that crosses the top of the Middle
 783 Miocene carbonate platform and delta drift south of Goidhoo Atoll in dip direction (for
 784 location see Fig. 1). B) Interpreted version showing mapped boundaries of the drift mega-
 785 sequences. DS1 is underlain by the prograding platform foresets, above in grey the sigmoidal
 786 clinoforms of the delta drift lobe. Marked with black lines are the clinoform roll-overs. Red
 787 arrows show reflector termination. Dashed black lines exhibit possible fault traces. CS: collapse
 788 structure. Dip indicator: average velocity applied 1600 m/s.

789 Fig. 5: Lithostratigraphic correlation of the IODP Expedition 359 Sites along the northern
 790 transect (see also Fig. 1). It crosses the platform top (U1465), the proximal upper slope
 791 (U1466) and the more distal middle slope (U1468) setting. Indicated are the boundaries of the
 792 drift sequences (DS). In color are DS1 to DS3, representing the deposits of the delta drift
 793 discussed in the text. Marked in green are the stratigraphically equivalent parts of DS1 and DS2
 794 in Sites U1466 and U1468, respectively. In addition, core and log total gamma radiation are
 795 plotted with the seismic data (distance 250 m) at the well location (red line) in the background.
 796 Depth is shown in mbsf.

797 Fig. 6: A) Detail of profile M74-65 (see Fig. 2 for location) showing aggradational asymmetric
 798 cyclic steps with steeper basinward flank (transparent black arrows indicate upslope
 799 migration). B) 3D view of cyclic steps at the base of drift sequence DS1. Color-coded is the dip

angle indicating a low upslope and a high downslope inclination of the margin-parallel sediment waves. It is characteristic for up-slope migrating asymmetrical cyclic steps after classification of Cartigny et al. (2011).

Fig. 7: Sketch of the Maldives palaeo-topography in the study area. View is to the west. A) Surface DS01 representing the initial “empty bucket” phase of the Inner Sea ca. 12.9 Ma ago. During this time the Inner Sea formed a semi-enclosed basin connected to the Indian Ocean via the northeast Kardiva Channel (Fig. 1). With the deposition of DS1 the platform top partially starts to drown and two proto-channels formed that connect the paleo-Inner Sea with the Indian Ocean. B) Top of delta drift sequence DS1 (DS02) demonstrating the lobe-shape infill of sequence DS1 attached to both waterways. The depocenter is deflected slightly southward by a south directed contour current (red arrow). Green arrows indicate possible sediment sources. C) Top of delta drift sequence DS2 (DS03) showing the progradation of sequence DS2 into the Inner Sea basin. DS2 continued to be deflected southward. Channel widening dominates over channel incision by platform backstepping. D) Top of delta drift sequence DS3 (DS04) revealing the final stage of delta drift formation ceasing at ca. 9.0 Ma. Indicated are the positions of the IODP Expedition 359 Sites. Isolines show depth in TWT. Red arrows indicate an assumed slope-parallel current, entering the Inner Sea from the deep eastern Kardiva Channel (see Fig. 1 for location). CS: collapse structure; E: excavation.

Fig. 8: A) Lithostratigraphic correlation of the IODP Expedition 359 Sites along the southern transect (see also Fig. 1). It crosses the platform top (U1470), the proximal upper slope (U1472) and the more distal middle slope (U1471) setting. Indicated are the boundaries of the drift sequences (DS). In color are DS1 to DS3, representing the deposits of the delta drift discussed in the text. In addition, core and log total gamma radiation are plotted with the seismic data (distance 250 m) at the well location (red line) in the background. Depth is shown in mbsf. B) $\delta^{18}\text{O}$ curve for the South Atlantic and Pacific after Cramer et al. (2011) with location of the delta drift sequences DS1 to DS3.

826 Fig. 9: A) Core section 61X (379.2 mbsf) and 70X (466.6 mbsf) of Site U1468 showing typical
 827 drift and platform slope strata, respectively (see Fig. 1 for location). Characteristic is the color
 828 change of the platform slope sequence. B) Bulk density and porosity measured on cores from
 829 Sites U1466 and U1468.

830 Figure 10: Facies of the delta drift (scale bar for core photographs =1 cm, scale bar for
 831 photomicrographs =1 mm). A) Core photograph of a fine-grained bioclastic packstone at Site
 832 U1468. B) Core photograph of an unlithified wackestone with intraclasts at Site U1468. C) Core
 833 photograph of a large benthic foraminifera-rich rudstone to grainstone with fining upward at
 834 Site U1468. Echinoid spines and intraclasts are common. D) Photomicrograph of the fine-
 835 grained packstone at Site U1466. E) Photomicrograph of the rudstone to grainstone facies at
 836 U1468 with abundant *Amphistegina* and intraclasts. See also figure 5 for location of samples.

837 Fig. 11: Summary of the delta drift characteristics, including a sketch showing the channel-
 838 related drift in strike and dip view. Gnstn: grainstone; pkstn: packstone; rdstn: rudstone;
 839 wkstn: wackestone.