



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Applicability of T1-weighted MRI in the assessment of forensic age based on the epiphyseal closure of the humeral head

Citation for published version:

Ekizoglu, O, Inci, E, Ors, S, Kacmaz, IE, Basa, CD, Can, IO & Kranioti, E 2018, 'Applicability of T1-weighted MRI in the assessment of forensic age based on the epiphyseal closure of the humeral head', *International journal of legal medicine*, pp. 1-8. <https://doi.org/10.1007/s00414-018-1868-7>

Digital Object Identifier (DOI):

[10.1007/s00414-018-1868-7](https://doi.org/10.1007/s00414-018-1868-7)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

International journal of legal medicine

Publisher Rights Statement:

The final publication is available at Springer via: <https://doi.org/10.1007/s00414-018-1868-7>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Applicability of T1-weighted MRI in the assessment of forensic age based on the epiphyseal closure of the humeral head

*Oguzhan Ekizoglu¹, Ercan Inci²,Suna Ors², Ismail Ozgur Can³, *Elena F. Kranioti^{4,5,6}

* Equal contribution to the paper.

¹Department of Forensic Medicine, Tepecik Training and Research
Hospital, Izmir, Turkey

²Department of Radiology, Bakirkoy Dr.Sadi Konuk Training and Research
Hospital, Izmir, Turkey

³Department of Forensic Medicine, Dokuz Eylul University, Faculty of Medicine,
Izmir, Turkey

⁴Edinburgh Unit for Forensic Anthropology, School of History, Classics and
Archaeology, University of Edinburgh, 4 Teviot Place, EH8 9AG.

⁵Forensic Pathology Division Crete, Hellenic Republic Ministry of Justice and
Human

Rights, Heraklion, Crete, Greece.

6Department of Medical Imaging, University Hospital of Heraklion, Heraklion, Crete 71110, Greece.

Corresponding author:

Oguzhan Ekizoglu

Department of Forensic Medicine, Tepecik Training and Research

Hospital, Izmir, Turkey

Güney mahallesi 1140/1 Yenisehir - Konak - Izmir, Turkey

E-mail: drekizoglu@gmail.com

Phone: +90 5324320139

Fax: +90 2125424491

Cite as:

Ekizoglu O, Inci E, Ors S, Eralp Kacmaz I, Doruk Basa C, Ozgur Can I, Kranioti EF. Applicability of T1-weighted MRI in the assessment of forensic age based on the epiphyseal closure of the humeral head. Int J Leg Med 2018 (in press).

Abstract

This work investigates the value of magnetic resonance imaging analysis of proximal epiphyseal fusion in research examining the growth and development of the humerus and its potential utility in establishing forensic age estimation.

In this study, 428 proximal humeral epiphyses (patient age, 12–30 years) were evaluated with T1-weighted turbo spin echo (T1 TSE) sequences in coronal oblique orientation on shoulder MRI images. A scoring system was created following a combination of the Schmeling and Kellinghaus methods.

Spearman's rank correlation analysis revealed a significant positive relationship between age and ossification stage of the proximal humeral epiphysis (all subjects: $\rho=0.664$, $p < 0.001$; males: 0.631 , $p < 0.001$; females: $\rho=0.651$, $p < 0.001$). The intra- and inter-observer reliability assessed using Cohen's kappa statistic was $\kappa=0.898$ and $\kappa=0.828$, respectively. The earliest age of epiphysis closure was 17 years for females and 18 years for males.

MRI of the proximal humeral epiphysis can be considered advantageous for forensic age estimation of living individuals in a variety of situations, ranging from monitoring public health to estimating the age of illegal immigrants/asylum seekers, minors engaged in criminal activities and illegal participants in competitive sports, without the danger of radiation exposure.

Keywords age estimation, proximal humeral epiphysis, magnetic resonance imaging

Introduction

Forensic age estimation (FAE) is a proxy for the biological maturation of an individual, which is important for a variety of demographic, clinical and forensic purposes. Every skeletal part of the human body develops in a predictable way, which enables for the estimation of skeletal age, given specific markers [1]. Skeletal development is subject to a combination of genetic and environmental factors; thus, recording the process in different parts of the world is of crucial importance for public health. The level of skeletal maturation is often requested by clinicians, such as endocrinologists, in order to be compared with chronological age in suspected developmental disorders in children, to monitor response to hormone therapy or to estimate stature in healthy subjects [1-4]. Estimating the age of living individuals is an important requirement for the birth registration of unknown persons [5-7]. Moreover, the age of a living individual is required in cases concerning migration and legal responsibility. Legal systems in different countries stipulate different age limits for legal assessment, particularly for those aged 10–21 years, with 14, 16, 18, and 21 years as notable thresholds [7-9]. Age assessment plays a critical role in the definitions of the civil rights of refugees and those seeking asylum and in the specification of the age of marriage and adoption [7,8,10]. Last, FAE is crucial in establishing entry thresholds for participants in competitive sports [11].

The primary application area of radiological evaluations for skeletal maturation is the epiphyseal union. Although skeletal maturation is driven by a

combination of intrinsic and extrinsic factors, it has been demonstrated that estrogen is primarily responsible for ultimate epiphyseal fusion [12]. In fact, it has been shown that prepubertal estrogen levels in girls are over seven-fold higher than in boys, which explains the five-fold more advanced skeletal maturation of girls in childhood [13]. It is thus possible that different estrogen levels between the sexes would impact the epiphyseal union time.

In long bones, the epiphyseal conversion to fatty marrow is noted within 6 months of the radiological appearance of the secondary ossification center. The fatty transformation continues in the diaphysis and moves toward the metaphysis. The last parts of the appendicular skeleton to convert are the proximal humeral and femoral metaphyses [14]. Epiphyseal closure has been studied to reveal minimal age limits for forensic age estimation using different articulations, imaging modalities and scoring systems [1,15-19]. The recommended methods for age estimation of living individuals by several forensic associations include physical and radiological examination of children and adolescents [6-7]. However, in recent years, radiation exposure in the pediatric age group has caused ethical concerns [20]. Therefore, developmental studies of living individuals have shifted to non-invasive modalities, such as ultrasonography [21-23] and magnetic resonance imaging (MRI) [15-17, 24-28].

MRI indeed offers a remarkable insight into the dynamic process of skeletal growth and maturation, as demonstrated by an increasing number of studies in children and adolescents in recent years [15-17, 24-40]. The vast majority of these studies focused on the clavicle, hand and wrist, producing large

reference databases for different populations. A direct application of this body of work was the introduction of an MRI examination of the wrist for age verification in international youth competitions (under 17 years old) by the Federation Internationale de Football Association (FIFA) in Nigeria in 2009 [41]. Despite the encouraging results of the MRI imaging in the investigation of age estimation, only one study focused on the development of the proximal humeral epiphysis but presented several limitations in methodology and sample size [39]. The current study aims to address these issues by investigating the epiphyseal union of the proximal humerus in a large sample ($N > 400$) with the aid of a hybrid staging system of previous methodologies. MRI imaging of the shoulder has a significant advantage in achieving detailed imaging of the non-ossified cartilaginous epiphysis, the secondary ossification centers, and the physis. MRI has also demonstrated to be highly efficient in detecting infections and tumors in young children or detecting sports-related injuries (intra-articular shoulder derangements and osseous abnormalities associated with shoulder dislocation) in older children and young adults [39]. The present work aims to evaluate the value of MRI analysis of the proximal epiphyseal fusion in the study of growth and development of the humerus and its potential utility in establishing FAE.

Materials and Methods

This cross-sectional retrospective study was performed at Bakırköy Dr Sadi Konuk Research and Training Hospital, İstanbul, Turkey from January 1, 2014 to January 1, 2016. We evaluated the left shoulder MRI of 449 patients who

were 12-30 years of age and who were admitted to the radiology department with traumatic and ligamentous disorders after clinical evaluation. Five patients with traumatic shoulder bone fracture, 3 patients with operation history, and 13 patients with insufficient age and sex information were excluded from the study. A total of 428 patients were included in the study. All medical documents and information on patient age and sex were obtained from the data-processing center of the hospital. Data were anonymized to comply with medical ethics at our institution.

All MRI examinations were performed with a 1.5-T whole body scanner (Avanto; Siemens, Erlangen, Germany) with comparable imaging properties using an extremity coil. For the analysis of the scans, T1-weighted turbo spin echo (T1 TSE) sequences in coronal oblique orientation were used (TR, 500 ms; TE, 15 ms; 150 mm; FoV, 150 mm; slice thickness, 2-4 mm; voxel size, 0.5x0.5x3.5 mm, and scan time 1 min 44 s). The ossification stage of the proximal humeral epiphysis was scored using a combination of the staging systems published by Schmeling et al. [18] and by Kellinghaus et al. [19]. Stages describe the ossification process of the proximal humeral epiphysis, as seen below, and applying the classification schemes, as shown in detail in Figure 1 and 2.

All MRI slices were evaluated in all cases for stage detection. The decision for stages 1,2a-2b-2c and 3a-3b-3c was based on the most-developed epiphysis in the slices. For stage 4, at least one epiphyseal scar should be observed in all sections. For stage 5, epiphyseal closure and no epiphyseal scar should be observed.




Figure 1. Ossification stages of the proximal humeral epiphysis; T1weighted turbo spin echo (T1 TSE) sequences in coronal oblique orientation. The length of the ossified epiphysis is over two thirds compared to the width of the metaphyseal ending (2c). Epiphyseal-metaphyseal fusion completes one third or less of the former gap between epiphysis and metaphysis. Arrow shows fused epiphyseal-metaphyseal part (3a). Epiphyseal-metaphyseal fusion completes between one third and two thirds of the former gap between epiphysis and metaphysis. Arrows show fused epiphyseal-metaphyseal part (3b). Epiphyseal-metaphyseal fusion completes over two thirds of the former gap between epiphysis and metaphysis. Arrow shows unfused epiphyseal-metaphyseal part (3c). The epiphyseal cartilage is fully ossified, and the epiphyseal scar is visible. Arrows show epiphyseal scar (4).




Figure 2. Schematic drawings of the stages of the proximal humeral epiphysis

Statistical Analysis

SPSS ver. 17 software (SPSS, Chicago, IL, USA) was used for the statistical analysis. Data were expressed as the mean or median with standard deviation (SDs), 95% confidence intervals, and ranges, as appropriate. Associations between age and ossification stage were evaluated via Spearman's

correlation analysis. Between-sex comparisons were performed using the Mann–Whitney U-test. A p -value < 0.05 was considered to be significant.

Two radiologists (R1 and R2) evaluated the MR images. R1 has 20 years of professional experience, while R2 has 6 years of experience. In addition, R1 has experience in forensic age estimation of living individuals via CT and MRI by measuring the ossification stage of the different epiphyseal areas. Cohen's kappa nonparametric test was used to evaluate intra- and inter-observer variabilities, and the κ statistic was calculated. The Altman [42] system was used to interpret the κ values: $\kappa < 0.20$, poor agreement; $\kappa = 0.21$ – 0.40 , fair agreement; $\kappa = 0.41$ – 0.60 , moderate agreement; $\kappa = 0.61$ – 0.80 , good agreement; and $\kappa = 0.81$ – 1.00 , very good agreement.

Results

We examined the left proximal humeral epiphysis of 428 patients aged 12–30 years (mean male age, 22.07 ± 4.37 years; mean female age, 22.10 ± 4.75 years; 240 males; 188 females; Table 1). Determination of the ossification stage of the proximal humeral epiphysis was possible in our sample. Figure 2 shows the MRI findings for ossification stages 2c, 3a, 3b, 3c and 4 that were observed. The remaining ossification stages were not found within the study population. Spearman's rank correlation analysis revealed a significant positive relationship between age and ossification stage of the proximal humeral epiphysis (all subjects: $\rho=0.664$, $p < 0.001$; males: 0.631 , $p < 0.001$; females: $\rho=0.651$, $p < 0.001$).

The intra- and inter-observer reliability for the proximal humeral epiphysis was $\kappa=0.898$ and $\kappa=0.828$, respectively. These values indicate good agreement.

Table 2 shows the minimum and maximum ages, mean \pm standard deviation and lower and upper quartiles of all parameters. No significant sex-related differences were found for all stages ($P>0.5$). The mean age at which stage 2c was seen was 13.44 ± 0.94 years in male and 12.95 ± 0.82 years in female patients. Stage 2c was first seen at 12.1 years of age among male patients and at 12.2 years among female patients. In male patients, the latest appearance of stages 2c was 15.3. In female patients, the latest appearance of stages 2c was 14.4 years. The mean age at which stage 4 was seen was 24.76 ± 3.47 years in male and 24.50 ± 3.22 years in female patients. Stage 4 was first seen at 18.6 years of age among male patients and at 17.2 years among female patients. In males, the latest appearance of stage 4 was 30.6, while in females, the latest appearance of stage 4 was 30.7 years.

Discussion

In recently published research, studies evaluating the development of the proximal humerus epiphysis are mostly based on direct (visual) or x-ray examination of skeletal collections (Table 3). In an anthropological study of proximal humeral epiphyses in the Lisbon skeletal collection, Cardoso [43] reported a complete union at the age of 17 years in both sexes. Another study on a Portuguese sample from Coimbra [44] noted that the minimum age at which

the epiphyseal union completes is two years later (at the age of 19) in males. In addition, compared to the Lisbon sample, the oldest individuals with open epiphyses were observed in the Coimbra sample and were 3 years older in both sexes. Both aforementioned studies were conducted in skeletal material using direct examination, the same staging system (3 stages), and similar size samples, which both derive from individuals who lived around the same period in Portugal (Table 3). Nonetheless, compared to the Lisbon study, the Coimbra study reports delayed development of approximately 3 years, which may reflect nutritional stress in the Coimbra sample.

Two more studies of the humeral epiphysis were conducted using direct analysis on skeletal remains from Bosnian and American soldiers [45-46]. For the two Bosnian samples, the minimum age of epiphyseal closure is 18, and the maximum age for which open epiphysis is described is 20 and 21 years, respectively. This finding is more consistent with the Cardoso study results [43]. Conversely, the maximum age for which open epiphysis in the McKern skeletal collection (derived from US soldiers) was 23 years, which indicates a two-year delay in skeletal maturation. These differences can be attributed to any number factors such as population differences or, possibly, different levels and durations of nutritional stress for the US soldiers compared to the Bosnians.

The most important study of the humeral epiphysis using x-rays was done on a large sample of US living patients [47]. This study reported complete union at the ages of 14 and 16 for males and females, respectively. This result is in agreement with two studies on Indian patients which, however, were conducted

using undefined protocols [48] or were limited in female patients [49]. The maximum age for which open epiphysis is noted is 21 for males and 20 for females. Interestingly, compared to the anthropological studies, all x-ray studies detected earlier ages of complete fusion (Table 3).

MRI studies of the proximal humerus are limited in the work of Kwong et al. [39] in a sample of 76 US patients. The authors attempted an overview of the humeral growth from 3 months to 17 years and focused on the detailed description of the MRI characteristics from the appearance of the humeral head secondary ossification centers to growth plate transformation, metaphyseal stripe and proximal metaphyseal marrow signal intensity. MRI examinations were performed on different MRI weighting and cross-sections. According to the results, the growth plate remained open until 14 years of age, showed partial closure in the 14-16 years age group (19 cases), and complete closure by 17 years (8 cases). The study had a limited sample with an upper limit at the age of 17 years old and no specifically defined staging system, and sex differences were not investigated. In addition, as stated by the authors, only 9/76 cases were healthy subjects, which could have seriously biased the results.

The current study used MRI, which is the imaging method of choice for children, to explore the developmental changes of the proximal epiphysis of the humerus in a large sample (N=428) balanced for both sexes, ranging in age from 12 to 30 years old. The study focused on the subtle differences in the epiphyseal line during development in an effort to capture the majority of developmental variation in the manifestation of skeletal age. We applied a combination of the

Schmeling [18] and Kellinghaus [19] staging system, and we found that the minimum age for closure of the epiphyseal union of the proximal humerus is 17.2 for females and 18.6 for males. These results agree with the standards presented for Bosnian males [46-47] but largely disagree with the standards for Americans [47]. These discrepancies can be attributed to differences in ethnicity, nutritional stress, dietary habits, environmental influences, secular changes and the modality used for data acquisition. The intra- and inter-observer reliability was $\kappa=0.898$ and $\kappa=0.828$, respectively, which is in agreement with other MRI studies in different anatomical regions (e.g., 33). Thus, discrepancies with other studies in the literature are unlikely to be due to human error.

This unequal distribution should be kept in mind as a substantive limitation in the evaluation of our work data. The fact that 345 out of 428 (ca. 80.6%) of our cases were above 18 years of age, and these cases only represented 2/3 (66%) of the total number of case, indicates potential selection bias. This study limitation is especially relevant, since the chance of documenting the lower extremes of a stage, i.e., the minimum age, is lowered. The unequal age distribution is a result of the availability of clinical images, as in any retrospective study of age estimation. In this respect, the results obtained with a balanced age distribution in future prospective studies will be important for demonstrating the applicability of the method.

It is important to define the sex differences for researching age-estimation studies. In our study, no significant difference was detected between the two sexes. At stage 4, however, the difference between the minimum ages (1.4 years) is remarkable. Unfortunately, there is no other proximal humerus MRI study to compare sex differences. Conversely, there are many different estimations of CT and MRI in different populations. In these studies, the differences between the sexes are variable, reaching 2 years for the stages in which the ossification is completed [15-17, 28, 30]. Thus, it is imperative to investigate other factors, such as the effect of socioeconomic status in the maturation of the humerus. Future studies may also reveal new data in discussing this difference.

The fact that stage 5 was not observed in our sample suggests that the minimum age for that stage is over the 30th year of life. Thus, stage 4 represents the final ossification stage for part of the population. There is a need for future studies in older age groups to determine the age at which the epiphyseal scar has completely disappeared, but this research would exceed the purposes of the current work, the focus of which is on the age estimation of living subjects to legal age thresholds.

Although the study was performed on one sectional plane and one MRI weighting, it was of paramount importance for recording the development of the proximal humeral epiphysis. The most prominent example of such limitation was observed in studies with different sectional planes and MRI weighting for the proximal tibial epiphysis, and the distal femoral epiphysis, creating important debates about the development of the methods [17, 24, 28, 30]. Assessing the

ossification of the epiphysis at different sectional planes may be important for the reliability of the results. In their original study, Dedouit et al. [17] reported that in addition to the coronal sections, sagittal sections were also evaluated in stages 2-3 and stages 3-4, but this practice was rarely encouraged. On the other hand, the axial and coronal sections of the clavicle CT study were used together for the same purpose in the staging, and Scharte et al. [50] reported a 35.6% difference between axial and coronal planes with clavicle CT. In our study, only coronal sections were used instead of a combination of coronal and sagittal sections. A follow-up study comparing the scores taken on both sectional planes for the proximal humeral epiphysis would enable us to establish a solid methodology for observation of the epiphyseal union using MRI modalities.

Another factor to take under consideration is that the sample derives from a single geographical area. Although this sample is the largest sample of MRI data from the proximal humeral epiphysis, one should stress that skeletal development is subject to many diverse influencing factors, which cannot be controlled in the current study and may have introduced bias to the estimates. For instance, the effects of socioeconomic status on the results should be acknowledged [45-46, 51, 52]. In fact, Schmeling and colleagues [50] suggested that socioeconomic status is a decisive factor for delays in skeletal maturation in contrast to ethnicity, which seems to have no effect. This issue is particularly relevant in all cases that require the assessment of skeletal age, such as for establishing legal responsibility. Franklin et al., [53] correctly stressed that “*the populations most in need of reliable age assessment standards (e.g., comprising*

individuals from localized geographic regions that are more likely to enter the criminal justice system – such as people smugglers) are disproportionately underrepresented in the published literature...” This property may be true for the present study, as well; thus, there is scope for further expanding the sample to include more socioeconomic and geographical groups.

In conclusion, this study focused on proximal humeral epiphysis using a non-ionizing MRI method in living individuals. We encountered difficulties comparing our data with previous research, as notably few studies have been conducted and different methods and populations have been used. Our data show that the MRI method can be very useful, as a supportive method for estimating skeletal age, as it provides a better assessment of bone and cartilage tissue due to its technical advantages and eliminates radiation exposure for living individuals. MRI of the proximal humeral epiphysis can be considered advantageous for skeletal age assessment of normal living subjects in a variety of situations ranging from illegal immigrants/asylum seekers, minor criminals to illegal participants in competitive sports, although it must be stressed that the age thresholds provided in this study are specific to a certain geographic region and should not be applied in a different population without further testing. Future directions of this work will include a prospective study on subjects of known socioeconomic status, nutritional habits and shoulder-related activities to fully exploit the investigative potential of MRI in the study of skeletal development and maturation.

Compliance with ethical standards:

Disclosure of potential conflicts of interest: The authors declare that they have no conflict of interest.

Research involving Human Participants and/or Animals: This article does not contain any studies with animals performed by any of the authors.

Informed consent : For this type of study formal consent is not required

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

References

- [1] Gilsanz V, Ratib O (2012) Hand Bone Age. A Digital Atlas of Skeletal Maturity. Berlin: Springer.
- [2] Pludowski P, Lebedowski M, Lorenc RS (2005) Evaluation of practical use of bone age assessments based on DXA-derived hand scans in diagnosis of skeletal status in healthy and diseased children. J Clin Densitom 8(1):48-56.
- [3] Ilich JZ, Hangartner TN, Skugor M, Roche AF, Goel PK, Matkovic V (1996) Skeletal age as a determinant of bone mass in preadolescent females. Skeletal radiology 25(5):431-9.

- [4] Morris EB, Shelso J, Smeltzer MP, Thomas NA, Karimova EJ, Li CS et al (2008) The use of bone age for bone mineral density interpretation in a cohort of pediatric brain tumor patients. *Pediatric radiology* 38(12):1285-92.
- [5] Ritz-Timme S, Cattaneo C, Collins MJ, Waite ER, Schutz HW, Kaatsch HJ et al (2000) Age estimation: the state of the art in relation to the specific demands of forensic practise. *Int J Legal Med* 113(3):129-36.
- [6] Schmeling A, Grundmann C, Fuhrmann A, Kaatsch HJ, Knell B, Ramsthaler F et al (2008) Criteria for age estimation in living individuals. *Int J Legal Med* 122(6):457-60.
- [7] Cunha E, Baccino E, Martrille L, Ramsthaler F, Prieto J, Schuliar Y et al (2009) The problem of aging human remains and living individuals: a review. *Forensic science international* 193(1-3):1-13.
- [8] Schmeling A, Olze A, Reisinger W, Geserick G (2004) Forensic age diagnostics of living people undergoing criminal proceedings. *Forensic science international* 144(2-3):243-5. doi:10.1016/j.forsciint.2004.04.059.
- [9] Janes L (2008) Criminal liability of minors and severity of penalties: European trends and developments. Howard League for Penal Reform http://www.europeanrights.eu/public/commenti/LauraJanes_en.pdf
Accessed: 21 Nov 2016

- [10] Jones VF (2012) Committee on Early Childhood A, Dependent C. Comprehensive health evaluation of the newly adopted child. *Paediatrics* 129(1):214-23.
- [11] Malina RM (2011) Skeletal Age and Age Verification in Youth Sport. *Sports Medicine* 41(11):925-47
- [12] Morishima A, Grumbach MM, Simpson ER, Fisher C, Qin K (1995) Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. *The Journal of clinical endocrinology and metabolism* 80(12):3689-98.
- [13] Grumbach MM (2000) Estrogen, Bone, Growth and Sex: A Sea Change in Conventional Wisdom. *Journal of Pediatric Endocrinology & Metabolism* 13:1439-55
- [14] Vogler JB, 3rd, Murphy WA (1988) Bone marrow imaging. *Radiology* 168(3):679-93.
- [15] Saint-Martin P, Rerolle C, Dedouit F, Bouilleau L, Rousseau H, Rouge D et al (2013) Age estimation by magnetic resonance imaging of the distal tibial epiphysis and the calcaneum. *Int J Legal Med* 127(5):1023-30.
- [16] Ekizoglu O, Hocaoglu E, Can IO, Inci E, Aksoy S, Bilgili MG (2015) Magnetic resonance imaging of distal tibia and calcaneus for forensic age estimation in living individuals. *Int J Legal Med* 129(4):825-31.

- [17] Dedouit F, Auriol J, Rousseau H, Rouge D, Crubezy E, Telmon N (2012) Age assessment by magnetic resonance imaging of the knee: a preliminary study. *Forensic science international* 217(1-3):232
- [18] Schmeling A, Schulz R, Reisinger W, Muhler M, Wernecke KD, Geserick G (2004) Studies on the time frame for ossification of the medial clavicular epiphyseal cartilage in conventional radiography. *Int J Legal Med* 118(1):5-8.
- [19] Kellinghaus M, Schulz R, Vieth V, Schmidt S, Pfeiffer H, Schmeling A (2010) Enhanced possibilities to make statements on the ossification status of the medial clavicular epiphysis using an amplified staging scheme in evaluating thin-slice CT scans. *Int J Legal Med* 124(4):321-5.
- [20] European Commission. Radiation protection. Medico-legal exposures, exposures with ionising radiation without medical indication. Proceedings of the International Symposium. 2002.

http://ec.europa.eu/energy/nuclear/radiation_protection/doc/publication/130.pdf. Accessed 4 April 2014.
- [21] Schmidt S, Schiborr M, Pfeiffer H, Schmeling A, Schulz R (2013) Age dependence of epiphyseal ossification of the distal radius in ultrasound diagnostics. *Int J Legal Med* 127(4):831-8.
- [22] Schmidt S, Schiborr M, Pfeiffer H, Schmeling A, Schulz R (2013) Sonographic examination of the apophysis of the iliac crest for forensic age

estimation in living persons. *Science & justice : journal of the Forensic Science Society* 53(4):395-401.

- [23] Schulz R, Schiborr M, Pfeiffer H, Schmidt S, Schmeling A (2013) Sonographic assessment of the ossification of the medial clavicular epiphysis in 616 individuals. *Forensic science, medicine, and pathology* 9(3):351-7.
- [24] Saint-Martin P, Rerolle C, Pucheux J, Dedouit F, Telmon N (2015) Contribution of distal femur MRI to the determination of the 18-year limit in forensic age estimation. *Int J Legal Med* 129(3):619-20.
- [25] Tangmose S, Jensen KE, Villa C, Lynnerup N (2014) Forensic age estimation from the clavicle using 1.0T MRI--preliminary results. *Forensic science international* 234:7-12.
- [26] Hillewig E, Degroote J, Van der Paelt T, Visscher A, Vandemaele P, Lutin B et al (2013) Magnetic resonance imaging of the sternal extremity of the clavicle in forensic age estimation: towards more sound age estimates. *Int J Legal Med* 127(3):677-89.
- [27] Vieth V, Schulz R, Brinkmeier P, Dvorak J, Schmeling A (2014) Age estimation in U-20 football players using 3.0 tesla MRI of the clavicle. *Forensic science international* 241:118-22.
- [28] Krämer JA, Schmidt S, Jürgens KU, Lentschig M, Schmeling A, Vieth V (2014) Forensic age estimation in living individuals using 3.0 T

MRI of the distal femur. *Int J Legal Med* 128(3):509-14. doi: 10.1007/s00414-014-0967-3.

- [29] Schmidt S, Muhler M, Schmeling A, Reisinger W, Schulz R (2007) Magnetic resonance imaging of the clavicular ossification. *Int J Legal Med* 121(4):321-4.
- [30] Ekizoglu O, Hocaoglu E, Inci E, Can IO, Aksoy S, Kazimoglu C (2016) Forensic age estimation via 3-T magnetic resonance imaging of ossification of the proximal tibial and distal femoral epiphyses: Use of a T2-weighted fast spin-echo technique. *Forensic science international* 260:102-7.
- [31] Wittschieber D, Vieth V, Timme M, Dvorak J, Schmeling A (2014) Magnetic resonance imaging of the iliac crest: age estimation in under-20 soccer players. *Forensic science, medicine, and pathology* 10(2):198-202.
- [32] Dvorak J, George J, Junge A, Hodler J (2007) Age determination by magnetic resonance imaging of the wrist in adolescent male football players. *British journal of sports medicine* 41(1):45-52.
- [33] George J, Nagendran J, Azmi K (2012) Comparison study of growth plate fusion using MRI versus plain radiographs as used in age determination for exclusion of overaged football players. *British journal of sports medicine* 46(4):273-8.

- [34] Terada Y, Kono S, Tamada D, Uchiumi T, Kose K, Miyagi R et al (2013) Skeletal age assessment in children using an open compact MRI system. *Magnetic resonance in medicine* 69(6):1697-702.
- [35] Tomei E, Battisti S, Martino M, Nissman D, Semelka RC (2014) Text-atlas of skeletal age determination: MRI of the hand and wrist in children. Hoboken: Wiley-Blackwell
- [36] Tomei E, Sartori A, Nissman D, Al Ansari N, Battisti S, Rubini A et al (2014) Value of MRI of the hand and the wrist in evaluation of bone age: preliminary results. *Journal of magnetic resonance imaging: JMRI* 39(5):1198-205.
- [37] Serinelli S, Panebianco V, Martino M, Battisti S, Rodacki K, Marinelli E et al (2015) Accuracy of MRI skeletal age estimation for subjects 12-19. Potential use for subjects of unknown age. *Int J Legal Med* 129(3):609-17.
- [38] Schmidt S, Vieth V, Timme M, Dvorak J, Schmeling A (2015) Examination of ossification of the distal radial epiphysis using magnetic resonance imaging. New insights for age estimation in young footballers in FIFA tournaments. *Science & justice: journal of the Forensic Science Society* 55(2):139-44.
- [39] Kwong S, Kothary S, Poncinelli LL (2014) Skeletal development of the proximal humerus in the pediatric population: MRI features. *AJR American journal of roentgenology* 202(2):418-25.

- [40] Zember JS, Rosenberg ZS, Kwong S, Kothary SP, Bedoya MA (2015) Normal Skeletal Maturation and Imaging Pitfalls in the Pediatric Shoulder. Radiographics: a review publication of the Radiological Society of North America, Inc 35(4):1108-22.

- [41] FIFA to introduce MRI screening at Nigeria 2009 to combat the fielding of over-age players. Federation Internationale de Football Association 2009.
<http://www.fifa.com/u17worldcup/news/y=2009/m=9/news=fifa-introduce-mri-screening-nigeria-2009-combat-the-fielding-over-age-1096817.html>.
 Accessed 3 Jan 2017

- [42] Altman DG (1991) Practical statistics for medical research. New York: Chapman & Hall

- [43] Cardoso HF (2008) Age estimation of adolescent and young adult male and female skeletons II, epiphyseal union at the upper limb and scapular girdle in a modern Portuguese skeletal sample. American journal of physical anthropology 137(1):97-105.

- [44] Coqueugniot H, Weaver TD (2007) Brief communication: infracranial maturation in the skeletal collection from Coimbra, Portugal: new aging standards for epiphyseal union. American journal of physical anthropology 134(3):424-37.

- [45] Schaefer MC (2008) A Summary of Epiphyseal Union Timings in Bosnian Males. Int J Osteoarchaeol 18(5):536-45.

- [46] Schaefer MC, Black SM (2005) Comparison of ages of epiphyseal union in north American and Bosnian skeletal material. *Journal of forensic sciences* 50(4):777-84.
- [47] Schaefer M, Aben G, Vogelsberg C (2015) A demonstration of appearance and union times of three shoulder ossification centers in adolescent and post-adolescent children. *J Forensic Radiology and Imaging* 3:49-56.
- [48] Jit I, Singh B (1971) A radiological study of the time of fusion of certain epiphyses in Punjabis. *J Anat Soc India* 20:1-27.
- [49] Tirpude B, Surwade V, Murkey P, Wankhade P, Meena S (2014) Age determination from epiphyseal union of bones at shoulder joint in girls of central India. *Journal of Forensic Medicine, Science and Law* 23(1)
- [50] Scharte P, Vieth V, Schulz R, Ramsthaler F, Püschel K, Bajanowski T, Pfeiffer H, Schmeling A, Schmidt S, Wittschieber D (2017) Comparison of imaging planes during CT-based evaluation of clavicular ossification: a multi-center study. *Int J Legal Med* doi: 10.1007/s00414-017-1615-5.
- [51] Schmeling A, Reisinger W, Loreck D, Vendura K, Markus W, Geserick G (2000) Effects of ethnicity on skeletal maturation: consequences for forensic age estimations. *Int J Legal Med* 113:253-258
- [52] Schmeling A, Schulz R, Danner B, Rosing FW (2006) The impact of economic progress and modernization in medicine on the ossification of hand and wrist. *Int J Legal Med* 120:121-126

- [53] Franklin D, Flavel A, Noble J, Swift L, Karkhanis S (2015) Forensic age estimation in living individuals: methodological considerations in the context of medico-legal practice. *Research and Reports in Forensic Medical Science* 5:53-66.

List of Tables

Table 1. Age distribution of male and female subjects.

| Age (years) | Male (N) | Female (N) |
|------------------------|---------------------|-----------------------|
| 12 | 2 | 6 |
| 13 | 5 | - |
| 14 | 6 | 12 |
| 15 | 10 | 2 |
| 16 | 12 | 13 |
| 17 | 8 | 7 |
| 18 | 12 | 11 |
| 19 | 16 | 8 |
| 20 | 25 | 13 |
| 21 | 25 | 24 |
| 22 | 18 | 11 |
| 23 | 24 | 14 |
| 24 | 16 | 9 |
| 25 | 17 | 16 |
| 26 | 9 | 9 |
| 27 | 4 | 7 |
| 28 | 10 | 8 |
| 29 | 13 | 9 |
| 30 | 8 | 9 |
| Total | 240 | 188 |

Table 2. Minimum and maximum ages, with means \pm SDs, lower and upper quartiles and medians, at all stages of proximal humeral epiphysis.

| Stage | Sex | N | Mean \pm SD | Min-Max | LQ; UQ;Median |
|-------|--------|-----|------------------|-----------|---------------------|
| 2c | Female | 8 | 12.95 \pm 0.82 | 12.2-14.4 | 12.37; 13.75; 12.65 |
| | Male | 10 | 13.44 \pm 0.94 | 12.1-15.3 | 12.92; 14.22; 13.25 |
| 3a | Female | 12 | 14.87 \pm 0.78 | 14.3-16.7 | 14.30; 14.90; 14.65 |
| | Male | 10 | 14.97 \pm 0.60 | 14.2-15.9 | 14.30; 15.37; 15.20 |
| 3b | Female | 8 | 16.58 \pm 1.15 | 15.2-18.4 | 15.72; 17.82; 16.25 |
| | Male | 12 | 16.56 \pm 1.02 | 15.3-18.2 | 15.80; 17.75; 16.35 |
| 3c | Female | 43 | 19.62 \pm 2.03 | 16.2-23.2 | 18.20; 21.30; 20.30 |
| | Male | 61 | 19.84 \pm 2.09 | 16.2-23.6 | 17.90; 21.30; 20.10 |
| 4 | Female | 117 | 24.76 \pm 3.47 | 17.2-30.7 | 22.05; 27.75; 25.10 |
| | Male | 147 | 24.50 \pm 3.22 | 18.6-30.6 | 21.80; 26.90; 24.90 |

Table 3. Comparative table of studies on the closure of the proximal humeral epiphysis.

| Population | Life span | Male | Female | Age range | Study protocol | Youngest fused | Oldest with open epiphysis | Reference |
|-----------------------------------------------------------------|----------------------------------------|------|--------|-------------------|------------------------------------|-----------------------------|----------------------------|---------------|
| Lisbon skeletal collection | 1887-1975 | 65 | 56 | 9-29 years | Direct analysis. Staging: 3 stages | Male 17 y, Female: 17 y | Male: 21 y Female:19 y | [43] |
| Coimbra skeletal collection | 1826-1938 | 69 | 68 | 7-29 years | Direct analysis. Staging: 3 stages | Male 19 y, Female: 17y | Male:24 y Female: 22y | [44] |
| Bosnian skeletal collection | 1962-1995 | 232 | 0 | 14-30 years | Direct analysis. Staging: 5 stages | Male: 18 y | Male: 21 y | [45] |
| Bosnian skeletal collection | 1965-1995 | 114 | 0 | 17-30 years | Direct analysis. Staging: 4 stages | Male:18 y | Male: 20 y | [46] |
| Mckern skeletal collection (US soldiers died in the Korean War) | 1924-1954 | 325 | 0 | 17-30 years | Direct analysis. Staging: 4 stages | Male: 17 y | Male: 23 y | [46] |
| USA-Living individuals/students | Examinations before 2015 | 258 | 179 | 10-21 years | X-ray staging: 4 stages | Male: 16 y, Female:14 y | Male:21 y Female:20 y | [47] |
| Indian-Living individuals | 1951-1971 | ? | ? | 14-19 years | X-Ray, staging undefined | Male:16 y, Female:15 y | Male 18 y, Female:17 y | [48] |
| Indian-Living individuals | Examinations before 2014 | 0 | 80 | 13-21 yeas | X-ray staging: 3 stages | Female: 15-16 y | Female:17-18 y | [49] |
| USA-Living individuals | Examinations between 2003-2012 | 44 | 32 | 3months -17 years | MRI: staging undefined | 15.5 y (sex not defined) | 17 y (sex not defined) | [39] |
| Turkish-Living individuals | Examinations from Jan 2014 to Jan 2016 | 240 | 188 | 12-31 years | MRI Staging: 5 stages | Male: 18.6 y, Female:17.2 y | Male:23.6 y, Female:23.2y | Present study |