



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Frightening and Traumatic Memories Early After Intensive Care Discharge

Citation for published version:

DESIST Investigators 2018, 'Frightening and Traumatic Memories Early After Intensive Care Discharge', *American Journal of Respiratory and Critical Care Medicine*. <https://doi.org/10.1164/rccm.201804-0699LE>

Digital Object Identifier (DOI):

[10.1164/rccm.201804-0699LE](https://doi.org/10.1164/rccm.201804-0699LE)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

American Journal of Respiratory and Critical Care Medicine

Publisher Rights Statement:

This is the author's peer-reviewed manuscript as accepted for publication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Frightening and Traumatic Memories Early After Intensive Care Discharge

Sarah Train¹, Kalliopi Kydonaki², Janice Rattray³, Jacqueline Stephen^{4,5}, Christopher J Weir^{4,5}, Timothy S Walsh^{1,5,6}; on behalf of the DESIST Investigators

¹ Anaesthetics, Critical Care and Pain Medicine, University of Edinburgh, Edinburgh, Scotland

²Edinburgh Napier University, Edinburgh, Scotland

³School of Nursing and Health Sciences, University of Dundee, Dundee, Scotland

⁴Edinburgh Clinical Trials Unit, University of Edinburgh, Scotland

⁵Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, Scotland

⁶Centre for Inflammation Research, University of Edinburgh, Edinburgh, Scotland

Corresponding Author

Professor Tim Walsh

Department of Anaesthesia, Critical Care & Pain Medicine Room S8208, 2nd Floor Royal Infirmary of Edinburgh, 51 Little France Crescent, Edinburgh EH16 4SA, Scotland

twalsh@staffmail.ed.ac.uk

0131 242 6395

Author Contributions

TSW, CK, and CJW conceived the DESIST study, secured funding, and managed the research. All authors conceived the current sub-study, contributed to analysis, wrote the manuscript, and approved the final version. All authors are accountable for the accuracy and integrity of the work.

Grants

The DESIST trial was supported by grants from the Chief Scientists Office, Scotland; and GE Healthcare (Unrestricted funding). ST was supported by a John Snow Anaesthesia Intercalated

bursary award for BMedSci students (Royal College of Anaesthetists/British Journal of Anaesthesia). CJW was supported in this work by NHS Lothian via the Edinburgh Clinical Trials Unit.

Running title: Early frightening and traumatic memories in ICU survivors

Descriptor: 4.06 ICU Management/Outcome

Word count: 948

References: 12

Anxiety, depression, and post-traumatic stress are prevalent among intensive care (ICU) survivors, can persist long-term, and are associated with low quality of life¹. Risk factors include frightening and delusional memories of ICU stay, especially for long-term post-traumatic stress^{2,3}. Frightening memories could relate to ICU stress, for example from pain, anxiety, and dreams. Delusional memories may originate from hallucinations and periods of delirium. These may all be influenced by sedation, analgesia and delirium management⁴. Few studies have described the nature and prevalence of frightening and traumatic memories early after ICU discharge. This transition of care is known to be stressful to both patients and family members, and is the time patients are starting to adjust to critical illness sequelae. Screening and intervention for psychological morbidity at this time might reduce long-term problems, especially if the patients at highest risk could be identified.

In a pre-planned sub-study within a published cluster randomised quality-improvement trial of sedation interventions in eight Scottish ICUs (the 'DESIST' trial^{5,6}), we administered two validated questionnaires to ICU survivors during their recovery on general wards early after ICU discharge. These were the Impact of Events Scale-Revised (IES-R) which measures traumatic symptomatology⁷, and the Intensive Care Experience Questionnaire (ICE-Q)⁸. The ICE-Q includes 31 items measuring patients' perceptions of their intensive care experience, grouped into four domains including a 'frightening experiences sub-score' (FESS; comprising six questions)⁸. The FESS demonstrates a Cronbach $\alpha > 0.7$ in previous studies, indicating validity, and previous studies have also found that ICE-Q scores could predict subsequent psychological morbidity^{8,9,10}.

Participants in the DESIST trial required mechanical ventilation (MV) with anticipated duration > 48 hours. When possible, surviving participants were approached after ICU discharge to complete the questionnaires and asked to anchor their responses to ICU memories. All patients provided written consent and the study was approved by the Scottish A Research Ethics Committee. These data were exploratory secondary outcomes, and were not reported in the main trial, but some results have been previously reported in the form of an abstract.¹¹ We report here a cohort analysis of all survivors who completed questionnaires.

We aimed to describe the frequency and nature of frightening and traumatic memories early after ICU discharge, while patients were still in hospital. For the ICE-Q questionnaire, we reported responses to each of the six questions comprising the FESS sub-score; the total FESS sub-score; and the proportion of patients with a FESS > 18 (a cut-off representing an average score > 3 ('neutral' response) across

contributing questions). For the IES-R we reported responses to all 22 questions; the IES-R total score; and the proportion of patients with a score >35 (indicating significant traumatic symptomatology⁷).

We also used responses to a separate ICE-Q question: 'I have no memory of intensive care' (five potential responses ranging from strongly agree to strongly disagree) to explore relationships between patients' overall memory of ICU and the FESS sub-score and IES-R score.

There were 1291 ICU survivors in the trial. The ICE-Q and IES-R were completed by 517 survivors (40%). Survivors who completed questionnaires had longer post-ICU hospital stays, and included lower proportions of general medical and higher proportions of general surgical patients than those who did not complete questionnaires (see table 1). There was also some imbalance in admission source. The median number of days to questionnaire completion was 7 days (IQR: 4-14 days) post ICU discharge. Responses to the individual questions are shown in figure 1. The median (IQR) sum score for the ICE-Q FESS was 15 (11-21), and for IES-R was 19 (7-36). There were 34.4% (95% Confidence Interval: 30.3 to 38.5%) of the patients with a FESS scores >18 and 25.1% (95% CI: 21.4 to 28.8%) with an IES-R score >35 . Patient responses to the two questionnaires correlated strongly indicating that patients who had more frightening memories also reported greater trauma symptomatology (Spearman Rho coefficient =0.62; $P<0.001$). Post-ICU memories were diverse: among the ICE-Q responses fear, helplessness, and anticipating death were frequent. Among the IES-R responses memories of numbness, unreality and sleep disturbance were most common.

In relation to patients' overall memory of ICU, we found those with less overall memory had fewer frightening memories (correlation with FESS scores: $\rho = -0.18$, $P<0.001$) and traumatic memories (correlation with IES-R scores: $\rho = -0.13$, $P=0.004$), although both associations were very weak.

To our knowledge, this is the largest reported study of frightening and traumatic memories in the early post-ICU discharge period. Our data suggest these occur in 25 to 35% of patients. Although our cohort was large, and from a general ICU population, both inclusion and response bias could have influenced these estimates. Patients who required no or only brief MV were excluded from the DESIST trial, and data were available for only 40% of participants. Missed responses were mainly due to limited trial resource; we found many patients were already discharged from hospital when ward-based follow-up occurred and later contact was not feasible. Missing cases were more likely to have general medical diagnoses, had shorter hospital stays, and were more often admitted directly from the emergency department. These and other unmeasured confounders could have biased the prevalence estimates.

However, our data are consistent with smaller studies describing ‘delusional’ memories early after ICU discharge and previous studies using the ICE-Q^{3,8-10}. Although delusion-like memories are known predictors of long-term psychological morbidity², controversy exists regarding whether factual memories, even if frightening, are protective^{3,12}. Our data illustrate the diversity, complexity and personalised nature of early post-ICU memories.

In conclusion, our data demonstrate prevalent early traumatic and frightening memories that might contribute to long-term psychological morbidity after ICU. This supports a hypothesis that systematic screening at this time may be useful as part of a strategy for personalised targeted care and interventions to reduce later psychological morbidity.

Table 1. Demographic data of participants who completed or did not complete the questionnaire follow-up. *P* value denotes comparison using Mann-Whitney U test for age, and duration variables; independent sample t test for APACHE II; Chi² for gender and case mix (medical or surgical). Unless stated, the variable contains no missing values. APACHE II: Acute Physiology and Chronic Health Evaluation II.

Characteristic		ICE-Q and IES-R	ICE-Q and IES-R	<i>p</i> value
		Completed	Not Completed	
		Valid (missing)	772 (2)	
Age, years	Median (IQR)	60.0 (48.5-69.0)	61.0 (47.0-71.0)	0.46
Male	n (%)	305/517 (59.0)	463 (59.9)	0.77
APACHE II Score	Valid (missing)	491 (26)	750 (24)	0.17
	Mean (SD)	20.58 (6.93)	21.16 (7.49)	
Length of ICU Stay days	Median (IQR)	6.74 (3.66-12.67)	6.76 (3.70-11.86)	0.43
Duration of Ventilation days	Median (IQR)	3.44 (1.70-8.40)	4.0 (1.86-8.08)	0.25
Length of Hospital Stay days	Valid (missing)	516 (1)	771 (3)	<0.001
	Median (IQR)	27.0 (15.0-48.75)	19.0 (10.0-34.0)	
Case mix:				
Proportion medical	n (%)	274 (52.9)	482 (62.4)	<0.001
Medical Specialty				
General Medicine		147 (28.9)	299 (38.7)	
Cardiology		33 (6.4)	44 (5.7)	
Gastrointestinal		29 (5.6)	32 (4.1)	
Respiratory		26 (5.0)	35 (4.5)	
Ear/Nose/Throat		11 (2.1)	13 (1.7)	
Neurology		5 (1.0)	18 (2.3)	
Renal		6 (1.2)	13 (1.7)	
Other		17 (3.3)	28 (3.6)	

Surgical Specialty


General Surgery	156 (30.2)	188 (24.3)
Vascular	24 (4.6)	43 (5.6)
Transplant	30 (5.8)	12 (1.6)
Orthopaedics	19 (3.7)	27 (3.5)
Cardiac	5 (1.0)	5 (0.6)
Neurosurgery	0 (0)	4 (0.5)
Other	9 (1.7)	11 (1.4)

Source of Admission

P = 0.02

Operating Room	171 (33.0)	198 (25.6)
Emergency Department	119 (23.0)	241 (31.2)
High Dependency Unit	101 (19.5)	129 (16.7)
Other Ward in hospital	69 (13.3)	100 (12.9)
Other Hospital	23 (4.4)	41 (5.3)
Other ICU	16 (3.1)	18 (2.3)
Other Immediate Care Area	12 (2.3)	18 (2.3)
Other	6 (1.2)	17 (2.2)

Figure 1. Individual responses to the questions that comprise the Frightening Experiences Sum Score (FESS) of the Intensive Care Experience Questionnaire (ICE-Q) and Impact of Event Scale Revised (IES-R) responses. Scores Represent:
 ICE-Q: 1 never/strongly disagree; 2 rarely/disagree; 3 sometimes/neither; 4 frequently/agree; 5 all the time
 IES-R: 1 not at all; 2 a little bit; 3 moderately; 4 quite a bit; 5 extremely.



References:

1. Ringdal M, Plos K, Lundberg D, Johansson L, Bergbom I. Outcome after injury: memories, health-related quality of life, anxiety, and symptoms of depression after intensive care. *J Trauma*. 2009;66(4):1226–33.
2. Parker A, Sricharoenchai T, Raparla S, Schneck K, Bienvenu O, Needham D. Posttraumatic stress disorder in critical illness survivors: A metaanalysis. *Crit Care Med* 2015;43(5):1121–9.
3. Kiekkas P, Theodorakopoulou G, Spyrtatos F, Baltopoulos G. Psychological distress and delusional memories after critical care: a literature review. *Int Nurs Rev* 2010;57: 288-296.
4. Aitken LM, Castillo MI, Ullman A, Engstrom A, Cunningham K, Rattray J. What is the relationship between elements of ICU treatment and memories after discharge in adult ICU survivors?. *Aust Crit Care* 2016;29(1):5–15.
5. Walsh TS, Kydonaki K, Antonelli J, Stephen J, Lee RJ, Everingham K, et al. Staff education, regular sedation and analgesia quality feedback, and a sedation monitoring technology for improving sedation and analgesia quality for critically ill, mechanically ventilated patients: a cluster randomised trial. *Lancet Respir Med* 2016;4(10):807–17.
6. Walsh TS, Kydonaki K, Antonelli J, Stephen J, Lee RJ, Everingham K, et al. Rationale, design and methodology of a trial evaluating three strategies designed to improve sedation quality in intensive care units (DESIST study). *BMJ Open* 2016; 6(3).
<http://bmjopen.bmj.com/content/6/3/e010148.abstract>
7. Bienvenu OJ, Williams JB, Yang A, Hopkins RO, Needham DM. Posttraumatic stress disorder in survivors of acute lung injury: Evaluating the impact of event scale-revised. *Chest* 2013;144(1):24–31.
8. Rattray J, Johnston M, Wildsmith JAW. The intensive care experience: Development of the ICE questionnaire. *J Adv Nurs*. 2004;47(1):64–73.
9. Alasad JA, Tabar NA, Ahmad MM. Patients' experience of being in intensive care units. *J Crit Care* 2015; 30: 859.e7–859.e11
10. Rattray J, Crocker C, Jones M, Connaghan J. Patients' perceptions of and emotional outcome

- after intensive care: results from a multicentre study. *Nursing in Critical Care* 2010;15:86-93
11. Train S , Rattray J, Antonelli J, Stephen J, Weir C, Walsh TS. A retrospective cohort study investigating prevalence and nature of frightening memories in adult ICU survivors and their association with ICU exposures. *J Intensive Care Soc* 2018;19(Suppl):e161
 12. Jones C, Griffiths RD, Humphris G, Skirrow PM Memory, delusions, and the development of acute posttraumatic stress disorder related symptoms after intensive care. *Critical Care Med* 2001;29(3):573–580.