THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Can Antarctic lichens acclimatize to changes in temperature?

Citation for published version:

Colesie, C, Büdel, B, Hurry, V & Green, TGA 2018, 'Can Antarctic lichens acclimatize to changes in temperature?', *Global Change Biology*, vol. 24, no. 3, pp. 1123-1135. <https://doi.org/10.1111/gcb.13984>

Digital Object Identifier (DOI):

[10.1111/gcb.13984](https://doi.org/10.1111/gcb.13984)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Global Change Biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 Title: Can Antarctic lichens acclimatise to changes in temperature?

2 Running head: Temperature acclimation in Antarctic lichens

3 List of authors:

4 Claudia Colesie¹, Burkhard Büdel², Vaughan Hurry¹ and T. G. Allan Green^{3,4}

5 1: Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural
6 Sciences; Umeå, Sweden

7 2: Department of Plant Ecology and Systematics, University of Kaiserslautern;
8 Kaiserslautern, Germany

9 3: Departamento de Biología Vegetal II, Facultad de Farmacia, Universidad Complutense,
10 Madrid, Spain

11 4: Department of Biological Sciences, University of Waikato, Hamilton, New Zealand

12

13 Corresponding author: Claudia Colesie

14 Telephone: 0046 70 333 1806

15 Fax: 0046 18 67 20 00

16 Email: Claudia.colesie@gmail.com

17 Keywords:

18 Thermal acclimation, climate warming, lichen, Antarctica, net photosynthesis, biological soil
19 crusts, *Usnea aurantiaco-atra*

20 Primary research article

21 Abstract

22 The Antarctic Peninsula, a tundra biome dominated by lichens and bryophytes, is an ecozone
23 undergoing rapid temperature shifts. Such changes may demand a high physiological
24 plasticity of the local lichen species in order for them to maintain their role as key drivers in
25 this pristine habitat. This study examines the response of net photosynthesis and respiration to
26 increasing temperatures for three Antarctic lichen species with different ecological response
27 amplitudes. We hypothesise that negative effects caused by increased temperatures can be
28 mitigated by thermal acclimation of respiration and/or photosynthesis. The fully controlled
29 growth chamber experiment simulated intermediate and extreme temperature increases over
30 the time course of six weeks. Results showed that, in contrast to our hypothesis, none of the
31 species was able to downregulate temperature-driven respiratory losses through thermal
32 acclimation of respiration. Instead, severe effects on photobiont vitality demonstrated that
33 temperatures around 15°C mark the upper limit for the two species restricted to the Antarctic,
34 and when mycobiont demands exceeded the photobiont capacity they could not survive within
35 the lichen thallus. In contrast, the widespread lichen species was able to recover its
36 homoeostasis by rapidly increasing net photosynthesis. We conclude that in order to
37 understand the complete lichen response, acclimation processes of both symbionts, the photo-
38 and the mycobiont, have to be evaluated separately. As a result, we postulate that any
39 acclimation processes in lichen are species specific. This, together with the high degree of
40 response variability and sensitivity to temperature in different species that co-occur spatially
41 close, complicates any predictions regarding future community composition in the Antarctic.
42 Nevertheless, our results suggest that species with a broad ecological amplitude may be
43 favoured with ongoing changes in temperature.

44 Introduction

45 The rates at which organisms process carbon and nutrients via biogeochemical cycling,
 46 photosynthesis (P) and respiration (R) are temperature sensitive. Understanding the biological
 47 mechanisms that regulate carbon exchange rates, and their response to climate change, are
 48 among the most urgent scientific challenges for ecosystem ecologists in order to assess
 49 terrestrial carbon cycle–climate feedbacks (Bardgett, Freeman, & Ostle, 2008). Many
 50 ecophysiological studies have addressed the question of how the absolute rates, and the
 51 balance between P and R, will change in response to climate change (e.g. Karhu et al., 2014;
 52 Wang et al., 2014; Atkin et al., 2015; Drake et al., 2016). It is known that the increases in
 53 temperature will directly affect photosynthesis and respiration and, traditionally, simplified
 54 climate models have assumed that both will rise exponentially with short-term changes in
 55 temperature (King, Gunderson, Post, Weston, & Wullschleger, 2006), generating a positive
 56 climate-ecosystem carbon feedback (Davidson & Janssens, 2006) with the potential to
 57 accelerate climate warming by up to 1.4 times (Cox, Betts, Jones, Spall, & Totterdell, 2000).
 58 In recent years, studies on vascular plants are leading to a re-evaluation of the model
 59 assumptions because many plants show physiological, structural and biochemical adjustments
 60 that mitigate the effects of temperature increases (Körner, 2006; Vanderwel, et al., 2015).
 61 This effect is referred to as thermal acclimation (Oechel et al., 2000; Luo, Wan, Hui, &
 62 Wallace, 2001; Davidson & Janssens, 2006). Acclimation of R to colder growth temperatures
 63 results in increased respiratory CO₂ release measured at a chosen standard temperature
 64 (R_mT). Conversely, acclimation to high growth temperature results in lower R_mT (Atkin,
 65 Bruhn, Hurry, & Tjoelker, 2005). Understanding and characterising the ecophysiological
 66 response of major contributors to ecosystem respiration is highlighted as an essential research
 67 priority to help predict accurately how warming will affect carbon efflux across different
 68 ecosystems (Heinemeyer et al., 2012). There is sound documentation of thermal acclimation

69 for vascular plants (e.g. Reich, Tjoelker, Machado, & Oleksyn, 2006), their mycorrhizal
70 symbionts (Heinemeyer, Ineson, Ostle, & Fitter, 2006) and free-living ectomycorrhizal fungi
71 grown in agar (Malcolm, López-Gutiérrez, Koide, & Eissenstat, 2008). For heterotrophic soil
72 microbes the topic is controversial (Carey et al., 2016; Crowther & Bradford, 2013; Min,
73 Lehmeier, Ballantyne, & Billings, 2016).

74 While these better understood global carbon players (vascular plants and their mycorrhizal
75 symbionts) have a major role in wet, moist and temperate terrestrial biomes, 35% of the
76 Earth's land mass is permanently or seasonally arid (accounting for the largest terrestrial
77 biome, Peel, Finlayson, & McMahon, 2007) and vascular plants are excluded or diminished
78 by low water availability or low temperatures. These environments are often dominated by
79 biological soil crusts (BSC, Pointing & Belnap, 2012). These inconspicuous communities,
80 composed of several poikilohydric organisms (lichens, bryophytes, cyanobacteria, algae,
81 bacteria and microfungi) have only recently been described to make a small but significant
82 (equal to annual anthropogenic carbon input) contribution to global CO₂ uptake (Elbert et al.,
83 2012; Porada, Weber, Elbert, Pöschl, & Kleidon, 2014). The habitats that are dominated by
84 BSC, hot and cold deserts, drylands, badlands, polar regions (Pointing & Belnap, 2012;
85 Belnap, Weber, & Büdel, 2016) are also suggested to be the first, and most severely affected,
86 by predicted temperature increases (IPCC report, 2014).

87 Lichens are a key component in late successional stage BSC (Rosentreter, Eldridge,
88 Westberg, Williams, & Grube, 2016). Compared to other BSC components, their
89 proportionate biomass is high, so that the ecophysiological response of a soil crust lichen can
90 be considered an appropriate proxy for the response of the entire crust (Lange, 2003). Lichens
91 are fungi (mycobiont) symbiotic with photosynthetic green algal or cyanobacterial partners
92 (photobiont). The mycobiont composes the major part of the lichen and contributes the
93 majority of the respired CO₂. In instantaneous measurements of lichen CO₂-exchange, the

respiration increases exponentially with increasing temperature, whilst gross photosynthesis increases up to about 30°C before beginning to decline. As a consequence, lichen *net* photosynthesis has an optimal temperature above which further increases of respiration depress net carbon gain, and it has been shown that net CO₂ exchange can become negative at moderately elevated temperatures (Green & Lange, 1994). Lethal temperatures for photosynthesis in active, hydrated lichens are not high, usually around 30 to 35°C (Lange, 1965; Smith, 1981; Chiarucci, Calderisi, Casini, & Bonini, 2008; Maphangwa, Musil, Raitt, & Zedda, 2014).

Because lichens, as poikilohydric organisms, often become hydrated overnight due to dew, fog or rain, the ecophysiological response to increased temperatures overnight are of special interest. While night-time hydration at *moderate* temperatures stimulated growth and resulted in thallus extension (Bidussi, Gauslaa, & Solhaug, 2013), it has been suggested that being hydrated during *warm* nights results in exceptionally poor carbon balance and that this may exclude lichens from some habitats (Lange, 2000). However, these assumptions are only valid if the instantaneous responses of P and R remain stable with respect to temperature and no acclimation occurs to mitigate these effects.

To date, the processes that underpin acclimation to increasing temperature are poorly understood for BSC communities and lichens (Green & Proctor, 2016). Larson and Kershaw (1975) reported species-specific acclimation with some species showing seasonal changes in the net photosynthetic (NP) capacity with constant respiration and others responding in a manner similar to the process of cold hardening found in higher plants (Larson & Kershaw, 1975). Therefore, the responses of the two processes (NP and R) should be considered separately to better understand the lichens response to changing climate. Although NP and R have different temperature sensitivities, both processes have been described to acclimate with changing seasons under natural conditions (Lange & Green, 2005; MacKenzie, MacDonald,

Dubois, & Campbell, 2001). While acclimation of lichen R seems to be species specific and can show full acclimation to temperature (Lange & Green, 2005), seasonal acclimation of lichen NP (electron transport rate and gross photosynthesis) is triggered by two factors, temperature and light availability (MacKenzie et al., 2001). The underlying physiological mechanisms are yet to be understood, and the response of lichens to environmental change is additionally confounded by their longevity through many seasonal cycles and by their slow growth rates (Lindsay, 1973; Sancho, Green, & Pintado, 2007).

In polar regions, lichens form a major part of the vegetation and are dominant in biological soil crusts (Williams et al., 2017a, Fig. 1a). Here, studies emphasising acclimation processes and the corresponding risk assessment are expected to be particularly useful because colder climates are considerably more responsive to increased ambient temperatures compared with warmer regions (Carey et al., 2016). The Antarctic Peninsula, especially, serves as an early warning system in understanding species and ecosystem responses to climate change because it recently experienced relatively fast regional climate changes (Turner et al., 2014). At present, temperatures are, at least temporary, declining (Turner et al., 2016) and this complicates the already complex response of the local biodiversity to a changing climate (Convey, 2011), for example, through “snowkill” as an additional threat to local lichen populations (Sancho et al., 2017). An important aspect of recent climate change scenarios overall is the increasing frequency of extreme events such as heat waves ($>5^{\circ}\text{C}$ above daily temperature for at least 5 consecutive days) (IPCC report, 2007). Such infrequent warming events might have significant and long-lasting impacts on local communities (Walther et al., 2002). In the Antarctic, for example, the extraordinarily warm summer 2001-2002 in Taylor Valley, continental Antarctica, had a disproportionately large impact on the local invertebrate community (Courtright, Wall, & Virginia, 2001), and provides a case study for projecting how above- and below-ground ecosystems may respond in the future (Wall, 2007). The most

drastic changes from this warming event were to water availability, with significant influences that persisted for several years (Barrett et al., 2008). This demonstrates the strong interconnection between the thermal and the hydric environment in the Antarctic and underlines the need for accurate experimental testing and monitoring.

This study aims to describe potential acclimation processes of R and NP to changing temperatures in polar lichens with special regard to differences in thermal acclimation within these symbiotic organisms. To isolate the temperature effect, we chose an experimental approach that allows maximum control and monitoring of conditions (water availability, light regime). The two following hypotheses are tested:

1: Lichens show thermal acclimation of respiration in a manner similar to patterns known from vascular plants, mitigating the effects of higher temperatures, while photosynthetic rates and the lichen thallus morphology remain more or less unaffected.

2: The degree of acclimation and the rates at which lichens acclimate to new temperatures will be species specific. We expect species with broader distribution patterns and ecophysiological amplitudes to acclimate both faster and more complete, than species with very specific physiological adjustments to their surrounding environment.

Materials and Methods

Species selection

We chose three different lichen species collected on Livingston Island in the maritime Antarctic. The lichens were selected to cover a variety of different growth forms, distribution patterns and photobionts with possible differences in their individual acclimation potential. For example, a cyanobacterial photobiont might contribute to a lichen's ability to adapt to temperature, as shown for the tropical lichen *Dictyonema glabratum* (Lange, Büdel, Zellner, Zotz, & Meyer, 1994), the epilithic lichen *Peltula capensis*, from South Africa (Wessels & Kappen, 1993) and *Collema tenax*, a typical soil-crust lichen in arid lands (Lange, Belnap, & Reichenberger, 1998). Because of the low photobiont diversity in the Antarctic, both for green algal photobionts (Domaschke, Fernández-Mendoza, García, Martín, & Printzen, 2012) and cyanobionts (Wirtz et al., 2003), we distinguish between these two functional groups rather than specific photobiont strains. To minimise covariation, each of the three traits (growth form, distribution pattern, photobiont) overlapped within two of the selected lichen species.

Stereocaulon alpinum Laurer is a member of the group of circumarctic-alpine lichens that are found bipolar and also in the alpine environments of the temperate regions (Øvstedal & Smith, 2001). *S. alpinum* also occurs in the dry cool boreal zone, where mean summer temperature reaches up to 13.8°C (Coxson & Marsh, 2001). *S. alpinum* is a fruticose lichen (Fig. 1b), circa 5-7-cm high, with cephalodia that contain cyanobacteria of the genus *Nostoc* as an additional cyanobiont, in addition to the trebouxoid primary green algal photobiont. Due to its broader distribution and its tripartite composition, this lichen is considered to have a relatively wide ecological amplitude. *Usnea aurantiaco-atra* (Jacq.) Bory, is a dominant component in vegetation communities of the maritime Antarctic and Alpine subantarctic regions (Øvstedal & Smith, 2001). *Usnea aurantiaco-atra* has a fruticose, erect growth form with many apothecia (Fig. 1c). It contains a trebouxoid green algal photobiont and can be

considered to be highly specialised to Antarctic climate conditions (Laguna-Defior, Pintado, Green, Blanquer, & Sancho, 2016). *Placopsis contortuplicata* I. M. Lamb, in contrast to the first two species, grows foliose to effigurate (Fig. 1d) but shares the feature of having cephalodia containing *Nostoc* as a cyanobiont with *S. alpinum*. The distribution of *P. contortuplicata* is restricted to the southernmost South America, the Subantarctic Islands and the Antarctic Peninsula (to at least 70°S), a distribution that it shares with *U. aurantiaco-atra*.

Sample collection

All lichen samples were collected in January 2015 in the vicinity of Juan Carlos I base (62°39 S; 60°23 W), which is located in the South Bay of Livingston Island, Antarctica. Mean annual temperatures are -2.8 °C with summer mean monthly temperatures above freezing, and the maximum mean monthly temperature is 4.3 °C. Mean annual precipitation is 444.5 mm, with 75 % falling in summer and autumn (Bañón, Justel, Velázquez, & Quesada, 2013). The bedrock of Livingston Island is a low-grade metamorphic turbidite sequence with volcanic to volcanoclastic rocks, intruded by igneous bodies (Arche, López-Martínez, & Martínez de Pisón, 1992; Moura, Francelino, Schaefer, Simas, & de Mendonça, 2012). Besides two native flowering plant species (*Deschampsia antarctica* Desv. and *Colobanthus quitensis* (Kunth) Bartl.), 110 lichen and 50 bryophyte species have been reported from the vicinity of Juan Carlos I base (Sancho, Schulz, Schroeter, & Kappen, 1999).

Four intact thalli of each species were collected: *S. alpinum*, *U. aurantiaco-atra* and *P. contortuplicata*. Identification was based on morphological and anatomical features using appropriate determination keys (Øvstedal & Smith, 2001). Samples were dried at room temperature, frozen at -20°C and transported to the laboratory, where they were stored in the frozen state until used. Frozen storage is described as being suitable for long-term storage of lichens for experimental studies (Honegger, 2003).

209 **Experimental design**

210 Because most biological processes in Antarctica operate at the scale of the organism and their
211 microclimate, we chose temperatures that are likely to occur under natural conditions in the
212 lichens microclimate. The overall design of this study was to incubate the lichens at three
213 different temperatures (one control plus two treatments with elevated temperatures) and to
214 track changes in photosynthesis and respiration rates over time. The control group is
215 represented by a set of samples incubated at 5°C as this temperature approximates the mean
216 temperature when the organisms are active under natural conditions (Schlensog, Green, &
217 Schroeter, 2013). The 15 °C treatment was considered to reflect moderately “increased”
218 temperatures as this temperature is 5°C above the recorded maximum thallus temperature
219 when the organisms were active at Livingston Island (Schroeter, Green, Pintado, Türk, &
220 Sancho, 2017). A 23°C treatment was chosen to reflect an “extreme” but still reasonable
221 change. Temperatures up to 26°C were recorded as maximum thallus temperature while the
222 organisms were active on Leonie Island, Antarctica (Schroeter et al., 2017). Our treatment
223 aims to increase the duration of exposure to such temperature extremes to simulate a “heat
224 wave” (De Boeck, Dreesen, Janssens, & Nijs, 2010). The treatments at the three temperatures
225 will be referred to as control (C₅), 15 degrees (T₁₅) and 23 degrees (T₂₃). Three replicates each
226 for the three selected species were used and, in order to avoid sample dependent presetting
227 (such as microhabitat dependent acclimation), each thallus was divided into 3 parts, with each
228 part allocated to a different temperature treatment.

229 After the start of the treatments, CO₂ exchange (NP, net photosynthesis and R, respiration)
230 was measured for all lichen samples at 5, 15 and 23°C and this was repeated at one week
231 intervals. A standardised label was allocated to each measured sample: eg. C_{5,5} = control
232 samples measured at 5°C, T_{23,15} = samples in the 23°C treatment measured at 15°C. The aim

was to detect any acclimation to the treatment temperature and the instantaneous response to the other two temperatures.

Sample treatment

Prior to the experiment, the intact lichen samples underwent a reactivation procedure composed of 2 days dry storage at 4°C in the dark and 24h at 4°C and 200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, before they were divided and fixed in CO₂-inert wire-mesh baskets. This procedure was found to be suitable for previous gas exchange studies on polar lichens and biological soil crusts (Colesie, Green, Haferkamp, & Büdel, 2014) and removes problems of water condensation on the sample and resaturation respiration that is known to differ both in amplitude and in time required to reach steady state after an initial burst in respiration (Sundberg, Ekblad, Näsholm, & Palmqvist, 1999). Initial test experiments showed that gas exchange rates were in the same order of magnitude as during field measurements from other studies (Green, Schroeter, Kappen, Seppelt, & Maseyk, 1998) indicating no physiological consequences from storage at -20°C. Nine baskets (3 species x 3 replicates) were put into a 30 cm x 20 cm plexiglass box with the lid slightly open, together with a temperature and humidity logger (HOBO, Onset). Three of these boxes were prepared and each of them allocated to a growth cabinet at 5 (control), 15 or 23°C (Total number of samples: 3 species x 3 replicates x 3 treatment temperatures). Each box was arranged in a way that 150-200 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ reached the lichen surface. For reactivation of lichen metabolism, the samples were sprayed with water until water saturation (external water droplets remaining on the lichens' surface). The activity of the lichens in the incubation boxes was monitored using an Imaging chlorophyll fluorometer (Imaging PAM, Walz, Germany). The lichens were then allowed to slowly desiccate in the boxes and once they had dried out and became inactive (Yield of PSII below 0.2) they were kept in this stage for 1 day until the next reactivation. This treatment was chosen to mimic natural conditions because lichens as poikilohydric

organisms often repeatedly undergo hydration-desiccation cycles under natural conditions (Green, Sancho, & Pintado, 2011) and similar treatments were shown to optimise lichen cultivation in growth chambers (Gauslaa, Alam, & Solhaug, 2016). Each hydration-desiccation cycle took about three to four days so that assays of photosynthesis and respiration rates were on a weekly basis. Total incubation time was six weeks.

Assays

Carbon dioxide gas exchange measurements were conducted under controlled laboratory conditions using a mini cuvette system (CMS400, Walz Company, Effeltrich, Germany). Relative humidity of the incoming air was adjusted using a cold trap and was kept stable at 90% for all measurements. Net photosynthesis (NP) was measured under saturating light levels at $500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ and rates of dark respiration (R) were obtained by shading the cuvette completely (until the ΔCO_2 signal had stabilised). Assay temperatures were adjusted to those of the treatments and each sample was measured at all three temperatures (5, 15 and 23 °C; total number of readings: 3 species x 3 replicates x 3 treatment temperatures x 3 assay temperatures). In order to minimise any effects due to the assay temperature being different to the treatment temperature the measurements at assay temperatures were made randomly and samples immediately replaced in their treatment temperature after the assays. The CO_2 exchange of the samples was related to chlorophyll content. Chlorophyll contents were determined by extracting the samples twice with dimethyl-sulfoxide (DMSO) at 60°C for 90 minutes and measuring the absorption at standard wavelengths (Ronen & Galun, 1984).

Microscopy

Visualisation of the internal thallus structure and anatomical properties was performed with a light microscope equipped with differential interference contrast (Axioskop, Carl Zeiss, Jena, Germany). Thin sections of the lichen thalli before and after the treatment were prepared

using a freezing microtome (Leitz, Wetzlar, Germany). Pictures were taken using the Axio-Vision software.

Calculations and statistics

The occurrence of acclimation over the time course of the experiment was investigated by presenting the results in three different ways.

First, the rates of net photosynthesis (NP), respiration (R) and the ratio of net photosynthesis to respiration (NP/R), measured at their respective incubation temperatures ($C_{5,5}$, $T_{15,15}$ and $T_{23,23}$), were plotted over the time course of the experiment with the objective of detecting changes over time (Fig. 3). The ratio of net photosynthesis to respiration (NP/R) was calculated in order to approximate whole lichen homeostasis. A value of 1 indicates that both processes compensate each other, while values below 1 indicate a high fraction of respiration compared to net photosynthesis and vice versa. Statistical testing was based on regression analysis using the Sigma Plot software (Systat Software GmbH, San Jose, USA). All linear regression lines are based on data that passed normality tests (Shapiro-Wilk) and tested for significance with $\alpha = 0.05$. For each plot, regression lines were fit to the data, the null hypothesis (slope equal to zero, $P > 0.05$) tested, and the coefficient of determination (r^2) calculated. Effects of temperature were analysed using a single factor GLM (General Linear Model) repeated measure procedure for each species separate. Effects of time were analysed using a single factor GLM repeated measure procedure for each species at each temperature separate. The species*temperature effect was tested with a two factor GLM repeated measure procedure (SPSS, IBM Analytics, New York). The low sample size did not permit the necessary degrees of freedom to test the interaction term between time and other main effects.

Second, in order to see if the instantaneous response to elevated temperature remains stable and whether *growing* at warmer temperatures reduces the negative effects resulting from this,

we assessed comparisons between $C_{5,5}$ and $C_{5,15}$, with those between $C_{5,5}$ and $T_{15,15}$. For the $C_{5,5}$ vs. $C_{5,15}$ comparison we expect respiration rates to increase when measured at a higher temperature, but net photosynthesis rates to decrease because 15°C is above the optimal temperature (Lange & Kappen, 1972). As a consequence, a line linking $C_{5,5}$ to $C_{5,15}$ would have a negative slope for both processes. Any acclimation to new, warmer growing conditions ($C_{5,5}$ vs. $T_{15,15}$ comparison) reduces the magnitude of this negative effect and results in a flattening of the negative slope. This means, that if acclimation occurs, the slope between $C_{5,5}$ and $T_{15,15}$ should be less than the slope between $C_{5,5}$ and $C_{5,15}$. Slopes were calculated from a linear equation at the beginning of the experiment and at the end. Means were compared by a two-way repeated measure ANOVA (SPSS, IBM Analytics, New York) using a significance level of $P < 0.05$ to check for differences between species and treatment temperature. Where ANOVA indicated significant results the treatment effect was assessed for each species.

Third, physiological rates (NP and R) of organisms grown at elevated temperature (15°C), but measured at the standard, control temperature ($T_{15,5}$) were analysed. Acclimation to higher growth temperatures results in decreasing rates when measured back at colder, standard temperatures. In order to demonstrate this decrease mean values from the beginning of the experiment were compared to those gathered after six weeks treatment and between the species by a two-way repeated measure ANOVA (SPSS, IBM Analytics, New York) using a significance level of $P < 0.05$.

Results

Incubation conditions

During the experiment, treatment temperatures remained stable with only a small discrepancy from the intended treatment temperatures (5, 15 and 23°C, Tab. 1). Active time was inversely proportional to the water vapour pressure deficit VPD (Fig. 2; $P = 0.0022$) and such relationships were previously described to appropriately simulate heat wave events (De Boeck et al., 2010).

Change in NP, R and their ratio over time

Control samples incubated and measured at 5°C showed stable net photosynthesis and respiration that did not change over the time course of the experiment (black lines, Fig. 3). All repeated measure GLMs showed no significant effects of time ($P > 0.05$). This stability under control conditions indicated that our control conditions were suitable for the stable maintenance of all 3 lichens selected for this study.

Net photosynthesis

For all species, the net photosynthetic rates were similar for the control and 15°C treatments at the start of monitoring (Fig. 3; 1.93 vs. 2.14 $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ s}^{-1}$ for *S. alpinum* ($P = 0.53$), 0.85 vs. 1.0 $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ s}^{-1}$ for *U. aurantiaco-atra* ($P=0.80$), 1.41 vs 1.69 $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ s}^{-1}$ for *P. contortuplicata* ($P = 0.58$)). However, during the treatments they changed significantly ($F_{3/4} = 8.25$ and $P = 0.035$ for *S. alpinum*; $F_{2/5} = 8.21$ and $P = 0.026$ for *P. contortuplicata*). At 15°C (red lines, Fig.3), NP increased for *S. alpinum* ($P = 0.0002$, $r^2 = 0.5925$, slope = 0.3229, $t = 4.8940$), remained stable for *U. aurantiaca-atra* ($F_{2/1} = 0.75$ and $P = 0.632$)) and decreased to near zero for *P. contortuplicata* ($P = 0.0103$, $r^2 = 0.3855$, slope = -0.4847, $t = -2.9637$). When exposed to extreme temperatures (23°C, green lines, Fig. 3), already at the start of the experiment, only *S. alpinum* showed NP rates similar to control and 15°C treatments. For *U.*

aurantiaco-atra and *P. contortuplicata* this was close to zero or even negative. Over time, under the extreme temperature regime, net photosynthesis declined significantly for *S. alpinum* ($P = 0.0044$, $r^2 = 0.6120$, slope = -1.3899, $t = -3.7740$) and *U. aurantiaco-atra* ($P = 0.0220$, $r^2 = 0.5509$, slope = -1.6229, $t = -2.9302$) with a similar trend for *P. contortuplicata* ($P = 0.0536$, $r^2 = 0.4342$, slope = -0.7583, $t = -2.3177$). However, all species ceased to show a NP response to light after 3 or 4 weeks indicating that the photobiont was dead and that treatment at 23°C exceeded their survival capacity at least within the local population.

Respiration

Respiration rates were significantly increased by treatment temperature ($F_{2/6} = 14.34$, $P = 0.005$ for *S. alpinum*, $F_{2/6} = 11.605$, $P = 0.009$ for *U. aurantiaco-atra*; $F_{2/4} = 105.99$, $P < 0.001$ for *P. contortuplicata*) and were at least double that of the controls (Fig. 3). Nevertheless, these rates did not change over the time course of the experiment ($F_{3/4} = 0.58$, $P = 0.65$ for *S. alpinum*, $F_{1/5} = 1.06$, $P = 0.344$ for *U. aurantiaco-atra*; $F_{2/5} = 0.18$, $P = 0.84$ for *P. contortuplicata*), indicating no thermal acclimation of these processes. Respiration rates were highest at 23°C for the first 3 or 4 weeks (green lines, Fig. 3) but after this period the 23°C samples did not show a reaction to changing light and the photobionts were therefore considered dead and the samples excluded from further analysis.

NP/R ratio

NP/R ratios were different between the different temperature treatments ($F_{6/8} = 67.220$, $P < 0.001$ for *S. alpinum*; $F_{2/6} = 35.76$, $P < 0.001$ for *U. aurantiaco-atra*; $F_{2/6} = 165.12$, $P < 0.001$ for *P. contortuplicata*). The NP/R ratio for the control groups of all three species indicated that NP rates at 5°C were at least double R rates during the whole experiment. At 15°C, the NP/R ratio was lower and close to 1 at the start of the treatments but as the experiment progressed (red lines, Fig.3), the ratio recovered to control levels for *S. alpinum* ($P = 0.0010$,

378 $r^2 = 0.7170$, slope = 0.2120, $t = 5.5859$), remained stable around 1 for *U. aurantiaca-atra* ($F_{2/1}$
 379 = 4.68 and $P = 0.311$)) and declined to below 1 after 3 weeks for *P. contortuplicata* ($P =$
 380 0.0325, $r^2 = 0.2868$, slope = -0.1881, $t = -2.3730$). At the extreme temperature (23°C, green
 381 lines, Fig. 3) NP/R ratios were below 1 for all samples and, as the experiment progressed,
 382 showed significant decreases for *S. alpinum* ($P = 0.0010$, $r^2 = 0.7170$, slope = -0.5331, $t = -$
 383 4.7751) and *U. aurantiaco-atra* ($P = 0.0002$, $r^2 = 0.8727$, slope = 0.4309, $t = -6.9282$). For *P.*
 384 *contortuplicata* NP/R ratio at 23 °C was stable around zero during the latter half of the
 385 experiment.

386 **Changes in the response to high temperatures**

387 According to our suggestion any acclimation to new, warmer growing conditions should
 388 entail a lesser negative slope at the end of the experiment (grey bars, Fig. 4) when compared
 389 to the start (black bars, Fig. 4). For the controls, no such decline occurred for any species,
 390 either for net photosynthesis (Fig. 4a, b, c) or for respiration (Fig. 4d, e, f), showing that the
 391 immediate response to increased temperatures was stable and consistent during the whole
 392 experiment for the controls. However, growth at elevated temperatures (15°C) had significant
 393 effects that were different between the species ($F_{2/3} = 95.03$, $P = 0.002$ for NP; $F_{2/3} = 88.88$, P
 394 = 0.002 for R). For NP of *S. alpinum* (Fig. 4a), positive slopes indicated that NP rates
 395 increased when measured at 15°C, which shows that the optimal temperature for this species
 396 was above 5°C from the beginning of the experiment. After growing at 15°C for 6 weeks the
 397 slope between $C_{5,5}$ and $T_{15,15}$ was significantly increased ($P = 0.039$), indicating that the
 398 temperature optimum had shifted to even higher temperatures and the species had acclimated
 399 to the new warmer growing temperature. No such changes occurred for respiration in *S.*
 400 *alpinum* (Fig. 4d) indicating that growing at elevated temperatures did not change the
 401 response of respiration to higher temperatures and there was no acclimation of R to the
 402 warmer growing temperature. For *U. aurantiaco-atra* (Fig. 4b, e) the treatment had no

significant effect on the slopes. For *P. contortuplicata* (Fig. 4c, f) the changes were most drastic. Here, net photosynthesis for $T_{15,15}$ at the end of the experiment was lower than it was at the start and compared to the control group, resulting in a significantly increased negative slope that indicated that these samples suffered from severe thermal stress and the 15°C treatment already exceeded their photobiont survival capacity. Respiration also increased significantly ($P = 0.019$, Fig. 4f) for the treatment at the end of the experiment, indicating that the lichens carbon balance tipped strongly into the negative.

Changes in response to control temperature

Net photosynthesis rates $T_{15,5}$ varied significantly between the species ($F_{2/1} = 345$, $P = 0.038$) with the highest rates for *S. alpinum*. Additionally, NP rates for all species decreased when compared between the beginning and the end of the experiment (Fig. 5; $F_{2/1} = 1470$, $P = 0.018$). This indicated that initial rates could not be maintained in the treatment and most possibly the temperature optimum had shifted for these species. It also implied that key traits (high NP rates at low temperatures) were lost during the experiment. For *P. contortuplicata* NP rates were close to zero and negative so that these were excluded from the analysis.

Respiration rates measured at the 5°C control temperature (Fig. 5) did not differ between the species ($F_{2/1} = 2.77$, $P = 0.39$) and also showed no change from the start to the end of the experiment ($F_{2/1} = 13.98$, $P = 0.186$). This indicated that the temperature response of respiration did not change during the experiment and initial rates were preserved.

Morphological changes

Visual effects of the incubation at 15°C varied drastically between the species. In untreated samples of *S. alpinum* the green algal photobiont was located in small bundles underneath the upper cortex (Fig. 6a). After six weeks of incubation at 15°C the photobiont layer appeared less constricted than before but remained vividly green (Fig. 6b). In *U. aurantiaco-atra* the

photobionts did not occur in a compact layer but were spread in small clusters between the outer cortex and the central string (Fig. 6c). These algal clusters could still be found after the 15°C treatment (Fig. 6d) but some of them only contained dead cell material (Fig. 6e). In *P. contortuplicata* the green algal photobiont originally formed a dense layer underneath the upper cortex (Fig. 6f) but after the incubation at 15°C only dead, brown cell material was present indicating that the photobiont inside this lichen species did not survive the treatment (Fig. 6g).

Discussion

In the present study, we have provided experimental evidence that polar macro lichens exposed to warmer growing conditions are unable to rapidly reduce their resulting respiration losses via thermal acclimation of respiration. For all tested species, an *extreme* increase in temperature exceeded their photobiont survival capacity at the latest after 3 or 4 weeks. At a more moderate *increased* temperature, we found a high degree of response variability and sensitivity between the species. Most interestingly, a widely distributed lichen species (*S. alpinum*) was capable of restoring its energy homoeostasis via an increase in net photosynthesis. In contrast, the specialised species, that were naturally growing in the same environment and spatially close, did not show this type of acclimation (*P. contortuplicata*), or showed it less obviously (*U. aurantiaco-atra*). Significant effects on photobiont vitality indicated that any acclimation processes in lichens are subject to complicated interplays between the two symbionts and strongly depend on their individual acclimation potential. Our finding emphasises species-specific sensitivity to changes in temperature in this pristine environment and underlines the fragility of the vegetation community composition.

Based on extensive studies in higher plants, thermal acclimation of respiration is a common biological feedback to long-term temperature increases (e.g. Atkin et al., 2015). If lichens have a similar capacity to acclimate their energy metabolism in response to changes in their

452 thermal environment, then we would expect that the respiration rates of lichens held at higher
453 temperatures would show a decrease in respiration losses over time. Our results show that, as
454 expected, respiration shows an immediate increase when all species were activated at higher
455 temperatures (Fig. 3,4). However, the increased R at the higher temperatures (Fig. 3) did not
456 show any downregulation with time over the 6 weeks of the experiment, suggesting that no
457 thermal acclimation of respiration occurred in any of three selected lichen species in this
458 study. We can support this assumption with three lines of evidence. Firstly, respiration rates
459 remained at the same level over the time course of the experiment (Fig. 3). Secondly, the
460 responses of respiration to higher temperatures remain stable during the treatment (Fig. 4).
461 Thirdly, organisms that were exposed to warmer growing conditions did not show any
462 downregulation of respiration when measured at the standard, control temperature (Fig. 5).
463 This finding is in line with a study on soil respiration in the Antarctic, where it has been
464 shown that both the biomass-specific respiration rate and the overall rate of SOC
465 mineralisation increased with temperature and this was interpreted as respiration by soil
466 micro-organisms not down-regulating relative to temperature (Laudicina et al., 2015). One
467 explanation for this finding might be that, unlike autotrophic counterparts, heterotrophic
468 organisms do not gain any evolutionary advantage from physiological downregulation in
469 response to increased temperature (Hartley, Heinemeyer, & Ineson, 2007). In agreement with
470 this, it is known that Antarctic invertebrates rely on life history traits that allow them to
471 remain dormant throughout most of the year whilst taking advantage of short-term favourable
472 (warmer) conditions (Convey, 1996; Convey, 1997). Such survival strategies enhance the
473 performance of the native biota under current climate conditions and are discussed to be an
474 important factor influencing soil invertebrate communities (Nielsen & Wall, 2013).
475 Nevertheless, this finding is unexpected, especially because lichens were previously described
476 to acclimate respiration rates within seasons under natural conditions (Lange & Green, 2005).

In contrast to the study from Lange and Green (2005), we applied drastic and abrupt changes rather than a continuous change in conditions. The severity of changes we applied was necessary in order to provoke significant responses in a reasonable amount of time and to simulate a heat wave stimulus. The advantage of this approach is that we have experimentally focused on one effect and can exclude factors that potentially cover temperature effects.

In addition to these negative effects of increased temperature on lichen respiration, one important factor in this study is the deleterious effect of the higher temperatures on the photobionts (Fig. 6). All three species showed a collapse in NP/R when incubated at 23°C (Fig. 3). This collapse appears to be due to photobiont death and is also shown at 15°C for *P. contortuplicata*, and to a lesser degree for *U. aurantico-atra* (Fig. 6). The death of the photobionts within the thallus has also previously been described for *Psora decipiens*, when cold and wet acclimated thalli were transplanted to hot desert conditions and vice versa (Williams et al., 2017 b). It indicates that 23°C is well above the survival temperature for all the photobionts in this study and 15°C is about the upper limit for the two highly specialised lichen species, which only occur with a narrow distribution range in the Antarctic. The finding is in line with other studies indicating a possible adaptation of Antarctic photobionts to colder growing conditions (Balarinová, Váczi, Barták, Hazdrová, & Forbelská, 2013). Photobiont death makes it difficult to interpret changes in NP/R, especially at 23°C, but also partly at 15°C for the two temperature-sensitive lichen species. Only *S. alpinum*, which is a lichen species with a wider distribution range, is robust enough to acclimate (Fig. 3).

Surprisingly, and in contrast to our hypothesis, the recovery of the NP/R ratio in *S. alpinum* resulted from increased net photosynthesis rates, rather than acclimation of respiration. This finding is clearly substantiated by a significantly lowered NPmT at the end of the experiment (Fig. 5) and the shift of the NP temperature optimum (Fig. 4). In lichens, there are two mechanisms available for acclimating photosynthesis to changing growth temperatures. The

first would be by changing the number of photobiont cells in the thallus (Tretiach, Bertuzzi, Carniel, & Virgilio, 2013). Domaschke, Vivas, Sancho, and Printzen (2013) demonstrated this option for *Cetraria aculeata*, where temperate populations of the same lichen species had a significantly higher NP and number of photobionts cell per mg dry weight than their polar counterparts. The second option would be through acclimation of the macromolecular composition to different environmental conditions within an existing photobiont cell (MacKenzie et al., 2001). It has been shown that within a nearly stable, non-dividing algal cell population in *Lobaria pulmonaria*, the key photosynthetic proteins showed significant seasonal acclimation (Schofield, Campbell, Funk, & MacKenzie, 2003). This second mechanism may be similar to findings in vascular plants, where the Rubisco enzyme content and activation is a key component of thermal acclimation of photosynthesis (e.g. Hurry, Strand, Tobiaeson, Gardestöm, & Öquist, 1995).

In order to estimate the relevance of acclimation processes in lichens under natural conditions, it is also important to know the actual rate at which such acclimation process occurs. Such an assessment can be done by checking when the NP/R homoeostasis was restored. For *S. alpinum* NP/R ratio of warm incubated samples already equals that of the controls after three weeks. This equates to only about 106 active hours and reflects a fast rate of acclimation, supporting the suggestion of acclimation via changes in the macromolecular composition, because this rate of change is higher than the described turn-over rate of photobionts in lichens (Hill, 1992). It also indicates how quickly lichens can respond to environmental temperature change and suggests that although lichens are slow-growing organisms, this does not mean that their metabolic processes are less responsive than in other organisms. For example, thermal acclimation by plants (Atkin & Tjoelker, 2003), animals (Seebacher, White, & Franklin, 2015) or fungi (Crowther & Bradford, 2013) manifests itself over time frames ranging from days to weeks.

Nevertheless, such acclimation of NP seems to be a species-specific trait. In general, broadly distributed, or species from variable temperature environments are likely to be more capable of acclimating than species experiencing a limited thermal range (Crowther & Bradford, 2013; Seebacher et al., 2015). If so, thermal acclimation as a species-specific trait in lichens can be suitable for environmental risk assessment and interpretation of the ecological spectrum of a species. Species with high acclimation potential are considered to be at less at risk than ones with narrow ecophysiological amplitudes. In our study the two species with restricted distribution in the Antarctic showed little acclimation potential, in fact, *P. contortuplicata*, did not show any signs of acclimation for R nor NP and died during the incubation at higher temperatures. *U. aurantiaco-atra* did not show increasing rates of NP with time but both the decreasing NPmT (Fig. 5) and the marginally lowered slope (Fig. 4b) points towards some potential acclimation processes. Therefore, we interpret the widely distributed lichen *S. alpinum* to be at less risk than highly adapted Antarctic restricted species such as *P. contortuplicata* or *U. aurantiaco-atra*.

Our findings provide mechanistic insight into why lichen biodiversity could decline and the lichen community composition shift to more dominant generalist species in the maritime Antarctic. This phenomenon is already described for the Arctic tundra (Lang et al., 2012). Such community shifts could lead to regional-scale biotic homogenisation, which is a threat for Antarctic ice-free habitats (Lee et al., 2017) and could alter ecosystem functioning and productivity (Clavel, Julliard, & Devictor, 2011). However, we stress that the response of individual organisms cannot fully reflect the entire community response for the selected ecozone. Future studies should address three important topics: the first concerns the potential for acclimation process in lichen photobionts. It is clear from our study that the photobionts in the studied lichens were not able to survive elevated temperatures that were maintained for extended periods (weeks). In one species even temperatures as low as 15°C were lethal in this

study. Differential adaptive and acclimative mechanisms appear to exist in phototrophic microorganisms residing in low-temperature environments, although these are also described to be understudied (Morgan-Kiss, Priscu, Pockock, Gudynaite-Savitch, & Huner, 2006). Secondly, the effects of increased temperature on the lichen and the biochemical mechanisms underlying this response should be studied in greater detail. Thirdly, future studies need to combine laboratory studies with in situ site performance.

Acknowledgements:

We are grateful to the head of the Spanish Antarctic Station, Dr. Jordi Felipe, and its entire staff for assistance in the field work and technical equipment as well as logistical support. The Spanish Antarctic Committee and Antonio Quesada also provided essential aid in travel to and from Livingston Island. Prof. Leo Sancho for overall support and assistance. Natalie Kunz for fruitful discussions about study design and data evaluation. Permits: sampling and research activities were approved by the German authorities (Biological soil crust algae from the Polar Regions; 24.09.2014). Financing and logistical support were kindly provided by the German Research Council in the framework of the Priority Programme ‘‘Antarctic Research’’ 1158 (DFG, BÜ666/17-1). CC gratefully acknowledges the Alexander von Humboldt Foundation for financial support via the Feodor Lynen Research Fellowship.

575

576 References:

577 Arche, A., López-Martínez, J., & Martínez de Pisón, E. (1992). Sedimentology of the Miers
578 Bluff formation, Livingston Island, South Shetland Islands. In Y. Yoshida, K. Kaminuma, &
579 K. Shiraishi (Eds.), *Recent progress in Antarctic Earth Science* (pp. 357–362). Tokyo: Terra
580 Scientific Publishing.

581 Atkin, O. K., & Tjoelker, M. G. (2003). Thermal acclimation and the dynamic response of
582 plant respiration to temperature. *Trends in Plant Science*, 8, 343-351.

583 Atkin, O. K., Bruhn, D., Hurry, V. M., & Tjoelker, M. G. (2005). Evans Review No. 2: The
584 hot and the cold: unravelling the variable response of plant respiration to temperature.
585 *Functional Plant Biology*, 32, 87-105.

586 Atkin, O. K., Bloomfield, K. J., Reich, P. B., Tjoelker, M. G., Asner, G. P., Bonal, D., ...
587 Creek, D. (2015). Global variability in leaf respiration in relation to climate, plant functional
588 types and leaf traits. *New Phytologist*, 206, 614-636.

589 Balarinová, K., Váczi, P., Barták, M., Hazdrová, J., & Forbelská, M. (2013). Temperature-
590 dependent growth rate and photosynthetic performance of Antarctic symbiotic alga *Trebouxia*
591 sp. cultivated in a bioreactor. *Czech Polar Reports*, 3, 19-27.

592 Bañón, M., Justel, A., Velázquez, D., & Quesada, A. (2013). Regional weather survey on
593 Byers Peninsula, Livingston Island, South Shetland Islands, Antarctica. *Antarctic Science*, 25,
594 146-156.

595 Bardgett, R. D., Freeman, C., & Ostle, N. J. (2008). Microbial contributions to climate change
596 through carbon cycle feedbacks. *The ISME Journal*, 2, 805-814.

- 597 Barrett, J. E., Virginia, R. A., Wall, D. H., Doran, P. T., Fountain, A. G., Welch, K. A., &
598 Lyons, W. B. (2008). Persistent effects of a discrete warming event on a polar desert
599 ecosystem. *Global Change Biology*, 14, 2249-2261.
- 600 Belnap, J., Weber, B., & Büdel, B. (2016). Biological Soil Crusts as an organizing principle in
601 drylands. In B. Weber, B. Büdel, & J. Belnap, (Eds.), *Biological soil crusts: An organizing*
602 *principle in drylands* (pp. 3-13), Ecological Studies 226. Springer International Publishing
603 Switzerland.
- 604 Bidussi, M., Gauslaa, Y., & Solhaug, K. A. (2013). Prolonging the hydration and active
605 metabolism from light periods into nights substantially enhances lichen growth. *Planta*, 237,
606 1359-1366.
- 607 Carey, J. C., Tang, J., Templer, P. H., Kroeger, K. D., Crowther, T. W., Burton, A. J., ...
608 Jiang, L. (2016). Temperature response of soil respiration largely unaltered with experimental
609 warming. *Proceedings of the National Academy of Sciences*, 113, 13797-13802.
- 610 Chiarucci, A., Calderisi, M., Casini, F., & Bonini, I. (2008). Vegetation at the limits for
611 vegetation: vascular plants, bryophytes and lichens in a geothermal field. *Folia Geobotanica*,
612 43, 19-33.
- 613 Clavel, J., Julliard, R., & Devictor, V. (2011). Worldwide decline of specialist species: toward
614 a global functional homogenization? *Frontiers in Ecology and the Environment*, 9, 222-228.
- 615 Colesie, C., Green, T.G.A., Haferkamp, I., & Büdel, B. (2014). Lichen dominated soil crusts
616 show changes in composition, CO₂ gas exchange, and carbon allocation as stress-related traits
617 across habitats of different severity. *The ISME Journal*, 8, 2104-2115.
- 618 Convey, P. (1996). The influence of environmental characteristics on life history attributes of
619 Antarctic terrestrial biota. *Biological Reviews*, 71, 191-225.

- 620 Convey, P. (1997). How are the life history strategies of Antarctic terrestrial invertebrates
621 influenced by extreme environmental conditions? *Journal of Thermal Biology*, 22, 429-440.
- 622 Convey, P. (2011). Antarctic terrestrial biodiversity in a changing world. *Polar Biology*, 34,
623 1629.
- 624 Courtright, E. M., Wall, D. H., & Virginia, R. A. (2001). Determining habitat suitability for
625 soil invertebrates in an extreme environment: the McMurdo Dry Valleys, Antarctica.
626 *Antarctic Science*, 13, 9-17.
- 627 Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., & Totterdell, I. J. (2000). Acceleration of
628 global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature*, 408, 184-
629 187.
- 630 Coxson, D. S., & Marsh, J. (2001). Lichen chronosequences (postfire and postharvest) in
631 lodgepole pine (*Pinus contorta*) forests of northern interior British Columbia. *Canadian*
632 *Journal of Botany*, 79, 1449-1464.
- 633 Crowther, T. W., & Bradford, M. A. (2013). Thermal acclimation in widespread heterotrophic
634 soil microbes. *Ecology Letters*, 16, 469-477.
- 635 Davidson, E. A., & Janssens, I. A. (2006). Temperature sensitivity of soil carbon
636 decomposition and feedbacks to climate change. *Nature*, 440, 165-173.
- 637 De Boeck, H. J., Dreesen, F. E., Janssens, I. A., & Nijs, I. (2010). Climatic characteristics of
638 heat waves and their simulation in plant experiments. *Global Change Biology*, 16, 1992-2000.
- 639 Domaschke, S., Fernández-Mendoza, F., A. García, M., Martín, M., & Printzen, C. (2012).
640 Low genetic diversity in Antarctic populations of the lichen-forming ascomycete *Cetraria*
641 *aculeata* and its photobiont. *Polar Research*, 31, 17353-17366.

- 642 Domaschke, S., Vivas, M., Sancho, L. G., & Printzen, C. (2013). Ecophysiology and genetic
643 structure of polar versus temperate populations of the lichen *Cetraria aculeata*. *Oecologia*,
644 173, 699-709.
- 645 Drake, J. E., Tjoelker, M. G., Aspinwall, M. J., Reich, P. B., Barton, C. V., Medlyn, B. E., &
646 Duursma, R. A. (2016). Does physiological acclimation to climate warming stabilise the ratio
647 of canopy respiration to photosynthesis? *New Phytologist*, 211, 850-863.
- 648 Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Büdel, B., Andreae, M. O., & Pöschl, U.
649 (2012). Contribution of cryptogamic covers to the global cycles of carbon and nitrogen.
650 *Nature Geoscience*, 5, 459-462.
- 651 Gauslaa, Y., Alam, M. A., & Solhaug, K. A. (2016). How to optimize lichen relative growth
652 rates in growth cabinets. *The Lichenologist*, 48(4), 305-310. doi:
653 10.1017/S002428291600030X
- 654 Green, T. G. A., & Lange, O. L. (1994). Photosynthesis in poikilohydric Plants: A
655 Comparison of Lichens and Bryophytes. In E. D. Schulze, & M. M. Caldwell, (Eds.),
656 *Ecophysiology of photosynthesis* (pp 319-337). Berlin: Springer.
- 657 Green, T. G. A., Schroeter, B., Kappen, L., Seppelt, R. D., & Maseyk, K. (1998). An
658 assessment of the relationship between chlorophyll a fluorescence and CO₂ gas exchange
659 from field measurements on a moss and lichen. *Planta*, 206, 611-618.
- 660 Green, T. G. A., Sancho, L. G., & Pintado, A. (2011). Ecophysiology of
661 desiccation/rehydration cycles in mosses and lichens. In U. Lüttge, E. Beck, & D. Barthels,
662 (Eds.), *Plant desiccation tolerance*, Ecological Studies 215, (pp. 89-120). Berlin: Springer.
- 663 Green, T. G. A. & Proctor, M. C. F. (2016). Physiology of photosynthetic organisms within
664 biological soil crusts: Their adaptation, flexibility, and plasticity. In B. Weber, B. Büdel, & J.

- 665 Belnap (Eds.), *Biological soil crusts: An organizing principle in drylands*, Ecological Studies
666 226, (pp 347-381). Springer International Publishing Switzerland.
- 667 Hartley, I. P., Heinemeyer, A., & Ineson, P. (2007). Effects of three years of soil warming and
668 shading on the rate of soil respiration: substrate availability and not thermal acclimation
669 mediates observed response. *Global Change Biology*, 13, 1761-1770.
- 670 Heinemeyer, A., Ineson, P., Ostle, N., & Fitter, A. H. (2006). Respiration of the external
671 mycelium in the arbuscular mycorrhizal symbiosis shows strong dependence on recent
672 photosynthates and acclimation to temperature. *New Phytologist*, 171, 159-170.
- 673 Heinemeyer, A., Wilkinson, M., Vargas, R., Subke, J. A., Casella, E., Morison, J. I., &
674 Ineson, P. (2012). Exploring the "overflow tap" theory: linking forest soil CO₂ fluxes and
675 individual mycorrhizosphere components to photosynthesis. *Biogeosciences*, 9, 79-95.
- 676 Hill, D. J. (1992). The coordination of development of symbionts in mutualistic symbiosis
677 with reference to the cell cycle of the photobiont in lichens. *Symbiosis*, 14, 325-333.
- 678 Honegger R. (2003). The impact of different long-term storage conditions on the viability of
679 lichen forming Ascomycetes and their green algal photobiont, *Trebouxia* spp. *Plant Biology*,
680 5, 324–330.
- 681 Hurry, V. M., Strand, Å., Tobiaeson, M., Gardeström, P., & Öquist, G. (1995). Cold
682 hardening of spring and winter wheat and rape results in differential effects on growth, carbon
683 metabolism, and carbohydrate content. *Plant Physiology*, 109, 697-706.
- 684 IPCC (2007). Contribution of Working Group I to the Fourth Assessment Report of the
685 Intergovernmental Panel on Climate Change. B., Metz, O. R., Davidson, P. R., Bosch, R.,
686 Dave & L. A. Meyer (Eds.), Cambridge and New York: Cambridge University Press.

- IPCC (2014). Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Core Writing Team, R.K. Pachauri & L.A. Meyer (Eds.), Cambridge and New York: Cambridge University Press.
- Karhu, K., Auffret, M. D., Dungait, J. A., Hopkins, D. W., Prosser, J. I., Singh, B. K., ... Gouriveau, F. (2014). Temperature sensitivity of soil respiration rates enhanced by microbial community response. *Nature*, 513, 81-84.
- King, A. W., Gunderson, C. A., Post, W. M., Weston, D. J. & Wullschlegel, S. D. (2006). Plant respiration in a warmer world. *Science*, 312, 536-537.
- Körner, C. (2006). Significance of Temperature in Plant Life. In J. I. L. Morison, & M D. Morecroft (Eds.), *Plant Growth and Climate Change* (pp 48-66). Oxford: Blackwell Publishing Ltd.
- Laguna-Defior, C., Pintado, A., Green, T. A., Blanquer, J. M., & Sancho, L. G. (2016). Distributional and ecophysiological study on the Antarctic lichens species pair *Usnea antarctica/Usnea aurantiaco-atra*. *Polar Biology*, 7, 1183-1195.
- Lang, S. I., Cornelissen, J. H., Shaver, G. R., Ahrens, M., Callaghan, T. V., Molau, U., ... Aerts, R. (2012). Arctic warming on two continents has consistent negative effects on lichen diversity and mixed effects on bryophyte diversity. *Global Change Biology*, 18, 1096-1107.
- Lange, O. L. (1965). Der CO₂-Gaswechsel von Flechten nach Erwärmung im feuchten Zustand. *Berichte der deutschen botanischen Gesellschaft*, 78, 441-454.
- Lange, O. L. & Kappen, L. (1972). Photosynthesis of lichens from Antarctica. In G. A. Llano (Ed.), *Antarctic Terrestrial Biology*. Washington D. C.: American Geophysical Union.

- 709 Lange, O.L., Büdel, B., Zellner, H., Zotz, G., Meyer, A. (1994). Field measurements of water
710 relations and CO₂-exchange of the tropical, cyanobacterial basidiolichen *Dictyonema*
711 *glabratum* in a Panamenian rainforest. *Botanica Acta*, 12, 279–290.
- 712 Lange, O.L., Belnap, B., & Reichenberger, H. (1998). Photosynthesis of the cyanobacterial
713 soil-crust lichen *Collema tenax* from arid lands in southern Utah, USA: role of water content
714 on light and temperature responses of CO₂ exchange. *Functional Ecology*, 12, 195-202.
- 715 Lange, O. L. (2000). Photosynthetic performance of a gelatinous lichen under temperate
716 habitat conditions: long-term monitoring of CO₂ exchange of *Collema cristatum*. *Bibliotheca*
717 *Lichenologica*, 75, 307-332.
- 718 Lange, O. L. (2003). Photosynthesis of soil-crust biota as dependent on environmental factors.
719 In J. Belnap, & O. L. Lange (Eds.), *Biological Soil Crusts: Structure, Function, and*
720 *Management*, Ecological Studies 150, Second edition, (pp. 217-241). Heidelberg, Berlin:
721 Springer.
- 722 Lange, O. L., & Green, T. G. A. (2005). Lichens show that fungi can acclimate their
723 respiration to seasonal changes in temperature. *Oecologia*, 142, 11-19.
- 724 Larson, D. W., & Kershaw, K. A. (1975). Acclimation in arctic lichens. *Nature*, 254, 421-423.
- 725 Laudicina, V. A., Benhua, S., Dennis, P. G., Badalucco, L., Rushton, S. P., Newsham, K. K.,
726 ... Hopkins, D. W. (2015). Responses to increases in temperature of heterotrophic micro-
727 organisms in soils from the maritime Antarctic. *Polar Biology*, 38, 1153-1160.
- 728 Lee, J. R., Raymond, B., Bracegirdle, T. J., Chadès, I., Fuller, R. A., Shaw, J. D., & Terauds,
729 A. (2017). Climate change drives expansion of Antarctic ice-free habitat. *Nature*, 547, 49- 55.
- 730 Lindsay, D. C. (1973). Estimates of lichen growth rates in the maritime Antarctic. *Arctic and*
731 *Alpine Research*, 5, 341-346.

- 732 Luo, Y., Wan, S., Hui, D., & Wallace, L. L. (2001). Acclimatization of soil respiration to
733 warming in a tall grass prairie. *Nature*, *413*, 622–625.
- 734 MacKenzie, T. D., MacDonald, T. M., Dubois, L. A., & Campbell, D. A. (2001). Seasonal
735 changes in temperature and light drive acclimation of photosynthetic physiology and
736 macromolecular content in *Lobaria pulmonaria*. *Planta*, *214*, 57-66.
- 737 Malcolm, G. M., López-Gutiérrez, J. C., Koide, R. T., & Eissenstat, D. M. (2008).
738 Acclimation to temperature and temperature sensitivity of metabolism by ectomycorrhizal
739 fungi. *Global Change Biology*, *14*, 1169-1180.
- 740 Maphangwa, K. W., Musil, C. F., Raitt, L., & Zedda, L. (2014). Will climate warming exceed
741 lethal photosynthetic temperature thresholds of lichens in a southern African arid region?
742 *African Journal of Ecology*, *52*, 228-236.
- 743 Min, K., Lehmeier, C. A., Ballantyne IV, F., & Billings, S. A. (2016). Carbon availability
744 modifies temperature responses of heterotrophic microbial respiration, carbon uptake affinity,
745 and stable carbon isotope discrimination. *Frontiers in Microbiology*, *7*, 2083- 2094.
- 746 Morgan-Kiss, R. M., Priscu, J. C., Pockock, T., Gudynaite-Savitch, L., & Huner, N. P. (2006).
747 Adaptation and acclimation of photosynthetic microorganisms to permanently cold
748 environments. *Microbiology and Molecular Biology Reviews*, *70*, 222-252.
- 749 Moura, P. A., Francelino, M. R., Schaefer, C. E. G., Simas, F. N., & de Mendonça, B. A.
750 (2012). Distribution and characterization of soils and landform relationships in Byers
751 Peninsula, Livingston Island, Maritime Antarctica. *Geomorphology*, *155*, 45-54.
- 752 Nielsen, U. N., & Wall, D. H. (2013). The future of soil invertebrate communities in polar
753 regions: different climate change responses in the Arctic and Antarctic? *Ecology Letters*, *16*,
754 409-419.

- 755 Oechel, W. C., Vourlitis, G. L., Hastings, S. J., Zulueta, R. C., Hinzman, L., & Kane, D.
756 (2000). Acclimation of ecosystem CO₂ exchange in the Alaskan Arctic in response to decadal
757 climate warming. *Nature*, 406, 978-981.
- 758 Øvstedal, D. O., & Smith, R. L. (2001). *Lichens of Antarctica and South Georgia: a guide to*
759 *their identification and ecology*. Cambridge University Press.
- 760 Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007). Updated world map of the Köppen-
761 Geiger climate classification. *Hydrology and earth system sciences discussions*, 4, 439-473.
- 762 Pointing, S. B., & Belnap, J. (2012). Microbial colonization and controls in dryland systems.
763 *Nature Reviews Microbiology*, 10, 551-562.
- 764 Porada, P., Weber, B., Elbert, W., Pöschl, U., & Kleidon, A. (2014). Estimating impacts of
765 lichens and bryophytes on global biogeochemical cycles. *Global Biogeochemical Cycles*, 28,
766 71-85.
- 767 Reich, P. B., Tjoelker, M. G., Machado, J. L., & Oleksyn, J. (2006). Universal scaling of
768 respiratory metabolism, size and nitrogen in plants. *Nature*, 439, 457-461.
- 769 Ronen, R., & Galun, M. (1984). Pigment extraction from lichens with dimethyl sulfoxide
770 (DMSO) and estimation of chlorophyll degradation. *Environmental and Experimental Botany*,
771 24, 239-245.
- 772 Rosentreter, R., Eldridge, D. J., Westberg, M., Williams, L., & Grube, M. (2016). Structure,
773 composition, and function of biocrust lichen communities. In B. Weber, B. Büdel, & J.
774 Belnap (Eds.), *Biological soil crusts: An organizing principle in drylands*, Ecological Studies
775 226 (pp. 121-138). Springer International Publishing Switzerland.

- 776 Sancho, L. G., Schulz, F., Schroeter, B. & Kappen, L. (1999). Bryophyte and lichen flora of
777 South Bay (Livingston Island: South Shetland Islands, Antarctica). *Nova Hedwigia*, 68, 301–
778 338.
- 779 Sancho, L. G., Green, T. G. A., & Pintado, A. (2007). Slowest to fastest: extreme range in
780 lichen growth rates supports their use as an indicator of climate change in Antarctica. *Flora-
781 Morphology, Distribution, Functional Ecology of Plants*, 202, 667-673.
- 782 Sancho, L. G., Pintado, A., Navarro, F., Ramos, M., De Pablo, M. A., Blanquer, J. M., ... &
783 Green, T. G. A. (2017). Recent warming and cooling in the Antarctic Peninsula Region has
784 rapid and large effects on lichen vegetation. *Nature Scientific Reports*, 7, 5689-5697.
- 785 Schlensog, M., Green, T.G.A., & Schroeter, B. (2013). Life form and water source interact to
786 determine active time and environment in cryptogams: an example from the maritime
787 Antarctic. *Oecologia*, 173, 59-72.
- 788 Schofield, S. C., Campbell, D. A., Funk, C., & MacKenzie, T. D. (2003). Changes in
789 macromolecular allocation in nondividing algal symbionts allow for photosynthetic
790 acclimation in the lichen *Lobaria pulmonaria*. *New Phytologist*, 159, 709-718.
- 791 Schroeter, B., Green, T. G. A., Pintado, A., Türk, R., & Sancho, L. G. (2017). Summer
792 activity patterns for mosses and lichens in Maritime Antarctica. *Antarctic Science*, online first.
793 doi: 10.1017/S095410201700027X
- 794 Seebacher, F., White, C. R., & Franklin, C. E. (2015). Physiological plasticity increases
795 resilience of ectothermic animals to climate change. *Nature Climate Change*, 5, 61-66.
- 796 Smith, C. W. (1981). Bryophytes and lichens of the Puhimau geothermal area, Hawaii
797 Volcanoes National Park. *The Bryologist*, 84, 457-466.

- 798 Sundberg, B., Ekblad, A., Näsholm, T. & Palmqvist, K. (1999). Lichen respiration in relation
799 to active time, temperature, nitrogen and ergosterol concentrations. *Functional Ecology*, 13,
800 119-125.
- 801 Tretiach, M., Bertuzzi, S., Carniel, F. C., & Virgilio, D. (2013). Seasonal acclimation in the
802 epiphytic lichen *Parmelia sulcata* is influenced by change in photobiont population density.
803 *Oecologia*, 173, 649-663.
- 804 Turner, J., Barrand, N. E., Bracegirdle, T. J., Convey, P., Hodgson, D. A., Jarvis, M., ...
805 Shanklin, J. (2014). Antarctic climate change and the environment: an update. *Polar Record*,
806 50, 237-259.
- 807 Turner, J., Lu, H., White, I., King, J. C., Phillips, T., Hosking, J. S., ... Deb, P. (2016).
808 Absence of 21st century warming on Antarctic Peninsula consistent with natural variability.
809 *Nature*, 535, 411-415.
- 810 Vanderwel, M. C., Slot, M., Lichstein, J. W., Reich, P. B., Kattge, J., Atkin, O. K., ...
811 Kitajima, K. (2015). Global convergence in leaf respiration from estimates of thermal
812 acclimation across time and space. *New Phytologist*, 207, 1026-1037.
- 813 Wall, D. H. (2007). Global change tipping points: above-and below-ground biotic interactions
814 in a low diversity ecosystem. *Philosophical Transactions of the Royal Society of London B:*
815 *Biological Sciences*, 362, 2291-2306.
- 816 Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J., ... Bairlein, F.
817 (2002). Ecological responses to recent climate change. *Nature*, 416, 389-395.
- 818 Wang, X., Liu, L., Piao, S., Janssens, I. A., Tang, J., Liu, W., ... Xu, S. (2014). Soil
819 respiration under climate warming: differential response of heterotrophic and autotrophic
820 respiration. *Global Change Biology*, 20, 3229-3237.

Wessels, D. & Kappen, L. (1993). Photosynthetic performance of rock-colonising lichens in the Mountain Zebra National Park, South Africa. *Koedoe*, 12, 27–48.

Williams, L., Borchhardt, N., Colesie, C., Baum, C., Komsic-Buchmann, K., Rippin, M., ... Büdel, B. (2017 a). Biological soil crusts of Arctic Svalbard and of Livingston Island, Antarctica. *Polar Biology*, 40, 399-411.

Williams, L., Colesie, C., Ullmann, A., Westberg, M., Wedin, M., & Büdel, B. (2017 b). Lichen acclimation to changing environments: Photobiont switching vs. climate-specific uniqueness in *Psora decipiens*. *Ecology and Evolution*, 7, 2560-2574.

Wirtz, N., Lumbsch, H. T., Green, T. G., Türk, R., Pintado, A., Sancho, L., & Schroeter, B. (2003). Lichen fungi have low cyanobiont selectivity in maritime Antarctica. *New Phytologist*, 160, 177-183.

Figure captions:

Figure 1: Study site and lichen species. a: Overview of the vegetation near Juan Carlos I Base, on Livingston Island. b: *Stereocaulon alpinum* in natural appearance, Glove as scale. c: *Usnea aurantiaco-atra*. d: *Placopsis contortuplicata*.

Figure 2: Active time of lichen samples in the experiment plotted against VPD in the experiment boxes.

Figure 3: Changes in net photosynthesis (upper graphs), respiration rates (middle graphs) and NP/R ratio (lower graphs) during incubation at different temperatures. Shown are mean values ($n = 3$) $\pm$ standard deviation of samples treated at different temperatures and measured at their

respective treatment temperature. • and black line = controls (5°C) measured at 5 °C assay temperature ($C_{5,5}$); • and red line = increased (15°C) measured at 15 °C assay temperature ($T_{15,15}$); ▲ and green line = extreme (23°C) measured at 23°C ($T_{23,23}$).

Figure 4: Responses to elevated temperature (15°C) for the control group and the 15°C treatment. Slopes from linear equations between $C_{5,5}$ vs. $C_{5,15}$ and $C_{5,5}$ vs. $T_{15,15}$ are compared from the beginning (black bars) and the end of the experiment (grey bars). Data are presented separately for changes in net photosynthesis (a,b,c) and respiration (d,e,f). Shown are mean values ($n = 3$) $\pm$ standard deviation and results from Post hoc tests, with different letters indicating significant differences between the means.

Figure 5: Rmt and NPmt. Net photosynthesis and respiration of organisms grown at elevated temperature (15°C), but measured back at the standard, control temperature ($T_{15,5}$) from the beginning (black bars) and the end of the experiment (grey bars). Shown are mean values $\pm$ standard deviation and results from post hoc tests, with different letters indicating significant differences between the means.

Figure 6: Microscopic comparison of lichen morphology. Pictures on the left show cross-sections of the lichen before the incubation (a: *S. alpinum*, c: *U. aurantiaco-atra*, f: *P. contortuplicata*). On the right side, pictures from after the 15°C incubation. b: *S. alpinum* the green algal photobiont in vivid green colour, d: trebuxioid photobiont in *U. aurantiaco-atra*, e: dead cell material in the photobiont layer of *U. aurantiaco-atra*, g: thalli of *P. contortuplicata* after the treatment.

Figure 1: Study site and lichen species. a: Overview of the vegetation near Juan Carlos 1 Base, on Livingston Island. b: *Stereocaulon alpinum* in natural appearance, Glove as scale. c: *Usnea aurantiaco-atra*. d: *Placopsis contortuplicata*.

306x206mm (300 x 300 DPI)

Figure 2: Active time of lichen samples in the experiment plotted against VPD in the experiment boxes.

157x122mm (300 x 300 DPI)

Figure 3: Changes in net photosynthesis (upper graphs), respiration rates (middle graphs) and NP/R ratio (lower graphs) during incubation at different temperatures. Shown are mean values ($n = 3$) $\pm$ standard deviation of samples treated at different temperatures and measured at their respective treatment temperature. • and black line = controls (5°C) measured at 5 °C assay temperature (C5,5); • and red line = increased (15°C) measured at 15 °C assay temperature ($T_{15,15}$); ▲ and green line = extreme (23°C) measured at 23°C ($T_{23,23}$).

281x211mm (300 x 300 DPI)

Figure 4: Responses to elevated temperature (15°C) for the control group and the 15°C treatment. Slopes from linear equations between $C_{5,5}$ vs. $C_{5,15}$ and $C_{5,5}$ vs. $T_{15,15}$ are compared from the beginning (black bars) and the end of the experiment (grey bars). Data are presented separately for changes in net photosynthesis (a,b,c) and respiration (d,e,f). Shown are mean values ($n = 3$) $\pm$ standard deviation and results from Post hoc tests, with different letters indicating significant differences between the means.

287x153mm (300 x 300 DPI)

Figure 5: Rmt and NPmt. Net photosynthesis and respiration of organisms grown at elevated temperature (15°C), but measured back at the standard, control temperature ($T_{15.5}$) from the beginning (black bars) and the end of the experiment (grey bars). Shown are mean values $\pm$ standard deviation and results from post hoc tests, with different letters indicating significant differences between the means.

105x140mm (300 x 300 DPI)

Figure 6: Microscopic comparison of lichen morphology. Pictures on the left show cross-sections of the lichen before the incubation (a: *S. alpinum*, c: *U. aurantiaco-atra*, f: *P. contortuplicata*). On the right side, pictures from after the 15°C incubation. b: *S. alpinum* the green algal photobiont in vivid green colour, d: trebuxioid photobiont in *U. aurantiaco-atra*, e: dead cell material in the photobiont layer of *U. aurantiaco-atra*, g: thalli of *P. contortuplicata* after the treatment.

448x513mm (300 x 300 DPI)

Table 1: Incubation conditions. Overall climatic conditions during the incubation and the count of hours that lichens were active after a hydration event. Given are mean values $\pm$ standard deviation.

Incubation Setup (code)	Air Temperature (°C)	Humidity when active (%)	Active time after hydration (h)
Control	5.9 $\pm$ 2.6	89.3 $\pm$ 13.9	42.2 $\pm$ 4.3
Increased	14.6 $\pm$ 3.4	80.4 $\pm$ 8.2	35.4 $\pm$ 2.7
Extreme	21.9 $\pm$ 4.3	79.1 $\pm$ 29.3	28.8 $\pm$ 1.9