



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Molecular apocrine tumours in EORTC 10994/BIG 1-00 phase III study: pathological response after neoadjuvant chemotherapy and clinical outcomes

Citation for published version:

Bonnefoi, H, MacGrogan, G, Poncet, C, Iggo, R, Pommeret, F, Grellety, T, Larsimont, D, Becette, V, Kerdraon, O, Bibeau, F, Ghnassia, J-P, Picquenot, J-M, Thomas, J, Tille, J-C, Slaets, L, Bodmer, A, Bergh, J & Cameron, D 2019, 'Molecular apocrine tumours in EORTC 10994/BIG 1-00 phase III study: pathological response after neoadjuvant chemotherapy and clinical outcomes', *British Journal of Cancer*.
<https://doi.org/10.1038/s41416-019-0420-y>

Digital Object Identifier (DOI):

[10.1038/s41416-019-0420-y](https://doi.org/10.1038/s41416-019-0420-y)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

British Journal of Cancer

Publisher Rights Statement:

This is a pre-copyedited, author-produced version of an article accepted for publication in "British Journal of Cancer" following peer review. The version of record "Molecular apocrine tumours in EORTC 10994/BIG 1-00 phase III study: pathological response after neoadjuvant chemotherapy and clinical outcomes." is available online at: DOI: 10.1038/s41416-019-0420-y / <https://www.nature.com/articles/s41416-019-0420-y>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Molecular apocrine tumours in EORTC 10994/BIG 1-00 phase III study: pathological response after neoadjuvant chemotherapy and clinical outcomes

Running title: Molecular apocrine breast cancers

Hervé Bonnefoi (1)

Gaetan MacGrogan (2)

Coralie Poncet (3)

Richard Iggo (4)

Fanny Pommeret (1)

Thomas Grellety (1)

Denis Larsimont (5)

Véronique Bécette (6)

Olivier Kerdraon (7)

Frédéric Bibeau (8)

Jean-Pierre Ghnassia (9)

Jean-Michel Picquenot (10)

Jeremy Thomas (11)

Jean-Christophe Tille (12)

Leen Slaets (3)

Alexandre Bodmer (13, 14)

Jonas Bergh (15, 16)

David Cameron (17,18)

and on behalf of the EORTC 10994/BIG 1-00 study investigators

Affiliations

- (1) Department of Medical Oncology, Institut Bergonié Unicancer, Univ. Bordeaux, INSERM U1218, INSERM CIC1401, Bordeaux, France
- (2) Department of BioPathology, Institut Bergonié Unicancer, INSERM U1218, Bordeaux, France
- (3) European Organisation for Research and Treatment of Cancer (EORTC) Headquarters, Brussels, Belgium
- (4) Institut Bergonié Unicancer, INSERM U1218, Bordeaux, France
- (5) Department of Pathology, Institut Jules Bordet, Brussels, Belgium
- (6) Department of Pathology, Institut Curie - Hôpital René Huguenin, Saint-Cloud, France
- (7) Department of Pathology, Centre René Gauducheau, Institut de Cancérologie de l'Ouest, Nantes, France
- (8) Department of Pathology, Institut de Cancérologie de Montpellier (ICM), Montpellier, France
- (9) Department of Pathology, Centre Paul Strauss, Strasbourg, France
- (10) Department of Pathology, Centre Henri Becquerel, Rouen, France
- (11) Department of Pathology, Edinburgh Cancer Centre, University of Edinburgh, Edinburgh, United Kingdom
- (12) Department of Pathologie, Hôpitaux Universitaires de Genève (HUG), Geneva, Switzerland
- (13) Swiss Group for Clinical Cancer Research (SAKK), Bern
- (14) Department of Oncology, Hôpitaux Universitaires de Genève (HUG), Geneva, Switzerland
- (15) Swedish Breast Cancer Group (SweBCG), Stockholm
- (16) Department of Oncology and Pathology, Karolinska Institutet, Stockholm, Sweden
- (17) Anglo-Celtic Cooperative Oncology Group (ACCOG), Edinburgh
- (18) Department of Medical Oncology, Edinburgh Cancer Centre, University of Edinburgh, Edinburgh, United Kingdom

Corresponding author

Prof Hervé Bonnefoi

Department of Medical Oncology, Institut Bergonié Unicancer, Univ. Bordeaux, INSERM

U1218, INSERM CIC1401, 229 Cours de l'Argonne, F-33000 Bordeaux, France.

Email: h.bonnefoi@bordeaux.unicancer.fr

Phone: +33 556 33 32 69 Fax: +33 556 33 33 30

Abstract

Background: We explored, within EORTC10994 study, the outcomes for patients with molecular apocrine (MA) breast cancer, defined immunohistochemically (IHC) as androgen-receptor (AR) positive, oestrogen (ER) and progesterone (PR) negative. We also assessed the concordance between IHC and gene expression arrays (GEA) in the identification of MA cancers.

Methods: Centrally-assessed biopsies for AR, ER, PR, HER2 and Ki-67 by IHC were classified into 6 subtypes: MA, triple negative (TN) basal-like, luminal A, luminal B HER2-negative, luminal B HER2-positive, and "other". The two main objectives were the pathological complete response (pCR) rates and survival outcomes in the overall MA subtype (and further divided by HER2 status) and the remaining 5 subtypes.

Results: IHC subtyping was obtained in 846 eligible patients. Ninety-three (11%) tumours were classified as MA subtype. Both IHC and GEA data were available for 64 patients. In this subset, IHC concordance was 88.3% in identifying MA tumours compared to GEA. Within the MA subtype, pCR was observed in 33.3% (95%CI: 29.4-43.9) and the 5-year recurrence-free interval was 59.2% (95%CI: 48.2-68.6). Patients with MA and TN basal-like tumours have lower survival outcomes.

Conclusions: Irrespective of their HER2 status, the prognosis for MA tumours remains poor and adjuvant trials evaluating anti-androgens should be considered.

Key words

Breast cancer, neoadjuvant chemotherapy, androgen receptor, molecular apocrine cancer, triple negative breast cancer

Introduction

Several gene expression array (GEA) studies have identified a breast cancer subtype characterized by the expression of the androgen receptor (AR), absence of the oestrogen receptor α (ER), and expression of many genes that are expressed in ER-positive luminal tumours (Doane AS, 2006; Farmer et al, 2005; Guedj et al, 2012). We named these tumours "molecular apocrine" (MA) as they have an increased androgen signalling expression profile and some, but not all, morphological hallmarks of apocrine tumours (Farmer et al, 2005). In approximately two-thirds of the cases, these tumours are human epidermal growth factor receptor 2 (HER2) positive; the importance of this HER2-positive AR-driven group of tumours has been recently highlighted (Daemen & Manning, 2018). The remainder are HER2-negative and are part of the heterogeneous triple negative breast cancer (TNBC) group.

In both HER2-positive and HER2-negative subgroups of MA tumours, prospective trials evaluating anti-androgens in patients with advanced breast cancer are ongoing. In these trials MA tumours are identified using an immunohistochemical (IHC) definition. In the HER2-negative sub-group, three prospective clinical trials demonstrated anti-tumour efficacy with anti-androgen treatment (Bonnefoi et al, 2016; Gucalp et al, 2013; Traina et al, 2018) and with long-term responders (Grellety et al, 2018). In the HER2-positive sub-group, encouraging preliminary results have been reported from a Simon 2-stage phase two study (Krop et al, 2016). These data have reinforced the interest in the MA subtype and the logical next step would be to evaluate these anti-androgen treatments in patients with MA early breast cancer, at least in the HER2-negative group.

Before considering adjuvant studies, there is a need to better understand the frequency of the MA subtype and its natural history. Previously published EORTC 10994/ BIG 1-00 study (Bonnefoi et al, 2011) offered an excellent opportunity to explore the outcomes for patients with MA tumours compared to other subtypes using an IHC definition. MA tumours were

identified using the following definition: AR-positive and ER-, progesterone receptor (PR)-negatives. Moreover, we categorized MA tumours into two sub-groups according to HER2 status. This IHC definition is commonly used to identify MA subtype in prospective therapeutic clinical trials in advanced breast cancer (Bonnetfoi et al, 2016; Gucaip et al, 2013; Traina et al, 2018). Other subtypes were defined in a similar way to the St Gallen 2011 simplified classification (Goldhirsch et al, 2011) with the exception of the basal-like subtype, which was by definition, AR negative (quadruple negative) in this study.

MA subtype was initially identified using GEA (Doane AS, 2006; Farmer et al, 2005; Guedj et al, 2012). For pragmatic reasons IHC is used to identify this subtype in prospective therapeutic trials. However the agreement between these two methods has never to our knowledge been assessed. Thus we determined in a subset of patients included in this sub-study, the concordance of IHC compared to this GEA classification in the identification of MA tumours. We used a biologically-based GEA classification of breast cancer recently developed to identify MA tumours (Iggo, 2018).

Materials and methods

Study design, eligibility and treatment

This was an unplanned analysis within the EORTC 10994/BIG 1-00 neoadjuvant phase III trial, in which 1856 patients were randomized in a 1:1 ratio between six cycles of fluorouracil, epirubicin, cyclophosphamide and a taxane-based regimen, docetaxel for three cycles followed by epirubicin+ docetaxel for three cycles, all administered prior to primary surgery as previously described (Bonnetfoi et al, 2011). Two frozen biopsies from the primary tumour were mandatory for research purposes. Formalin fixed paraffin embedded (FFPE) biopsies were performed for diagnostic purposes. Eligible patients for the EORTC 10994/BIG 1-00 trial were women <71 years with histologically-proven invasive breast cancer suitable for neoadjuvant

chemotherapy, with any large operable or locally advanced/inflammatory breast cancer. At completion of chemotherapy, locoregional treatment was planned in accordance with the guidelines described in the protocol. Treatment was completed with hormonal therapy according to each centre's policy. Patients with HER2-positive tumours were allowed to participate in adjuvant clinical trials assessing trastuzumab or to receive this treatment in the adjuvant setting once it became standard practice, but none received neoadjuvant trastuzumab. The trial was registered with ClinicalTrials.gov number NCT00017095 and approved by national and/or local ethics committees in all participating centres. Before registration, all patients signed an informed consent for the trial and for mandatory p53 gene assessment on tumour samples. In addition patients were asked to consent for optional biological research on their tumour samples.

For the sub-study that is the subject of this report, a subgroup of the initial population of 1856 patients was selected based on the following criteria: (i) patients eligible for the main EORTC 10994/BIG 1-00 trial; (ii) patients who received at least one cycle of neo-adjuvant chemotherapy and who did not receive radiotherapy before surgery; (iii) patients who agreed to consent for optional biological research on their tumour samples; (iv) patients with sufficient tumour in their pre-treatment core biopsies and whose tumour subtype was identified based on the central analysis of their biopsies included in the ancillary tissue microarray (TMA) study.

Histopathological assessment

Histological type and grade were assessed locally by pathologists at each participating centre and the data collected on case report forms in the context of the EORTC 10994 trial. Pathological response was assessed by local pathologists after completion of the neoadjuvant chemotherapy. No central pathology review was performed either for histological type and grade at diagnosis or pathological response at surgery.

Construction of tissue microarrays

Breast cancer FFPE core biopsies taken at diagnosis before neoadjuvant chemotherapy were retrospectively collected and sent to Institut Bergonié by different participating centres. All core biopsies were reviewed on H and E-stained sections, and representative tumour areas were selected for TMA construction. For each case, three 0.6 mm-diameter tumour cores were used. The TMA was constructed using a tissue micro-arrayer (Alphelys France). Evaluation of the entire section was performed by a board certified pathologist (GMG).

Immunohistochemical and dual detection in situ hybridisation methods and interpretation

Tumour phenotype concerning AR, ER, PR, HER2 status, and proliferation status (Ki-67) were defined on TMA. AR was scored as positive if $\geq 10\%$ of tumour cell nuclei showed a positive signal. This is the commonly used cut-off (Bonnetfoi et al, 2016; Gucaip et al, 2013; Traina et al, 2018). ER and PR were scored negative if $< 1\%$ of tumour cells were positive. For Ki-67, results were given by % of positive cells. The threshold used to define high Ki-67 expression was $\geq 14\%$ (Cheang et al, 2009). For HER2, results were given by % of positive cells and intensity of staining. Final HER2 status was scored according to the ASCO/CAP recommendations (Wolff et al, 2013). An IHC3+ score was considered positive. An IHC2+ score was considered equivocal. It was then retested by silver in situ hybridisation (SISH). Cases with ≥ 6 *HER2* copies per cell nucleus were considered positive. Details for IHC staining of ER, PR, HER2, Ki67 and AR are provided in the supplementary table 1.

Simplified breast cancer molecular subtypes classification

Tumours were classified into 6 subtypes: MA, triple negative basal-like (as named in the first gene expression arrays classification) (Perou et al, 2000), luminal A, luminal B HER2-negative, luminal B HER2-positive, and non-luminal non-MA HER2-positive. This classification is detailed in table 1. The MA subtype was further divided in two subgroups according to HER2 status: positive or negative. The luminal group (ER and/or PR positive, any HER2 status, any Ki67) was further divided in two subgroups according to AR status: positive or negative.

TP53 status

TP53 status from frozen biopsies was assessed using a yeast functional test as previously described (Flaman et al, 1995; Waridel et al, 1997).

Gene expression array analysis

Microarray data from our previous studies (Farmer et al, 2009; Farmer et al, 2005) was downloaded from the NCBI GEO database using accession numbers GSE1561 and GSE6861. A biology-based classification of breast cancer was developed using a mammary lineage model (supplementary figure 1) (Iggo, 2018). The first step in the classification splits tumours into hormone sensing tumours (10 transcripts including ESR1, AR, FOXA1) and secretory cell tumours (9 transcripts including ELF5, FOXC1, KLF5). The second step splits hormone sensing tumours into classic ER+ luminal tumours and molecular apocrine tumours. To separate luminal from molecular apocrine tumours, 30 genes were selected based on correlation with ESR1 expression, half showing positive (luminal) and half negative (molecular apocrine) correlation. Using these 30 genes luminal and apocrine scores were created. This classification uses a total of 49 preselected transcripts (supplementary figure 1). Of note ERBB2 is not in the list. The tumours were assigned to luminal, molecular apocrine and basal (LAB) classes as described (Iggo, 2018). The LAB classification includes a fourth category, "unknown" or "non-interpretable", for tumours that are too close to the thresholds separating classes to be assigned any particular class confidently. Full details of this classification can be found in the publication (Iggo, 2018).

Objectives and end-points definitions

The two main objectives were to describe the pathological complete response (pCR) rates and to report the survival outcome measures, recurrence-free interval (RFI), distant recurrence-free interval (DRFI) and overall survival (OS): (i) in the MA subtype (in the overall MA population and in the two subgroups according to HER2 status); (ii) in the remaining 5

subtypes, (iii) within the luminal group (3 subtypes: luminal A, luminal B HER2 negative, luminal B HER2 positive), in the two subgroups according to AR status (any HER2 and Ki67 status). As an additional aim we also assessed in a subset of patients the agreement of IHC to identify MA cancers compared to the gold standard GEA.

pCR was defined as no evidence of residual invasive cancer (or very few scattered tumour cells left) with or without residual ductal carcinoma in situ (DCIS) and negative axillary lymph nodes (ypT0/is ypN0). Patients whose tumour progressed on neoadjuvant chemotherapy and patients who did not undergo surgery or with missing information on the surgical pathology report were considered as having "no pCR".

The survival endpoints were defined according to the standardised definitions for efficacy endpoints (STEEP) system (Hudis et al, 2007). RFI was measured as time from randomization to progression on chemotherapy, ipsilateral invasive breast (local) recurrence, regional recurrence (chest wall and regional nodes: axillary, internal mammary, infraclavicular, and supraclavicular nodes), distant recurrence or death due to breast cancer and/or treatment toxicity, whichever came first. DRFI was calculated as the time from randomization to distant recurrence or death due to breast cancer and/or treatment toxicity, whichever came first. OS was calculated as the time from randomization to death due to any cause. In the EORTC 10994/BIG 1-00 trial, both first locoregional recurrence and first distant metastasis were registered. Events diagnosed within two months were considered as simultaneous and we chose to declare the site of first event as the one with the worst prognosis. Patients who did not present with any of the events mentioned above during their follow-up were censored at the time of their last follow-up. Contralateral breast cancer and second primary invasive cancer (non-breast) were not considered as primary events.

Statistical analysis

A statistical analysis plan was prospectively defined. All the statistical analyses were performed using SAS version 9.4 (SAS Institute Inc., Cary, NC, USA).

pCR analysis

Logistic regression model was used to estimate the effect of subtype on the odds of having a pCR. The associated exact 95% Clopper-Pearson confidence interval and p-value based on the Wald method were also presented. Three logistic models were conducted: (i) comparing the six simplified subtypes using the luminal A as reference group, (ii) within the MA subtype, comparing HER2 positive to HER2 negative subgroups, (iii) within the luminal group (3 subtypes as mentioned before), comparing AR-positive to AR-negative subgroups.

Survival outcomes

Time to event end-points were analysed per Kaplan Meier method reporting 5-year estimate and corresponding Kaplan-Meier curve. P-values were based on the logrank test. Hazard ratios were estimated from a Cox proportional hazards model and the corresponding 95% confidence intervals (CI) (Wald method) were added.

Concordance between IHC and GEA subtype classification

In the subset of patients with both subtype classifications based on IHC or GEA methods, the proportion of concordant subtype classification and Kappa agreement coefficient as well as their 95% confidence intervals were estimated.

Results

Of the 1856 patients originally randomized, core biopsies of 1092 eligible patients were centralized in Bordeaux and available for the TMA construct. A total of 846 patients with a tumour classified in one of the 6 IHC-based subtypes (table 1) based on this TMA were included in this sub-study. The reasons for ineligibility are shown in the Consort diagram (supplementary figure 2). Baseline characteristics and treatment of patients included in this analysis (eligible) and those excluded are presented on supplementary table 2 (significant p-

values below 0.05 are indicated). The median follow-up of the patients included in this sub-study was 56 months from the date of randomization. We will first describe the MA population.

Molecular apocrine tumours

A total of 93/846 (11.0%) eligible tumours were classified in the MA subtype. Baseline characteristics and treatment are reported in supplementary table 4. Median age was 54.1 years. Ki67 was high in 81.5% (75/92) and TP53 status was mutated in 72.1% (49/68) patients. Approximately one-third of MA tumours were HER2-negative (32/92) and two thirds were HER2-positive (59/92) and one case was equivocal. MA HER2-negative tumours represented 25.4% of all triple negative breast cancers (32/126). Patient and tumour characteristics were compared between HER2-positive and HER2-negative groups. MA HER2-positive tumours presented more frequently with a high Ki67. All the other characteristics were similar between the two groups except nodal status and Ki67.

A pCR was observed in 31 of 93 (33.3%, [95%CI: 23.9-43.9]) patients with MA tumours (table 2). pCR rates were not significantly different between HER2-negative and positive subgroups (Odds ratio HER2-positive versus HER2-negative 1.31 ([95% CI: 0.51-3.36]; p =0.57).

The RFI, DRFI and OS curves are shown in figure 1 and supplementary figures 3 and 4. The 5-year estimate of RFI rate was 59.2% (95%CI: 48.2-68.6) (figure 1 and supplementary table 3). Within the MA subtype, survival outcomes measures were not statistically different between HER2-positive and negative sub-groups (figure 1 and supplementary table 5). Approximately one-third of first events in MA cancers were loco-regional recurrences (supplementary table 6). Patterns of distant relapses are reported in supplementary table 7.

Other molecular subtypes and comparison with molecular apocrine subtype

Baseline characteristics and treatment are reported in supplementary table 9. Patients with MA tumours when compared with other molecular subtypes were older and more often postmenopausal.

pCR rates differed significantly ($p < 0.001$) across intrinsic subtypes, with the lowest rate for luminal A (8.7%, [95%CI: 5.3-13.2]) and the highest rates for MA and triple negative basal-like (33.3%, [95%CI: 23.9-43.9] and 34.0%, [95%CI: 24.6-44.5], respectively) (table 2). The pCR rate of HER2-positive non-luminal and non-MA tumours was high (42.9%, [95%CI: 10.0-81.6]) but the number of patients in this group is very small.

Patients with MA and triple negative basal-like tumours showed the lowest 5-year RFI, DRFI and OS estimates (figure 2, supplementary table 3, supplementary figures 5 and 6).

Within luminal subtypes, 8.6% (luminal A) to 18.2% (luminal B HER2-negative) experienced loco-regional recurrence as first event contributing to RFI. In non-luminal subtypes including MA tumours, one-third of patients experienced loco-regional recurrence as first event contributing to RFI (supplementary table 6).

Patterns of distant relapses by simplified breast cancer subtypes are reported in supplementary table 7. Compared with patients with luminal tumours, patients with MA tumours presented more often with visceral metastasis ($p = 0.0343$) and less often with bone metastasis ($p = 0.0006$) (supplementary table 8).

Analysis of the ER and/or PR positive group by AR status

Within the luminal group, 93.7% (599/639) were AR-positive. pCR rates were not statistically different by AR status (Odds ratio 0.62, [95%CI: 0.27-1.39]; $p = 0.242$) (supplementary table 10). RFI, DRFI and OS were not statistically different by AR status (supplementary table 11).

Concordance of immunohistochemistry with gene expression to identify molecular apocrine cancers

We compared the gene expression-based LAB classification (Iggo, 2018) with IHC classification in a subset of 64 patients for whom GEA and IHC data were available. Note that the selection of cases for GEA was not random: it was enriched for ER-negative tumours. By definition the LAB classification splits tumours into only three groups: luminal, MA and basal-like (table 1). Hence, to compare it to IHC classification we assigned cases to three IHC groups (luminal, MA and basal-like) based on IHC for ER, PR and AR. In four patients the gene expression values were too close to the thresholds for the tumours to be assigned confidently to any particular group, leaving 60 samples for comparison with the IHC data. The concordance was 88.3% (95%CI: 80.2-96.5). The Kappa agreement coefficient between IHC and GEA methods to identify the LAB MA subtype was 0.82 (95%CI: 0.694-0.945) (table 3).

HER2 is not used in the LAB classification because it is commonly expressed by both luminal and MA tumours (our hypothesis is that HER2 promotes apocrine metaplasia of luminal cells leading to a high frequency of HER2 amplification in the MA group). To illustrate the potential limitations of using HER2 to identify MA tumours we plotted HER2 against ESR1 in figure 3A (the tumours are labelled according to the LAB classification in three molecular groups). Tumours expressing high levels of HER2 were indeed classified as MA but several tumours with high levels of HER2 were luminal (upper right quadrant) or even basal-like, and one MA tumour expressed a low level of HER2. Figure 3B shows the distribution of AR and ESR1 expression in the three molecular groups. The tumours fall into the three expected groups: MA tumours (AR high and ESR1 low) in the upper left quadrant, basal-like tumours (AR low and ESR1 low) in the lower left quadrant, and luminal tumours (AR high and ESR1 high) in the upper right quadrant.

Supplementary figures 7 and 8 highlight the seven discordant cases between the IHC and GEA classifications. Five discordant cases lie close to the thresholds separating the tumour types and can readily be explained by slightly differing placement of the thresholds by the two approaches. For example, tumours 250 and 337 are classified as MA by gene expression but

luminal by IHC; they express AR well but they also express ESR1 at a level almost exactly at the cut-off separating luminal from MA tumours (supplementary figure 8). The two remaining discordant cases are outliers (tumours 335 and 856). In these two cases possible explanations for the discordance are, post-transcriptional modifications, tumour heterogeneity or even a sample labelling issue.

In summary, the overall agreement between classification by IHC and gene expression was good, with the disagreements concentrated near the thresholds.

Moreover we compared LAB classification and a simplified PAM50 classification (Parker et al, 2009). We used 43 genes out of the 50 genes which constitute PAM50 successfully mapped to the Affymetrix dataset (7 genes were not present on the U133A chip). We excluded from the comparison tumours which were incomparable (classified as Normal by PAM or unknown by LAB) leaving 59 tumours to compare. There was a perfect agreement between the two classifications for the basal tumours. Of note 89.3% (25/28) of those classified as luminal by PAM were classified as luminal by LAB; three luminal tumours by PAM were MA by LAB. Moreover 78.6% (11/14) of those classified as molecular apocrine by LAB were classified as HER2-enriched by PAM (supplementary table 12).

Discussion

With a total of 93 MA cancer patients, this is the largest series from a prospective neoadjuvant trial assessing clinicopathological characteristics, frequency, chemosensitivity and prognosis of this subtype.

In this series, patients diagnosed with MA were older (median age 54.1) and were more often postmenopausal (62.4%) compared to other subtypes. One-third of the first relapses were locoregional. This proportion is similar to that observed in triple negative basal-like subtype. Two-thirds of MA tumours were HER2-positive and the remainder, HER2-negative. TP53

mutation rate was high (72.1%) and was similar to the one observed in triple negative basal-like cancers (73.3%).

Regarding frequency, 11% of cancers were classified in the MA subtype. Within the TNBC group, approximately one-quarter were MA (32/126). This information is potentially important when estimating the feasibility of a prospective trial in this molecular subtype in early breast cancer particularly in the HER2 negative subgroup where no targeted therapy can be offered. In the literature, based on IHC, the frequency of MA tumours in the TNBC group ranges from 21.6% (24/111) in the GBG Gepartrio sub-study (Loibl et al, 2011) to 35.9% (122/339) in the Nurses' Health study (Collins et al, 2011). Based on GEA, using the TNBCtype classification, the frequency of MA tumours is 11.1% (65/587) in a first analysis of 21 publicly available breast cancer GEA data sets performed by the Vanderbilt University group (Lehmann et al, 2011). In a second Vanderbilt University analysis, the authors simplified their classification from six into four subtypes (TNBCtype-4). Using this refined TNBCtype-4 classification, the frequency is 16% (50/316) in a second analysis performed by this group combining 5 publicly available GEA data sets of patients treated with neoadjuvant chemotherapy (Lehmann et al, 2016). Using a different classification algorithm, the frequency of MA is 17.7% (35/198) in a series of tumours collected from U.S. and European sites, with IHC triple negative status centrally reviewed and GEA analysed in Houston (Burstein et al, 2015).

The difference in the frequency of AR-positive TNBCs, whether IHC or GEA was used, is difficult to explain given that there is 88.3% concordance between IHC and GEA to identify MA subtype in our EORTC series. We believe that this high concordance rate validates the IHC approach taken in this study. Although not perfect, IHC has the advantage of being inexpensive and routinely available in diagnostic histopathology departments. In addition, IHC is commonly used to identify MA cancers in therapeutic trials assessing anti-androgen treatments from which a significant proportion of patients benefited (Bonnetfoi et al, 2016; Gucalp et al, 2013; Traina et al, 2018). We suspect, however, that real progress in the identification of these

tumours will come not from analysis of arbitrary signatures or IHC profiles but rather from a deeper understanding of the underlying biological entity so we can devise tests that identify that entity on the basis of its essential properties. It was for this reason that we developed the LAB classification (Iggo 2018).

There are few data reported in the literature regarding chemosensitivity of MA tumours in particular, in the HER2-negative subgroup. In the GBG series, the authors used IHC to identify MA tumours. In the TNBC group (n=111), the pCR rates of AR-positive (n=24) and negative (n=87) tumours were similar, 29.2% and 33.3%, respectively (Loibl et al, 2011). In our series, within the TNBC group (n=126), the pCR rates of AR-positive (n=32) and negative (n=94) tumours were also similar. In the first Vanderbilt University analysis of TNBC, a total of 42 patients included in 2 trials received neoadjuvant chemotherapy (Lehmann et al, 2011). The pCR rates were 14.3% (1/7) and 63.2% (12/19) in MA named luminal AR and basal-like subtypes, respectively (Lehmann et al, 2011). The MD Anderson Cancer Center (MDACC) group used a similar approach with GEA in a series of 130 evaluable patients with TNBC treated with neoadjuvant chemotherapy (Masuda et al, 2013). In luminal AR, basal-like 1 and basal-like 2 subtypes, the pCR rates were 10% (2/20), 52.4% (11/21) and 0% (0/8), respectively. In the second Vanderbilt University analysis, the authors used their simplified TNBCtype-4 classification and assessed the pCR rates in each subtype using data from 4 publicly available GEA data sets (including the MDACC cohort) corresponding to a total of 306 patients with TNBC (Lehmann et al, 2016). In this publication the pCR rate in the luminal AR was 29% (15/52) which is similar to the results observed in IHC series. It is difficult to explain these apparently different results in pCR rates observed whether IHC or GEA with TNBCtype or GEA with TNBCtype-4 classifications were used to identify MA tumours. Our interpretation is that, as shown in the LAB classification (Iggo, 2018), it is easy to separate basal-like from luminal and MA tumours by gene expression but far more difficult to differentiate luminal and MA tumours. Hence, a possible explanation for the divergent results in pCR rates is that, when using the initial TNBCtype classification tool in the Vanderbilt University and MD Anderson

studies (Lehmann et al, 2011; Masuda et al, 2013) a low pCR rate in the LAR group was observed because they may have included within this group some classic ER-positive luminal tumours, a subgroup known to have lower pCR rates. By their own admission, the Vanderbilt group acknowledge that they included from 55 to 82% of luminal A or B tumours (identified using the published intrinsic 306-gene set or the PAM50) in the LAR group (Lehmann et al, 2011; Lehmann et al, 2016).

In our series, patients with MA and triple negative basal-like tumours had the worst outcome. Although the distribution of recurrences during the first 3 years was very similar in the 2 groups, it becomes different after 3 years. In the triple negative basal-like group (n=94) a plateau was observed but not in the MA group (n=93) (figure 2). This plateau is a classic observation in basal-like series. For example, in a French study using GEA, the authors applied their molecular subtype classifier model to a large Affymetrix validation set comprising 2291 breast cancers. On the metastasis-free survival curves a plateau was observed in the basal-like group (n=264) but not in the MA group (n=146). More than 40% of patients with MA tumours relapsed within 5 years and survival outcomes were not statistically different between HER2-negative and positive sub-groups. However a numerical difference was observed for patients with MA HER2-positive tumours. This difference could be explained by the fact that only 1/3 of patients received adjuvant trastuzumab (EORTC10994/BIG 1-00 accrual period extended from April 2001 to November 2006; Herceptin became standard practice at the end of 2005 which explains why only 1/3 of patients received this treatment). As far as MA HER2-negative tumours are concerned, the risk of relapse at 5 years in the EORTC study was more than one-third (34.5%). In the two Vanderbilt publications, using GEA based classifications for TNBCs, the risk of relapse at 5 years of MA cancers was 50% in both series of 62 and 50 MA tumours (Lehmann et al, 2011; Lehmann et al, 2016).

Our study has some strengths and limitations. This is the largest series from a prospective trial assessing the frequency and the prognosis of MA tumours, in particular, in TNBCs. The main

weakness is that only 45.6% of patients included in EORTC 10994 study were included in this sub-study. In EORTC study two frozen biopsies were mandatory but we did not plan to prospectively collect FFPE blocks prior to neoadjuvant chemotherapy. Hence, these samples were collected retrospectively. However, the characteristics of patients included in this sub-study and those who were excluded were similar (supplementary table 2). In addition there was no central assessment of pathological response. Lastly we did not use a transcriptomic signature to identify MA tumours and other subtypes. However, as mentioned before, both IHC and GEA methods have high concordance in the identification of MA tumours (88.3%).

In conclusion, this study demonstrates that the prognosis of MA breast cancers is very poor despite their acceptable rate of pCR after neoadjuvant chemotherapy. Moreover the MA subtype is frequent, representing approximately 11% of all breast cancers and 25% within the TNBC group. This specific molecular subtype should be considered as an unmet need particularly in the HER2 negative subgroup where no targeted therapy can be offered. In the advanced setting, three clinical trials in patients with AR-positive TNBCs have demonstrated efficacy of anti-androgen treatments (Bonnefoi et al, 2016; Gucalp et al, 2013; Traina et al, 2018). Based on the data reported in these publications anti-androgen treatments should be evaluated in the adjuvant setting in patients with AR-positive TNBCs.

Additional information

Ethics approval and consent to participate

The trial was registered with ClinicalTrials.gov number NCT00017095 and approved by national and/or local ethics committees in all participating centres. Before registration, all patients signed an informed consent for the trial and for mandatory p53 gene assessment on tumour samples. In addition patients were asked to consent for optional biological research on their tumour samples. The study was performed in accordance with the Declaration of Helsinki.

Availability of data and material

Data can be accessed through the EORTC data sharing platform. Data request form is available on EORTC website: <http://www.eortc.org/data-sharing/>

Conflict of interest

The authors declare no conflict of interest.

Funding

EORTC sponsored, designed and coordinated the trial. The funding sources had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the study data and the final responsibility for the decision to submit for publication. Trial design, conduct, and analysis were done at the EORTC headquarters independently from all funding bodies.

Authorship

HB, LS, DC conceived and designed the work, acquired data and played an important role in interpreting the results. GMG, CP, RI designed the work, acquired data and played an important role in interpreting the results. FP, TG designed the work and acquired data. DL, VB, OK, FB, JPG, JMP, JT, JCT, AT, JB acquired data. All authors drafted or revised the manuscript, approved the final version and agreed to be accountable for all aspects of the work.

Acknowledgements

We thank the patients, doctors and nurses involved in the EORTC 10994/BIG 1-00 study for their generous participation. We thank Valérie Velasco for technical assistance in constructing the TMA. We also thank the data managers from the EORTC, the Anglo- Celtic Cooperative Oncology Group (ACCOG) at the Information and Statistics Division of the Scottish NHS, the Swedish Breast Cancer Group (SweBCG) and the Swiss Group for Clinical Cancer Research

(SAKK). We thank the Fondation pour la lutte contre le cancer et pour des recherches medico-biologiques and SIRIC BRIO (Site de Recherche Intégrée sur le Cancer – Bordeaux Recherche Intégrée Oncologie) for financial support [Grant INCa-DGOS-Inserm 6046]. The authors would also like to thank Ravi Nookala of Institut Bergonié for medical writing service. This publication was supported by the EORTC Cancer Research Fund.

References

- Bonnefoi H, Grellety T, Tredan O, Saghatchian M, Dalenc F, Mailliez A et al. A phase II trial of abiraterone acetate plus prednisone in patients with triple-negative androgen receptor positive locally advanced or metastatic breast cancer (UCBG 12-1). *Ann Oncol* 2016; **27**: 812-818
- Bonnefoi H, Piccart M, Bogaerts J, Mauriac L, Fumoleau P, Brain E et al. TP53 status for prediction of sensitivity to taxane versus non-taxane neoadjuvant chemotherapy in breast cancer (EORTC 10994/BIG 1-00): a randomised phase 3 trial. *Lancet Oncol* 2011; **12**: 527-539
- Burstein MD, Tsimelzon A, Poage GM, Covington KR, Contreras A, Fuqua SA et al. Comprehensive genomic analysis identifies novel subtypes and targets of triple-negative breast cancer. *Clin Cancer Res* 2015; **21**: 1688-1698
- Cheang MC, Chia SK, Voduc D, Gao D, Leung S, Snider J et al. Ki67 index, HER2 status, and prognosis of patients with luminal B breast cancer. *J Natl Cancer Inst* 2009; **101**: 736-750
- Collins LC, Cole KS, Marotti JD, Hu R, Schnitt SJ, Tamimi RM Androgen receptor expression in breast cancer in relation to molecular phenotype: results from the Nurses' Health Study. *Mod Pathol* 2011; **24**: 924-931
- Daemen A, Manning G. HER2 is not a cancer subtype but rather a pan-cancer event and is highly enriched in AR-driven breast tumors. *Breast Cancer Res* 2018; **20**: 8
- Doane AS, Danso M, Lal P, Donaton M, Zhang L, Hudis C et al. An estrogen receptor-negative breast cancer subset characterized by a hormonally regulated transcriptional program and response to androgen. *Oncogene* 2006; **25**: 1-15
- Farmer P, Bonnefoi H, Anderle P, Cameron D, Wirapati P, Becette V et al. A stroma-related gene signature predicts resistance to neoadjuvant chemotherapy in breast cancer. *Nat Med* 2009; **15**: 68-74
- Farmer P, Bonnefoi H, Becette V, Tubiana-Hulin M, Fumoleau P, Larsimont D et al. Identification of molecular apocrine breast tumours by microarray analysis. *Oncogene* 2005; **24**: 4660-4671
- Flaman JM, Frebourg T, Moreau V. A simple p53 functional assay for screening cell lines, blood, and tumors. *Proc Natl Acad of Sci USA* 1995; **92**: 3963-3967
- Goldhirsch A, Wood WC, Coates AS, Gelber RD, Thurlimann B, Senn HJ. Strategies for subtypes--dealing with the diversity of breast cancer: highlights of the St. Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2011. *Ann Oncol* 2011; **22**: 1736-1747
- Grellety T, MacGrogan G, Chakiba C, Kind M, Bonnefoi H. Long term complete response of an androgen receptor-positive triple-negative metastatic breast cancer to abiraterone acetate. *JCO Precision Oncology* 2018 doi: 10.1200/PO.17.00223
- Gucalp A, Tolaney S, Isakoff SJ, Ingle JN, Liu MC, Carey LA et al. Phase II trial of bicalutamide in patients with androgen receptor-positive, estrogen receptor-negative metastatic Breast Cancer. *Clin Cancer Res* 2013; **19**: 5505-5512

Guedj M, Marisa L, de Reynies A, Orsetti B, Schiappa R, Bibeau F et al. A refined molecular taxonomy of breast cancer. *Oncogene* 2012; **31**: 1196-1206

Hudis CA, Barlow WE, Costantino JP, Gray RJ, Pritchard KI, Chapman JA et al. Proposal for standardized definitions for efficacy end points in adjuvant breast cancer trials: the STEEP system. *J Clin Oncol* 2007; **25**: 2127-2132

Iggo R. Classification of breast tumours into molecular apocrine, luminal and basal groups based on an explicit biological model. *BioRxiv* 2018 dx.doi.org/10.1101/270975

Krop I, Cortes J, Miller K, Huizing MT, Provencher L, Gianni L, Chan S et al. A single-arm phase 2 study to assess clinical activity, efficacy and safety of enzalutamide with trastuzumab in HER2+ AR+ metastatic or locally advanced breast cancer. *Breast Cancer Res (supplement)* 2017; **77**: 4, abstract P4-22-08

Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y et al. Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest* 2011; **121**: 2750-2767

Lehmann BD, Jovanovic B, Chen X, Estrada M, Johnson K, Shyr Y et al. Refinement of Triple-Negative Breast Cancer Molecular Subtypes: Implications for Neoadjuvant Chemotherapy Selection. *PLoS One* 2016; **11**: e0157368.

Loibl S, Muller BM, von Minckwitz G, Schwabe M, Roller M, Darb-Esfahani S et al. Androgen receptor expression in primary breast cancer and its predictive and prognostic value in patients treated with neoadjuvant chemotherapy. *Breast Cancer Res Treat* 2011; **130**: 477-487

Masuda H, Baggerly KA, Wang Y, Zhang Y, Gonzalez-Angulo AM, Meric-Bernstam F et al. Differential response to neoadjuvant chemotherapy among 7 triple-negative breast cancer molecular subtypes. *Clin Cancer Res* 2013; **19**: 5533-5540

Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol* 2009; **27**: 1160-1167

Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA et al. Molecular portraits of human breast tumours. *Nature* 2000; **406**: 747-752

Traina TA, Miller K, Yardley DA, Eakle J, Schwartzberg LS, O'Shaughnessy J et al. Enzalutamide for the Treatment of Androgen Receptor-Expressing Triple-Negative Breast Cancer. *J Clin Oncol* 2018; **36**: 884-890

Waridel F, Estreicher A, Bron L, Flaman JM, Fontolliet C, Monnier P et al. Field cancerisation and polyclonal p53 mutation in the upper aero-digestive tract. *Oncogene* 1997; **14**: 163-169

Wolff AC, Hammond ME, Hicks DG, Dowsett M, McShane LM, Allison KH et al. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *J Clin Oncol* 2013; **31**: 3997-4013

Figures titles and legends

Figure 1. Recurrence-free interval in the molecular apocrine subtype (any HER2 status, HER2 positive and HER2 negative subgroups)

Abbreviations:

HR: hazard ratio; CI: confidence interval; MA: molecular apocrine; HER2-: human epidermal growth factor receptor 2 negative; HER2+: human epidermal growth factor receptor 2 positive.

Figure 2. Recurrence-free interval in the six subtypes

Abbreviations:

HR: hazard ratio; CI: confidence interval; lum A: luminal A; lum B: luminal B; HER2-: human epidermal growth factor receptor 2 negative; HER2+: human epidermal growth factor receptor 2 positive; MA: molecular apocrine; TN: triple negative.

Figure 3. HER2, AR and ESR1 gene expression of individual tumours labelled by LAB class. A, HER2 and ESR1; B, AR and ESR1.

Legend:

The points are coloured according to the LAB classification. The gene expression units are arbitrary Affymetrix signal intensities after normalisation with the rma algorithm.

Abbreviations:

HER2: human epidermal growth factor receptor 2; AR: androgen receptor; ESR1: estrogen receptor 1; L: luminal; B: basal; MA: molecular apocrine.

Table 1. Simplified breast cancer molecular subtypes classifications (including the molecular apocrine subtype)					
Classification in 6 IHC subtypes	Classification in 3 IHC subtypes for the comparison to GEA	AR(1)	ER/PR(2)	HER2(3)	Ki67(4)
MA	MA	Positive	Both negative	Any	Any
Luminal A	Luminal	Any	ER and/or PR positive	Negative	Low
Luminal B HER2-negative		Any	ER and/or PR positive	Negative	High
Luminal B HER2-positive		Any	ER and/or PR positive	Positive	Any
Triple negative basal-like	Basal-like	Negative	Both negative	Negative	Any
Non-luminal and non-MA HER2-positive		Negative	Both negative	Positive	Any
<p>Abbreviations: IHC: immunohistochemical; GEA: gene expression array; AR: androgen receptor; ER: oestrogen receptor; PR: progesterone receptor; HER2: human epidermal growth receptor 2; MA: molecular apocrine-like subtype. Legend: (1) AR positive $\geq 10\%$; (2) ER and PR negative $< 1\%$; (3) HER2 positive: immunohistochemistry (IHC) 3+ or IHC2+ and Dual Detection In Situ Hybridisation (DDISH); (4) Ki67 high $\geq 14\%$.</p>					

Table 2. pCR rates by simplified breast cancer molecular subtype					
	Patients (N=846)	No pCR (%)	No data on residual tumour ^a (%)	pCR (%) [95%CI]	Odds ratio (95% CI)
MA					
Any HER2-status ^b	93	58 (62.4)	4 (4.3)	31 (33.3) [23.9, 43.9]	5.26 (2.78, 9.96)
<i>HER2-negative</i>	32	21 (65.6)	2 (6.3)	9 (28.1)	
<i>HER2-positive</i>	59	37 (62.7)	2 (3.5)	20 (33.9)	
Triple negative basal-like	94	55 (58.5)	7 (7.4)	32 (34.0) [24.6, 44.5]	5.43 (2.88, 10.25)
Luminal A	219	199 (90.9)	1 (0.5)	19 (8.4) [5.3, 13.2]	1.00
Luminal B HER2-negative	323	279 (86.4)	3 (0.9)	41 (12.7) [9.2, 16.8]	1.53 (0.86, 2.72)
Luminal B HER2-positive	110	77 (70.0)	3 (2.7)	30 (27.3) [19.2, 36.6]	3.95 (2.10, 7.41)
Non-luminal and non-MA HER2-positive	7	4 (57.1)	0 (0.0)	3 (42.9) [10.0, 81.6]	7.89 (1.64, 37.91)
P-value ^c					<0.001
Legend: ^a No surgery performed (progression on neoadjuvant chemotherapy); considered as No pCR in the logistic regression model ^b Two patients with equivocal or missing HER2, not included in the subgroups by HER2-status ^c P value for Wald test of a difference between the 6 subtypes using a logistic regression model.					
Abbreviations: MA: molecular apocrine; pCR: pathological complete response; CI: confidence interval; HER2: human epidermal growth factor receptor 2.					

Table 3. Molecular subtypes identified by gene expression array and immunohistochemistry for oestrogen, progesterone and androgen receptors				
	GEA classification			Total (N=60)
	Luminal (N=25)	MA (N=14)	Basal-like (N=21)	
	N (%)	N (%)	N (%)	N (%)
IHC classification				
Luminal (*)	23 (38.3)	2 (3.3)	2 (3.3)	27 (45.0)
MA (**)	1 (1.7)	12 (20.0)	1 (1.7)	14 (23.3)
Basal-like (***)	1 (1.7)	0 (0.0)	18 (30.0)	19 (31.7)
Abbreviations: GEA: gene expression array; IHC: immunohistochemistry; MA: molecular apocrine Legend: (*) ER-positive and/or PR-positive (**) AR-positive, ER- and PR-negatives (***) ER-, PR- and AR-negatives				