THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Factors associated with acute fatigue in primary care

Citation for published version:

Power, M, Chalder, T & al, E 2010, 'Factors associated with acute fatigue in primary care', *Psychological Medicine*, vol. 40, pp. 1289-1295.

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Psychological Medicine

Publisher Rights Statement:

T. Chalder, J. Neeleman, S. E. Reme, M. Power and S. Wessely
Psychological Medicine / Volume 40 / Issue 08 / August 2010, pp 1289 - 1295
DOI: 10.1017/S0033291709992327, Published online: 27 January 2010

Link to this article: http://journals.cambridge.org/abstract_S0033291709992327

T. Chalder, J. Neeleman, S. E. Reme, M. Power and S. Wessely (2010). Factors associated with acute fatigue in primary care. *Psychological Medicine*, 40, pp 1289-1295 doi:10.1017/S0033291709992327

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Psychological Medicine

<http://journals.cambridge.org/PSM>

Additional services for *Psychological Medicine*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)

Factors associated with acute fatigue in primary care

T. Chalder, J. Neeleman, S. E. Reme, M. Power and S. Wessely

Psychological Medicine / Volume 40 / Issue 08 / August 2010, pp 1289 - 1295

DOI: 10.1017/S0033291709992327, Published online: 27 January 2010

Link to this article: http://journals.cambridge.org/abstract_S0033291709992327

How to cite this article:

T. Chalder, J. Neeleman, S. E. Reme, M. Power and S. Wessely (2010). Factors associated with acute fatigue in primary care. Psychological Medicine, 40, pp 1289-1295 doi:10.1017/S0033291709992327

Request Permissions : [Click here](#)

Factors associated with acute fatigue in primary care

T. Chalder*, J. Neeleman, S. E. Reme, M. Power and S. Wessely

Department of Psychological Medicine and Department of Primary Care, King's College Hospital, London, UK

Background. To examine the role of psychological distress, negative life events, social support and lack of fitness (using breathlessness on exertion as a proxy) in the development of new onset fatigue in a primary care population.

Method. Adults between the ages of 18 and 45 years who were registered with five general practices in South East England were asked to complete a fatigue questionnaire and the 12-item General Health Questionnaire. Between 1 and 12 months later, subjects who visited the general practitioner (GP) with a suspected viral infection were recruited to the study and asked to complete measures of fatigue, psychological distress, life events, social support and allergies (stage 2). The next person to present to the GP with a complaint other than a viral illness was recruited as a control. Factors assessed at stage 2 that were associated with the development of fatigue were examined with stepwise logistic regression.

Results. Acute fatigue was not associated with a viral illness. Negative life events and breathlessness on exertion (interpreted as lack of fitness) were associated with incident cases of fatigue. However, when controlling for concurrent psychological distress, the independent association of negative life events disappeared.

Conclusions. Psychological distress was strongly associated with new onset fatigue and hence emphasizes the significance of psychological distress as a concomitant complaint in fatigue. Further, the salient association between breathlessness and fatigue may indicate the need to recommend exercise as a therapeutic strategy to improve physical fitness in the primary care setting.

Received 14 September 2009; Revised 30 November 2009; Accepted 14 December 2009; First published online 27 January 2010

Key words: Fatigue, life events, primary care.

Introduction

Fatigue is common and experienced by many but prevalence rates vary according to the setting (Kroenke & Price, 1993; Pawlikowska *et al.* 1994; Eriksen *et al.* 1998; Van't Leven *et al.* 2009). It is distributed as a continuous variable in the community (Pawlikowska *et al.* 1994). Moving from normal tiredness through mild to severe fatigue, the controversial syndrome of chronic fatigue syndrome (CFS) lies at the severest end of this spectrum. Although ubiquitous, fatigue is not a trivial problem and is often associated with disability comparable with that found in other illnesses (Kroenke *et al.* 1988). In the primary care setting it is commonly associated with psychological distress and not disease as some might assume (Ridsdale *et al.* 1993). Despite the rise in interest in fatigue over the last few years there has been little prospective research into its aetiology.

Complaints of fatigue were in a recent study found to affect as much as one-third of the general population, with 5% short-term fatigue (<6 months), 31% chronic fatigue (>6 months), and 1% CFS-like fatigue (Van't Leven *et al.* 2009). Common characteristics between all three groups of fatigued subjects were an unhealthy life-style that included smoking, unhealthy food, little physical activity and more use of analgesia, antidepressants and sedatives (Van't Leven *et al.* 2009). This indicates that life-style is a crucial factor for the whole dimension of fatigue, and might also be important for prolongation of fatigue. Other prospective population-based studies have found that previous psychiatric disorder (Harvey *et al.* 2008) or pre-morbid emotional instability (also known as neuroticism) and self-reported stress are important predictors for developing chronic fatigue (Kato *et al.* 2006). Pre-morbid fatigue score was the only significant predictor of chronic fatigue when examined prospectively in another population-based study (Lawrie *et al.* 1997). The authors found a cross-sectional correlation between fatigue and psychological distress, but not a predictive relationship, indicating that psychiatric morbidity is associated with, but not necessarily a risk factor for, chronic fatigue (Lawrie *et al.* 1997). This is

* Address for correspondence: T. Chalder, Department of Psychological Medicine, Guy's, King's and St Thomas' School of Medicine, 103 Denmark Hill, London SE5 8AZ, UK.
(Email: trudie.chalder@kcl.ac.uk)

Fig. 1. Patient flow and patients analysed in this study. FQ, Fatigue questionnaire; GHQ, General Health Questionnaire; GP, general practitioner.

not quite in agreement with other studies that have found psychological distress to be a predictor of CFS (Hotopf *et al.* 1996) and post-infectious fatigue. Several outcome studies have further suggested that psychiatric and psychological rather than physical measures predict long-term disability in CFS (Sharpe *et al.* 1992; Lawrie & Pelosi, 1994; Wilson *et al.* 1994; Cairns & Hotopf, 2005).

Long-term fatigue is frequently reported after various viral illnesses (Hotopf *et al.* 1996; White *et al.* 1998; Buchwald *et al.* 2000). Viral illnesses have therefore been suggested to be a risk factor for CFS (Cope *et al.* 1994), although our previous study found no evidence of CFS being caused by common viral infections (Wessely *et al.* 1995). Several retrospective studies have reported the presence of stressful life events and difficulties before onset of CFS (Stricklin *et al.* 1990; Masuda *et al.* 1994; Theorell *et al.* 1999; Hatcher & House, 2003), including childhood adversity (Heim *et al.* 2006). One of the studies suggests that a particular kind of stressor precedes the onset of CFS, namely irresolvable dilemmas which were detected in nearly a third of the CFS patients but not in the controls (Hatcher & House, 2003). However, other studies have not been able to find this difference (Lewis *et al.* 1994). Another factor purported to maintain fatigue is reduced exercise capacity or deconditioning (Berelowitz *et al.* 1995; White *et al.* 2001). The purpose of this study was to test the hypothesis that negative life events, lack of social support and lack of fitness were independently associated with incident

fatigue, irrespective of the presence of a viral infection and concurrent psychological distress.

Method

Design

A longitudinal cohort study was conducted in UK primary care. At stage 1, 31 651 men and women aged between 18 and 45 years and registered with five general practices were mailed a questionnaire, designed to measure subjective physical and mental fatigue (Chalder *et al.* 1993), and the 12-item General Health Questionnaire (GHQ-12; Goldberg *et al.* 1997).

Recruitment procedures

Between 1 and 12 months later, patients between the ages of 18 and 45 years (stage 2) visiting the general practitioner (GP) with an acute viral illness were recruited to the study. Most patients presented with 'flu-like' episodes or infections of the upper respiratory tract. An additional group of patients, presenting with complaints unrelated to viral infections, were also recruited as controls. More detailed descriptions of the sample and recruitment procedures are reported elsewhere (Wessely *et al.* 1995; Chalder *et al.* 1996). Those with anxiety and depression were not excluded. After complete description of the study, written informed consent was obtained from all subjects. They were then asked to complete a series of questionnaires (see Fig. 1).

Measures

Chalder Fatigue Scale

The Chalder Fatigue Scale is an 11-item scale measuring physical and mental fatigue during the last month. There are four response options available, ranging from 'less than usual' to 'much more than usual'. The scale's total score can range from 0 to 33, but a bimodal score ranging from 0 to 11 can also be obtained. With a cut-off score of ≥ 4 , the bimodal score is considered indicative of excessive fatigue. The scale is both reliable and valid (Chalder *et al.* 1993).

GHQ

The short version of the GHQ (GHQ-12) has been extensively used as a short screening instrument to assess psychological distress during the past few weeks. It has been shown to be both reliable and valid, and a good case detector when compared with diagnostic interviews (Goldberg *et al.* 1997). The GHQ has also been used extensively in primary care.

Life Events Inventory

The Life Events Inventory is a checklist of both positive and negative life events (Cochrane & Robertson, 1973). The checklist includes 48 life events and has spaces for subjects to indicate which, if any, of the events have happened to them during the last 6 months. The questionnaire has two subscales: positive life events and negative life events. The score used in this report is a summary of negative life events only.

Significant Others Scale (social support)

The Significant Others Scale measures different functional resources of social support that may be provided by a number of significant role relationships within an individual's social network. We used a shortened version of the Significant Others Scale which yields levels of actually received and ideally desired levels of support and the discrepancy between them, further called 'perceived lack' (Power *et al.* 1988).

Allergy Scale

An allergy questionnaire designed specifically for this study covered self-reported atopic and allergic symptoms. A question called 'breathlessness on exertion' was included and assumed to assess perceived lack of physical fitness. The exact wording of the question was: 'Have you ever been excessively breathless after minor exercise or after exposure to smoke, fumes or similar factors?' Responses were dichotomous (0 = no, 1 = yes).

Social class was measured using the Registrar General's Classification System.

Statistical analysis

All analyses were performed using SPSS version 15 (SPSS Inc., USA). The primary outcome used was fatigue cases (defined as those scoring ≥ 4 out of a maximum of 11 on the fatigue scale). Odds ratios, obtained by logistic regression, were used to compare associations between fatigue cases and non-cases. Marital status, social class, presence of viral illness and breathlessness on exertion were treated as categorical variables (e.g. married/co-habiting *versus* other; professional/intermediate/skilled *versus* part-skilled/unskilled/unemployed).

To obtain an incident fatigue sample, subjects scoring ≥ 4 on the fatigue questionnaire or who had missing data at stage 1 were excluded from the analysis, leaving 1219 subjects. First, crude associations between fatigue caseness at stage 2 and the various risk factors were obtained. Second, we examined whether these associations differed by the presence of viral illness, using the likelihood ratio test for interaction. Third, a full model was specified containing all variables and the interaction terms between them. The time lag between stages 1 and 2 (1–12 months) was controlled for throughout. A final model was obtained using a backward stepwise approach.

Results

There were 15 283 responders at stage 1, an overall response rate of 48%. The response rate adjusted for inaccuracies in the inner-city practice registers was 67% (Pawlikowska *et al.* 1994). At stage 2, 2376 subjects were recruited, 1199 with a viral illness and 1177 without a viral illness. All or nearly all of the questionnaires were completed by 97% of those with ($n=1167$) and 98% of those without viral illness ($n=1160$). A total of 1873 subjects completed questionnaires at stages 1 and 2. Subjects with missing data at either stage were more likely to be male (181/723 = 25%) than female (302/1633 = 18%) ($p < 0.001$), and unmarried (187/456 = 41%) than married (880/1841 = 47.8%) ($p = 0.009$). Social class did not differ between responders and non-responders. More information about response rates is reported elsewhere (Wessely *et al.* 1995; Chalder *et al.* 1996). There were 502 cases of incident fatigue at stage 2.

Negative life events, psychological distress, perceived lack of practical and emotional support, and breathlessness on exertion were crudely associated with incident fatigue, all of which were assessed at stage 2. None of the demographic variables was

Table 1. Univariate and multivariate associations (ORs) between fatigue caseness and other variables at stage 2

	Univariate associations		Multivariate model	
	OR (95% CI)	P	OR (95% CI)	p
Sex	1.11 (0.86–1.42)	0.423	1.00 (0.76–1.32)	0.978
Age	1.00 (0.99–1.02)	0.957	1.00 (0.98–1.02)	0.906
Marital status	1.19 (0.95–1.50)	0.137	1.02 (0.77–1.35)	0.891
Social class	0.92 (0.72–1.18)	0.504	0.80 (0.61–1.06)	0.120
Perceived lack of practical support	1.21 (1.07–1.37)	0.002	1.04 (0.88–1.23)	0.650
Perceived lack of emotional support	1.22 (1.08–1.38)	0.002	1.10 (0.93–1.29)	0.274
Negative life events	1.47 (1.28–1.69)	0.000	1.51 (1.28–1.78)	0.000
Breathlessness on exertion	2.01 (1.54–2.61)	0.000	1.76 (1.31–2.35)	0.000

OR, Odds ratio; CI, confidence interval.

Table 2. Univariate and multivariate associations (ORs) between fatigue caseness and other variables, with psychological distress at stage 2 included

Associated variables	Univariate associations		Multivariate model	
	OR (95% CI)	p	OR (95% CI)	p
Sex	1.11 (0.86–1.42)	0.423	0.96 (0.72–1.27)	N.S.
Age	1.00 (0.99–1.02)	0.957	1.01 (0.99–1.03)	N.S.
Marital status (co-habiting or not)	1.19 (0.95–1.50)	0.137	1.12 (0.83–1.51)	N.S.
Social class	0.92 (0.72–1.18)	0.504	0.80 (0.61–1.06)	N.S.
Perceived lack of practical support	1.21 (1.07–1.37)	0.002	0.88 (0.52–1.50)	N.S.
Perceived lack of emotional support	1.22 (1.08–1.38)	0.002	0.82 (0.46–1.46)	N.S.
Negative life events	1.47 (1.28–1.69)	0.000	1.04 (0.97–1.11)	N.S.
Breathlessness on exertion	2.01 (1.54–2.61)	0.001	1.66 (1.23–2.25)	0.001
Psychological distress, GHQ	5.60 (4.32–7.27)	0.000	5.60 (4.25–7.37)	0.000

OR, Odds ratio; CI, confidence interval; N.S., non-significant; GHQ, General Health Questionnaire.

associated with fatigue and the crude associations did not differ between subjects with or without a viral illness. Thus, viral illness is not considered in the subsequent analyses. Past psychological distress (measured at stage 1) did not predict new incidents of fatigue (at stage 2) and was therefore not included in the multivariate model. The best model for new onset fatigue included negative life events and breathlessness on exertion. None of the other variables contributed significantly. The final model including negative life events and breathlessness correctly classified 63.0% of cases and non-cases of fatigue compared with 63.8% in the full model with all variables included (see Table 1).

However, when concurrent psychological distress (from stage 2) was added to the multivariate model, the effect of negative life events disappeared (see Table 2). The final model including psychological

distress and breathlessness now correctly classified 71.5% of cases and non-cases of fatigue compared with 71.6% in the full model with all variables included (see Table 2). The final model explained between 16% and 21% of the variance.

Discussion

As hypothesized, this study found that negative life events and lack of fitness were associated with incident fatigue. However, when concurrent psychological distress was included in the multivariate model, the independent association of negative life events disappeared. Perceived lack of support, whether it be emotional or practical, was not independently associated with incident fatigue. Concurrent psychological distress was the strongest factor in determining fatigue.

The results could be interpreted in a number of ways. First, the association between negative life events and fatigue disappeared when controlling for psychological distress, which could indicate that psychological distress mediates the effect of negative life events on fatigue. However, the direction could also go the other way, as the data were collected simultaneously; chronic fatigue has for instance been found to predict psychiatric disorders in another community-based sample (Taylor *et al.* 2003). Second, our findings are in agreement with Lawrie *et al.* (1997) where cross-sectional correlations between fatigue and psychological distress were found but no predictive relationships when adjusted for baseline fatigue. The authors hence concluded that psychiatric morbidity is associated with, but not necessarily a risk factor for, chronic fatigue. Our results, on the other hand, cannot eliminate the possibility of psychological distress as a predictor of chronic fatigue, as this study only investigates new onset of fatigue. Given the evidence of emotional instability (Kato *et al.* 2006) and psychological distress as predictors of chronic fatigue (Hotopf *et al.* 1996; Russo *et al.* 1998) and post-infectious fatigue (Wessely *et al.* 1995), it still seems reasonable to assume that psychological distress is involved in prolongation of fatigue.

Breathlessness on exertion was the other independent association with fatigue in this study. Breathlessness is a symptom of anxiety. However, given that this question specifically asked about breathlessness on exertion it seems unlikely that anxiety was the sole explanatory factor. Second, breathlessness could have been associated with asthma. However, both univariate and multivariate statistics were carried out excluding subjects with self-reported asthma. The effect remained. In keeping with previous research, it seems plausible therefore that breathlessness on exertion is indeed an indication of lack of fitness or deconditioning. Lack of activity or exercise is commonly associated with fatigue (Haines, 1974; Greenleaf & Kozlowski, 1982; White *et al.* 2001), and the negative effects of inactivity on exercise physiology are well established (Convertino *et al.* 1997), as is post-exertional malaise which has been associated with prolongation of fatigue (Taylor *et al.* 2002). Finally, breathlessness on exertion as an indication of little physical activity is in agreement with Van't Leven *et al.* (2009) who argued that an unhealthy life-style is significant in the whole spectrum of fatigue.

In this study, we asked about people's perception of their fitness, which could differ from their actual level of physical fitness. Future studies should therefore look at both subjective and objective measures of physical fitness. In addition, the latter part of the question (... exposure to smoke, fumes or similar factors) might

conflate sensitivity to exposure and perceived sensitivity to exposure, which may in turn be associated with neuroticism and distress (Van den Bergh *et al.* 2004). On balance, however, we think that the most likely interpretation of this finding is level of fitness in accordance with previous studies showing that physical activity is associated with reduced feelings of fatigue (Puetz, 2006).

There did not appear to be any interactions between possible viral infection and the other risk factors such as psychological distress, life events, social support or breathlessness on exertion. The multivariate analysis was therefore carried out without entering viral status as a covariate. This does not necessarily exclude a role for viruses in a model of understanding chronic fatigue. It does seem reasonable to assume that in some, the virus acts as a trigger for the fatigue in a vulnerable individual but does not diminish the effect of other predictor variables such as psychological distress and breathlessness on exertion. We were interested in the fact that social support was not retained in the final model and so examined the correlation between social support and psychological distress and found small but significant associations ($-0.22, p < 0.01$). It is possible that although social support is not central to the development of fatigue it may be more important in determining chronicity.

The main methodological problem with this study is the fact that most of the risk factors were assessed at the same time as the outcome, introducing the possibility of reporting bias. However, although this may have inflated the univariate associations, this is less likely to have affected the multivariate model because all estimates would have been inflated more or less to a similar extent. The salient associations found in this study should therefore be investigated further in order to determine their significance as risk factors for new onset fatigue.

In conclusion, psychological distress did not predict new onset of fatigue, but was strongly associated with fatigue and hence emphasizes the significance of psychological distress as a concomitant complaint in fatigue. Further, the salient association between breathlessness and fatigue could indicate a need to recommend exercise as a therapeutic strategy to improve physical fitness in the primary care setting, although these findings need replication before such therapeutic strategies are recommended.

Acknowledgements

We are grateful to Vivian Hobbs, Julie Dennison, Dawn Baker, Sue Fox, Marjorie Peers and Brenda Hinton for helping with the data collection. We would also like to thank the GPs for their assistance. The

study was supported by the Linbury Trust. T.C. and S.W. acknowledge financial support from the Department of Health via the National Institute for Health Research (NIHR) Specialist Biomedical Research Centre for Mental Health award to South London and Maudsley NHS Foundation Trust (SLaM) and the Institute of Psychiatry at King's College London.

Declaration of Interest

Trudie Chalder is an author of books on chronic fatigue.

References

- Berelowitz GJ, Burgess AP, Thanabalasingham T, Murray-Lyon IM, Wright DJ (1995). Post-hepatitis syndrome revisited. *Journal of Viral Hepatitis* **2**, 133–138.
- Buchwald DS, Rea TD, Katon WJ, Russo JE, Ashley RL (2000). Acute infectious mononucleosis: characteristics of patients who report failure to recover. *American Journal of Medicine* **109**, 531–537.
- Cairns R, Hotopf M (2005). A systematic review describing the prognosis of chronic fatigue syndrome. *Occupational Medicine* **55**, 20–31.
- Chalder T, Berelowitz G, Pawlikowska T, Watts L, Wessely S, Wright D, Wallace EP (1993). Development of a fatigue scale. *Journal of Psychosomatic Research* **37**, 147–153.
- Chalder T, Power MJ, Wessely S (1996). Chronic fatigue in the community: 'a question of attribution'. *Psychological Medicine* **26**, 791–800.
- Cochrane R, Robertson A (1973). The life events inventory: a measure of the relative severity of psycho-social stressors. *Journal of Psychosomatic Research* **17**, 135–140.
- Convertino VA, Bloomfield SA, Greenleaf JE (1997). An overview of the issues: physiological effects of bed rest and restricted physical activity. *Medicine and Science in Sports and Exercise* **29**, 187–190.
- Cope H, David A, Pelosi A, Mann A (1994). Predictors of chronic 'postviral' fatigue. *Lancet* **344**, 864–868.
- Eriksen HR, Svendsrød R, Ursin G, Ursin H (1998). Prevalence of subjective health complaints in the Nordic European countries in 1993. *European Journal of Public Health* **8**, 294–298.
- Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O, Rutter C (1997). The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychological Medicine* **27**, 191–197.
- Greenleaf JE, Kozlowski S (1982). Physiological consequences of reduced physical activity during bed rest. *Exercise and Sport Sciences Reviews* **10**, 84–119.
- Haines RF (1974). Effect of bed rest and exercise on body balance. *Journal of Applied Physiology* **36**, 323–327.
- Harvey SB, Wadsworth M, Wessely S, Hotopf M (2008). The relationship between prior psychiatric disorder and chronic fatigue: evidence from a national birth cohort study. *Psychological Medicine* **38**, 933–940.
- Hatcher S, House A (2003). Life events, difficulties and dilemmas in the onset of chronic fatigue syndrome: a case-control study. *Psychological Medicine* **33**, 1185–1192.
- Heim C, Wagner D, Maloney E, Papanicolaou DA, Solomon L, Jones JF, Unger ER, Reeves WC (2006). Early adverse experience and risk for chronic fatigue syndrome: results from a population-based study. *Archives of General Psychiatry* **63**, 1258–1266.
- Hotopf M, Noah N, Wessely S (1996). Chronic fatigue and minor psychiatric morbidity after viral meningitis: a controlled study. *Journal of Neurology, Neurosurgery and Psychiatry* **60**, 504–509.
- Kato K, Sullivan PF, Evengard B, Pedersen NL (2006). Premorbid predictors of chronic fatigue. *Archives of General Psychiatry* **63**, 1267–1272.
- Kroenke K, Price RK (1993). Symptoms in the community. Prevalence, classification, and psychiatric co-morbidity. *Archives of Internal Medicine* **153**, 2474–2480.
- Kroenke K, Wood DR, Mangelsdorff AD, Meier NJ, Powell JB (1988). Chronic fatigue in primary care. Prevalence, patient characteristics, and outcome. *Journal of the American Medical Association* **260**, 929–934.
- Lawrie SM, Manders DN, Geddes JR, Pelosi AJ (1997). A population-based incidence study of chronic fatigue. *Psychological Medicine* **27**, 343–353.
- Lawrie SM, Pelosi AJ (1994). Chronic fatigue syndrome: prevalence and outcome. *British Medical Journal* **308**, 732–733.
- Lewis S, Cooper CL, Bennett D (1994). Psychosocial factors and chronic fatigue syndrome. *Psychological Medicine* **24**, 661–671.
- Masuda A, Nozoe SI, Matsuyama T, Tanaka H (1994). Psychobehavioral and immunological characteristics of adult people with chronic fatigue and patients with chronic fatigue syndrome. *Psychosomatic Medicine* **56**, 512–518.
- Pawlikowska T, Chalder T, Hirsch SR, Wallace P, Wright DJ, Wessely SC (1994). Population based study of fatigue and psychological distress. *British Medical Journal* **308**, 763–766.
- Power MJ, Champion LA, Aris SJ (1988). The development of a measure of social support: the Significant Others (SOS) Scale. *British Journal of Clinical Psychology* **27**, 349–358.
- Puetz TW (2006). Physical activity and feelings of energy and fatigue: epidemiological evidence. *American Journal of Sports Medicine* **36**, 767–780.
- Ridsdale L, Evans A, Jerrett W, Mandalia S, Osler K, Vora H (1993). Patients with fatigue in general practice: a prospective study. *British Medical Journal* **307**, 103–106.
- Russo J, Katon W, Clark M, Kith P, Sintay M, Buchwald D (1998). Longitudinal changes associated with improvement in chronic fatigue patients. *Journal of Psychosomatic Research* **45**, 67–76.
- Sharpe M, Hawton K, Seagroatt V, Pasvol G (1992). Follow up of patients presenting with fatigue to an

- infectious diseases clinic. *British Medical Journal* **305**, 147–152.
- Stricklin A, Sewell M, Austad C** (1990). Objective measurement of personality variables in epidemic neuromyasthenia patients. *South African Medical Journal* **77**, 31–34.
- Taylor RR, Jason LA, Curie CJ** (2002). Prognosis of chronic fatigue in a community-based sample. *Psychosomatic Medicine* **64**, 319–327.
- Taylor RR, Jason LA, Jahn SC** (2003). Chronic fatigue and sociodemographic characteristics as predictors of psychiatric disorders in a community-based sample. *Psychosomatic Medicine* **65**, 896–901.
- Theorell T, Blomkvist V, Lindh G, Evengard B** (1999). Critical life events, infections, and symptoms during the year preceding chronic fatigue syndrome (CFS): an examination of CFS patients and subjects with a nonspecific life crisis. *Psychosomatic Medicine* **61**, 304–310.
- Van den Bergh O, Winters W, Devriese S, van Diest I, Vos G, de Peuter S** (2004). Accuracy of respiratory symptom perception in persons with high and low negative affectivity. *Psychology and Health* **19**, 9.
- Van't Leven M, Zielhuis GA, van der Meer JW, Verbeek AL, Bleijenberg G** (2009). Fatigue and chronic fatigue syndrome-like complaints in the general population. *European Journal of Public Health*. Published online: 18 August 2009. doi:10.1093/eurpub/ckp113.
- Wessely S, Chalder T, Hirsch S, Pawlikowska T, Wallace P, Wright DJ** (1995). Postinfectious fatigue: prospective cohort study in primary care. *Lancet* **345**, 1333–1338.
- White PD, Thomas JM, Amess J, Crawford DH, Grover SA, Kangro HO, Clare AW** (1998). Incidence, risk and prognosis of acute and chronic fatigue syndromes and psychiatric disorders after glandular fever. *British Journal of Psychiatry* **173**, 475–481.
- White PD, Thomas JM, Kangro HO, Bruce-Jones WD, Amess J, Crawford DH, Grover SA, Clare AW** (2001). Predictions and associations of fatigue syndromes and mood disorders that occur after infectious mononucleosis. *Lancet* **358**, 1946–1954.
- Wilson A, Hickie I, Lloyd A, Hadzi-Pavlovic D, Boughton C, Dwyer J, Wakefield D** (1994). Longitudinal study of outcome of chronic fatigue syndrome. *British Medical Journal* **308**, 756–759.