THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Parasite variation and the evolution of virulence in a *Daphnia*-microparasite system

Citation for published version:

Little, TJ, Chadwick, W & Watt, K 2008, 'Parasite variation and the evolution of virulence in a *Daphnia*-microparasite system', *Parasitology*, vol. 135, no. 3, pp. 303-308.
<https://doi.org/10.1017/S0031182007003939>

Digital Object Identifier (DOI):

[10.1017/S0031182007003939](https://doi.org/10.1017/S0031182007003939)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Parasitology

Publisher Rights Statement:

RoMEO green

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Parasite variation and the evolution of virulence in a *Daphnia*-microparasite system

T. J. LITTLE*, W. CHADWICK and K. WATT

Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Kings Buildings, West Mains Road, Edinburgh EH9 3JT, UK

(Received 21 June 2007; revised 24 September and 2 October 2007; accepted 3 October 2007; first published online 16 November 2007)

SUMMARY

Understanding genetic relationships amongst the life-history traits of parasites is crucial for testing hypotheses on the evolution of virulence. This study therefore examined variation between parasite isolates (the bacterium *Pasteuria ramosa*) from the crustacean *Daphnia magna*. From a single wild-caught infected host we obtained 2 *P. ramosa* isolates that differed substantially in the mortality they caused. Surprisingly, the isolate causing higher early mortality was, on average, less successful at establishing infections and had a slower growth rate within hosts. The observation that within-host replication rate was negatively correlated with mortality could violate a central assumption of the trade-off hypothesis for the evolution of virulence, but we discuss a number of caveats which caution against premature rejection of the trade-off hypothesis. We sought to test if the characteristics of these parasite isolates were constant across host genotypes in a second experiment that included 2 *Daphnia* host clones. The relative growth rates of the two parasite isolates did indeed depend on the host genotype (although the rank order did not change). We suggest that testing evolutionary hypotheses for virulence may require substantial sampling of both host and parasite genetic variation, and discuss how selection for virulence may change with the epidemiological state of natural populations and how this can promote genetic variation for virulence.

Key words: natural selection, co-evolution, trade-off, genetic variation, pathogen, parasite, specificity, *Pasteuria*.

INTRODUCTION

A common model for understanding parasite or pathogen evolution suggests that pathogens will be selected to balance their rate of transmission with their rate of host exploitation (Bremermann and Pickering, 1983; May and Anderson, 1983; Bull, 1994; Frank, 1996; Andre *et al.* 2003; Choo *et al.* 2003; Day and Proulx, 2004). The crux of this optimality argument is that parasites will maximize their fitness through host exploitation, but excessive host exploitation might kill hosts too quickly, thereby reducing parasites' transmission opportunities. This trade-off model for the evolution of virulence offers a general explanation for why parasites do not always evolve towards benevolence (e.g. Ebert, 1998), and has provided insight regarding how pathogen evolution may thwart intervention strategies (Gandon *et al.* 2001).

The virulence trade-off hypothesis assumes that higher within-host replication rates lead to higher virulence, and this has been supported in some (Lipsitch and Moxon, 1997; Ebert, 1998; Mackinnon and Read, 1999), but not all systems (Weiss, 2002; Escriu *et al.* 2003; Sacristán *et al.* 2005; Stewart *et al.* 2005; Pagan, 2007). Some workers have highlighted

how other factors, such as competition between parasites, need to be incorporated into the trade-off model (e.g. de Roode *et al.* 2005). Theoretical treatments of within-host competition have generally indicated that within-host selection will favour higher virulence, because more rapidly growing parasite genotypes may kill the host or competitively exclude prudent genotypes before the latter have achieved transmission (reviewed by Read and Taylor, 2001). However, a number of nuances are imaginable in competition, for example, chemical warfare between combatants could alter virulence evolution if parasite genotypes suppress each other's growth. Alternatively, immune-mediated competition can alter outcomes when one genotype attracts the majority of attention from the immune system (Almogy *et al.* 2002). Testing the generality of the trade-off or other hypotheses requires further data on genetic relationships amongst the key life-history traits of parasites, as well as an understanding of the likelihood and consequences of within-host competition.

Our work concerns the bacterium *Pasteuria ramosa*, a specialist parasite of the crustacean *Daphnia magna*. This is a naturally co-occurring host-parasite species pair for which past studies have revealed genetic variation for resistance in hosts, genetic differences among parasite isolates, and substantial genetic specificity (e.g. Little and Ebert, 2000;

* Corresponding author. Tel: +0131 650 7781. Fax: +0131 650 6564. E-mail: tom.little@ed.ac.uk

Carius *et al.* 2001; Little *et al.* 2006). *Pasteuria ramosa* is an ideal organism for estimating parasite fitness because transmission spores are only released upon host death, allowing for very accurate measurement of transmission potential or lifetime reproductive success. This feature of the system has been exploited in previous studies, and consequently *P. ramosa*–*D. magna* infections have provided rare empirical evidence of parasite fitness being maximized at an intermediate level of virulence, supporting a version of the virulence trade-off model (Jensen *et al.* 2006).

Parasite isolates used in many previous studies on this system have simply been the parasites collected from a single wild-caught female (as opposed to a true clone based on a single bacterial cell, which is not possible to isolate because *P. ramosa* can only be grown *in vivo*). Therefore, in the absence of molecular markers, it was not known if wild females harboured multiple parasite strains. Jensen *et al.* (2006) performed experiments indicating the presence of strain variation within single infections, and thus the potential for within-host competition in natural populations of *D. magna* and *P. ramosa*. The present work sought to associate this variation with host fecundity and mortality to shed further light on the potential for within-host competition and its likely impact on the evolution of virulence. By also measuring parasite growth and transmission potential we tested the assumption of a positive correlation between virulence and the within-host replication rate.

MATERIALS AND METHODS

Organisms and preparation of material

Daphnia magna is a planktonic crustacean found in still, freshwater bodies in temperate regions. Like most *Daphnia*, *D. magna* is a cyclical parthenogen where reproduction is usually the apomictic (i.e. clonal) production of all-female broods, but sex is occasionally induced by particular environmental factors (changes in food levels, crowding and photoperiod, among others). The product of sex is a resting egg that requires a period of diapause before hatching. Thus, by controlling environmental conditions and discarding any resting eggs produced, genetically identical clonal lines of *Daphnia* can be maintained in the laboratory. *Pasteuria ramosa* is a bacterial spore-forming, obligate endoparasite of *D. magna* that greatly reduces host fecundity (Ebert *et al.* 1996). Transmission is horizontal, but only achieved at host death, when spores are released from the decomposing cadavers (Ebert *et al.* 1996). *Pasteuria ramosa* passes through a series of well-defined developmental stages, ending in the production of morphologically distinct transmission spores that are straightforward to quantify using a

haemocytometer. The *D. magna* and *P. ramosa* used in this study were from a single pond near the village of Gaarzerfeld in northern Germany.

Pasteuria ramosa was collected from a single wild-caught infected *D. magna* that produced viable offspring before the parasite curtailed host reproduction. These clonally produced *Daphnia* offspring were collected and propagated to provide new hosts. The original infected host was also kept alive in the laboratory, and upon death her cadaver was crushed and used to infect new hosts. To accumulate parasites for experimentation, this propagation was repeated twice. Each new round of infections was performed in 200 ml jars with 10 hosts per jar and there were 10 replicate jars per passage. Infections were initiated with 1×10^6 spores per host and infection periods were 7 days. Infected hosts were grown until they died, at which point they were pooled and frozen for use in the next passage. In the last passage, dead infected hosts were separated and stored in 2 categories: hosts that had died relatively early (days 25–37; termed early killers) or those that died relatively late (days 55–67; termed late killers).

Main experiment

An experiment was conducted using ‘early killing’ spores or ‘late killing’ spores, each applied at 3 doses (1×10^4 , 1×10^5 or 1×10^6 spores per jar). We established 60 replicates, each replicate being a single *Daphnia* maintained alone in a 60 ml jar. These replicates were kept as such for 3 generations prior to experimentation, and during this time water was changed twice in the first 10 days of life, and thereafter with every clutch (clutches are produced every 3–4 days). *Daphnia* were fed 5×10^6 cells of the algae *Scenedesmus* daily. Six newborn from the third clutch of third generation females were split into the 6 treatments (2 spore types, 3 doses each) and parasite spores were added to each jar. *Daphnia* were left in contact with the spores for 5 days, after which they were transferred to fresh jars and water. From here on, *Daphnia* were transferred to fresh water each time they had a clutch (for hosts that did not become infected) or every 3 days (for hosts that became infected and thus had few or no clutches).

For 65 days, we counted the number of offspring produced by each *Daphnia* and recorded mortality and infection status. Whenever a host died, it was stored frozen. All individuals that survived until day 65 were censored. Additionally, a group of 60 hosts, evenly drawn from the 6 treatments, were sacrificed on day 25 (and recorded as censored) and used to gain an estimate of parasite growth in the earlier stages of the experiment. Each dead host was crushed in 200 μ l of distilled water and 8 μ l of this was placed in a haemocytometer for counting of transmission spores at $40\times$ magnification. We performed 4 transmission spore counts for each host and used a mean of these

Table 1. Summary statistics for the main infection experiment involving *Daphnia magna* and the parasite *Pasteuria ramosa*

(N, D indicates numerator and denominator degrees of freedom.)

Proportion infected	N, D d.f.	L.R. χ^2	P
Spore type	1, 343	8.84	0.0038
Dose	2, 343	40.1	<0.0001
Mortality	D.F.	L.R. χ^2	P
Spore type	1	22.8	<0.0001
Dose	2	4.55	0.10
Spore counts day 25	N, D d.f.	F	P
Spore type	1, 54	5.85	0.019
Dose	2, 54	7.15	0.002
Spore counts day 65	N, D d.f.	F	P
Spore type	1, 97	5.92	0.017
Dose	2, 97	0.87	0.42
Reproduction (yes/no)	N, D d.f.	F	P
Spore type	1, 279	2.49	0.12
Dose	2, 279	15.62	0.0004
Number of offspring	N, D d.f.	F	P
Spore type	1, 26	0.01	0.97
Dose	2, 26	0.68	0.68
Age at first reproduction	D.F.	L.R. χ^2	P
Spore type	1	0.51	0.48
Dose	2	1.98	0.16

for analysis. The counting procedure also allowed us to confirm infection status.

The explanatory variables for analyses were spore type (early killing or late killing) and dose (low, medium or high). We used logistic regression to analyse if the proportion infected differed by treatment. For all other analyses, we focussed on the group of infected hosts (the fitness effects of being exposed to parasites but resisting infection are reported elsewhere (Little and Killick, 2007)). Proportional hazards were used to analyse time to death. ANOVA was used to study counts of parasite transmission spores. For analysis of transmission spores we studied 2 data sets: hosts sacrificed on day 25 and hosts sacrificed on day 65. All analyses were performed in SAS version 9.0 (SAS Institute Inc, 2002).

Reproduction was difficult to analyse because very few (11%) infected hosts reproduced. We scored each host as either reproducing or not, and used a logistic regression to relate this to dose and spore type. We also created a separate data set including only those hosts that did reproduce and used this to compare the number of (square root transformed) offspring among treatments (via ANOVA), as well as to

Fig. 1. Mortality of *Daphnia magna* hosts infected with 2 spore types of the bacterium *Pasteuria ramosa*.

investigate age at first reproduction, which was analysed via proportional hazards because age at first reproduction is a time to event variable.

Experiment 2

The above experiment was repeated (with some minor methodological differences) but included 2 different host genotypes (the same one used above, plus a different host clone). The second host clone was isolated from the same population at the same time as the first, and is known to differ from the first both at allozyme loci and in terms of its resistance characteristics (Carius *et al.* 2001), although both clones show high overall susceptibility (these clones are labelled as '3' and '4' in Carius *et al.* (2001)). We focussed this experiment on testing for a host clone by spore type interaction, and the design was such that we had the most power to do so for early parasite growth. Hosts were prepared as above for 3 generations and there were 5 replicates of each host genotype (here, however, 5 *Daphnia* were kept in each jar, and jar size was 200 ml, but jar is still the largest unit of independent replication). On the day the experiment began, 30 newborn from each replicate were isolated and groups of 5 were placed in 200 ml jars corresponding to the 6 treatments (early and late killing spores and 3 spore doses of each, as above). When hosts died, their bodies were frozen, and on day 30 all surviving host were frozen for later counting of parasite spores.

RESULTS

Main experiment

Across treatments, 82% of hosts became infected and both dose and spore type (odds ratio, early killing spores = 0.48) influenced this proportion. Mortality among infected hosts was not influenced by dose, but was strongly affected by spore type (Table 1; Fig. 1). Parasite spore production at both day 25 and day 65

Fig. 2. Parasite (*Pasteuria ramosa*) spore production after 25 (left graph) or 65 days (right graph) of growth in the host *Daphnia magna*.

was influenced by spore type (Table 1; Fig. 2), while dose influenced spore counts only at the earlier time-point. In no case did we observe a dose by spore type interaction.

In the high, low and medium dose treatments, only 1.4, 4.2 and 5.6% of hosts, respectively, reproduced, but this was a significant difference (Table 1). Spore type (early killing or late killing) did not explain differences in whether hosts reproduced or not (Table 1). Among the 32 hosts that reproduced, the number of offspring they had was not explained by spore type or dose (Table 1; early killers, average number of offspring = 16.3 (S.E. = 5.3), late killers, average number of offspring = 19.4 (S.E. = 6.3)), and neither dose nor spore type explained age at first reproduction (Table 1; early killers, average age at first reproduction day 19.1 (S.E. = 1.6), late killers, average age at first reproduction day 21.4 (S.E. = 1.9)). For all reproduction traits, dose by spore type interactions were not significant.

That the number of host offspring did not differ between hosts suffering a fast-growing versus a slow growing parasite might at first seem to contradict previous studies (Ebert *et al.* 2004) showing an overall negative correlation between parasite growth and host reproduction. However, adding parasite spore production as a covariate (nested within spore type) to the analysis of the 32 hosts that reproduced reveals such a negative relationship (spore production [spore type], $F_{2,28} = 5.64$, $P < 0.01$ $r^2 = 0.29$). Thus, while differences between parasites do not account for host reproduction (presumably this is because the strong mortality differences between spore types obscures the parasite growth–host fecundity relationship), within each spore type there is a negative correlation between parasite growth and host reproduction.

Experiment 2

As in the main experiment, spore counts of late killing spores at a defined time-point were significantly

Fig. 3. The proportion of 2 clones of *Daphnia magna* becoming infected in relation to 2 spore types of the bacterium *Pasteuria ramosa*. Clone 1 was used in the main experiment, clone 2 was the additional host clone used in Exp. 2.

larger (spore type main effect: $F_{1,46} = 62.57$, $P < 0.001$), and there was also a spore type by host clone interaction (Fig. 3; $F_{1,46} = 6.95$, $P = 0.014$). Regarding the other response variables, this experiment confirmed the results of the first, since early and late killing spores showed the expected differences in the proportion of hosts becoming infected ($\chi^2 = 10.18$, $P = 0.0014$) and mortality ($\chi^2 = 6.4$, $P = 0.014$). Although the two host clones differed in terms of the mortality suffered (clone effect: $\chi^2 = 10.17$, $P = 0.0014$) and the proportion that became infected (clone effect: $\chi^2 = 11.56$, $P = 0.0007$), neither of these variables showed a significant spore type by host clone interaction.

DISCUSSION

The two parasite isolates studied here have markedly different life-history traits and were originally found to co-infect a single host line, indicating that multiple infections and hence within-host competition are possible in the wild (although we cannot exclude that one strain is a mutant which emerged between field sampling and experimentation). The isolates were separated based on the mortality they cause: one kills hosts relatively quickly, the other kills relatively slowly. These isolates showed additional differences, in particular the slow killing isolate showed greater infectivity and, surprisingly, a greater rate of replication. Clearly, within-host replication rate can be decoupled from mortality. While this challenges a key assumption of the virulence trade-off hypothesis, this result should be interpreted cautiously because of the small number of parasite strains studied. The trade-off theory assumes a genetic relationship between within-host parasite growth and virulence,

but fully assessing this will require more substantial sampling of the genetic variation in the parasite population.

Additionally, for *Daphnia-Pasteuria* infections, mortality and virulence (parasite-induced fitness losses in hosts) are not strictly equivalent. The trade-off model essentially states that the rate at which parasites extract resources from their hosts equates to virulence, typically taken to be mortality. *Pasteuria ramosa* should follow this trade-off: prudent parasites might be out-competed while those that grow fast and rapidly extract host resources might kill their hosts before achieving maximal transmission potential. However, even living hosts suffer great parasite-induced costs because *Pasteuria* infections sterilize hosts. Virulence in this system can be thus sensibly accounted for by studying host fecundity instead of mortality, although both perspectives predict that parasite growth is positively related to virulence. Fecundity, however, will be determined jointly by mortality (early-dying hosts will have had less time to reproduce) and the amount of reproduction living hosts achieve prior to sterilization. The isolates under study presently actually do not differ in the virulence they cause from the perspective of host reproduction, although it is notable that only a very small fraction of hosts reproduced and so the power to detect reproductive differences was limited. Sterilization benefits *Pasteuria* because it liberates resources that the host might otherwise put into reproduction (Ebert *et al.* 2004). Thus while *Pasteuria* does, as expected, follow a trade-off regarding mortality (Jensen *et al.* 2006), selection for parasites of intermediate virulence is not expected for fecundity: parasites benefit by sterilizing hosts as early as possible.

We can only speculate as to the proximate cause of higher mortality in slow-growing infections. This isolate may produce high levels of toxins. Another possibility is that the early-killing isolate has a greater propensity to induce fecundity compensation in their hosts, and it is this extra effort towards early reproduction in hosts that leads to higher mortality, but we could find no evidence of this (age at first reproduction did not differ between spore types). Conceivably, because our parasite isolates may not be pure strains, interactions between bacterial strains within a spore type could mitigate virulence; for example the late-killing spore suspension may contain mitigating strains the early-killing suspension lacks. (Unfortunately, this possibility is untestable until more advanced microbiological techniques can be applied to *Pasteuria*.) A further possibility is that the quick-killing isolate stimulates a stronger immune response, leading to immunopathology or a cost associated with the energetic demands of launching a strong immune response. If this were the case, we might expect hosts that did not become infected (assuming that resistance involves an

immune response) to show higher mortality when resisting the quick-killing, more immunogenic strain. However, we also found no evidence that differences in survivorship between uninfected hosts were explained by spore type (data not shown).

The existence of a quick-killing, poorly infecting and slow-growing parasite isolate appears puzzling. It may be that this parasite isolate is simply a maladaptive form in decline when originally isolated. Alternatively, it may be that this slow-growing isolate does not show these features on all host genotypes. Parasite isolate by host genotype interactions are prevalent in the *D. magna-P. ramosa* system (Carius *et al.* 2001). The Carius *et al.* (2001) study on specificity was based on measures of infectivity and host fecundity. Part of the current result indicates that genetic specificity can be extended to include parasite growth rate, although the relationship between strains was qualitatively similar across host genotypes (it was the magnitude of the difference that changed). However, again, the full effect of parasite genotype by host genotype effects can only be assessed with substantial sampling of the genetic variation in both interactors. This raises the point that fully assessing evolution of virulence models (as well as models of co-evolution) might invariably require extensive sampling of both host and parasite genetic variation (see Grech *et al.* 2006; Lefebvre *et al.* 2007).

Alternatively, a quick-killing strategy could be successful under particular epidemiological conditions, for instance, during periods of high host density or population growth where the most successful parasites kill as quickly as possible and move on to new hosts. During summer gerbil-leishmania epidemics in Russia, for example, when the possibilities for transmission are high, a large proportion of infections are virulent, while a lower percentage of infections are virulent during other times of the year, when transmission opportunities are comparatively few (Dye and Davies, 1990). In the laboratory, *Leishmania* strains that cause high virulence out-compete strains that cause less virulence, but it is thought that the former will be selected against during the low transmission periods in the wild because of the excessive mortality they cause. This highlights the fact that the frequencies of parasite strains that cause different levels of virulence may fluctuate with transmission possibilities (see also Herre, 1995) and the epidemic state of the population, possibly leading to periods where slow-growing strains predominate. *Daphnia* population sizes fluctuate dramatically, and these appear to be associated with summer parasite epidemics (Duncan *et al.* 2006; Duncan and Little, 2007). We propose that quick-killing strains may only have an advantage during periods of high population growth and a spreading epidemic, and insight into this could be had by isolating parasites from a range of

time-points. Generally, associations between virulence and patterns of epidemiological spread (which are essentially genotype by environment interactions) are a topic deserving of further study.

We thank Jaap Deroode for comments and advice. This work was supported by grants from NERC and the Wellcome Trust.

REFERENCES

- Almog, G., Cohen, N., Stocker, S. and Stone, L.** (2002). Immune response and virus population composition: HIV as a case study. *Proceedings of the Royal Society of London, B* **269**, 809–815.
- Andre, J. B., Ferdy, J. B. and Godelle, B.** (2003). Within-host parasite dynamics, emerging trade-off, and evolution of virulence with immune system. *Evolution* **57**, 1489–1497.
- Bremermann, H. J. and Pickering, J.** (1983). A game-theoretical model of parasite virulence. *Journal of Theoretical Biology* **100**, 411–426.
- Bull, J. J.** (1994). Virulence. *Evolution* **48**, 1423–1437.
- Carius, H.-J., Little, T. J. and Ebert, D.** (2001). Genetic variation in a host–parasite association: Potential for coevolution and frequency dependent selection. *Evolution* **55**, 1136–1145.
- Choo, K., Williams, P. D. and Day, T.** (2003). Host mortality, predation and the evolution of parasite virulence. *Ecology Letters* **6**, 310–315.
- Day, T. and Proulx, S. R.** (2004). A general theory for the evolutionary dynamics of virulence. *American Naturalist* **163**, E40–E63.
- De Roode, J. C., Pansini, R., Cheesman, S. J., Helinski, M. E. H., Huijben, S., Wargo, A. R., Bell, A. S., Chan, B. H. K., Walliker, D. and Read, A. F.** (2005). Virulence and competitive ability in genetically diverse malaria infections. *Proceedings of the National Academy of Sciences, USA* **102**, 7624–7628.
- Duncan, A., Mitchell, S. E. and Little, T. J.** (2006). Parasite-mediated selection in *Daphnia*: the role of sex and diapause. *The Journal of Evolutionary Biology* **19**, 1183–1189.
- Duncan, A. B. and Little, T. J.** (2007). Parasite-driven genetic change in a natural population of *Daphnia magna*. *Evolution* **64**, 796–803.
- Dye, C. and Davies, C. R.** (1990). Glasnost and the great gerbil: virulence polymorphisms in the epidemiology of leishmaniasis. *Trends in Ecology and Evolution* **5**, 237–238.
- Ebert, D.** (1998). Experimental evolution of parasites. *Science* **282**, 1432–1435.
- Ebert, D., Carius, H.-J., Little, T. J. and Decaestecker, E.** (2004). The evolution of virulence when parasites cause host castration and gigantism. *American Naturalist* **164**, s19–s32.
- Ebert, D., Rainey, P., Embley, T. M. and Scholz, D.** (1996). Development, life cycle, ultrastructure and phylogenetic position of *Pasteuria ramosa* Metchnikoff 1888: rediscovery of an obligate endoparasite of *Daphnia magna* Straus. *Philosophical Transactions of the Royal Society of London, B* **351**, 1689–1701.
- Escriu, F., Fraile, A. and García-Arenal, F.** (2003). The evolution of virulence in a plant virus. *Evolution* **57**, 755–765.
- Frank, S. A.** (1996). Models of parasite virulence. *Quarterly Review of Biology* **71**, 37–78.
- Gandon, S., Mackinnon, M. J., Nee, S. and Read, A. F.** (2001). Imperfect vaccines and the evolution of pathogen virulence. *Nature, London* **414**, 751–755.
- Grech, K., Watt, K. and Read, A. F.** (2006). Host–parasite interactions for virulence and resistance in a malaria model system. *Journal of Evolutionary Biology* **19**, 1620–1630.
- Herre, E. A.** (1995). Factors affecting the evolution of virulence: nematode parasites of fig wasps as a case study. *Parasitology* **111** (Suppl.), S179–S191.
- Jensen, K. N., Little, T. J., Skorpington, A. and Ebert, D.** (2006). Empirical support for an optimal virulence in a castrating parasite. *Plos Biology* **4**, e197. doi:10.1371/journal.pbio.0040197.
- Lefebvre, T., Sanchez, M., Ponton, F., Hughes, D. and Thomas, F.** (2007). Virulence and resistance in malaria: who drives the outcome of infection. *Trends in Parasitology* **23**, 299–302.
- Lipsitch, M. and Moxon, E. R.** (1997). Virulence and transmissibility of pathogens: What is the relationship? *Trends in Microbiology* **5**, 31–37.
- Little, T. J. and Ebert, D.** (2000). The cause of parasitic infection in natural populations of *Daphnia*: the role of host genetics. *Proceedings of the Royal Society of London, B* **267**, 2037–2042.
- Little, T. J. and Killick, S. C.** (2007). Evidence for a cost of immunity when the crustacean *Daphnia magna* is exposed to the bacterial pathogen *Pasteuria ramosa*. *Journal of Animal Ecology* **76**, 1202–1207.
- Little, T. J., Watt, K. and Ebert, D.** (2006). Parasite-host specificity: experimental studies on the basis of parasite adaptation. *Evolution* **60**, 31–38.
- Mackinnon, M. J. and Read, A. F.** (1999). Genetic relationships between parasite virulence and transmission in the rodent malaria *Plasmodium chabaudi*. *Evolution* **53**, 689–703.
- May, R. M. and Anderson, R. M.** (1983). Epidemiology and genetics in the coevolution of parasites and hosts. *Proceedings of the Royal Society of London, B* **219**, 281–313.
- Pagan, I.** (2007). The relationship of within-host multiplication and virulence in a plant-virus system. *PLoS ONE* **2**, e786. doi:10.1371/journal.pone.000786.
- Read, A. F. and Taylor, L. H.** (2001). The ecology of genetically diverse infections. *Science* **292**, 1099–1102.
- Sacristán, S., Fraile, A., Malpica, J. M. and García-Arenal, F.** (2005). An analysis of host adaptation and its relationship with virulence in cucumber mosaic virus. *Phytopathology* **95**, 827–833.
- Sas Institute Inc** (2002). *SAS, Version 9.0*. SAS Institute Inc., Cary, NC, USA.
- Stewart, A. D., Logsdon, J. M. and Kelley, S. E.** (2005). An empirical study of the evolution of virulence under both horizontal and vertical transmission. *Evolution* **59**, 730–739.
- Weiss, R. A.** (2002). Virulence and pathogenesis. *Trends in Microbiology* **10**, 314–317.