THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Differential distribution of WC1(+) gamma delta TCR(+) T lymphocyte subsets within lymphoid tissues of the head and respiratory tract and effects of intranasal M. bovis BCG vaccination

Citation for published version:

Price, S, Davies, M, Villarreal-Ramos, B & Hope, J 2010, 'Differential distribution of WC1(+) gamma delta TCR(+) T lymphocyte subsets within lymphoid tissues of the head and respiratory tract and effects of intranasal M. bovis BCG vaccination', *Veterinary Immunology and Immunopathology*, vol. 136, no. 1-2, pp. 133-137. <https://doi.org/10.1016/j.vetimm.2010.02.010>

Digital Object Identifier (DOI):

<http://dx.doi.org/10.1016/j.vetimm.2010.02.010>

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Early version, also known as pre-print

Published In:

Veterinary Immunology and Immunopathology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Short communication

Differential distribution of WC1⁺ $\gamma\delta$ TCR⁺ T lymphocyte subsets within lymphoid tissues of the head and respiratory tract and effects of intranasal *M. bovis* BCG vaccination

Sally Price, Marc Davies, Bernardo Villarreal-Ramos, Jayne Hope*

Institute for Animal Health, Compton, High Street, Newbury, Berkshire RG20 7NN, United Kingdom

ARTICLE INFO

Article history:

Received 18 December 2008

Received in revised form 29 January 2010

Accepted 12 February 2010

Keywords:

Bovine

Gamma delta T cell

*Mycobacterium bovis*IFN γ

ABSTRACT

BCG vaccination of neonatal calves induces significant protection against tuberculosis. The enhanced protection observed in neonatal calves may be linked to an enhanced capacity for IFN γ production by innate cells, including WC1⁺ $\gamma\delta$ T cells, which constitute a major population in young cattle. Intranasal BCG vaccination of mice induces high levels of IFN γ in the lungs, which may enhance protection against subsequent challenge with virulent strains of mycobacteria. We used an intranasal BCG vaccination model in calves to study the effect on the distribution of WC1⁺ $\gamma\delta$ T cells expressing two alternate forms of WC1: WC1.1 and WC1.2. These subsets of WC1⁺ $\gamma\delta$ T cells have previously been shown to have a differential capacity for IFN γ secretion. Our results indicate that there is a selective expansion/recruitment of $\gamma\delta$ T cells expressing the IFN γ -associated WC1.1 isoform in tissues of the lungs and upper respiratory tract following intranasal BCG vaccination.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

T cells expressing the alternate receptor consisting of γ and δ chains (often referred to as TCR1) are found in all vertebrate species, though the relative numbers of $\gamma\delta$ -expressing T cells varies. For example, in humans and mice, $\gamma\delta$ T cells comprise ~10% of the circulating T cells, while in ruminants they may constitute up to 50–60%. There are two distinct subsets of bovine $\gamma\delta$ T cells, based on expression of workshop cluster antigen-1 (WC1). WC1 is a 220 kDa glycoprotein, encoded by a complex family of genes in cattle, sheep, goats, pigs and camelids (Mackay et al., 1986, 1989; Clevers et al., 1990; Carr et al., 1994) and also in humans and mice, although in these the antigen is not expressed and the gene family is less complex. In cattle,

13 members of the WC1 gene family have been identified, on two loci on chromosome 5 (Herzig and Baldwin, 2009). The extra-cellular domains of WC1 share homology with the scavenger receptor cysteine-rich (SRCR) family, though its function remains enigmatic. It has been postulated that WC1 may be a pattern recognition receptor or a regulator of lymphocyte receptor signalling (Herzig and Baldwin, 2009).

Distinct functional sub-populations of bovine WC1⁺ $\gamma\delta$ T cells have been identified, characterised by the expression of either the WC1.1 or WC1.2 isoform (Rogers et al., 2005a; Rogers et al., 2005b; Chen et al., 2004). Within the WC1⁺ population, the WC1.1⁺ sub-population is the principal IFN γ -secreting population, while the WC1.2-expressing cells are more responsive to mitogen stimulation. It is likely that these two sub-types of WC1⁺ $\gamma\delta$ T cell play discrete roles in the innate immune response, and may display distinct behaviour in response to infection or vaccination.

Potential roles for $\gamma\delta$ T cells in innate immune responses have been investigated in cattle. Depletion of WC1⁺ $\gamma\delta$ from *Mycobacterium bovis* infected calves increased levels of IL-

Abbreviations: WC, workshop cluster; FCM, flow cytometry.

* Corresponding author. Tel.: +44 01635 577239;

fax: +44 01635 577263.

E-mail address: jayne.hope@bbsrc.ac.uk (J. Hope).

4, decreased IgG2 and, importantly, decreased secretion of innate IFN γ (Kennedy et al., 2002) suggesting that these cells have a role in directing Th1 bias. WC1⁺ $\gamma\delta$ T cells are among the first cells to accumulate at sites of infection and concurrently reduce in number in the peripheral blood, implying a migration from the blood to the tissues (Pollock et al., 1996). Bovine WC1⁺ $\gamma\delta$ T cells may play a significant role in shaping the adaptive immune response through early interactions with dendritic cells, resulting in secretion of significant amounts of IFN γ (Price et al., 2007).

Bovine tuberculosis caused by *M. bovis* is a major economic and potential zoonotic problem in the UK. Use of the BCG vaccine in cattle has been hampered by the fact that it has not been possible to distinguish between infected and vaccinated animals by current diagnostic tests, and that BCG vaccination may have variable efficacy as seen in humans (Suazo et al., 2003). Experimentally, BCG vaccination is effective against bovine TB (Hope and Vordermeier, 2005) with BCG vaccination of neonatal calves eliciting significant protection against challenge with *M. bovis* (Hope et al., 2005; Buddle et al., 2003). Several studies in mice and cattle have indicated that enhanced protection against experimental infection with virulent mycobacteria may be achieved by delivering BCG via the intratracheal route (Buddle et al., 1995; Falero-Diaz et al., 2000). This protective effect is linked to rapid production of IFN γ production in the lungs in mice (Lyadova et al., 2001). In addition, intranasal infection of mice with *M. bovis* BCG was shown to induce increased levels of $\gamma\delta$ T cells by 7 days post-infection, and $\gamma\delta$ T cells isolated from the lungs of infected mice showed high levels of IFN γ production and cytotoxic activity against infected macrophages (Dieli et al., 2003). These studies indicate that effects on sub-populations of lymphocytes post-vaccination are of key importance in the development of protective immune responses.

In order to investigate potential roles for WC1⁺ $\gamma\delta$ T cells in vaccine-induced responses in calves we studied the distribution of WC1⁺ $\gamma\delta$ T cells in the head and respiratory tract following intranasal BCG vaccination. Differences in the distribution of cells expressing the alternate isoforms of WC1 were also assessed. We demonstrated functionally relevant alterations in WC1⁺ cells in calves 1 week following intranasal BCG vaccination.

2. Materials and methods

2.1. Intranasal BCG vaccination

Four 5–10-month-old Friesian calves were selected on the basis of their lack of response to mycobacterial antigens (purified protein derivatives from *M. avium* (PPD-A) and *M. bovis* (PPD-B) as previously described (Hope et al., 2005)). Calves were vaccinated intranasally by introduction of a 6" long catheter into the left nostril. Approximately 1.5×10^7 cfu of *Mycobacterium bovis* Bacille Calmette Guerin strain Pasteur (BCG Pasteur) suspended in 2 ml of sterile PBS was delivered into the nostril, followed by 2 ml of PBS alone. The calves showed no adverse effects of vaccination. After 7 days, the animals were killed by captive bolt. Tissues from three age-matched unvaccinated calves were used for control samples. All animal procedures used were approved

Table 1

Tissues taken post-mortem.

Tissue	Location	Abbreviation
Cardiac (medial) lung (right)	Lung	RC
Cardiac (medial) lung (left)	Lung	LC
Intermediate lung	Lung	I
Diaphragmatic lung (right)	Lung	RD
Diaphragmatic lung (left)	Lung	LD
Pallatine tonsil (right)	Distal palate	Pall R
Pallatine tonsil (left)	Distal palate	Pall L
Retropharyngeal lymph node (right)	Pharynx	RPLN R
Retropharyngeal lymph node (left)	Pharynx	RPLN L
Pharyngeal tonsil	Pharynx	Ph ton
Cervical lymph node	Draining distal trachea	CLN

by the local ethics committee, consisting of expert and lay members.

2.2. Isolation of peripheral blood mononuclear cells

PBMC were isolated from blood collected at the time of slaughter, as previously described (Price et al., 2007). Cells were re-suspended at 2×10^5 cells per ml in PBS with 0.1% BSA and 0.01% sodium azide (PBS/BSA/Azide) for immunostaining.

2.3. Isolation of mononuclear cells from tissues

Pieces of tissue approximately 2.5 cm³ and selected peripheral lymph nodes were removed at post-mortem (Table 1) and washed thoroughly in sterile PBS. The lymph nodes were sliced open and flushed repeatedly with PBS to liberate cells from within the tissue. Other tissue samples were homogenised in a pestle and mortar containing sterile PBS. The resulting cell suspensions were passed through a 40 μ m nylon cell strainer (Becton Dickinson, Oxford, UK). The mononuclear cells were then isolated by density gradient centrifugation over Histopaque 1083 (Sigma Aldrich, Gillingham, UK) as described (Price et al., 2007). Cells were re-suspended at 2×10^5 cells per ml for immunostaining.

2.4. Flow cytometric analysis

Cells suspended at 2×10^5 per ml were incubated for 10 min with 10% normal mouse serum. Cells were then incubated with either an isotype-matched control antibody or antibody to the bovine $\gamma\delta$ T cell receptor TCR1 (mAb GB21a; IgG2b, kindly provided by Dr W. Davis, WSU, Pullman, USA) in conjunction with either anti-pan WC1 (CC39-Alexa Fluor⁶⁴⁷ IgG1; AbD-Serotec, Oxford, UK) anti-WC1.1 (mAb BAQ159A; IgG1, W. Davis) or anti-WC1.2 (mAb CACTB32A; IgG1, W. Davis) antibodies. Isotype-specific goat anti-mouse secondary reagents were then added (goat anti-mouse IgG2b tri-colour or IgG2b FITC; goat anti-mouse IgG1 phycoerythrin; Southern Biotechnology Associates, Birmingham, USA). Flow cytometric acquisition was carried out on a FACSCalibur flow cytometer (Becton Dickinson, Oxford, UK) and the data was analysed

using FCS Express version 3 (DeNovo Software, Ontario, Canada). A minimum of 200,000 events were collected.

2.5. Statistical analyses

Statistical analyses were carried out with Minitab release 15.0 (Minitab Inc, Coventry, UK) using a Mann

Whitney non-parametric test and p values of <0.05 were considered significant.

3. Results and discussion

In order to establish a potential role for $\gamma\delta$ TCR⁺ T lymphocytes in BCG vaccination induced immune responses

Fig. 1. Percentages of $\gamma\delta$ T cell subsets in control, non-vaccinated, cattle. The percentage of cells expressing the $\gamma\delta$ TCR (TCR1; a) and percentages of TCR1⁺ T cells expressing WC1 (b), WC1.1 (c) or WC1.2 (d) were assessed in tissues of the respiratory tract and head (see Table 1 for abbreviations) by FCM. The ratio of WC1.1:WC1.2 cells is shown in (e). Individual animal data is shown with means (solid bars). Statistically significant differences were assessed using Mann Whitney non-parametric analysis. *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$.

we measured the percentage of T cells expressing the $\gamma\delta$ TCR, and subsets of $\gamma\delta$ T cells expressing isoforms of the WC1 receptor, in lymphoid tissues of the head and respiratory tract of cattle. Control, non-vaccinated calves were compared with calves that had been vaccinated intranasally with BCG 7 days prior to analysis.

In control, non-vaccinated calves, significant differences in the percentages of $\gamma\delta$ T cell subsets were noted between tissues (Fig. 1). There were significantly higher numbers of T lymphocytes expressing the $\gamma\delta$ TCR (TCR1) in the lung lobes compared to the lymphoid tissues of the head (Fig. 1a; $p < 0.01$), but the number of $\gamma\delta$ TCR⁺ T cells was lower in all tissues than in the peripheral blood ($p < 0.001$). Within the $\gamma\delta$ TCR-expressing T lymphocyte population there were

also tissue specific differences noted in the percentage of cells expressing the WC1 receptor (Fig. 1b). Notably, there was a significantly higher number of $\gamma\delta$ T cells expressing WC1 in the lung lobes and blood, when compared to lymphoid tissues of the head ($p < 0.001$). The percentage of $\gamma\delta$ T cells expressing WC1.1 (Fig. 1c) or WC1.2 (Fig. 1d) also differed between tissues. Significantly greater percentages of WC1.1 and WC1.2 expressing cells were present in the lung lobes and peripheral blood compared to head lymphoid tissues ($p < 0.001$). Assessment of the relative ratio of WC1.1:WC1.2 expressing $\gamma\delta$ T cells within tissues (Fig. 1e) revealed a greater ratio of WC1.1:WC1.2 within the lymphoid tissues of the head ($p < 0.05$). The ratio in the lung and blood was less than 1 indicating a predominance of WC1.2

Fig. 2. Comparison of $\gamma\delta$ T cell subsets in control and BCG vaccinated cattle. The percentage of cells expressing the $\gamma\delta$ TCR (TCR1; a) and percentages of TCR1⁺ T cells expressing WC1 (b), WC1.1 (c) or WC1.2 (d), WC1.1 or WC1.2 were assessed as for Fig. 1. Mean $\pm$ SE is illustrated. Differences between control (open symbols) and vaccinated (closed symbols) calves were assessed by Mann Whitney non-parametric statistical analysis. * $p < 0.05$.

expressing $\gamma\delta$ T cells. By comparison, within the lymphoid tissues of the head the WC1.1:WC1.2 ratio was approximately 1 suggesting a more balanced population of cells expressing either of the two isoforms of WC1. Given the suggested functional differences associated with expression of either WC1.1 or WC1.2, this may have implications for immune responses being induced in different tissues (Rogers et al., 2005a, b).

The percentage of $\gamma\delta$ TCR-expressing cells and the relative proportions which expressed WC1 and the isoforms WC1.1 and WC1.2 was subsequently assessed in calves that had been intranasally vaccinated with BCG (Fig. 2). A significant increase in the percentage of $\gamma\delta$ TCR-expressing T cells (Fig. 2a) and of WC1⁺ $\gamma\delta$ TCR⁺ T cells (Fig. 2b) was observed in the pharyngeal tonsil ($p < 0.05$) of BCG vaccinated calves (Fig. 2; closed symbols). Significant differences in the percentage of $\gamma\delta$ TCR⁺ T cells expressing WC1.1 (Fig. 2c), but not WC1.2 (Fig. 2d) were observed in each of the lung regions assessed and in the pharyngeal tonsil ($p < 0.05$). Increased percentages of WC1.1 expressing cells were observed in each of the tissues assessed although, due to large animal to animal variation, this did not reach statistical significance for the head lymph nodes, or blood. The ratio of WC1.1:WC1.2 was significantly increased within the draining pharyngeal tonsil (Fig. 2e; $p < 0.05$).

These data indicate that intranasal BCG vaccination induces a predominance of WC1.1⁺ cells in the tissues of the respiratory tract. Selective recruitment and/or expansion of WC1.1⁺ $\gamma\delta$ T cells, which have been shown to have capacity for high level IFN γ secretion, may play an important role in the induction of the protective immune response induced by BCG vaccination in cattle.

Conflict of interest

None of the authors have any conflict of interest.

Acknowledgments

This work was funded by the Biotechnology and Biological Sciences Research Council (BBSRC). Jayne C. Hope is a Jenner Investigator. The authors would like to thank Nazneen Siddiqui for assistance with post-mortems and tissue preparation. We also gratefully acknowledge the animal services staff at IAH for care of the cattle used within these experiments.

References

Buddle, B.M., Keen, D., Thomson, A., Jowett, G., McCarthy, A.R., Heslop, J., De Lisle, G.W., Stanford, J.L., Aldwell, F.E., 1995. Protection of cattle from

- bovine tuberculosis by vaccination with BCG by the respiratory or subcutaneous route, but not by vaccination with killed *Mycobacterium vaccae*. Res. Vet. Sci. 59, 10–16.
- Buddle, B.M., Wedlock, D.N., Parlange, N.A., Corner, L.A., De Lisle, G.W., Skinner, M.A., 2003. Revaccination of neonatal calves with *Mycobacterium bovis* BCG reduces the level of protection against bovine tuberculosis induced by a single vaccination. Infect. Immun. 71, 6411–6419.
- Carr, M.M., Howard, C.J., et al., 1994. Expression on porcine gamma delta lymphocytes of a phylogenetically conserved surface antigen previously restricted in expression to ruminant gamma delta T lymphocytes. Immunology 81, 36–40.
- Chen, L., Wang, J., et al., 2004. Single intranasal mucosal *Mycobacterium bovis* BCG vaccination confers improved protection compared to subcutaneous vaccination against pulmonary tuberculosis. Infect. Immun. 72, 238–246.
- Clevers, H., MacHugh, N.D., Bensaid, A., Dunlap, S., Baldwin, C.L., Kaushal, A., Iams, K., Howard, C.J., Morrison, W.I., 1990. Identification of a bovine surface antigen uniquely expressed on CD4⁺CD8⁺ T cell receptor gamma/delta⁺ T lymphocytes. Eur. J. Immunol. 20, 809–817.
- Dieli, F., Ivanyi, J., Marsh, P., Williams, A., Naylor, I., Sireci, G., Caccamo, N., Di Sano, C., Salerno, A., 2003. Characterization of Lung {gamma}{delta} T Cells Following Intranasal Infection with *Mycobacterium bovis* Bacillus Calmette–Guerin. J. Immunol. 170, 463–469.
- Falero-Diaz, G., Challacombe, S., Banerjee, D., Douce, G., Boyd, A., Ivanyi, J., 2000. Intranasal vaccination of mice against infection with *Mycobacterium tuberculosis*. Vaccine 18, 3223–3229.
- Herzig, C.T., Baldwin, C.L., 2009. Genomic organization and classification of the bovine WC1 genes and expression by peripheral blood gamma delta T cells. BMC Genom. 10, 191.
- Hope, J.C., Thom, M.L., Villarreal-Ramos, B., Vordermeier, H.M., Hewinson, R.G., Howard, C.J., 2005. Vaccination of neonatal calves with *Mycobacterium bovis* BCG induces protection against intranasal challenge with virulent *M. bovis*. Clin. Exp. Immunol. 139, 48–56.
- Hope, J.C., Vordermeier, H.M., 2005. Vaccines for bovine tuberculosis: current views and future prospects. Expert. Rev. Vaccines 4, 891–903.
- Kennedy, H.E., Welsh, M.D., Bryson, D.G., Cassidy, J.P., Forster, F.I., Howard, C.J., Collins, R.A., Pollock, J.M., 2002. Modulation of immune responses to *Mycobacterium bovis* in cattle depleted of WC1(+) gamma delta T cells. Infect. Immun. 70, 1488–1500.
- Lyadova, I.V., Vordermeier, H.M., Eruslanov, E.B., Khaidukov, S.V., Apt, A.S., Hewinson, R.G., 2001. Intranasal BCG vaccination protects BALB/c mice against virulent *Mycobacterium bovis* and accelerates production of IFN-gamma in their lungs. Clin. Exp. Immunol. 126, 274–279.
- Mackay, C.R., Beya, M.F., et al., 1989. Gamma/delta T cells express a unique surface molecule appearing late during thymic development. Eur. J. Immunol. 19, 1477–1483.
- Mackay, C.R., Maddox, J.F., et al., 1986. Three distinct subpopulations of sheep T lymphocytes. Eur. J. Immunol. 16 (1), 19–25.
- Pollock, J.M., Pollock, D.A., Campbell, D.G., Girvin, R.M., Crookard, A.D., Neill, S.D., Mackie, D.P., 1996. Dynamic changes in circulating and antigen-responsive T-cell subpopulations post-*Mycobacterium bovis* infection in cattle. Immunology 87, 236–241.
- Price, S.J., Sopp, P., Howard, C.J., Hope, J.C., 2007. Workshop cluster 1⁺ gammadelta T-cell receptor T cells from calves express high levels of interferon-gamma in response to stimulation with interleukin-12 and -18. Immunology 120, 57–65.
- Rogers, A.N., VanBuren, D.G., Hedblom, E., Tilahun, M.E., Telfer, J.C., Baldwin, C.L., 2005a. Function of ruminant [gamma][delta] T cells is defined by WC1.1 or WC1.2 isoform expression. Vet. Immunol. Immunopathol. 108, 211–217.
- Rogers, A.N., Vanburen, D.G., Hedblom, E.E., Tilahun, M.E., Telfer, J.C., Baldwin, C.L., 2005b. Gammadelta T cell function varies with the expressed WC1 coreceptor. J. Immunol. 174, 3386–3393.
- Suazo, F.M., Escalera, A.M.A., Torres, R.M.G., 2003. A review of *M. bovis* BCG protection against TB in cattle and other animals species. Prevent. Vet. Med. 58, 1–13.