THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The spatial distribution of campylobacteriosis in New Zealand, 1997-2005

Citation for published version:

Rind, E & Pearce, J 2010, 'The spatial distribution of campylobacteriosis in New Zealand, 1997-2005', *Epidemiology & Infection*, vol. 138, no. 10, pp. 1359-1371. <https://doi.org/10.1017/S095026881000018X>

Digital Object Identifier (DOI):

[10.1017/S095026881000018X](https://doi.org/10.1017/S095026881000018X)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Epidemiology & Infection

Publisher Rights Statement:

Published by Cambridge University Press (2010)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The spatial distribution of campylobacteriosis in New Zealand, 1997–2005

E. RIND^{1,3*} AND J. PEARCE^{1,2}

¹ GeoHealth Laboratory, Department of Geography, University of Canterbury, New Zealand

² Institute of Geography, School of GeoSciences, University of Edinburgh, UK

³ School of Environmental Sciences, University of East Anglia, Norwich, UK

(Accepted 4 January 2010; first published online 9 February 2010)

SUMMARY

In New Zealand *Campylobacter* infection rates have increased steadily since 1980, reaching a peak in 2003 (396/100 000 population). Compared to other nations, disease rates are unfavourably high (e.g. Australia 117/100 000 population, UK 85/100 000 population, USA 13/100 000 population). This ecological study investigated spatial variations in *Campylobacter* infection rates across New Zealand's Territorial Local Authorities (TLAs, $n = 73$) for the period 1997–2005. Applying multiple linear regression, we examined whether geographical factors such as socio-demographic characteristics, climate, land use, water and the food environment were associated with local differences in the occurrence of *Campylobacter* infection rates. The results suggested significant variations in campylobacteriosis across TLAs (average annual rates ranging from 97 to 526/100 000 population), with higher rates in the South Island. Disease rates were associated with lower socio-economic deprivation ($P < 0.01$), the proportion of the population aged 25–44 years ($P < 0.01$) and fresh food outlet density ($P < 0.76$). The results underline the role of area-level characteristics in explaining the spatial distribution of campylobacteriosis in New Zealand. In particular, the findings draw attention to the relatively unexplored role of fresh food outlets as a potential risk factor for increased *Campylobacter* notifications.

Key words: Campylobacteriosis, ecological study, geographical determinants, New Zealand.

INTRODUCTION

Although different surveillance systems complicate international comparisons, New Zealand's rates of campylobacteriosis are among the highest reported rates worldwide (Fig. 1). Rates in New Zealand have increased from 14/100 000 population in 1980, when the disease became notifiable, to 396/100 000 population in 2003 [1]. For this increase no conclusive

reasons have yet been clarified. While the disease is usually self-limiting within 3–7 days, severe sequelae can arise, including reactive arthritis (1%), respiratory and severe neurological dysfunctions (0.1%), or death (0.1%) [2, 3]. Hospitalization arises in about 13% of cases [2]. The total annual economic costs (medical and non-medical) of *Campylobacter* infections in New Zealand have been estimated at NZ\$40 million, which is about 73% of the country's total economic cost of foodborne infectious diseases [4].

Regulated by the Health Act 1956 and the Tuberculosis Act 1948, any suspected or diagnosed case of campylobacteriosis is required to be reported

* Author for correspondence: E. Rind, School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK.
(Email: E.Rind@uea.ac.uk)

Fig. 1. Notification rates for selected affluent nations in 2003. (Adapted from Baker *et al.* 2006 [1].)

to the local public health service. On suspicion of a case, clinicians and general practitioners (GPs) request laboratory tests and send a notification message to the respective public health unit (PHU) on confirmation of an infection. Since 2007 laboratories are required to report test results directly to one of the 20 PHU offices across New Zealand where the national notifiable disease database (EpiSurv) is maintained [5, 6]. EpiSurv provides the basis for surveillance activities including the identification and monitoring of campylobacteriosis, the investigation of risk factors, and the evaluation of morbidity and mortality trends.

Across New Zealand, there are distinct temporal and spatial variations in the distribution of *Campylobacter* rates. One recent national study identified three dominant spatial and seasonal patterns: low summer incidence and low inter-seasonal variation in rural areas mainly across the North Island; higher summer incidence and greater degree of seasonality in urban areas mainly across the North Island; and highest summer incidence and greatest inter-seasonal variation in urban areas mainly across the South Island [7]. These observed trends are generally consistent with those reported from other affluent nations such as the UK, Denmark and Finland [8–10]. However, the explanation for the spatial distribution of campylobacteriosis in New Zealand is poorly understood, and study designs adopting a spatial approach to consider the aetiology of campylobacteriosis are rare. Earlier research can be grouped into (i) studies which have examined underlying environmental factors, including factors affecting the distribution of *Campylobacter* (e.g. climate, land use patterns), and (ii) work concerned with characteristics of the population and associated reporting or surveillance

mechanisms (e.g. demographic characteristics, deprivation) [11, 12].

In order to explain the geographical distribution of *Campylobacter*, rainfall and different water sources have received some attention [9, 13, 14]. Heavy rainfall often initiates extreme run-off events which can result in a significant increase of microbiological loads in surface waters. In the environment, the pathogen occurs more frequently during the winter months because the organism grows well in water below 10 °C. However, freezing, drying, acid conditions (pH ≤ 5), UV light, and salinity reduce its growth [8, 9, 15, 16].

Further, drinking water and recreational water have been considered to be significant transmission routes from the environment to humans, and the faecal indicator organism *E. coli* can be used to estimate the possible presence of *Campylobacter* in different water sources [17]. A study in Sweden investigated associations between the spatial distribution of *Campylobacter* incidence and environmental factors related to water and livestock, and water was concluded to be an important transmission route for *Campylobacter* infections [11]. In New Zealand, Savill *et al.* [18] found *Campylobacter* prevalent in shallow groundwater (75%), river water (60%), roof water (38%), and drinking water (29%). In particular, private wells and roof-water supplies in rural areas are at risk of being faecally contaminated from cattle and wild birds, respectively, as these water supplies remain predominantly untreated [14, 19, 20]. A recent study investigating the microbiological quality of roof-collected rainwater of private dwellings in New Zealand concluded that at least 30% of the samples collected were faecally contaminated [21].

Pastoral farming is another key aspect in determining the survival and reproduction of *Campylobacter* in the environment [20]. This factor is likely to be particularly pertinent in New Zealand where about 40% of the land cover is pastoral (e.g. high- and low-producing grassland, scrub/tussock) [22, 23]. Studies investigating the appearance of *Campylobacter* in rural areas of New Zealand [24, 25] identified the pathogen in human as well as in animal faeces, including faeces from dairy cattle, beef cattle, sheep, pigs and ducks. The results indicate that for the examined rural population a range of transmission routes can be related to infected animals. In particular, farm and meat-processing workers, abattoir and meat freezing-plant workers, and veterinarians are at risk of becoming infected [26, 27].

Other risk factors include diet, particularly the consumption of fresh poultry and fresh fruits or vegetables from retail outlets or restaurants, ready-to-eat chicken products and other fast foods [28].

Various characteristics of the exposed population are also significant in understanding the spatial distribution of campylobacteriosis. In New Zealand, the average annual *Campylobacter* incidence for the period 2001–2003 was significantly higher in males (326/100 000 population) than in females (296/100 000 population), as well as in children (aged <15 years) and young adults (aged <45 years). The reasons for children and young males being particularly susceptible to developing campylobacteriosis have not been clarified. Explanations range from underlying immunological differences between the genders, and children being more likely to see a doctor and thus their infection being notified more frequently. In New Zealand, Europeans show the highest notification rates (329/100 000 population) compared to all other ethnic groups (Māori 94/100 000 population; Pacific people 70/100 000 population; other ethnicities 193/100 000 population) which is linked frequently to socio-economic deprivation. It has been suggested that some of the ethnic differences are associated with poorer access to primary care and health services resulting in lower notification rates for non-European ethnicities compared to Europeans [1]. Prior studies from New Zealand found, for example, a negative relationship between disease rates and unsafe water supplies (e.g. outdated sanitary installations, less monitored water supplies) occurring more frequently in socio-economically deprived areas compared to less deprived areas [20, 29]. Moreover, differences in laboratory procedures (e.g. detection techniques) or

reporting practices between GPs may partially account for the observed spatial patterns in New Zealand [1, 30]. Although underestimation of absolute *Campylobacter* rates has not been quantified, a study from Auckland found that only 49% of the actual cases of giardiasis were notified which indicates that the notified *Campylobacter* cases might only capture a fraction of the true prevalence of the disease [31].

This ecological study investigates the spatial distribution of campylobacteriosis in New Zealand for the period 1997–2005. The study has two objectives. First, we identify areas with particularly high and low rates of the disease, and examine whether there are parts of the country where rates have increased more than in others. Second, we identify whether geographical factors such as climate, land use, water and the food retail environment as well as local socio-demographic characteristics are associated with the spatial distribution of campylobacteriosis. Approaching this topic with a spatial focus will contribute to a better understanding of the mechanisms affecting the distribution of campylobacteriosis in New Zealand, a basic requirement for reducing rates and the costs involved.

DATA AND METHODS

Data sources and data manipulation

For all Territorial Local Authorities (TLAs), excluding the Chatham Islands ($n=73$), the campylobacteriosis notification data (cases/100 000 population per TLA=dependent variable) were obtained from the notifiable disease database EpiSurv for each of the years 1997–2005 [32]. TLAs are statistical units with a median population of 27 393 in 2001 (Fig. 2). Due to confidentiality constraints, no individual-level information was available.

For each TLA, a range of potential area-level explanatory variables were developed (see Table 1). The climate data [33] were aggregated to the TLA level. An index for the degree of seasonal change in *Campylobacter* rates was calculated for each TLA using data published elsewhere [7]. To emphasize higher summer incidence and seasonal variation in certain TLAs, this index was weighted, using the three types of seasonal grouping developed by Hearnden *et al.* [7]. To ensure the comparability of different loads of livestock as a measurement of pastoral intensity, the total number of livestock (cattle, horses, pigs, sheep, goats, deer, poultry) were transformed into

Fig. 2. Territorial Local Authorities in New Zealand.

stock units (stock units/km², stock units/capita) (e.g. 1 dairy cattle = 1 stock unit; 1 sheep = 0.2 stock units) [23, 34]. In order to estimate the possibility of becoming infected from contaminated food, it was assumed that the risk of exposure increased with better access to food retailing. For each TLA, the density of restaurants, fast food outlets, fresh food outlets and supermarkets was calculated (proportion of food outlets and supermarkets/100 000 population) [35].

As outlined above, children are particularly at risk of becoming infected [1]. Both of the <5 and 5–14

years age groups were significantly related with the *Campylobacter* rate (results not shown) and combined as one age group (<5–14 years) to obtain a greater population. To represent socio-economic characteristics of the affected population, a TLA-level measure was derived by aggregating the 2001 New Zealand Deprivation index from the small Census Area Units [36]. The deprivation index is a population-weighted area-level aggregation of nine variables from the 2001 New Zealand Census including, for example, unemployment and low professional education. The

higher the index score, the greater the level of socioeconomic disadvantage [36]. For all other variables, the original data were either available at or aggregated to the TLA level.

Analysis

In order to determine whether there was an increase or decrease in *Campylobacter* rates within the study period, the percentage change in the number of incidents reported for 2005 compared to 1997 was calculated per TLA. A standard deviation (s.d.) classification scheme was used to identify areas deviating significantly from the national mean, calculated for the entire study period. The spatial autocorrelation statistic, Moran's *I*, was calculated to assess statistical significance of spatial clustering which is indicated when the results of the test are >0 . For these analyses ArcMap™ 9.1 (ESRI Inc., USA) was used.

Linear regression analysis was applied to investigate whether each of the potential explanatory variables were associated with the aggregated TLA-level rates of campylobacteriosis (1997–2005), using SPSS 13.0 (SPSS Inc., USA). Our *a priori* assumptions were that all but one of the explanatory variables showed positive associations with the *Campylobacter* rate; for the variable 'temperature' an inverse relationship was expected. The independent variables were entered in a predetermined order based on theoretical considerations [37, 38]. First, 23 univariate models were derived. The subsequent multivariate modelling steps explored how the effects of each of the variables changed with increasing complexity of the regression models within the predefined variable categories, excluding those coefficients for which effects of multicollinearity were observed. Multicollinearity was indicated when high correlation coefficients occurred between the independent variables. Tabachnick & Fidell [39] recommend not to include two variables with a bivariate correlation close to $r \geq 0.7$. Modelling step 2 incorporated five models (climate/seasonal change, land use, water, food, demographic characteristics of the affected population), excluding the categories of 'rurality/occupation', 'social deprivation' and 'artefacts' since each of those included only one variable. Modelling step 3 consisted of two models incorporating those variables classified into the categories 'conditions affecting the distribution of the organism' and 'exposure'. The fourth modelling step consisted of two models describing the associations

between the variables classified in the categories 'underlying environmental factors' and 'surveillance characteristics'. The fifth modelling step combined all of the variables. Subsequently, the standardized residuals of the final model were tested for the assumption of normality of the error term (one-sample Kolmogorov–Smirnov procedure) and spatial interdependency (Durbin–Watson statistic).

RESULTS

The spatial distribution of campylobacteriosis

The average annual rate across all TLAs for the entire study period was 285/100 000 population. The rates of campylobacteriosis for the period 1997–2005 were not uniform across New Zealand, ranging from 97/100 000 population (Opotiki District) to 526/100 000 population (Wellington City). Over the study period, there has been an increase of *Campylobacter* rates in 85% of the TLAs (62/73 TLAs) (Fig. 3). The highest increase was in Nelson City (+377%), followed by the Wairoa District (+363%), and the Central Hawke's Bay District (+270%). Of the 11 TLAs with a rate reduction, the largest decrease was in the Banks Peninsula District (−71%), followed by the Opotiki District (−42%), and the Ruapehu District (−36%).

Higher and lower rates (s.d. $> \pm 1$) appeared to be slightly clustered with a concentration of higher rates in the South Island. Wellington had very high rates of *Campylobacter* notifications (s.d. $> +2$), followed by the Selwyn District, Queenstown-Lakes District and the South Wairarapa District (s.d. $> +1$ to $+2$), whereas rates in Kawereau, Far North and the Opotiki District were particularly low (s.d. < -1 to -2). The results of the autocorrelation statistic confirmed low, but statistically significant clustering for the *Campylobacter* rate 1997–2005 (Moran's $I=0.11$, $P=0.01$).

Results of the regression analysis

The results of the univariate modelling showed that for 16/23 explanatory variables the associations were in the anticipated direction, and eight coefficients were significantly related with the *Campylobacter* rate (Table 2). Multicollinear effects were observed for the variables pastoral and stock units/km², stock units/capita and occupation, stock units/capita and males, occupation and males, population aged <15 years and Europeans, population aged <15 years and

Table 1. *Campylobacteriosis in New Zealand, 1997–2005, classification and description of the independent variables included in the regression analysis*

Independent variables						
Main category	Subcategory 1	Subcategory 2	Name	Unit per TLA	Description	Data source
Underlying environmental factors	Conditions affecting the distribution of the organism	Climate/seasonal change	Rainfall	mm	Monthly mean maximum total rainfall	Landcare Research, 2002 [33]
			Temperature	°C	Monthly mean maximum daily temperatures	Landcare Research, 2002 [33]
			WIS	Index	Weighted index of interseasonal change	Calculated from: Hearnden <i>et al.</i> 2003 [7]
			Pastoral	%	Land use primarily pastoral (high- and low-producing grassland)	New Zealand Landcover Database (LCDB2), Ministry of the Environment/Terralink International, 2000–2002 [22]
	Exposure	Land use	Scrub/tussock	%	Land use: scrub (e.g. fernland, gorse, broom, subalpine shrubland); tussock grassland (e.g. low-producing and depleted grassland)	New Zealand Landcover Database (LCDB2), Ministry of the Environment/Terralink International, 2001–2002 [22]
			Stock units/km ²	Stock units/km ²	Total stock units/km ² , including dairy, beef, sheep, deer, pigs, goats, horses, poultry	MAF, 2002 (http://www.maf.govt.nz/statistics/pastoral/) [23]
		Water	<i>E. coli</i>	%	Intensity of drinking-water quality monitoring: % of <i>E. coli</i> exceedances in drinking water	NZPHO, 2002 (http://www.nzpho.org.nz/data_available.asp) [32]
			Rainwater supply	%	Population with rainwater supply	ESR, MoH, 2006 (http://www.drinkingwater.co.nz/default.asp) [49]
			Other water supply	%	Population not on registered water supply	NZPHO, 2002 (http://www.nzpho.org.nz/data_available.asp) [32]
			Measurements	<i>per capita</i>	Number of drinking water measurements	NZPHO, 2002 (http://www.nzpho.org.nz/data_available.asp) [32]
		Rurality/occupation	Stock units/ <i>per capita</i>	<i>per capita</i>	Total stock units per average population 1997/2001, including dairy, beef, sheep, deer, pigs, goats, horses, poultry	Statistics NZ, 2001 Census of Population and Dwellings [34]; MAF, 2002 (http://www.maf.govt.nz/statistics/pastoral/) [23]
			Rural population	%	Population living in highly rural/remote area	Statistics NZ, 2001 Census of Population and Dwellings [34]
			Occupation	per 1000	Residence by occupation for the employed usually president population count, aged ≥ 15 yr working in dairy- and meat-related industries	Statistics NZ, 2001 Census of Population and Dwellings [34]
		Food	Restaurants	per 100 000	Number of restaurants, bars, taverns, clubs with alcohol licence	Pearce <i>et al.</i> 2006 [35]
			Fast food	per 100 000	Number of fast food outlets	Pearce <i>et al.</i> 2006 [35]
			Fresh food	per 100 000	Number of fresh food outlets	Pearce <i>et al.</i> 2006 [35]
			Supermarkets	per 100 000	Number of supermarkets	Pearce <i>et al.</i> 2006 [35]

Table 1 (cont.)

Independent variables						
Main category	Subcategory 1	Subcategory 2	Name	Unit per TLA	Description	Data source
Surveillance/ characteristics of the affected population	Demography	Demographic characteristics of the affected population particularly at risk	Age < 15 yr	%	People aged < 15 yr	Statistics NZ, 2001 Census of Population and Dwellings [34]
			Age 25–44 yr	%	People aged 25–44 yr	Statistics NZ, 2001 Census of Population and Dwellings [34]
			Males	%	Males	Statistics NZ, 2001 Census of Population and Dwellings [34]
			Europeans	%	Europeans	Statistics NZ, 2001 Census of Population and Dwellings [34]
	Socioeconomic factors	Social deprivation	Deprivation	index	New Zealand Deprivation Score, 2001	Salmond & Crampton 2001 [36]
	Notification	Artefacts	GP density	per 100 000	Number of GPs	Pearce <i>et al.</i> 2006 [35]

deprivation, Europeans and deprivation; hence, five variables were excluded from the subsequent models.

The second modelling step showed that 11/15 associations were in the anticipated direction, and four of the variables were statically significantly related to the *Campylobacter* rate (scrub/tussock, stock units/km², rainwater supply, and population aged 25–44 years). Modelling steps 3 and 4 showed changes in the direction of associations for six variables, including, for example, rainfall and fast food outlet density. There were no major changes in the levels of statistical significance; only the association between the disease rate and the variable ‘scrub/tussock’ became statistically insignificant.

The results of the final multivariate model indicated that higher rates of campylobacteriosis occurred in TLAs with a high proportion of people aged 25–44 years (age 25–44: $P < 0.01$) and a low deprivation index (deprivation: $P < 0.01$) (Table 3). The variable representing a high density of fresh food outlets was close to reaching statistical significance (fresh food outlets: $P = 0.076$). The final model’s residuals resembled a normal distribution and indicated a low tendency of negative autocorrelation; consequently, there was no serious violation of the assumption of normality of the error term.

Evaluating the entire modelling procedure, 5/23 independent variables showed expected and stable associations throughout the regression analysis with no change of the direction of association; those were variables representing seasonality (WIS), pastoral land use (pastoral), rurality (rural population), fresh food outlet density (fresh food outlets), and the 25–44 years age group. Five explanatory variables showed stable, but inverse associations with the disease rate (*E. coli*, rainwater supply, restaurants, deprivation, GP density); the other variables had unstable associations throughout the modelling procedure.

DISCUSSION

This study examined the spatial distribution of campylobacteriosis in New Zealand. We found that the observed spatial patterns were in line with previous national studies [5, 7]. Further, our analysis gained insight into the relative importance of plausible area-level determinants of this disease. Some of the variables relating to the underlying environmental factors (e.g. stock units/km², rural population) and characteristics of the affected population (25–44 years age group) showed expected associations throughout the

Fig. 3. Percentage change in *Campylobacter* rates, New Zealand, 1997–2005.

modelling process, indicating that these variables were independently related to the disease rate [40, 41].

Interestingly, this analysis provides statistical evidence for a relationship between campylobacter rates and socio-economic deprivation. For New Zealand, there has been little evidence of a distinct association between socio-economic deprivation and campylobacteriosis [20, 29], and our results showed a consistent and statistically significant negative relationship throughout the modelling procedure. This finding is contrary to most other health outcomes as worse

health (e.g. higher morbidity/mortality rates) is commonly found in areas of lower socio-economic status [42]. Potential differences in food consumption patterns between areas stratified by socio-economic deprivation may explain this relationship. In terms of campylobacteriosis, Evans *et al.* [43] and Eberhart-Phillips *et al.* [40] identified the consumption of fresh vegetables, drinking bottled water and eating out as independent risk factors of becoming infected. These patterns of consumption tend to be associated with higher socio-economic status [44–46]. Consequently,

Table 2. Associations from the univariate and multivariate regression models predicting *Campylobacter* rates in New Zealand, 1997–2005

Models ...	Univariate models			Multivariate models											Final multivariate model				
Modelling step ...	1			2				3				4				5			
Variable	Association		<i>P</i> value*	Subcategory 2	Association		<i>P</i> value*	Subcategory 1	Association		<i>P</i> value*	Main category	Association		<i>P</i> value*	Association	<i>P</i> value*		
Rainfall	+	Expected	0.435	Climate/ seasonal change	–	Inverse	0.913	Conditions affecting the distribution of the organism	–	Inverse	0.960	Underlying environmental factors	+	Expected	0.969	–	Inverse	0.664	
Temperature	–	Expected	0.036		–	Expected	0.120		–	Expected	0.250		–	Expected	0.338	+	Inverse	0.868	
WIS	+	Expected	0.030		+	Expected	0.113		+	Expected	0.439		+	Expected	0.402	+	Expected	0.144	
Pastoral	+	Expected	0.747	Land use															
Scrub/tussock	+	Expected	0.032		+	Expected	0.003		+	Expected	0.047		+	Expected	0.105	–	Inverse	0.651	
Stock units/km ²	+	Expected	0.289		+	Expected	0.014		+	Expected	0.028		+	Expected	0.022	+	Expected	0.296	
<i>E. coli</i>	–	Inverse	0.939	Water	–	Inverse	0.912	Exposure	–	Inverse	0.249	Surveillance/ characteristics of the affected population	–	Inverse	<i>0.065</i>	–	Inverse	0.883	
Rainwater supply	–	Inverse	0.003		–	Inverse	0.003		–	Inverse	0.003		–	Inverse	0.023	–	Inverse	0.143	
Other water supply	–	Inverse	0.467		+	Expected	0.256		+	Expected	0.418		–	Inverse	0.912	–	Inverse	0.616	
Measurements	+	Expected	0.330		+	Expected	0.457		–	Inverse	0.824		+	Expected	0.958	+	Expected	0.986	
Stock units/capita	+	Expected	0.239	Rurality/ occupation															
Rural population		Expected	0.195						+	Expected	0.129		+	Expected	<i>0.066</i>	+	Expected	0.193	
Occupation	+	Expected	0.291																
Restaurants	–	Inverse	0.307	Food	–	Inverse	<i>0.053</i>		–	Inverse	0.140		–	Inverse	0.271	–	Inverse	0.455	
Fast food	+	Expected	0.686		+	Expected	0.557		+	Expected	0.811		–	Inverse	0.570	–	Inverse	0.247	
Fresh food	+	Expected	0.170		+	Expected	0.206		+	Expected	0.139		+	Expected	0.091	+	Expected	<i>0.076</i>	
Supermarkets	+	Expected	0.260		+	Expected	0.336		+	Expected	0.888		–	Inverse	0.407	+	Expected	0.600	
Age < 15 yr	–	Inverse	0.000	Demographic characteristics of the affected population particularly at risk															
Age 25–44 yr	+	Expected	0.000		+	Expected	0.000							+	Expected	0.008	+	Expected	0.005
Males	+	Expected	0.504		+	Expected	0.641							–	Inverse	0.679	–	Inverse	0.184
Europeans	+	Expected	0.001																
Deprivation	–	Inverse	0.000	Social deprivation										–	Inverse	0.000	–	Inverse	0.007
GP density	–	Inverse	0.123	Artefacts										–	Inverse	0.526	–	Inverse	0.212

* **Significant P values**; P values are very close to being significant at the <0.05 level.

Shading in cells:

Dark grey, the relationships are stable and show the expected associations throughout the modelling process.

Light grey, the relationships are stable and show inverse associations throughout the modelling process.

White, the relationships are unstable throughout the modelling process.

Table 3. Regression coefficients from the final multivariate regression model predicting *Campylobacter* rates in New Zealand, 1997–2005

Final model	Subcategory 2	Variable	<i>b</i> coefficient	<i>P</i> value
$R^2=0.583$	Climate/seasonal change	Rainfall	−0.205	0.664
		Temperature	12.991	0.868
Adjusted		WIS	2.721	0.144
$R^2=0.441$	Land use	Scrub/tussock	−4.439	0.651
		Stock units/km ²	1.680	0.296
Intercept = 12 978, 552	Water	<i>E. coli</i>	−6.921	0.883
Standard error of the estimate = 638.669		Rainwater supply	−65.361	0.143
		Other water supply	−4.283	0.616
		Measurements	0.794	0.986
	Rurality	Rural population	12.206	0.193
	Food	Restaurants	−0.436	0.455
		Fast food	−4.211	0.247
		Fresh food	4.107	0.076***
		Supermarkets	4.174	0.600
	Demographic characteristics	Age 25–44 yr	129.410	0.005*
		Males	−143.890	0.184
	Social deprivation	Deprivation	−7.105	0.007*
	Artefacts	GP density	−10.693	0.212

* Significant at the 0.01 level; *** close to significance at the <0.05 level.

there could be a higher risk of becoming infected from the consumption of fresh fruits and vegetables in areas of higher socio-economic status. Linking the inverse association identified here between socio-economic deprivation and the *Campylobacter* rate to the positive association with the variable fresh food outlet density, this study suggests statistical evidence for an inter-connection between socio-economic status, the distribution of fresh food outlets and *Campylobacter* notifications.

Several of the other explanatory variables, however, did not have a consistent or expected relationship with the disease rate. For example, national and international findings have demonstrated a significant underestimation of *Campylobacter* cases related to contaminated water supplies [11, 21, 47], but our study showed little evidence that water is a significant independent contributor to notification rates in New Zealand. However, there is some evidence that higher rates of campylobacteriosis occur during the summer months when outdoor activities such as swimming may increase the probability of infections [41]. Further, the inverse association between *E. coli* presence and *Campylobacter* notifications observed here may be due to regional differences in detection procedures [17]. Frequently, the concentration of *E. coli* is too low to detect faecal contamination of water. Even if the indicator organism is not detected, the water

might be contaminated nevertheless. Conversely, a high level of *E. coli* exceedances in water is not a conclusive indicator for contamination with *Campylobacter* [18].

This study has three key limitations. First, applying an ecological study design, the evaluation of the results is confronted inherently with issues related to the ecological fallacy, and it is not possible to infer causality from the observed associations. Second, there are several reasons for inconsistencies between the reported and actual prevalence of campylobacteriosis between regions. People might not seek medical attention for various reasons (e.g. unequal access to healthcare facilities, different view of sickness); hence, their illness is not notified which in turn results in an undercount of cases. Further, the disease might not specifically be diagnosed as campylobacteriosis, but more generally as gastrointestinal infection [21].

Finally, the fully adjusted regression model did not account for all of the assumed impacts of different land-use categories, occupational effects, ethnicity and age. Future research could concentrate on identifying those unspecified determinants of campylobacteriosis and on minimizing the multicollinear effects detected here. New data sources and the improvement of the national notification system, including the EpiSurv messaging system provide enhanced surveillance sources. For example,

SurvINZ, including the databases EpiSurv and WINZ (Water Information New Zealand), is a new platform where surveillance activities have been integrated to provide more comprehensive and timely information to analyse morbidity trends and to investigate mechanisms affecting the distribution of campylobacteriosis in New Zealand [6]. In particular, it would be interesting to investigate the possibly underestimated role of water in the circle of infection with *Campylobacter* since the different dimensions of this relationship have not been quantified on a broader scale (e.g. impact of recreational water contact, consumption of tap or bottled water, cross-contamination of drinking water). It might also be useful to study areas where disease rates are lower than expected in order to identify and understand protective factors potentially limiting the increase in rates in some areas. Moreover, it is important to investigate determinants of campylobacteriosis separately for urban and rural areas (e.g. differences in consumption patterns, the exposure to risk factors such as infected cattle or contaminated water supplies), and to study the aetiology of outbreaks and sporadic cases, respectively.

For reducing high rates of campylobacteriosis in New Zealand, banning of fresh poultry from sale and improving consumer education are the most discussed public health implications. In Belgium, for example, the *Campylobacter* rate dropped significantly when fresh poultry potentially contaminated with the chemical pollutant dioxin was banned from sale for 4 weeks during 1999. Interventions to change consumer behaviour include, for example, information about the thorough cleaning of kitchen utensils, cutting boards, taps, and the rinsing of fresh fruits and vegetables [48].

In conclusion, area-level characteristics exerted an influence on *Campylobacter* rates in New Zealand over the period 1997–2005, and the observed plausible associations between fresh food outlet density, higher socio-economic status and an increased disease rate cast light on currently relatively unexplored dimensions of campylobacteriosis. Infections with *Campylobacter* present a complex, persistent, increasing and expensive public health issue. Regarding the high rates of campylobacteriosis, current public health interventions appear to be insufficient, and further research is necessary to effectively target particular causes of the infection. Socio-economic conditions might be a key factor in explaining spatial differences in *Campylobacter* notifications across New Zealand.

ACKNOWLEDGEMENTS

This research was funded by the GeoHealth Laboratory, a joint initiative between the University of Canterbury and the Ministry of Health, New Zealand.

DECLARATION OF INTEREST

None.

REFERENCES

1. Baker M, Sneyd E, Wilson NA. Is the major increase in notified campylobacteriosis in New Zealand real? *Epidemiology and Infection* 2006; **135**: 163–170.
2. Environmental Science and Research. Campylobacter (<http://www.nzfsa.govt.nz/science/data-sheets/campylobacter.pdf>). Accessed 2 August 2006.
3. WHO. Campylobacter (<http://www.who.int/mediacentre/factsheets/fs255/en/>). Accessed 8 August 2006.
4. Scott WG, *et al.* Economic cost to New Zealand of foodborne infectious disease. *New Zealand Medical Journal* 2000; **113**: 281–284.
5. Environmental Science and Research. Notifiable and other diseases in New Zealand. Annual Report 2005. Wellington: Ministry of Health, 2006. Client Report FW 0621.
6. Ministry of Health. Direct Laboratory Notification of Communicable Diseases. National Guidelines (<http://www.surv.esr.cri.nz/LabSurv/Documents/dln-national-guidelines-dec07.pdf>). Accessed 3 August 2009.
7. Hearnden M, *et al.* The regionality of campylobacteriosis seasonality in New Zealand. *International Journal of Environmental Health Research* 2003; **13**: 337–348.
8. Kovats RS, *et al.* Climate variability and *Campylobacter* infection: an international study. *International Journal of Biometeorology* 2005; **49**: 207–214.
9. Louis VR, *et al.* Temperature-driven *Campylobacter* seasonality in England and Wales. *Applied and Environmental Microbiology* 2005; **71**: 85–92.
10. Nylen G, *et al.* The seasonal distribution of *Campylobacter* infection in nine European countries and New Zealand. *Epidemiology and Infection* 2002; **128**: 383–390.
11. Nygård K, *et al.* Association between environmental risk factors and *Campylobacter* infections in Sweden. *Epidemiology and Infection* 2004; **132**: 317–325.
12. Sandberg MM, *et al.* Incidence trend and risk factors for *Campylobacter* infections in humans in Norway. *BMC Public Health* 2006; **6**: 179–205.
13. Curriero FC, *et al.* The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948–1994. *American Journal of Public Health* 2001; **91**: 1194–1199.
14. Hänninen ML, *et al.* Detection and typing of *Campylobacter jejuni* and *Campylobacter coli* and analysis

- of indicator organisms in three waterborne outbreaks in Finland. *Applied and Environmental Microbiology* 2003; **69**: 1391–1396.
15. **Altekruse SF, et al.** *Campylobacter jejuni* – an emerging foodborne pathogen. *Emerging Infectious Diseases* 1999; **5**: 28–35.
 16. **Newell DG, et al.** The ecology of *Campylobacter jejuni* in avian and human hosts and in the environment. *International Journal of Infectious Diseases* 2002; **6**: 3S16–3S21.
 17. **Khan R, Voice T, Phillips D.** Developing National Environmental Health Indicators for New Zealand. Christchurch, Wellington: Environmental Science and Research, Ministry of Health, 2005. Client Report FW 0570.
 18. **Savill MG, et al.** Enumeration of *Campylobacter* in New Zealand recreational and drinking waters. *Journal of Applied Microbiology* 2001; **91**: 38–46.
 19. **Gallay A, et al.** A large multi-pathogen waterborne community outbreak linked to faecal contamination of a groundwater system, France, 2000. *Clinical Microbiology and Infection* 2006; **12**: 561–570.
 20. **National Health Committee.** The Health of People and Communities. The effect of environmental factors on the health of New Zealanders. Report to the Minister of Health from the Public Health Advisory Committee. Wellington: Public Health Advisory Committee, 2002. HP: 3574.
 21. **Abbott SE, Douwes J, Caughley BP.** A survey of the microbiological quality of roof-collected rainwater of private dwellings in New Zealand (http://www.water2006.com/manuscripts_06/index.htm). Accessed 28 April 2007.
 22. **Ministry for the Environment.** The New Zealand land cover database (<http://www.mfe.govt.nz/issues/land/land-cover-dbase/>). Accessed 21 January 2007.
 23. **Ministry of Agriculture and Forestry.** Livestock by Territorial Authority (<http://www.maf.govt.nz/statistics/pastoral/>). Accessed 5 August 2006.
 24. **Devane ML, et al.** The occurrence of *Campylobacter* subtypes in environmental reservoirs and potential transmission routes. *Journal of Applied Microbiology* 2005; **98**: 980–990.
 25. **Savill M, et al.** Elucidation of potential transmission routes of *Campylobacter* in New Zealand. *Water Science and Technology* 2003; **47**: 33–38.
 26. **Kapperud G, et al.** Factors associated with increased and decreased risk of *Campylobacter* infection: a prospective case-control study in Norway. *American Journal of Epidemiology* 2003; **158**: 234–242.
 27. **TeckLok W, On SLW, Michie H.** *Campylobacter* in New Zealand: reservoirs, sources and the labyrinth of transmission routes. *New Zealand Journal of Environmental Health* 2006; **29**: 1–6.
 28. **Baker M, et al.** Potential transmission routes of *Campylobacter* from environment to humans. Christchurch, Wellington: Environmental Science and Research, Ministry of Health, 2002. Client Report FW0246.
 29. **Hales S, et al.** Social deprivation and the public health risks of community drinking water supplies in New Zealand. *Journal of Epidemiology and Community Health* 2003; **57**: 581–583.
 30. **Scott KM, Marwick JC, Crampton PR.** Utilization of general practitioner services in New Zealand and its relationship with income, ethnicity and government subsidy. *Health Services Management Research* 2003; **16**: 45–55.
 31. **Hoque ME, et al.** Under-notification of giardiasis in Auckland, New Zealand: a capture-recapture estimation. *Epidemiology and Infection* 2005; **133**: 71–79.
 32. **New Zealand Public Health Observatory.** Notifiable disease and environmental health indicator dataset (http://www.nzpho.org.nz/data_available.asp). Accessed 7 July 2006.
 33. **Leathwick JR, Wilson G, Stephens RTT.** Climate surfaces for New Zealand. Landcare Research Contract Report: LC9798/126 (http://www.landcareresearch.co.nz/databases/lenz/downloads/Climate_Surfaces_for_New_Zealand_revised.pdf). Accessed 19 February 2007.
 34. **Statistics New Zealand.** Census of Population and Dwellings 2001 (<http://www.stats.govt.nz/census/default.htm#2001Census>). Accessed 12 January 2007.
 35. **Pearce J, Witten K, Bartie P.** Neighbourhoods and health: a GIS approach to measuring community resource accessibility. *Journal of Epidemiology and Community Health* 2006; **60**: 389–395.
 36. **Salmond C, Crampton P.** NZDep2001 Index of Deprivation. User's Manual ([http://www.moh.govt.nz/moh.nsf/Files/phi-users-manual/\\$file/phi-users-manual.pdf](http://www.moh.govt.nz/moh.nsf/Files/phi-users-manual/$file/phi-users-manual.pdf)). Accessed 14 July 2006.
 37. **Blakely T, et al.** No association of neighbourhood volunteerism with mortality in New Zealand: a national multilevel cohort study. *International Journal of Epidemiology* 2006; **35**: 981–989.
 38. **Duncan C, Jones K, Moon G.** Context, composition and heterogeneity: using multilevel models in health research. *Social Science and Medicine* 1998; **46**: 97–117.
 39. **Tabachnik BG, Fidell LS.** *Using Multivariate Statistics*, 4th edn. Boston, MA: Allyn & Bacon, 2001.
 40. **Eberhart-Phillips J, et al.** *Campylobacteriosis* in New Zealand: results of a case-control study. *Journal of Epidemiology and Community Health* 1997; **51**: 686–691.
 41. **Eyles R, et al.** Spatial and temporal patterns of *Campylobacter* contamination underlying public health risk in the Taieri River, New Zealand. *Journal of Environmental Quality* 2003; **32**: 1820–1828.
 42. **Pearce J, Dorling D.** Increasing geographical inequalities in health in New Zealand, 1980–2001. *International Journal of Epidemiology* 2006; **35**: 597–603.
 43. **Evans MR, Ribeiro CD, Salmon RL.** Hazards of healthy living: bottled water and salad vegetables as risk factors for *Campylobacter* infection. *Emerging Infectious Diseases* 2003; **9**: 1219–1225.
 44. **Giskes K, et al.** Socio-economic differences in fruit and vegetable consumption among Australian adolescents

- and adults. *Public Health Nutrition* 2002; **5**: 663–669.
45. **Lindstrom M, et al.** Socioeconomic differences in the consumption of vegetables, fruit and fruit juices. The influence of psychosocial factors. *European Journal of Public Health* 2001; **11**: 51–59.
 46. **Shohaimi S, et al.** Residential area deprivation predicts fruit and vegetable consumption independently of individual educational level and occupational social class: a cross sectional population study in the Norfolk cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk). *Journal of Epidemiology and Community Health* 2004; **58**: 686–691.
 47. **Wilson N.** A systematic review of the aetiology of human campylobacteriosis in New Zealand (<http://www.nzfsa.govt.nz/science/research-projects/campy-aetiol/campy-aetiol.pdf>). Accessed 8 August 2006.
 48. **Baker MG, et al.** Regulation of chicken contamination urgently needed to control New Zealand's serious campylobacteriosis epidemic. *New Zealand Medical Journal* 2006; **119**: 1–8.
 49. **Environmental Science and Research, Ministry of Health.** Drinking water for New Zealand. Water information New Zealand (WINZ). (<http://www.drinkingwater.co.nz/default.asp>). Accessed 17 October 2006.