THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Primary palliative care needs urgent attention

Citation for published version:

Mitchell, S, Tan, A, Moine, S, Dale, J & Murray, SA 2019, 'Primary palliative care needs urgent attention', *BMJ (Clinical research ed.)*, vol. 365, pp. l1827. <https://doi.org/10.1136/bmj.l1827>

Digital Object Identifier (DOI):

[10.1136/bmj.l1827](https://doi.org/10.1136/bmj.l1827)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

BMJ (Clinical research ed.)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

EDITORIALS

Primary palliative care needs urgent attention

Primary care teams are vital to the initiation and delivery of effective palliative care

Sarah Mitchell *general practitioner and NIHR doctoral research fellow*¹, Amy Tan *family physician and associate professor of family medicine*², Sebastien Moine *general practitioner and visiting fellow*^{3, 4}, Jeremy Dale *professor of primary care*¹, Scott A Murray *professor of primary palliative care*³

¹Unit of Academic Primary Care, Warwick Medical School, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK; ²Dept. of Family Medicine, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada; ³Primary Palliative Care Research Group, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Teviot Place, Edinburgh, UK; ⁴Health Education and Practices Laboratory, University of Paris 13, Bobigny, France

Growing numbers of people, both old and young, have complex chronic conditions and multimorbidity. Their disease trajectories are highly unpredictable; sudden health deteriorations, and the possibility of dying, are a constant risk for many. This presents huge challenges for primary and community healthcare services. Patients and families often turn to emergency healthcare services at times of acute crisis, which can lead to emergency admission to hospital. In the United Kingdom, up to one third of hospital inpatients are in their last year of life.¹

Palliative care improves the quality of life of people with serious illnesses, or life threatening conditions, or both, by assessment and management of their physical, psychological, social, and spiritual needs. Substantial inequalities exist in provision, however, and patients with multimorbidity, non-malignant disease, or dementia are less likely to receive such care.² Many patients who need palliative care are not included in UK general practice palliative care registers, but those with cancer are more likely to be included than those with non-malignant conditions.³

Similar concerns exist in other developed nations. In Canada, only 15-30% of people have access to publicly funded specialist palliative care services, and they often access them only in the last days or weeks of life, mainly in acute care settings. Those who do receive palliative care services in the community are 2.5 times more likely to die at home.⁴

In 2018, 40 years after the landmark Alma Ata declaration formally recognised primary healthcare as integral to achieving health for all,⁵ the World Health Organization affirmed that providing palliative care to patients, families, and communities is core to the role and identity of primary care clinicians.⁶ General practitioners or family physicians have the skills to deliver palliative and end-of-life care in every encounter in the community—when their working environment allows it and they are adequately supported by specialist colleagues.^{7, 8}

Trusted, longstanding relationships between primary care professionals and their patients create an ideal context for early identification of palliative care needs, including for those who are frail, those with multimorbidity, and those at risk of unwanted admissions to hospital. These relationships are valued by patients and families.⁹ Primary care, however, is under unprecedented pressure in the UK, Canada, and elsewhere. Barriers include a lack of time with patients, conflicting clinical, administrative, and managerial demands, compromised continuity of care (including out of hours), and depleted community nursing services.¹⁰

The BMJ drew attention to the pressing need to develop primary palliative care to realise the expectations of dying people 15 years ago.⁸ Some progress has been achieved since then with research, national policies, educational resources, frameworks for care,^{11, 12} and the European Association of Palliative Care toolkit.¹³ But the integration of palliative care into primary care continues to need attention, alongside the development of specialist services, including specialist outreach, which, although appropriate for many patients, is neither needed nor wanted by others.

Palliative care is needed by populations all over the world,¹⁴ and primary care has the potential to deliver this person centred care. As we approach the first high level meeting on universal health coverage at the United Nations General Assembly in September 2019, the message for policy makers and primary healthcare professionals is clear. Developing primary palliative care alongside specialist palliative care services is vital to fully realise the benefits of early identification of need. These benefits include improved coordination of care and symptom control, support for families, and less time in hospital.¹⁵

Frameworks to improve care^{10, 11} and models of palliative care integration between primary, secondary, and tertiary services

already exist.¹⁶ Educational resources to support the management of patients with life threatening, complex multimorbid conditions in primary care should also be improved, in keeping with the changing needs of the population.

Future policy, and those charged with implementing it, should place greater emphasis on the role of general practitioners, family physicians, and community nurses in palliative care. They must recognise that primary care teams have a key role in the initiation and delivery of effective palliative care for patients and families and must ensure that training, support and a working environment and culture conducive to primary palliative care is provided for all.

Competing interests: We have read and understood BMJ policy on declaration of interests and declare no competing interests.

Provenance and peer review: Not commissioned, peer reviewed.

- 1 Clark D, Armstrong M, Allan A, Graham F, Camon A, Isles C. Imminence of death among hospital inpatients: Prevalent cohort study. *Palliat Med* 2014;28:474-9. 10.1177/0269216314526443 24637342
- 2 Care Quality Commission. A different ending. Addressing inequalities in end of life care. Overview report. Newcastle upon Tyne, CQC, 2016. https://www.cqc.org.uk/sites/default/files/20160505%20CQC_EOLC_OVERVIEW_FINAL_3.pdf
- 3 Harrison N, Cavers D, Campbell C, Murray SA. Are UK primary care teams formally identifying patients for palliative care before they die? *Br J Gen Pract* 2012;62:e344-52. 10.3399/bjgp12X641465 22546594
- 4 The Way Forward, Aller De L'Avant. The palliative approach: improving care for Canadians with life-limiting illnesses. 2018. <http://hpcintegration.ca/media/38753/TWF-palliative-approach-report-English-final2.pdf>
- 5 World Health Organization. Declaration of Alma-Ata International Conference on Primary Health Care, Alma-Ata, USSR, 6-12 September 1978. 1978. https://www.who.int/social_determinants/tools/multimedia/alma_ata/en/

- 6 World Health Organization, United Nations International Children's Fund. Declaration of Astana. Global Conference on Primary Health Care. Astana, Kazakhstan, October 25-26, 2018. 2018. <https://www.who.int/docs/default-source/primary-health/declaration/gcphc-declaration.pdf>
- 7 Brazil K, Sussman J, Bainbridge D, Whelan T. Who is responsible? The role of family physicians in the provision of supportive cancer care. *J Oncol Pract* 2010;6:19-24. 10.1200/JOP.091060 20539726
- 8 Murray SA, Boyd K, Sheikh A, Thomas K, Higginson IJ. Developing primary palliative care. *BMJ* 2004;329:1056-7. 10.1136/bmj.329.7474.1056 15528593
- 9 Bulsara C, Ward AM, Joske D. Patient perceptions of the GP role in cancer management. *Aust Fam Physician* 2005;34:299-300, 302.15861760
- 10 Mitchell S, Loew J, Millington-Sanders C, Dale J. Providing end-of-life care in general practice: findings of a national GP questionnaire survey. *Br J Gen Pract* 2016;66:e647-53. 10.3399/bjgp16X686113 27381483
- 11 Millington-Sanders C, Noble B. New UK general practice core standards for advanced serious illness and end of life care. *Br J Gen Pract* 2018;68:114-5. 10.3399/bjgp18X694913 29472205
- 12 Shaw KL, Clifford C, Thomas K, Meehan H. Review: improving end-of-life care: a critical review of the gold standards framework in primary care. *Palliat Med* 2010;24:317-29. 10.1177/0269216310362005 20156934
- 13 Murray SA, Firth A, Schneider N, et al. Promoting palliative care in the community: production of the primary palliative care toolkit by the European Association of Palliative Care Taskforce in primary palliative care. *Palliat Med* 2015;29:101-11. [https://www.eapconet.eu/Portals/0/adam/Tables/d8SJmTWv1Ey7keWyC6b5yQ/Col2/PM201529\(2\)Murray.pdf](https://www.eapconet.eu/Portals/0/adam/Tables/d8SJmTWv1Ey7keWyC6b5yQ/Col2/PM201529(2)Murray.pdf). 10.1177/0269216314545006 25395577
- 14 Murray SA, Kendall M, Mitchell G, Moine S, Ambias-Novellas J, Boyd K. Palliative care from diagnosis to death. *BMJ* 2017;356:j878. 10.1136/bmj.j878 28242747
- 15 Knaul FM, Farmer PE, Krakauer EL, et al. Lancet Commission on Palliative Care and Pain Relief Study Group. Alleviating the access abyss in palliative care and pain relief—an imperative of universal health coverage: the Lancet Commission report. *Lancet* 2018;391:1391-454. 10.1016/S0140-6736(17)32513-8 29032993
- 16 Kreitzer MJ, Monsen KA, Nandram S, de Blok J. Buurtzorg Nederland: a global model of social innovation, change, and whole-systems healing. *Glob Adv Health Med* 2015;4:40-4. 10.7453/gahmj.2014.030 25694851

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>