THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Antenatal glucocorticoid treatment

Citation for published version:

Reynolds, RM & Seckl, JR 2012, 'Antenatal glucocorticoid treatment: are we doing harm to term babies?', *The Journal of Clinical Endocrinology & Metabolism (JCEM)*, vol. 97, no. 10, pp. 3457-3459.
<https://doi.org/10.1210/jc.2012-3201>

Digital Object Identifier (DOI):

[10.1210/jc.2012-3201](https://doi.org/10.1210/jc.2012-3201)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

The Journal of Clinical Endocrinology & Metabolism (JCEM)

Publisher Rights Statement:

Copyright © 2012 by The Endocrine Society

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Antenatal Glucocorticoid Treatment: Are We Doing Harm to Term Babies?

Rebecca M. Reynolds and Jonathan R. Seckl

Endocrinology Unit, University/British Heart Foundation Centre for Cardiovascular Science, University of Edinburgh, Queen's Medical Research Institute, Edinburgh EH16 4TJ, United Kingdom

Antenatal glucocorticoids are given to 7–10% of women in Europe and North America at threat of preterm labor. It is undisputed that this therapy significantly reduces neonatal morbidity and mortality in infants born before 34 wk gestation (1, 2). Based on outcomes from multiple clinical trials, current obstetric practice is to administer either a single 2-d course of betamethasone or dexamethasone (*i.e.* two 12-mg doses of betamethasone im at 24-h intervals or four 6-mg doses of dexamethasone im at 12-h intervals) to women in preterm labor between 24 and 34 wk gestation (3). The use of repeat courses of antenatal glucocorticoids is more controversial, although some evidence suggests that a repeat course, if the mother does not deliver but remains at risk of preterm labor, may yield neonatal benefit (2). However, because early clinical diagnosis of preterm labor is difficult and there are no reliable laboratory tests, there are a substantial number of women who receive antenatal glucocorticoid treatment yet subsequently deliver babies at full term, *i.e.* after 37 wk gestation. Little is known about the potential risks or benefits of synthetic glucocorticoid exposure to these infants.

It is now well recognized that exposure to an adverse *in utero* environment during development has long-term effects on physiology and later risk of adult disease (4). Overexposure of the developing fetus to excess glucocorticoids is a key mechanism thought to underlie such “developmental programming” (4). Numerous preclinical studies have shown that administration of dexamethasone to pregnant mothers leads to long-term changes in the offspring, notably permanently altering cardiometabolic and neuropsychiatric functions as well as hypothalamic-pituitary-adrenal (HPA) axis activity (4). The latter may

contribute to some of the broader pathogenesis associated with experimental programming.

Both betamethasone and dexamethasone have high affinity for glucocorticoid receptors but are poor substrates for the placental enzyme 11- β hydroxysteroid dehydrogenase type 2, which normally acts as a partial barrier, protecting the fetus from glucocorticoids by breaking down endogenous cortisol into inactive cortisone (4). These synthetic glucocorticoids readily cross the placenta, potentially exposing the fetus to excess glucocorticoids. In animal models, the developing HPA axis appears particularly vulnerable to excess glucocorticoids (4), yet the evidence for detrimental effects of antenatal glucocorticoids on human HPA axis activity has been less clear-cut. This is partly because most studies have included preterm infants, who often have coexisting medical complications and are exposed to stressors and painful procedures while in the neonatal intensive care unit, factors that can *per se* alter HPA axis function (5). Likewise, the type, dose, and timing of administration of glucocorticoid therapy often differ between studies, complicating interpretation. Furthermore, studies have included limited assessments of the offspring HPA axis, often with a single measurement of cortisol levels. Given these caveats, the limited existing data, which include preterm babies in whom the HPA axis may be immature, suggest that among infants exposed to antenatal glucocorticoids, baseline cortisol levels are suppressed during the first postnatal week, and cortisol responses to the painful stress of the heel-stick procedure are suppressed for up to 8 wk (6), suggesting some persistent suppressive effects on HPA axis regulation, but hardly the programming of increased (or reduced) HPA activity seen throughout the lifespan in animal models.

ISSN Print 0021-972X ISSN Online 1945-7197

Printed in U.S.A.

Copyright © 2012 by The Endocrine Society

doi: 10.1210/jc.2012-3201 Received August 28, 2012. Accepted August 29, 2012.

Abbreviation: HPA, Hypothalamic-pituitary-adrenal.

For article see page 3538

In the article by Alexander *et al.* (7) in this issue of the *JCEM*, the authors report on the first study to examine the impact of antenatal synthetic glucocorticoid exposure on endocrine stress reactivity in term-born children aged 6–10 yr. Maternal stress has itself been implicated as a programming factor (4), and hospitalized women receiving antenatal glucocorticoids are likely to be more worried and “stressed.” Alexander *et al.* (7) therefore introduced an additional control group of stressed mothers who had been admitted to hospital with pregnancy complications but had not received antenatal glucocorticoids. This careful study design goes some way to disentangle the potential confounding effect of maternal stress from the direct effects of antenatal glucocorticoid exposure.

The children exposed to antenatal glucocorticoids were smaller at birth and were born slightly earlier than the control groups; these variables were included as covariates in the statistical analysis. The main finding was that children exposed to antenatal glucocorticoid treatment had significantly increased cortisol reactivity to a standardized laboratory stressor, the Trier Social Stress Test for Children (TSST-C), compared with both control groups. Although these data are the most comprehensive to date, they are still cross-sectional and observational. Prospective data with longitudinal follow-up are necessary to confirm the findings and overcome any possible bias in recruitment/retention. Nevertheless, these findings are consistent with a recent study showing a more pronounced cortisol response to the heel-stick procedure among term infants exposed to betamethasone *in utero*, compared with an unexposed control group (8), and suggest long-lasting effects of antenatal glucocorticoid exposure on HPA axis reactivity. Alexander *et al.* (7) also found that cortisol reactivity to the TSST-C was more pronounced in girls than boys, a finding that is consistent with the sexually dimorphic responses in animal studies where females are generally more susceptible to programming of HPA axis stress responsivity (9). Increased HPA axis activity during the morning diurnal peak has also been reported in children of mothers who voluntarily consumed higher quantities of the 11- β hydroxysteroid dehydrogenase type 2 inhibitor licorice in pregnancy (10).

What are the implications of increased HPA axis stress reactivity in these children? From an evolutionary perspective, such an adaptation to an intrauterine insult is thought to be beneficial in the short-term and linked to increased chance of offspring survival in a predicted stressful environment, and hence increased chances of survival to reproductive age (11). However, such adaptations may be detrimental in the longer term. For example, in adult men and women, activation of the HPA axis has been linked to increased risk of cardiovascular risk factors com-

prising the metabolic syndrome including higher glucose, blood pressure, and triglycerides (12), as well as ischemic heart disease, cognitive decline, and depression (12) in later life. Indeed, in preterm infants, antenatal glucocorticoid exposure has been associated with an increase in behavioral disorders at age 3 yr (13), higher systolic and diastolic blood pressures in adolescence (14), and increased insulin resistance, a precursor of type 2 diabetes, at age 30 yr (15). Whether or not the HPA axis hyperactivity observed by Alexander *et al.* (7) contributes to such effects remains to be determined.

There is no doubt that antenatal glucocorticoid treatment for women in preterm labor has benefits for preterm babies, and this justifies the small risk of long-term side effects. Yet with this emerging evidence of potential harm for term infants and no evidence of benefits, it is crucial that we develop ways to accurately identify those women at the greatest risk of preterm labor. Approaches might employ biological markers such as fetal fibronectin as a test of exclusion of preterm labor (16), or the research tool of measurement of cervical length by transvaginal ultrasonography (17). In addition, in 2011, the U.S. Food and Drug Administration approved the use of progesterone supplementation (hydroxyprogesterone caproate) during pregnancy to reduce the risk of recurrent preterm birth in women with a history of at least one prior spontaneous preterm delivery (18). Such strategies may avoid unnecessary glucocorticoid treatment of those women who go on to deliver beyond 37 wk gestation. However, recent Royal College of Obstetrics and Gynaecology guidelines in the United Kingdom recommend routine administration of antenatal glucocorticoids for all women undergoing elective cesarean section at less than 39 wk gestation (19). Although endogenous glucocorticoids rise dramatically in pregnancy, peaking in the third trimester at three times nonpregnant levels (20), the long-term effects of exogenous glucocorticoid exposure in late pregnancy for these children are unknown. A further approach is to consider minimizing the doses of glucocorticoids used. Although Alexander *et al.* (7) observed no differences in childhood cortisol stress responses according to the type of glucocorticoid used, data in a nonhuman primate model suggest dose-associated programming of cardiometabolic parameters by dexamethasone (21), implying that the lowest dose of glucocorticoid possible should be used to minimize long-term adverse effects. This principle is well-established in adult glucocorticoid therapy and might beneficially be incorporated into obstetric thinking as well.

Acknowledgments

Address all correspondence and requests for reprints to: Dr. Rebecca Reynolds, Endocrinology Unit, University/British Heart Foundation Centre for Cardiovascular Science, University of Edinburgh, Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh EH16 4TJ, United Kingdom. E-mail: R.Reynolds@ed.ac.uk.

We acknowledge the support of the British Heart Foundation.

Disclosure Summary: Both authors have no conflicts of interest to declare.

References

1. Roberts D, Dalziel S 2006 Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev* 3:CD004454
2. McKinlay CJ, Crowther CA, Middleton P, Harding JE 2012 Repeat antenatal glucocorticoids for women at risk of preterm birth: a Cochrane Systematic Review. *Am J Obstet Gynecol* 206:187–194
3. 1995 Effect of corticosteroids for fetal maturation on perinatal outcomes. NIH Consensus Development Panel on the Effect of Corticosteroids for Fetal Maturation on Perinatal Outcomes. *JAMA* 273:413–418
4. Seckl JR 2004 Prenatal glucocorticoids and long-term programming. *Eur J Endocrinol* 151(Suppl 3):U49–U62
5. Grunau RE, Holsti L, Haley DW, Oberlander T, Weinberg J, Solimano A, Whitfield MF, Fitzgerald C, Yu W 2005 Neonatal procedural pain exposure predicts lower cortisol and behavioural reactivity in preterm infants in the NICU. *Pain* 113:293–300
6. Waffarn F, Davis EP 13 June 2012 Effects of antenatal corticosteroids on the hypothalamic-pituitary-adrenocortical axis of the fetus and newborn: experimental findings and clinical considerations. *Am J Obstet Gynecol* doi: 10.1016/j.ajog.2012.06.012
7. Alexander N, Rosenlocher F, Stalder T, Linke J, Distler W, Morgner J, Kirschbaum C 2012 Impact of antenatal synthetic glucocorticoid exposure on endocrine stress reactivity in term born children. *J Clin Endocrinol Metab* 97:3538–3544
8. Davis EP, Waffarn F, Sandman CA 2011 Prenatal treatment with glucocorticoids sensitizes the HPA axis response to stress among full-term infants. *Dev Psychobiol* 53:175–183
9. Weinstock M, Matlina E, Maor GI, Rosen H, McEwen BS 1992 Prenatal stress selectively alters the reactivity of the hypothalamic-pituitary-adrenal system in the female rat. *Brain Res* 595:195–200
10. Räikkönen K, Seckl JR, Heinonen K, Pyhälä R, Feldt K, Jones A, Pesonen AK, Phillips DI, Lahti J, Järvenpää AL, Eriksson JG, Matthews KA, Strandberg TE, Kajantie E 2010 Maternal prenatal licorice consumption alters hypothalamic-pituitary-adrenocortical axis function in children. *Psychoneuroendocrinology* 35:1587–1593
11. Storm JJ, Lima SL 2010 Mothers forewarn offspring about predators: a transgenerational maternal effect on behavior. *Am Nat* 175:382–390
12. Reynolds RM 2012 Nick Hales Award Lecture 2011: glucocorticoids and the early life programming of cardiometabolic disease. *J Dev Origins Health Dis* 3:309–314
13. Newnham JP, Moss TJ, Nitsos I, Sloboda DM 2002 Antenatal corticosteroids: the good, the bad and the unknown. *Curr Opin Obstet Gynecol* 14:607–612
14. Doyle LW, Ford GW, Davis NM, Callanan C 2000 Antenatal corticosteroid therapy and blood pressure at 14 years of age in preterm children. *Clin Sci* 98:137–142
15. Dalziel SR, Walker NK, Parag V, Mantell C, Rea HH, Rodgers A, Harding JE 2005 Cardiovascular risk factors after antenatal exposure to betamethasone: 30-year follow-up of a randomised controlled trial. *Lancet* 365:1856–1862
16. Honest H, Bachmann LM, Gupta JK, Kleijnen J, Khan KS 2002 Accuracy of cervicovaginal fetal fibronectin test in predicting risk of spontaneous preterm birth: systematic review. *BMJ* 325:301
17. Honest H, Bachmann LM, Coomarasamy A, Gupta JK, Kleijnen J, Khan KS 2003 Accuracy of cervical transvaginal sonography in predicting preterm birth: a systematic review. *Ultrasound Obstet Gynecol* 22:305–322
18. Norwitz ER, Caughey AB 2011 Progesterone supplementation and the prevention of preterm birth. *Rev Obstet Gynecol* 4:60–72
19. 2010 Antenatal corticosteroids to reduce neonatal morbidity and mortality. Greentop Guideline no 7. London: Royal College of Obstetricians and Gynaecologists
20. Jung C, Ho JT, Torpy DJ, Rogers A, Doogue M, Lewis JG, Czajko RJ, Inder WJ 2011 A longitudinal study of plasma and urinary cortisol in pregnancy and postpartum. *J Clin Endocrinol Metab* 96:1533–1540
21. de Vries A, Holmes MC, Heijnis A, Seier JV, Heerden J, Louw J, Wolfe-Coote S, Meaney MJ, Levitt NS, Seckl JR 2007 Prenatal dexamethasone exposure induces changes in nonhuman primate offspring cardiometabolic and hypothalamic-pituitary-adrenal axis function. *J Clin Invest* 117:1058–1067