THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Multiple feeding strategies observed in the cold-water coral *Lophelia pertusa*

Citation for published version:

Murray, F, De Clippele, L, Hiley, A, Wicks, L, Roberts, J & Hennige, S 2019, 'Multiple feeding strategies observed in the cold-water coral *Lophelia pertusa*', *Journal of the Marine Biological Association of the UK*, vol. 99, no. 6, pp. 1281-1283. <https://doi.org/10.1017/S0025315419000298>

Digital Object Identifier (DOI):

[10.1017/S0025315419000298](https://doi.org/10.1017/S0025315419000298)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of the Marine Biological Association of the UK

Publisher Rights Statement:

COPYRIGHT: © Marine Biological Association of the United Kingdom 2019

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

*Title: Multiple feeding strategies observed in the cold-water coral *Lophelia pertusa**

Authors: Fiona Murray^{1}, Laurence H. De Clippele¹, Alexandra Hiley^{1,2}, Laura Wicks³, J. Murray Roberts¹, Sebastian Hennige^{1*}*

Author affiliations:

¹School of GeoSciences, The Grant Institute, James Hutton Road, King's Buildings, University of Edinburgh, EH9 3FE, UK

²*present address:* Nova Southeastern University, Halmos College of Natural Science and Oceanography in Dania Beach, Florida

³Heriot-Watt University, Riccarton, Edinburgh, EH14 4AS, UK

*Corresponding authors: fiona.murray25@gmail.com, s.hennige@ed.ac.uk

Abstract

Cold-water coral reefs are biodiversity hotspots of the deep sea. The most dominant reef building cold-water coral in the Atlantic is *Lophelia pertusa* which builds vast and structurally complex habitats. Studying the behaviours of deep-sea species is challenging due to the technological difficulties in making prolonged observations *in situ* so little is known about the behavioural ecology of this important species. Observations in laboratory studies can help to enhance our understandings of the range of behaviours these species have. Here we present video evidence that the cold-water coral *Lophelia pertusa* is capable of producing mucus nets as part of their feeding strategy. This finding suggests that have a more diverse range of feeding strategies than previously thought.

24 *Introduction*

25 Cold-water coral reefs are biodiversity hotspots of the deep sea. In contrast to their shallow, tropical
26 counterparts, these reefs are dominated by only a handful of reef framework-forming scleractinian
27 corals, yet these few species build vast and structurally complex reefs that support up to an estimated
28 1,300 species (Roberts et al. 2009). Since the discovery of the ecological importance of these diverse,
29 deep-sea communities, and the rapid technological advancement for exploring the deep sea at the
30 end of the last century, interest in cold-water corals has grown rapidly. For the North Atlantic, *Lophelia*
31 *pertusa* is the best studied, arguably most significant, and most widespread reef building cold-water
32 coral (e.g. Rogers 1999, Freiwald & Roberts 2005). Despite this attention, observations of the
33 behavioural ecology of *L. pertusa* remain limited due to the inaccessibility of their remote, deep-sea
34 homes. However, in contrast to many deep-sea species, *L. pertusa* has an extensive depth range (39
35 m - 3380 m, Mortensen et al. 2001) and is able to survive collection and transport to marine
36 laboratories where they can be maintained for months to years (e.g. Hennige et al. 2015). This allows
37 an insight into their behaviours that is beyond the scope of our current capabilities *in situ*. Here we
38 report on feeding behaviours recorded in laboratory mesocosms that suggest that the feeding
39 strategies of *L. pertusa* are more diverse than previously thought.

40

41 It has been established that the diet of *L. pertusa* consists predominantly of zooplankton and
42 phytodetritus (Carlier et al. 2009, Mueller et al. 2014), and previous laboratory observations reported
43 that polyps caught food items through nematocyst adhesion (Mortensen, 2001): that is, they capture
44 items that come into contact with their tentacles. Polyps then transfer particles to the centre of the
45 oral disc and into the pharynx for consumption. Mortensen (2001) noted that small amounts of mucus
46 were also excreted when potential prey had come into contact with the tentacles but had
47 subsequently escaped. Our understanding of the production of mucus in relation to feeding was
48 previously limited to Mortensen's observations and it was thought that *L. pertusa* was limited to

consuming food items that came into direct contact with its tentacles. Indeed, mucus excretion has been predominantly reported as a disturbance response (Mortensen 2001), an antifouling strategy (Reitner 2005) and to have a role in skeletal growth (Reitner 2005). However, we have now produced video evidence on a freshly collected *L. pertusa* specimen that *L. pertusa* is able to construct mucus nets that it casts into the water column to capture food.

Methods

Colonies of *Lophelia pertusa* were collected using a modified video assisted van-Veen grab³ from 141 – 167 m depth at the Mingulay Reef Complex, Outer Hebrides, UK (56°49'N, 7°23'W, see Fig. 1 in Hennige et al. 2014), in June 2011 during the RRS *Discovery* D366/7 Cruise. Upon return to the surface, corals were placed in a holding tank at ambient seabed temperature conditions (9.5 °C) for 2 days, to recover from collection, at which time polyps were extended and feeding and mucus was no longer visibly being produced (Hennige et al. 2014). Corals were then carefully fragmented into smaller pieces (Hennige et al. 2014). These fragments had 5-20 polyps, and were taken from the top of sampled colonies to ensure that relatively young polyps were used consistently, as polyp age can determine physiological response (Maier et al. 2009). Coral fragments were transferred to tanks in a 10 °C temperature-controlled room, fed cultured *Artemia salina*, and acclimatised for five days, (comparable to Naumann et al. 2014).

The fragment presented here was filmed for 1 hour using a Canon PowerShot G9. The video was edited using Final Cut Pro X (version 10.3) and sped up 5 times.

Results and Discussion

73

74 Following the introduction of *A. salina*, small quantities of mucus were released creating two distinct
75 and separate mucus nets. Subsequently the nets and captured *A. salina* were consumed (Fig. 1). The
76 entire process from net production to consumption lasted approximately 18 minutes, far longer than
77 an ROV (remotely operated vehicle) generally spends on a single patch of coral during deep-sea
78 expeditions, making it difficult to observe such behaviours *in situ*. The key sequence is presented in
79 still images (Fig. 1) and video clips are included as a digital supplement (Video S1). Whilst corals
80 habitually produce mucus as a stress response when collected, our acclimation procedures and
81 observations that the mucus nets were only produced in the presence of food allows us to conclude
82 that this is a feeding behaviour.

83

84 These observations suggest that *L. pertusa* has a more diverse range of feeding strategies than
85 previously thought. The prevalence and frequency of the use of mucus nets remains unknown,
86 however other benthic species use a similar strategy and have been more comprehensively studied.
87 Polychaetes (e.g. *Hediste diversicolor*, Riisgård 1991) and gastropods (e.g. *Dendropoma maximum*,
88 Rybak et al. 2005) for example, produce mucus nets as part of their suspension feeding strategies. In
89 both cases, environmental cues, such as food availability, have been shown to influence choice of
90 feeding method.

91

92 Scleractinian coral mucus is functionally important and a prominent source of dissolved organic matter
93 (DOM) in both warm and cold-water coral reefs (Rix et al. 2016). It plays a key role in nutrient cycling
94 (Wild et al. 2008) and drives the “sponge loop”- the key trophic pathway to transfer DOM, the most
95 abundant nutrient rich resource on tropical coral reefs, to higher trophic levels (de Goeij et al. 2013).

Our observations suggest that the functions of mucus for the coral itself may also be more diverse and important than first appears.

Acknowledgements

This paper is a contribution to the UK Ocean Acidification Research Programme supported by Natural Environment Research Council NE/H017305/1, NE/K009028/1, NE/K009028/2.

References

- Carrier A., Le Guilloux, E., Olu K., Sarrazin J., Mastrototaro F., Taviani M., Clavier J. (2009) Trophic relationships in a deep Mediterranean cold-water coral bank (Santa Maria di Leuca, Ionian Sea). *Mar. Ecol. Prog. Ser.* 397: 125-137.
- de Goeij, J.M., van Oevelen, D., Vermeij, M.J.A., Osinga, R., Middelburg, J.J., de Goeij, A.F.P.M., Admiraal, W. (2013). Surviving in a marine desert: the sponge loop retains resources within coral reefs. *Science*. 642:6154 108-110.
- Freiwald, A., & Roberts, J. M. (Eds.). (2005). Cold-water corals and ecosystems. *Springer Science & Business Media*.
- Hennige, S. J., Wicks, L.C., Kamenos, N.A., Bakker, D.C.E., Findlay, H.S., Dumousseaud, C., Roberts, J.M. (2014). Short-term metabolic and growth responses of the cold-water coral *Lophelia pertusa* to ocean acidification. *Deep. Res. Part II Top. Stud. Oceanogr.* 99, 27–35.
- SJ Hennige, S.J., Wicks, L.C., Kamenos, N.A., Perna, G., Findlay, H.S., Roberts, J.M. (2015). Hidden impacts of ocean acidification to live and dead coral framework. *Proc. R. Soc. B* 282 (1813), 20150990.
- Maier, C., Hegeman, J., Weinbauer, M. G. & Gattuso, J.-P. Calcification of the cold-water coral *Lophelia pertusa* under ambient and reduced pH. *Biogeosciences* 6, 1671–1680 (2009).
- Mortensen, P.B. (2001) Aquarium observations on the deep-water coral *Lophelia pertusa* (L., 1758) (scleractinia) and selected associated invertebrates, *Ophelia*, 54:2, 83-104, DOI:10.1080/00785236.2001.10409457.
- Mueller, C.E., Larsson, A.I., Veuger, B., Middelburg, J.J., van Oevelen, D. (2014). Opportunistic feeding on various organic food sources by the cold-water coral *Lophelia pertusa*. *Biogeosciences*, 11: 123-133.
- Naumann, M. S., Orejas, C. & Ferrier-Pagès, C. (2014). Species-specific physiological response by the cold-water corals *Lophelia pertusa* and *Madrepora oculata* to variations within their natural temperature range. *Deep. Res. Part II Top. Stud. Oceanogr.* 99, 36–41.

- Reitner, J. (2005). Calcifying extracellular mucus substances (EMS) of *Madrepora oculata*—a first geobiological approach. *Cold-water corals and ecosystems*, 731-744.
- Riisgård, H.U. (1991). Suspension feeding in the polychaete *Nereis diversicolor*. *Marine Ecology Progress Series*, 29-37.
- Rix, L., de Goeij, J.M., Mueller, C.E., Struck, U., Middelburg, J.J., van Duyl, F.C., Al-Horani, F.A., Wild, C., Naumann, M.S., van Oevelen, D. (2016). Coral mucus fuels the sponge loop in warm- and cold-water coral reef ecosystems. *Scientific Reports*. 6.
- Roberts, J. M., Wheeler, A. J., Freiwald, A., Cairns, S. D. (2009). Cold-Water Corals: The Biology and Geology of Deep-Sea Coral Habitats. *Cambridge University Press*.
- Rogers, A. D. (1999). The Biology of *Lophelia pertusa* (Linnaeus 1758) and Other Deep-Water Reef-Forming Corals and Impacts from Human Activities. *International review of hydrobiology*, 84(4), 315-406.
- Ribak, G., Heller, J., & Genin, A. (2005). Mucus-net feeding on organic particles by the vermetid gastropod *Dendropoma maximum* in and below the surf zone. *Marine Ecology Progress Series*, 293, 77-87.
- Wild, C., Mayr, C., Wehrmann, L., Schöttner, S., Naumann, M., Hoffmann, F., & Rapp, H. T. (2008). Organic matter release by cold water corals and its implication for fauna–microbe interaction. *Marine Ecology Progress Series*, 372, 67-75.

Figure legend

Fig. 1: Production and consumption of mucus net: (a) free swimming *A. salina*, (b) production of mucus net and trapping of plankton, (c-d) pulling in of mucus net towards oral disc, (g) absence of mucus net after consumption. The black dots approximately represent visible *A. salina*.

Appendix

Video S1: *L. pertusa* catching and consuming *A. salina* using a mucus net.