



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Common Genetic Variants Associate with Serum Phosphorus Concentration

Citation for published version:

Kestenbaum, B, Glazer, NL, Koettgen, A, Felix, JF, Hwang, S-J, Liu, Y, Lohman, K, Kritchevsky, SB, Hausman, DB, Petersen, A-K, Gieger, C, Ried, JS, Meitinger, T, Strom, TM, Wichmann, HE, Campbell, H, Hayward, C, Rudan, I, de Boer, IH, Psaty, BM, Rice, KM, Chen, Y-DI, Li, M, Arking, DE, Boerwinkle, E, Coresh, J, Yang, Q, Levy, D, van Rooij, FJA, Dehghan, A, Rivadeneira, F, Uitterlinden, AG, Hofman, A, van Duijn, CM, Shlipak, MG, Kao, WHL, Witteman, JCM, Siscovick, DS & Fox, CS 2010, 'Common Genetic Variants Associate with Serum Phosphorus Concentration', *Journal of the American Society of Nephrology*, vol. 21, no. 7, pp. 1223-1232. <https://doi.org/10.1681/ASN.2009111104>

Digital Object Identifier (DOI):

[10.1681/ASN.2009111104](https://doi.org/10.1681/ASN.2009111104)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of the American Society of Nephrology

Publisher Rights Statement:

Copyright © 2010 by the American Society of Nephrology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Common Genetic Variants Associate with Serum Phosphorus Concentration

Bryan Kestenbaum,* Nicole L. Glazer,[†] Anna Köttgen,[‡] Janine F. Felix,^{§||} Shih-Jen Hwang,[¶] Yongmei Liu,^{**} Kurt Lohman,^{††} Stephen B. Kritchevsky,^{‡‡} Dorothy B. Hausman,^{§§} Ann-Kristin Petersen,^{|||} Christian Gieger,^{|||} Janina S. Ried,^{|||} Thomas Meitinger,^{¶¶} Tim M. Strom,^{¶¶} H. Erich Wichmann,^{|||} Harry Campbell,^{***} Caroline Hayward,^{†††} Igor Rudan,^{***†††} Ian H. de Boer,* Bruce M. Psaty,^{†§§§|||¶¶¶} Kenneth M. Rice,^{†****} Yii-Der Ida Chen,^{††††} Man Li,[‡] Dan E. Arking,^{††††} Eric Boerwinkle,^{§§§§|||} Josef Coresh,^{†¶¶¶¶****} Qiong Yang,[¶] Daniel Levy,[¶] Frank J.A. van Rooij,^{§||} Abbas Dehghan,^{§||} Fernando Rivadeneira,^{||†††††} André G. Uitterlinden,^{||†††††} Albert Hofman,^{§||} Cornelia M. van Duijn,^{§||} Michael G. Shlipak,^{†††††} W.H. Linda Kao,[‡] Jacqueline C.M. Witteman,^{§||} David S. Siscovick,^{†§§§|||} and Caroline S. Fox^{¶§§§§§}

Author affiliations are listed at the end of the manuscript

ABSTRACT

Phosphorus is an essential mineral that maintains cellular energy and mineralizes the skeleton. Because complex actions of ion transporters and regulatory hormones regulate serum phosphorus concentrations, genetic variation may determine interindividual variation in phosphorus metabolism. Here, we report a comprehensive genome-wide association study of serum phosphorus concentration. We evaluated 16,264 participants of European ancestry from the Cardiovascular Health Study, Atherosclerosis Risk in Communities Study, Framingham Offspring Study, and the Rotterdam Study. We excluded participants with an estimated GFR <45 ml/min per 1.73 m² to focus on phosphorus metabolism under normal conditions. We imputed genotypes to approximately 2.5 million single-nucleotide polymorphisms in the HapMap and combined study-specific findings using meta-analysis. We tested top polymorphisms from discovery cohorts in a 5444-person replication sample. Polymorphisms in seven loci with minor allele frequencies 0.08 to 0.49 associate with serum phosphorus concentration ($P = 3.5 \times 10^{-16}$ to 3.6×10^{-7}). Three loci were near genes encoding the kidney-specific type IIa sodium phosphate co-transporter (*SLC34A1*), the calcium-sensing receptor (*CASR*), and fibroblast growth factor 23 (*FGF23*), proteins that contribute to phosphorus metabolism. We also identified genes encoding phosphatases, kinases, and phosphodiesterases that have yet-undetermined roles in phosphorus homeostasis. In the replication sample, five of seven top polymorphisms associate with serum phosphorus concentrations ($P < 0.05$ for each). In conclusion, common genetic variants associate with serum phosphorus in the general population. Further study of the loci identified in this study may help elucidate mechanisms of phosphorus regulation.

J Am Soc Nephrol 21: 1223–1232, 2010. doi: 10.1681/ASN.2009111104

Received November 2, 2009. Accepted March 2, 2010.

Published online ahead of print. Publication date available at www.jasn.org.

B.K., N.L.G., A.K., J.F.F., and S.-J.H. contributed equally to this work.

W.H.L.K., J.C.M.W., D.S.S., and C.S.F. contributed equally to this work.

Correspondence: Dr. Bryan Kestenbaum, University of Washington, Kidney Research Institute, Division of Nephrology, Department of Medicine, Harborview Medical Center, Room 10EH11, Box 359764, Seattle, WA 98104-2499. Phone: 206-731-4029; Fax: 206-731-2252; E-mail: brk@u.washington.edu

Copyright © 2010 by the American Society of Nephrology

Phosphorus is an essential mineral that is responsible for maintaining cellular energy, mineralizing the skeleton, and protecting nonosseous tissue from calcification.¹ In humans, the majority of phosphorus resides within bone and teeth as hydroxyapatite and within cells as a component of nucleic acids and phospholipid membranes. A small proportion of phosphorus (approximately 1%) circulates in the serum under tight regulation by the complex actions of specialized ion transporters and regulatory hormones, which balance gastrointestinal phosphorus absorption, bone uptake, cellular flux, and excretion through the kidneys.²

Dysregulated phosphorus metabolism may lead to adverse clinical consequences. In the setting of chronic kidney disease, higher serum phosphorus concentrations are associated with vascular calcification and mortality.^{3,4} Higher serum phosphorus concentrations within the normal laboratory range are also associated with incident cardiovascular events among individuals without known kidney disease.^{5,6} In cell culture models, phosphorus directly transforms vascular smooth muscle tissue into osteoblast-like cells that calcify the medial vessel wall.^{7,8}

Several lines of evidence suggest that interindividual differences in the steady-state serum phosphorus concentration may be partly heritable. Rare phosphorus wasting disorders, such as hypophosphatemic rickets, are caused by specific mutations in genes within phosphorus metabolic pathways.^{9,10} Genetic disruption of hormones that regulate phosphorus alters circulating phosphorus concentrations in animal models.^{11,12} Moreover, dietary phosphorus intake is only weakly associated with the serum phosphorus concentration in humans.¹³

We conducted the first large-scale genome-wide association study (GWAS) to investigate common genetic variants associated with serum phosphorus concentrations in the general population. We conducted a meta-analysis of results from 16,264 individuals of European ancestry participating in the Cardiovascular Health Study, the Atherosclerosis Risk in Communities Study, the Framingham Heart Study, and the Rotterdam Study.

RESULTS

Genome-wide Associations for the Serum Phosphorus Concentration

A total of 16,264 study participants from four cohorts were available for meta-analysis (Table 1). Seven genetic loci located on chromosomes 1, 3, 5, 6, and 12 met the prespecified meta-analysis statistical significance threshold for the association with serum phosphorus (Figure 1). The strongest statistical association was observed for locus 1p36.13 ($P = 3.5 \times 10^{-16}$). The top single-nucleotide polymorphism (SNP) within this locus (rs1697421) does not reside within any known gene, but lies within 10 kb of genes encoding for tissue-nonspecific alkaline phosphatase (*ALPL*) and neuroblastoma breakpoint family member 3 (*NBPF3*; Table 2, Figure 2A).

On chromosome 3, the top SNP within locus 3q21.1 (rs17265703) was in strong linkage disequilibrium ($R^2 = 0.93$)

with a second SNP (rs1801725), which represents a known alanine (A) to serine (S) polymorphism of amino acid 986 in the calcium-sensing receptor (*CASR*) gene (Figure 2B).¹⁴

On chromosome 5, locus 5q35.3 contained several significant SNPs that were associated with serum phosphorus (Figure 2C). The top SNP was located within an intron of *RGS14*, a regulator of G-protein signaling that lies directly adjacent to *SLC34A1*, which encodes for the kidney-specific type IIa sodium phosphate co-transporter (Npt2a).

Three distinct loci on chromosome 6 (6p21.31, 6q23.2, and 6q23.1) were each associated with differences in the circulating serum phosphorus concentration (Figure 2, D through F). Loci on 6q23.1 and 6q23.2 are presumed independent because they are >4000 kb apart and are not correlated ($R^2 < 0.001$), and simultaneous adjustment for both SNPs did not alter the results for either. Locus 6p21.31 contains two intronic SNPs (rs6942022 and rs9469578) in strong linkage disequilibrium within the inositol hexakisphosphate kinase 3 (*IHPK3*) gene. The top significant SNP at locus 6q23.2 is located near the gene encoding for phosphodiesterase 7B (*PDE7B*). The top significant SNP at locus 6q23.1 is located within an intron of the ectonucleotide pyrophosphatase/phosphodiesterase 3 (*ENPP3*) gene, which lies adjacent to the *ENPP1* gene.

On chromosome 12, locus 12p13.32 lies within an open reading frame that is adjacent to RAD51-associated protein (*RAD51AP1*) and fibroblast growth factor 6 (*FGF6*). The gene encoding for the phosphaturic factor fibroblast growth factor 23 (*FGF23*) is located 133 kb upstream from this top SNP; however, there is a recombination point between them (Figure 2G). Several SNPs within the *FGF23* gene were close to meeting genome-wide statistical significance for the association with serum phosphorus concentration.

Replication

There were 5444 individuals available for the replication meta-analysis (Table 3). For all seven SNPs measured in replication, regression coefficients were in the same direction as those from the discovery sample. On the basis of β coefficients adjusted for bias in the discovery cohort and minor allele frequencies for the top seven SNPs, statistical power for replication was limited in some cases. For five of the top SNPs, replication associations were significant at the $P = 0.05$ level, and three were associated at the Bonferroni-corrected $P = 0.007$ level (Table 3).

Interactions by Sex and Estimated Kidney Function

Associations of the seven top SNPs with the serum phosphorus concentration were similar in magnitude comparing men with women, and no significant interactions were found (all P for interaction >0.1). Results also remained similar after restricting the analysis to individuals who had an estimated GFR (eGFR) ≥ 60 ml/min per 1.73 m².

Gene Score Model

A gene score model that encompassed the sum of risk alleles from the top seven SNPs was significantly associated with the

Table 1. Characteristics of the discovery and replication cohorts

	Discovery Cohorts				Replication Cohorts			
	CHS (n = 1761)	FHS (n = 2865)	ARIC (n = 8048)	RS (n = 3516)	KORA-F3 (n = 1599)	KORA-F4 (n = 1779)	Health ABC (n = 1507)	Vis (n = 659)
Age, years	71.0 (4.6)	43.6 (9.8)	54.3 (5.7)	69.4 (9.1)	62.5 (10.1)	60.9 (8.9)	73.7 (2.8)	56.8 (15.3)
Women	69.6%	52.1%	52.9%	59.4%	50.5%	51.3%	46.9%	60.1%
eGFR, ml/min	82.3 (21.4)	103.4 (35.9)	89.7 (17.7)	80.7 (14.9)	80.1 (18.1)	81.9 (18.0)	72.4 (13.4)	95.7 (34.7)
eGFR 45 to 60 ml/min	14.3%	3.2%	2.7%	4.8%	11.5%	7.6%	21.5%	10.2%
Serum phosphorus								
Mean, mg/dl	3.60	3.15	3.41	3.68	3.30	3.39	3.51	3.50
(SD)	(0.49)	(0.44)	(0.48)	(0.60)	(0.55)	(0.50)	(0.48)	(0.54)
Median, mg/dl	3.6	3.2	3.4	3.7	3.3	3.4	3.5	3.5
(IQR)	(3.3, 3.9)	(2.9, 3.4)	(3.1, 3.7)	(3.3, 4.1)	(2.9, 3.7)	(3.1, 3.7)	(3.2, 3.8)	(3.1, 3.9)

CHS, Cardiovascular Health Study; FHS, Framingham Health Study; ARIC, Atherosclerosis Risk in Communities Study; RS, Rotterdam Study; eGFR, estimated glomerular filtration rate; IQR, interquartile range.

serum phosphorus concentration in the discovery sample ($P = 2.6 \times 10^{-57}$; unadjusted for regression to the mean). The majority of individuals had gene scores between 3 and 12 (Figure 3). In aggregate, the top seven SNPs identified in this study explained approximately 1.5% of the variation in serum phosphorus concentrations (model $R^2 = 0.0147$). Serum phosphorus concentrations ranged from 3.24 to 3.58 mg/dl across the spectrum of risk allele dosage.

DISCUSSION

In this first genome-wide association study of the serum phosphorus concentration, we identified seven genetic loci, located on chromosomes 1, 3, 5, 6, and 12, that were associated with the circulating serum phosphorus concentration. In replica-

tion meta-analysis, five of these loci were associated with serum phosphorus at the $P = 0.05$ level and three were associated at the $P = 0.007$ level. Three loci identified in the discovery analysis were near genes that encode for proteins known to be involved in phosphorus metabolism: the sodium phosphate co-transporter type IIa, the calcium-sensing receptor, and *FGF23*. Other candidate genes identified in this study include a phosphatase, a kinase, and two phosphodiesterases that have a yet-undetermined role in the regulation of phosphorus balance.

Some of the genes identified in this study support previously understood biology of phosphorus metabolism. The sodium phosphate co-transporter type IIa (*Npt2a*) is expressed in the proximal tubule of the kidney and plays a major role in determining urinary phosphorus excretion.² Serum phosphorus excess leads to the removal of *Npt2a* transporters from the cell-surface, resulting in diminished phosphorus re-absorption and subsequent elimination of phosphorus in the urine.¹⁵ Conversely, a low serum phosphorus concentration stimulates upregulation of *Npt2a*, which enhances phosphorus re-absorption in the kidney. Deletion of the *Npt2a* gene in animal models leads to marked urinary phosphorus wasting and impaired skeletal development.¹¹ Heterozygous mutations in *Npt2a* have been described in humans with nephrolithiasis and urinary phosphorus leak.¹⁶

The calcium-sensing receptor can influence phosphorus metabolism through connections with key hormones that regulate both calcium and phosphorus.¹⁷ For example, a low serum calcium level detected by the calcium-sensing receptor stimulates release of parathyroid hormone, which increases urinary phosphorus excretion via downregulation of *Npt2a*.¹⁸ A low serum calcium level also stimulates the production of calcitriol, which enhances absorption of both calcium and phosphorus through the gastrointestinal tract via actions on their selective ion channels.^{19,20}

The SNP rs2970818 is located within 50 kb of *FGF6* and within 133 kb of *FGF23*. *FGF6* is a paracrine factor that plays a




Figure 1. Single nucleotide polymorphisms on chromosomes 1, 3, 5, 6, and 12 associate with the serum phosphorus concentration. Log (P values) for individual SNPs plotted for each chromosome in the discovery sample. Horizontal line represents genome-wide significance level of 4×10^{-7} .

Table 2. Genetic loci associated with the serum phosphorus concentration in the discovery sample

Locus	Top SNP ID	Position	Major/Minor Allele (MAF)	Meta		In Gene	Closest Genes ^b
				β^a	P		
1p36.13	rs1697421	21695879	G/A (0.49)	+0.044	3.47×10^{-16}	—	ALPL NBPF3
3q21.1	rs17265703	123531334	A/G (0.15)	-0.040	6.24×10^{-08}	CSTA	CASR CCDC58
5q35.3	rs4074995	176729949	G/A (0.28)	-0.032	6.25×10^{-08}	RGS14	SLC34A1 PFN3 LMAN2 F12
6p21.31	rs9469578	33814457	C/T (0.08)	-0.064	5.15×10^{-10}	IHPK3	ITPR3 LEMD2 MLN
6q23.1	rs453639	132091350	A/C (0.36)	+0.036	3.59×10^{-07}	ENPP3	
6q23.2	rs947583	136175352	T/C (0.29)	+0.035	2.19×10^{-09}	—	PDE7B
12p13.32	rs2970818	4476429	T/A (0.09)	+0.052	4.04×10^{-08}	—	FGF6 RAD51AP1 FGF23 ^c

MAF, minor allele frequency; ALPL, alkaline phosphatase, liver/bone/kidney; NBPF3, neuroblastoma breakpoint family, member 3; CSTA, cystatin A; CASR, calcium-sensing receptor; CCDC58, coiled-coil domain-58; RGS14, regulator of G-protein signaling 14; SLC34A1, solute carrier family 34 (sodium phosphate); PFN3, profilin 3; LMAN2, lectin, mannose-binding 2; F12, coagulation factor XII (Hageman factor); IHPK3, inositol hexakisphosphate kinase 3; ITPR3, inositol 1,4,5-triphosphate receptor, type 3; LEMD2, LEM domain containing 2; MLN, motilin; ENPP3, ectonucleotide pyrophosphatase/phosphodiesterase 3; PDE7B, phosphodiesterase 7B; FGF6, fibroblast growth factor 6; RAD51AP1, RAD51 associated protein 1; FGF23, fibroblast growth factor 23.

^aAssociation with serum phosphate concentration in mg/dl.

^bWithin 50 kb.

^cDistance = 133 kb.

role in myoblast proliferation and muscle differentiation;²¹ associations of FGF6 with phosphorus metabolism have not been reported. Another fibroblast growth factor, FGF23, is an endocrine factor that is considered to be central to the regulation of phosphorus balance.²² Originally identified as the cause of tumor-induced phosphorus wasting,²³ FGF23 binds to its receptor within the kidney to potentially increase the urinary fractional excretion of phosphorus.²⁴ The hormone klotho is a key co-factor for FGF23; deletion of FGF23 or klotho in animal models yields a similar phenotype characterized by hyperphosphatemia, short-life span, vascular calcification, and osteoporosis.^{25,26} Mutations in FGF23 are the cause of autosomal dominant hypophosphatemic rickets,⁹ and elevated FGF23 levels are recently identified as the cause of phosphorus wasting in kidney transplant recipients.²⁷

The most robust association in this study was for SNP RS1697421, which is located adjacent to the tissue-nonspecific alkaline phosphatase gene. Alkaline phosphatase is a membrane-bound enzyme that plays a key role in mineralizing the skeleton. In osteoblasts, alkaline phosphatase hydrolyzes pyrophosphate into two phosphate molecules, which are used to synthesize hydroxyapatite. Although this process will raise phosphorus levels within cells, the effect on the serum phosphorus concentration is unknown. Mutations in the alkaline phosphatase gene are linked with hypophosphatasia, a rare inherited condition characterized by failure to mineralize the teeth and bone, and excess circulating levels of pyrophosphate, a natural calcification inhibitor.²⁸ However, to our knowledge

differences in the serum phosphorus concentration among individuals with hypophosphatasia have not been reported. Mutations in the ENPP1 gene are linked with generalized arterial calcification of infancy, and polymorphisms in this gene are associated with insulin resistance and obesity.^{29,30} Inositol hexakisphosphate kinase-3 is suspected to convert inositol hexakisphosphate to diphosphoinositol pentakisphosphate.³¹

Associations of individual genetic loci with the serum phosphorus concentration were modest. In aggregate, the seven top genetic loci identified in this study contributed to an approximate 0.4 mg/dl difference in the serum phosphorus concentration among the discovery sample. This variation is similar to that naturally observed between men and women and may be relevant for cardiovascular health. For example, each 0.5 mg/dl higher serum phosphorus concentration was linearly associated with an approximate 15% greater risk of cardiovascular events in the Framingham Heart Study and the Cholesterol And Recurrent Events study.^{5,6}

Known intraindividual variation in serum phosphorus concentrations may have precluded identification of additional loci. Serum phosphorus concentrations vary throughout the day and from day to day, although they do not appear to be associated with fasting status.³² In one study of 1878 participants in the National Health and Nutrition Examination Survey (NHANES) who underwent serum phosphorus measurements approximately 2 weeks apart, the correlation between serum phosphorus concentrations was 0.63.¹³ In future studies, multiple measurements of serum phosphorus within an




Figure 2. Seven loci meet genome wide significance level for the association with the serum phosphorus concentration. Observed log (*P* values) by chromosome position for genetic regions surrounding each top SNP in the discovery sample. Recombination rates are shown in red on the right-sided Y-axis. Correlation of the top SNP with surrounding SNPs indicated by shading—darker boxes indicate greater correlation (scale in top right-hand corner). (A) Locus 1p36.13; (B) locus 3q21.1; (C) locus 5q35.3; (D) locus 6p21.31; (E) locus 6q21.3; (F) locus 6q23.2; (G) locus 12p13.32.

individual would help reduce this source of variation. Our study was limited to Caucasian subjects; therefore, results do not apply to other race/ethnicities. The SNPs identified in this

study do not imply a direct causal relationship with the serum phosphorus concentration; fine mapping and re-sequencing studies are needed to pinpoint potential causal variants, and

Table 3. Meta-analysis of the replication cohorts

SNP ID	Replication <i>n</i>	Power ^a	Discovery β	Replication β	Replication <i>P</i>	Combined β^b	Combined <i>P</i> ^b
rs1697421	5536	0.999	+0.044	+0.070	2.08×10^{-14}	+0.050	1.14×10^{-27}
rs17265703	5543	0.730	-0.040	-0.026	0.015	-0.036	4.32×10^{-9}
rs4074995	5497	0.734	-0.032	-0.010	0.168	—	—
rs9469578	5544	0.978	-0.064	-0.046	0.004	-0.059	1.11×10^{-11}
rs453639	5544	0.409	+0.036	+0.014	0.096	—	—
rs947583	5543	0.921	+0.035	+0.034	3.94×10^{-4}	+0.035	3.45×10^{-12}
rs2970818	5544	0.760	+0.052	+0.034	0.025	+0.047	4.38×10^{-9}

^aPower to replicate findings from the discovery cohorts.

^bCombined results shown for replication *P* < 0.05.




Figure 3. Mean serum phosphorus concentrations associate with the number of SNPs individually associated with serum phosphorous. Mean serum phosphorus concentration as a function of the number of risk alleles in the discovery sample; number of participants with the risk alleles shown below the graph.

functional studies are needed to understand how the specific genetic variation may be realized at the protein level. Because of known selection bias in selecting SNPs on the basis of genome-wide statistical significance and relatively low allele frequencies, study power to replicate associations for loci 3q21.1, 5q35.3, 6q23.1, and 12p13.32 was inadequate. β coefficients for SNPs within these loci were smaller in magnitude, but in the same direction as those from the discovery cohort. Finally, this genome-wide association study is limited to detecting common genetic variants and is therefore unable to identify associations for rare alleles, such as those responsible for phosphorus wasting disorders in humans.

In summary, we demonstrate that common genetic variants are associated with the serum phosphorus concentration using meta-analysis of study-specific GWAS from four large adult populations. Follow-up studies are needed to identify potential causal genetic loci in or near the candidate genes identified in this GWAS. Candidate genes may be explored in more comprehensive metabolic studies of phosphorus metabolism and in translational animal models that could shed new light on the mechanisms and clinical implications of phosphorus homeostasis.

CONCISE METHODS

CHARGE Consortium

The Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium was developed to facilitate the conduct and replication of genetic association studies across established cohort studies.³³ Each participating study approved guidelines for collaboration, and the institutional review boards for each study approved the consent procedures, data security measures, and collection of genetic material. All study participants provided written informed consent for genetic research.

Discovery Study Populations

Serum phosphorus measurements were available from four participating discovery cohorts within the CHARGE consortium: the Cardiovascular Health Study (CHS), the Framingham Heart Study (FHS), the Atherosclerosis Risk in Communities Study (ARIC), and the Rotterdam Study (RS).

CHS is a community-based study of cardiovascular disease among adults ages 65 years and older.³⁴ CHS recruited 5201 ambulatory older adults in 1989 to 1990 from four U.S. communities: Forsyth County, NC; Sacramento County, CA; Washington County, MD; Pittsburgh, PA. Subjects were excluded if they were institutionalized, required a wheelchair in the home, or were undergoing treatment for cancer. Serum phosphorus concentrations and genomic data were available for 2337 CHS participants without cardiovascular disease at the time of 1992 to 1993 exam.

FHS is a prospective cohort study initiated in 1948 to investigate risk factors for cardiovascular disease. Children and spouses of children from the original cohort were recruited in 1972 (Framingham Offspring Study) and followed prospectively with clinic examinations every 4 years.³⁵ Serum phosphorus concentrations and genomic data were available for 2865 participants during the second examination cycle in 1979 to 1982.⁵

ARIC is a community-based, prospective cohort study of atherosclerosis that recruited 15,792 Caucasian and African-American individuals aged 45 to 64 years from four U.S. communities: Forsyth County, NC, Washington County, MD, Minneapolis, MN, and Jackson, MS.³⁶ Participants underwent three follow-up examinations at 3-year intervals. Serum phosphorus concentrations and genomic data were available for 8122 ARIC participants from the baseline examination in 1987 to 1989.

RS is a community-based cohort study to assess determinants of chronic diseases among the elderly that recruited 7983 participants (78% of the eligible population) aged ≥ 55 years living in a well-defined suburb of the city of Rotterdam, The Netherlands, in 1990 to 1993.³⁷ Home visits were used to collect health status information. Participants were subsequently examined at the research center in 1990 to 1993 and every 3 to 4 years thereafter. Serum phosphorus concentrations and genomic data were available for 3516 RS participants.

Because of the relatively small number of non-Caucasian participants, we limited analyses to participants of primarily European (Caucasian) ancestry. Previous studies have demonstrated that serum phosphorus concentrations increase markedly in advanced stages of chronic kidney disease.³⁸ We therefore excluded participants who had an estimated GFR < 45 ml/min per 1.73 m² to focus on potential factors that regulate phosphorus under normal conditions. We estimated GFR using the four-variable Modification of Diet in Renal Disease equation.³⁹

Replication Study Populations

KORA S3 is an independent population-based sample from the general population living in the region of Augsburg, Southern Germany, in 1994 to 1995. A total of 3006 subjects participated in a follow-up examination in 2004 to 2005 (KORA F3). The KORA S4 survey (4261 participants) is an independent population-based sample from the same region and was conducted in 1999 to 2001.⁴⁰ Serum phosphorus concentrations and genomic data were available for 1599 participants of KORA F3 and for 1779 participants of KORA F4. The Vis study was conducted in 2003 to 2004 among 986 unselected, 18- to 93-year-old Croats recruited from the villages of Vis and Komiza, on the Dalmatian island of Vis. These settlements have unique population histories and have preserved isolation from other villages and from the outside world for centuries. Health ABC is a prospective cohort study designed to investigate the effect of health conditions on age-related functional changes. Participants aged 70 to 79 years were recruited from the metropolitan areas surrounding Pittsburgh, PA, and Memphis, TN. Eligibility criteria were no difficulty walking one-quarter of a mile, climbing 10 steps, or performing basic activities of daily living.

Serum Phosphorus Measurements

Blood specimens were collected in the morning, centrifuged, and either frozen at -70°C for storage or run daily; not all participants were fasting. Serum phosphorus concentrations were quantified using an automated platform (Beckman-Coulter for CHS and ARIC, Roche for FHS, and a Kone Diagnostica reagent kit and a Kone autoanalyzer for RS) in which inorganic phosphorus reacts with ammonium molybdate in an acidic solution to form a colored phosphomolybdate complex. The reported intra-assay coefficients of variation for these assays are 5.8 and 5.6%, respectively. Intraindividual biologic variation in serum phosphorus measurements are available from a previous population-based cohort study in which 1878 participants underwent repeat serum phosphorus measurements a median of 16 days apart.¹³ The correlation between measurements was 0.63, and the SD for the difference in measurements was 0.31 mg/dl.

Genotyping

CHS.

Genotyping was performed at the General Clinical Research Center's Phenotyping/Genotyping Laboratory at Cedars-Sinai using the Illumina 370CNV BeadChip system. Genotypes were called using the Illumina BeadStudio software. Samples were excluded for genotypic sex mismatch, discordance with prior genotyping, or call rate $< 95\%$. Genotyping was successful in 96%. The following exclusions were then applied: call rate $< 97\%$, Hardy Weinberg Equilibrium $P < 10^{-5}$, duplicate error or Mendelian inconsistency, heterozygote frequency of approximately 0, or SNP not found in dbSNP. SNPs were further excluded from analysis if the ratio of the variance of the allele dosage to the variance expected under Hardy Weinberg Equilibrium was < 0.01 .

ARIC.

Genotyping was performed using the Affymetrix 6.0 array and genotypes were called using the Birdsees software. Of 8861 samples, 734 were excluded because of discordance with previous genotype data, mismatch of reported and genotypic sex, first-degree relative of an included individual based on genotype data, or genetic outliers as assessed by average allele sharing and principal components analyses. Before imputation, SNPs with call rate $< 95\%$, minor allele frequency $< 1\%$, and Hardy-Weinberg equilibrium $P < 10^{-5}$ were excluded to ensure good quality for imputation. After imputation, no SNPs were excluded.

FHS.

Genotyping was conducted using the Affymetrix 500K mapping array and the Affymetrix 50K gene-focused macrophage inflammatory protein array. Individuals were excluded when their call rate across all SNPs was $< 97\%$. After exclusion of SNPs with genotype call rate $< 95\%$ or Hardy-Weinberg equilibrium $P < 10^{-6}$, there were 503,551 SNPs available for analysis.

RS.

Genotyping was done with the Illumina 550K array (Illumina, San Diego, CA, USA) in self-reported Caucasian individuals. Individuals were excluded for an overall call rate $< 97.5\%$, excess autosomal heterozygosity, mismatch between genotypic and phenotypic sex, or outliers identified by the identity-by-state clustering analysis; genotyping was successful in 93% of RS. SNPs were excluded when the minor allele frequency was $\leq 1\%$, Hardy-Weinberg equilibrium $P < 10^{-6}$, or SNP call rate $\leq 98\%$, resulting in 530,683 directly measured SNPs used for imputation.

Imputation

In all studies, genotypes were imputed to approximately 2.5 million SNPs in HapMap, using the Phase II CEU individuals as a reference panel. For imputation software, ARIC, FHS, and RS used MACH and CHS used BAMBAM.^{41,42} Imputation results are summarized as an "allele dosage" (a fractional value between 0 and 2), defined as the expected number of copies of the minor allele at that SNP.

Statistical Analysis

We evaluated serum phosphorus concentrations as a continuous variable, in mg/dl. The distribution of serum phosphorus concentrations

was not strongly skewed and was analyzed without transformation. We did not exclude any individuals based on their serum phosphorus concentration.

Genome-wide analyses were conducted within each cohort. Using an additive genetic model, we used linear regression to evaluate the association between the allele dosage and serum phosphorus concentration, quantifying the regression slope (β) and standard error [SE(β)]. We adjusted analyses for age, sex, and study site. We accounted for relatedness of individuals in FHS using random effects to account for the covariance between family members with the specific covariance structure determined by the degree of relatedness between each relative pair.³³ Within each study, genomic control was used to adjust each study's standard errors for potential effects of population stratification; the genomic control λ values ranged from 1.01 to 1.04 across the discovery and replication cohorts. In all cohorts except for Health ABC, principal components were not associated with the serum phosphorus concentration. In Health ABC, principal components were associated with serum phosphorus; therefore, we adjusted for principal components in Health ABC to account for population substructure. We combined within-study associations by meta-analysis, using inverse variance weighting. After meta-analysis, we filtered results on weighted minor allele frequency <0.005 . We selected an *a priori* genome-wide significance threshold of 4×10^{-7} , which corresponds to a ≤ 1 expected false-positive result for 2.5 million tests. The validity of this boundary is not affected by correlation between test statistics.⁴³

To assess interactions, we used a genotype-sex interaction parameter within the same linear modeling and meta-analysis framework. To summarize the main effects from the discovery sample, we constructed an additive gene score model using the number of copies of each risk allele from each SNP (possible values 0 to 14) and used linear regression to evaluate the association of gene score with serum phosphorus concentration.

We estimated power to replicate observed associations from the discovery phase in a replication sample size of 5544, an observed SD for phosphorus of 0.5, and a one-sided α of 0.05. Allele frequencies were based on the discovery cohorts and effect sizes were adjusted for the "winners curse" selection bias.⁴⁴ Because of this adjustment and relatively low minor allele frequencies, power was limited to replicate findings in loci 3q21.1, 5q35.3, 6q23.1, and 12p13.32.

ACKNOWLEDGMENTS

We acknowledge the individual participating studies and investigators of the CHARGE (Cohorts for Heart and Aging Research in Genome Epidemiology).

Cardiovascular Health Study.

The CHS research reported in this article was supported by contracts N01-HC-85079 through N01-HC-85086, N01-HC-35129, N01-HC-15103, N01-HC-55222, N01-HC-75150, and N01-HC-45133, grants U01-HL080295 and R01 HL087652, and R01 AG027002 from the National Heart, Lung, and Blood Institute, with additional contribution from the National Institute of Neurological Disorders and

Stroke. A full list of principal CHS investigators and institutions can be found at <http://www.chs-nhlbi.org/pi.htm>. DNA handling and genotyping was supported in part by the National Center for Research Resources grant M01RR00069 to the Cedars-Sinai General Clinical Research Center Genotyping core and National Institute of Diabetes and Digestive and Kidney Diseases grant DK063491 to the Southern California Diabetes Endocrinology Research Center. Dr. Kestenbaum received grant funding for the serum phosphate measurements under National Institutes of Health R01 HL084443.

Atherosclerosis Risk in Communities Study.

The Atherosclerosis Risk in Communities Study is carried out as a collaborative study supported by National Heart, Lung, and Blood Institute contracts N01-HC-55015, N01-HC-55016, N01-HC-55018, N01-HC-55019, N01-HC-55020, N01-HC-55021, N01-HC-55022, R01HL087641, R01HL59367, and R01HL086694; National Human Genome Research Institute contract U01HG004402; and National Institutes of Health contract HHSN268200625226C. The authors thank the staff and participants of the ARIC study for their important contributions. Infrastructure was partly supported by grant UL1RR025005, a component of the National Institutes of Health and NIH Roadmap for Medical Research.

Framingham Heart Study.

This research was conducted in part using data and resources from the Framingham Heart Study of the National Heart, Lung, and Blood Institute of the National Institutes of Health and Boston University School of Medicine. The analyses reflect intellectual input and resource development from the Framingham Heart Study investigators participating in the SNP Health Association Resource (SHARe) project. This work was partially supported by the National Heart, Lung, and Blood Institute's Framingham Heart Study (contract N01-HC-25195) and its contract with Affymetrix, Inc., for genotyping services (contract N02-HL-6-4278). A portion of this research utilized the Linux Cluster for Genetic Analysis (LinGA-II) funded by the Robert Dawson Evans Endowment of the Department of Medicine at Boston University School of Medicine and Boston Medical Center.

Rotterdam Study.

The Rotterdam Study is supported by the Erasmus MC and Erasmus University Rotterdam; the Netherlands Organization for Scientific Research; the Netherlands Organization for Health Research and Development (ZonMw); the Research Institute for Diseases in the Elderly; the Netherlands Heart Foundation; the Ministry of Education, Culture and Science; the Ministry of Health, Welfare and Sports; the European Commission; and the Municipality of Rotterdam. Support for genotyping was provided by the Netherlands Organization for Scientific Research (NWO) (175.010.2005.011, 911.03.012) and Research Institute for Diseases in the Elderly (RIDE). This study was further supported by the Netherlands Genomics Initiative (NGI)/Netherlands Organisation for Scientific Research (NWO) project 050-060-810. We thank P. Arp, M. Jhamai, M. Moorhouse, M. Verkerk, and S. Bervoets for their help in creating the Rotterdam database and M. Struchalin for his contributions to the imputations of the Rotterdam data.

KORA Study.

The MONICA/KORA Augsburg studies were financed by the Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany, and supported by grants from the German Federal Ministry of Education and Research (BMBF). Part of this work was financed by the German National Genome Research Network (NGFN) and supported within the Munich Center of Health Sciences (MC Health) as part of LMUinnovativ. We gratefully acknowledge the contribution of P. Lichtner, G. Eckstein, and all other members of the Helmholtz Zentrum München genotyping staff in generating and analyzing the SNP dataset and G. Fischer for data management. We thank all members of field staffs who were involved in the planning and conduct of the MONICA/KORA Augsburg studies.

Vis Study.

The Vis study in the Croatian island of Vis was supported by grants from the Medical Research Council UK and from the Ministry of Science, Education and Sports of the Republic of Croatia and by funds from EUROSPAN (European Special Populations Research Network) supported by European Commission FP6 STRP grant 018947 (LSHG-Court-2006-01947). The Vis authors collectively thank a large number of individuals for their help in organizing, planning, and carrying out the field work related to the project and data management: Professor Pavao Rudan and the staff of the Institute for Anthropological Research in Zagreb, Croatia (organization of the field work, anthropometric and physiological measurements, and DNA extraction); Professor Ariana Vorko-Jovic and the staff and medical students of the Andrija Stampar School of Public Health of the Faculty of Medicine, University of Zagreb, Croatia (questionnaires, genealogy reconstruction, and data entry); Dr. Branka Salzer from the biochemistry lab "Salzer", Croatia (measurements of biochemical traits); local general practitioners and nurses (recruitment and communication with the study population); and the employees of several other Croatian institutions who participated in the field work, including but not limited to the University of Rijeka and Split, Croatia; Croatian Institute of Public Health; Institutes of Public Health in Split and Dubrovnik, Croatia. SNP genotyping of the Vis samples was carried out by the Genetics Core Laboratory at the Wellcome Trust Clinical Research Facility, WGH, Edinburgh, Scotland.

DISCLOSURES

None.

AUTHOR AFFILIATIONS

*Division of Nephrology, Department of Medicine, University of Washington, Kidney Research Institute, Seattle, Washington;

†Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, Washington; Departments of

‡Epidemiology and ¶Biostatistics, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland;

§Epidemiology and ¶¶Internal Medicine, and

||Netherlands Consortium for Healthy Ageing, Erasmus Medical

Center, Rotterdam, The Netherlands; ¶¶Framingham Heart Study, National Heart, Lung, and Blood Institute, Framingham, Massachusetts; Departments of **Epidemiology and Prevention and ¶¶Biostatistical Sciences, and ¶¶Sticht Center on Aging, Wake Forest University School of Medicine, Winston-Salem, North Carolina; §§Department of Foods and Nutrition, University of Georgia, Athens, Georgia; Institutes of ¶¶Epidemiology and ¶¶Human Genetics, Helmholtz Zentrum München, Neuherberg, Germany; ***Public Health Sciences Institute of Genetics and Molecular Medicine, University of Edinburgh, Scotland, UK; ¶¶Human Genetics Unit, University of Edinburgh Western General Hospital, Edinburgh, Scotland, UK; ¶¶Croatian Centre for Global Health, University of Split Medical School, Split, Croatia; Departments of §§§Medicine, ¶¶Epidemiology, and ¶¶Biostatistics, University of Washington, Seattle, Washington; ¶¶Group Health Research Institute, Group Health, Seattle, Washington; ¶¶Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, California; ¶¶¶McKusick-Nathans Institute of Genetic Medicine and ¶¶¶Department of Medicine, Johns Hopkins Medical Institutions, Baltimore, Maryland; §§§§Human Genetics Center and ¶¶¶Division of Epidemiology, University of Texas, Houston, Texas; ¶¶¶¶General Internal Medicine Section, Veterans Affairs Medical Center, University of California San Francisco, San Francisco, California; §§§§§Harvard Medical School, Boston, Massachusetts

REFERENCES

- Hruska K, Slatopolsky E: Disorders of phosphorous, calcium, and magnesium metabolism. In: *Diseases of the kidney*, edited by Schrier R, Gottschalk C, London, Little, Brown, and Company, 1996, pp 2477–2526
- Kronenberg HM: NPT2a—The key to phosphate homeostasis. *N Engl J Med* 347: 1022–1024, 2002
- Adeney KL, Siscovick DS, Ix JH, Seliger SL, Shlipak MG, Jenny NS, Kestenbaum BR: Association of serum phosphate with vascular and valvular calcification in moderate CKD. *J Am Soc Nephrol* 20: 381–387, 2009
- Block GA, Hulbert-Shearon TE, Levin NW, Port FK: Association of serum phosphorus and calcium x phosphate product with mortality risk in chronic hemodialysis patients: a national study. *Am J Kidney Dis* 31: 607–617, 1998
- Dhingra R, Sullivan LM, Fox CS, Wang TJ, D'Agostino RB Sr, Gaziano JM, Vasan RS: Relations of serum phosphorus and calcium levels to the incidence of cardiovascular disease in the community. *Arch Intern Med* 167: 879–885, 2007
- Tonelli M, Sacks F, Pfeffer M, Gao Z, Curhan G: Relation between serum phosphate level and cardiovascular event rate in people with coronary disease. *Circulation* 112: 2627–2633, 2005
- Jono S, McKee MD, Murry CE, Shioi A, Nishizawa Y, Mori K, Morii H, Giachelli CM: Phosphate regulation of vascular smooth muscle cell calcification. *Circ Res* 87: E10–E17, 2000
- Giachelli CM: Vascular calcification: in vitro evidence for the role of inorganic phosphate. *J Am Soc Nephrol* 14: S300–S304, 2003
- Autosomal dominant hypophosphataemic rickets is associated with mutations in FGF23. *Nat Genet* 26: 345–348, 2000
- Bastepe M, Juppner H: Inherited hypophosphatemic disorders in children and the evolving mechanisms of phosphate regulation. *Rev Endocr Metab Disord* 9: 171–180, 2008
- Beck L, Karaplis AC, Amizuka N, Hewson AS, Ozawa H, Tenenhouse HS: Targeted inactivation of Npt2 in mice leads to severe renal phosphate wasting, hypercalciuria, and skeletal abnormalities. *Proc Natl Acad Sci U S A* 95: 5372–5377, 1998

12. Shimada T, Kakitani M, Yamazaki Y, Hasegawa H, Takeuchi Y, Fujita T, Fukumoto S, Tomizuka K, Yamashita T: Targeted ablation of Fgf23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *J Clin Invest* 113: 561–568, 2004
13. de Boer IH, Rue TC, Kestenbaum B: Serum phosphorus concentrations in the third National Health and Nutrition Examination Survey (NHANES III). *Am J Kidney Dis* 53: 399–407, 2009
14. Marz W, Seelhorst U, Wellnitz B, Tiran B, Obermayer-Pietsch B, Renner W, Boehm BO, Ritz E, Hoffmann MM: Alanine to serine polymorphism at position 986 of the calcium-sensing receptor associated with coronary heart disease, myocardial infarction, all-cause, and cardiovascular mortality. *J Clin Endocrinol Metab* 92: 2363–2369, 2007
15. Biber J, Forgo J, Murer H: Modulation of Na⁺-Pi cotransport in opossum kidney cells by extracellular phosphate. *Am J Physiol* 255: C155–C161, 1988
16. Prie D, Huart V, Bakouh N, Planelles G, Dellis O, Gerard B, Hulin P, Benque-Blanchet F, Silve C, Grandchamp B, Friedlander G: Nephrolithiasis and osteoporosis associated with hypophosphatemia caused by mutations in the type 2a sodium-phosphate cotransporter. *N Engl J Med* 347: 983–991, 2002
17. Brown EM, Gamba G, Riccardi D, Lombardi M, Butters R, Kifor O, Sun A, Hediger MA, Lytton J, Hebert SC: Cloning and characterization of an extracellular Ca(2+)-sensing receptor from bovine parathyroid. *Nature* 366: 575–580, 1993
18. Kempson SA, Lotscher M, Kaissling B, Biber J, Murer H, Levi M: Parathyroid hormone action on phosphate transporter mRNA and protein in rat renal proximal tubules. *Am J Physiol* 268: F784–F791, 1995
19. Birge SJ, Miller R: The role of phosphate in the action of vitamin D on the intestine. *J Clin Invest* 60: 980–988, 1977
20. Bouillon R, Van Cromphaut S, Carmeliet G: Intestinal calcium absorption: Molecular vitamin D mediated mechanisms. *J Cell Biochem* 88: 332–339, 2003
21. Armand AS, Pariset C, Laziz I, Launay T, Fiore F, Della Gaspera B, Birnbaum D, Charbonnier F, Chanoine C: FGF6 regulates muscle differentiation through a calcineurin-dependent pathway in regenerating soleus of adult mice. *J Cell Physiol* 204: 297–308, 2005
22. Liu S, Quarles LD: How fibroblast growth factor 23 works. *J Am Soc Nephrol* 18: 1637–1647, 2007
23. White KE, Jonsson KB, Carn G, Hampson G, Spector TD, Mannstadt M, Lorenz-Depiereux B, Miyauchi A, Yang IM, Ljunggren O, Meitinger T, Strom TM, Juppner H, Econs MJ: The autosomal dominant hypophosphatemic rickets (ADHR) gene is a secreted polypeptide overexpressed by tumors that cause phosphate wasting. *J Clin Endocrinol Metab* 86: 497–500, 2001
24. Yu X, Ibrahim OA, Goetz R, Zhang F, Davis SI, Garringer HJ, Linhardt RJ, Ornitz DM, Mohammadi M, White KE: Analysis of the biochemical mechanisms for the endocrine actions of fibroblast growth factor-23. *Endocrinology* 146: 4647–4656, 2005
25. Sitara D, Razzaque MS, Hesse M, Yoganathan S, Taguchi T, Erben RG, Juppner H, Lanske B: Homozygous ablation of fibroblast growth factor-23 results in hyperphosphatemia and impaired skeletogenesis, and reverses hypophosphatemia in PheX-deficient mice. *Matrix Biol* 23: 421–432, 2004
26. Kuro-o M, Matsumura Y, Aizawa H, Kawaguchi H, Suga T, Utsugi T, Ohyama Y, Kurabayashi M, Kaname T, Kume E, Iwasaki H, Iida A, Shiraki-Iida T, Nishikawa S, Nagai R, Nabeshima YI: Mutation of the mouse *klotho* gene leads to a syndrome resembling ageing. *Nature* 390: 45–51, 1997
27. Evenepoel P, Naesens M, Claes K, Kuypers D, Vanrenterghem Y: Tertiary ‘hyperphosphatoniism’ accentuates hypophosphatemia and suppresses calcitriol levels in renal transplant recipients. *Am J Transplant* 7: 1193–1200, 2007
28. Whyte MP: Hypophosphatasia and the role of alkaline phosphatase in skeletal mineralization. *Endocr Rev* 15: 439–461, 1994
29. Rutsch F, Ruf N, Vaingankar S, Toliat MR, Suk A, Hohne W, Schauer G, Lehmann M, Roscioli T, Schnabel D, Epplen JT, Knisely A, Superti-Furga A, McGill J, Filippone M, Sinaiko AR, Vallance H, Hinrichs B, Smith W, Ferre M, Terkeltaub R, Nurnberg P: Mutations in ENPP1 are associated with ‘idiopathic’ infantile arterial calcification. *Nat Genet* 34: 379–381, 2003
30. Meyre D, Bouatia-Naji N, Tounian A, Samson C, Lecoer C, Vatin V, Ghousaini M, Wachter C, Herberg S, Charpentier G, Patsch W, Pattou F, Charles MA, Tounian P, Clement K, Jouret B, Weill J, Maddux BA, Goldfine ID, Walley A, Boutin P, Dina C, Froguel P: Variants of ENPP1 are associated with childhood and adult obesity and increase the risk of glucose intolerance and type 2 diabetes. *Nat Genet* 37: 863–867, 2005
31. Saiardi A, Nagata E, Luo HR, Snowman AM, Snyder SH: Identification and characterization of a novel inositol hexakisphosphate kinase. *J Biol Chem* 276: 39179–39185, 2001
32. Isakova T, Gutierrez O, Shah A, Castaldo L, Holmes J, Lee H, Wolf M: Postprandial mineral metabolism and secondary hyperparathyroidism in early CKD. *J Am Soc Nephrol* 19: 615–623, 2008
33. Psaty B, O'Donnell C, Gudnason V, Lunetta K, Folsom A, Rotter J, Uitterlinden A, Harris T, Witteman JCM, Boerwinkle E: Cohorts for heart and aging research in genomic epidemiology (CHARGE) consortium: design of prospective meta-analyses of genome-wide association studies from 5 cohorts. *Circ Cardiovasc Genet* 2: 73–80, 2009
34. Fried LP, Borhani NO, Enright P, Furberg CD, Gardin JM, Kronmal RA, Kuller LH, Manolio TA, Mittelmark MB, Newman A, O'Leary D, Psaty B, Rautaharju PM, Tracy R: The Cardiovascular Health Study: design and rationale. *Ann Epidemiol* 1: 263–276, 1991
35. Feinleib M, Kannel WB, Garrison RJ, McNamara PM, Castelli WP: The Framingham Offspring Study. Design and preliminary data. *Prev Med* 4: 518–525, 1975
36. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *Am J Epidemiol* 129: 687–702, 1989
37. Hofman A, Breteler MM, van Duijn CM, Janssen HL, Krestin GP, Kuipers EJ, Stricker BH, Tiemeier H, Uitterlinden AG, Vingerling JR, Witteman JC: The Rotterdam Study: 2010 objectives and design update. *Eur J Epidemiol* 24: 553–572, 2009
38. Kestenbaum B, Sampson JN, Rudser KD, Patterson DJ, Seliger SL, Young B, Sherrard DJ, Andress DL: Serum phosphate levels and mortality risk among people with chronic kidney disease. *J Am Soc Nephrol* 16: 520–528, 2005
39. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D: A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 130: 461–470, 1999
40. Wichmann HE, Gieger C, Illig T: KORA-gen-resource for population genetics, controls and a broad spectrum of disease phenotypes. *Gesundheitswesen* 67 Suppl 1: S26–S30, 2005
41. Li Y, Ding J, Abecasis G: Mach 1.0: Rapid haplotype reconstruction and missing genotype inference. Presented at the 2006 Annual Meeting of the American Society of Human Genetics, New Orleans, LA, October 9–13, 2006
42. Servin B, Stephens M: Imputation-based analysis of association studies: candidate regions and quantitative traits. *PLoS Genet* 3: 1296–1308, 2007
43. Gordon A, Glazko G, Qiu X, Yakovlev A: Control of the mean number of false discoveries, Bonferroni and stability of multiple testing. *Ann Appl Stat* 1: 179–190, 2007
44. Zhong H, Prentice RL: Bias-reduced estimators and confidence intervals for odds ratios in genome-wide association studies. *Biostatistics* 9: 621–634, 2008