THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Lymphoma with Mott cell differentiation and validation of immunohistochemical lymphoid markers in an African pygmy hedgehog (*Atelerix albiventris*)

Citation for published version:

Cazzini, P, Richardson, J, Smith, N, Lodzinska, J, Robinson, AL & Philbey, A 2019, 'Lymphoma with Mott cell differentiation and validation of immunohistochemical lymphoid markers in an African pygmy hedgehog (*Atelerix albiventris*)', *Veterinary Clinical Pathology: An International Journal of Laboratory Medicine*.
<https://doi.org/10.1111/vcp.12816>

Digital Object Identifier (DOI):

[10.1111/vcp.12816](https://doi.org/10.1111/vcp.12816)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Veterinary Clinical Pathology: An International Journal of Laboratory Medicine

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Title: Lymphoma with Mott cell differentiation and validation of immunohistochemical**
2 **lymphoid markers in an African pygmy hedgehog (*Atelerix albiventris*)**

4 Running header: Lymphoma with Mott cell differentiation in a hedgehog

6 Paola Cazzini^{1*}, Jenna Richardson², Nicola Smith¹, Joanna Lodzinska³, Amy L. Robinson⁴,
7 Adrian W. Philbey¹

9 ¹ Easter Bush Pathology, Royal (Dick) School of Veterinary Studies and The Roslin
10 Institute, The University of Edinburgh, Easter Bush, Edinburgh EH25 9RG, UK

11 ² Hospital for Small Animals, Rabbit and Exotic Animal Practice, Royal (Dick) School of
12 Veterinary Studies and The Roslin Institute, The University of Edinburgh, Easter Bush,
13 Edinburgh EH25 9RG, UK

14 ³ Hospital for Small Animals, Diagnostic Imaging Department, Royal (Dick) School of
15 Veterinary Studies and The Roslin Institute, The University of Edinburgh, Easter Bush,
16 Edinburgh EH25 9RG, UK

17 ⁴ Royal (Dick) School of Veterinary Studies and The Roslin Institute, The University of
18 Edinburgh, Easter Bush EH25 9RG, Edinburgh, UK

21 *Correspondence: Paola Cazzini; current address: Easter Bush Pathology, Royal (Dick)
22 School of Veterinary Studies and The Roslin Institute, The University of Edinburgh, Easter
23 Bush, Edinburgh EH25 9RG, UK. E-mail: paola.cazzini@ed.ac.uk

Key words: African pygmy hedgehog, lymphoma, Mott cell, plasma cell, immunohistochemistry, cytology

Abstract

A 2 year old, female entire African pygmy hedgehog was presented for diagnostic investigation of a 2-month reduction in appetite, with weight loss, and recent vomiting. Clinical examination revealed a large, firm mass originating from the left cranial abdomen. Ultrasound guided fine needle aspirates of the mass, the liver and the mesenteric lymph nodes revealed a population of pleomorphic round cells, some of which contained variable numbers of round, clear vacuoles, consistent with a diagnosis of lymphoma with Mott cell differentiation. At post-mortem examination, there was marked diffuse splenic enlargement, with infiltration by a soft tissue mass. There were multiple coalescing masses in the liver, pallor of the kidneys, and enlargement of the mesenteric lymph nodes. On histological examination, the spleen, lymph nodes, and masses in the liver were extensively infiltrated by proliferating lymphoid cells with plasmacytoid and Mott cell differentiation. Cells with Mott cell morphology had accumulation of periodic acid-Schiff positive material in cytoplasmic inclusions, and were positive for cytoplasmic nucleic acids when stained with methyl green pyronin. In the population of neoplastic lymphoid cells, a majority of cells expressed the transcription factor Pax5, which drives B cell differentiation, and a minority expressed transcription factor IRF4/MUM-1, which drives plasma cell differentiation, consistent with a B cell lymphoma with plasmacytoid differentiation.

Case presentation

A 2 year old, female entire, African pygmy hedgehog (*Atelerix albiventris*) was presented to the Rabbit and Exotic Animal Practice at the Royal (Dick) School of Veterinary Studies,

Edinburgh, in February 2017, for diagnostic investigation of a 2-month period of weight loss and recent episodes of vomiting. On initial observations, the hedgehog was ambulatory, but with a subdued demeanor and exercise intolerance, when allowed to explore the mammal ward floor. The respiratory rate was elevated and there was increased respiratory effort and noise.

Clinical examination was performed under general anesthesia. The hedgehog weighed 275 g and had a reduced body condition score at 2 out of 5. Abdominal palpation revealed a large, well-defined mass in the cranial abdomen. On ultrasound examination, a large, heterogeneous mass was identified; the mass occupied most of the peritoneal cavity, causing a mass effect on all the abdominal organs. The mass appeared to be of splenic origin because of its location and appearance; no normal splenic parenchyma was visualized. The displaced abdominal organs, including the liver, both kidneys and the mesenteric lymph nodes, were also involved. The liver was moderately heterogeneous, with multiple hypoechoic nodules. Both kidneys were hyperechoic and had decreased definition of the corticomedullary junction. The abdominal lymph nodes were enlarged and heterogeneous. A moderate quantity of anechoic peritoneal effusion was evident.

The presumptive diagnosis from ultrasound investigation was splenic neoplasia, most likely lymphoma, with metastasis to the liver, kidneys and abdominal lymph nodes. Fine needle aspirates (FNAs) were taken from the abdominal mass, liver and lymph nodes, routinely stained by the May-Grünwald-Giemsa method, and examined microscopically. The abdominal mass FNAs were highly cellular and heavily hemodiluted. A population of pleomorphic round cells predominated. They ranged from large, round cells (approximately 20 µm in diameter), with a high nuclear to cytoplasmic ratio (N:C), deeply basophilic cytoplasm, occasionally with perinuclear clearing and some clear vacuoles, and a round to

76 irregular nucleus, with finely stippled chromatin, to smaller, oval cells, with a more abundant
77 cytoplasm, filled with clear vacuoles, and a peripherally located nucleus with clumped
78 chromatin (Figure 1 A & B). Mitotic figures were frequent and occasionally were bizarre
79 (Figure 1B). Similar cells were present in the mesenteric lymph nodes and in the liver.
80 Unstained lymph node smears were stained with periodic acid-Schiff (PAS), which indicates
81 the presence of carbohydrates, and methyl green pyronin (MGP), a histochemical stain used
82 to visualise RNA and DNA. The cytoplasmic inclusions of the neoplastic cells were strongly
83 positive for PAS (Figure 1 C), suggesting the presence of antibodies that have been
84 glycosylated in the endoplasmic reticulum,¹ while the cells had marked cytoplasmic
85 pyroninophilia (Figure 1 D), demonstrating high cytoplasmic nucleic acid (RNA) content.

86
87 A diagnosis of lymphoma with Mott cell differentiation was established. The hedgehog
88 was euthanized on welfare grounds without additional work-up (e.g. hematology and
89 biochemistry) and submitted for post-mortem examination. The spleen was markedly
90 enlarged, weighing 76 g, and was mottled, dark red to purple, with multilobular,
91 coalescent, pale yellow-brown masses; fibrin strands and haemorrhage were present on the
92 surface (Figure 2). The liver was moderately enlarged, weighing 25 g, and was diffusely
93 pale yellow-brown, with rounded borders and an accentuated lobular pattern. Multiple,
94 raised, soft, pale yellow-brown nodules, approximately 5 mm in diameter, were present in
95 the right lateral and right medial lobes of the liver. The mesenteric lymph nodes were
96 enlarged and pale yellow-brown. The kidneys were slightly enlarged and pale brown. The
97 brain was unremarkable on external examination; on sectioning, a 3 to 4 mm diameter
98 white mass was evident, involving the ventral cerebellum and adjacent medulla oblongata.
99 The lungs were moderately, diffusely oedematous. There were 6 mL of clotted blood in the
100 peritoneal cavity. The bone marrow was pale yellow. The gross findings were consistent
101 with infiltrative neoplasia of the spleen, liver, kidneys, and brain.

102

103 On histological examination, affected areas of the spleen, liver, kidneys, and lymph nodes
104 were infiltrated, expanded and effaced by dense sheets of large, round, moderately
105 pleomorphic, neoplastic lymphoid cells (Figure 3 A). The infiltrating cells had large, round
106 to ovoid, sometimes indented, nuclei, clumped chromatin, large, prominent, eosinophilic
107 nucleoli, and moderate amounts of eosinophilic to amphophilic cytoplasm; in some cells,
108 there was a clear perinuclear area, consistent with a plasmacytoid morphology. Mott cells
109 with eosinophilic cytoplasmic inclusions, consistent with Russell bodies, were scattered
110 through the tissue. There were six to eight mitoses per high power field (400x
111 magnification). Apoptosis was evident and there was multifocal, locally extensive necrosis
112 and haemorrhage. Infiltrates of neutrophils were present in some areas. The bone marrow
113 was hypercellular with a myeloid predominance. Mildly increased numbers of plasma cells
114 and Mott cells were present; however, it was unclear if these and some of the larger
115 immature cells present represented an early neoplastic infiltrate.

116

117 Immunohistochemistry was performed on the spleen; healthy splenic tissue from an
118 unrelated African pygmy hedgehog was retrieved from our archive and used as a control
119 for validation of the immunohistochemical stain. Sections (4µm thickness) of formalin-
120 fixed, paraffin wax-embedded tissue were placed on SuperFrost® Plus coated slides
121 (Thermo Electron, Runcorn, Cheshire, UK), dewaxed, hydrated, and rinsed in distilled
122 water. To block non-specific endogenous peroxidase activity, sections were treated with
123 blocking agent (Dako REAL blocking agent S202386) for 10 minutes at room temperature.
124 Antibodies were diluted in antibody diluent (S0809, Dako) at 1/200 for detection of CD3
125 (mouse monoclonal anti-CD3; Novocastra, NCL-L-CD3), 1/40 for detection of PAX 5
126 (mouse monoclonal anti-PAX 5; Becton & Dickinson, P67320) and 1/40 for detection of
127 MUM 1/IRF4 (interferon regulatory factor 4; rabbit polyclonal anti-MUM1; Thermofisher,

PA5-32511). Antigen retrieval was performed using 0.01 M citrate buffer pH 6.0 at 110 °C for 5 minutes (Histos 5 microwave processor), then sections were incubated with the primary antibody for 30 minutes at room temperature (RT) following antigen retrieval. Following incubation with primary antibody, the sections were incubated with secondary antibody (Envision anti-mouse HRP; Dako K4007) and visualized with DAB+ chromogen (Dako K3468). Sections were counterstained with Harris haematoxylin. All washings between steps were carried out using Tris-buffered saline with Tween (Thermo Fisher Scientific TA-999-TT).

On immunohistochemical examination of the splenic mass in the African pygmy hedgehog with lymphoma, a majority of neoplastic lymphoid cells (60-70%) expressed the transcription factor Pax5 (Figure 3 B), whilst a minority (10-20%) expressed the transcription factor MUM-1 (Figure 3 C). This was consistent with the majority of the neoplastic cells being morphologically compatible with lymphocytes, and with lower numbers being morphologically consistent with Mott cells. Moderate numbers of CD3 positive tumor infiltrating lymphocytes were present within the areas of neoplastic infiltration, representing 5% of the total cell population (Figure 3 D). In samples of spleen from a control African pygmy hedgehog that died of unrelated disease, there was positive immunostaining for Pax5, CD3, and MUM1 in expected lymphoid tissue zones (Supplementary Figures 4 – 6).

Discussion

Neoplasia is reported commonly in African pygmy hedgehogs (*Atelerix albiventris*).² In a study on captive African hedgehogs, 53% of animals between 2 and 5.5 years of age had at least one type of tumor, while 8.6% of animals had more than one type of tumor.² In another retrospective study of 14 African hedgehogs, the prevalence of neoplasia was 29%.³

The most commonly reported neoplasms are carcinomas and lymphomas, and malignant neoplasia is more frequent.²

Mott cells are plasma cells which have retained immunoglobulins packed in vesicles, known as Russell bodies, giving these cells a distinctive appearance.^{4, 5} Due to the carbohydrate component of the immunoglobulins, Russell bodies are also positive when stained with PAS (Figure 1 C).^{1, 6, 7} Plasmacytoid cells producing immunoglobulin and immunoblasts contain high quantities of rough endothelial reticulum and their cytoplasm is therefore positive on staining with methyl green pyronin, which highlights nucleic acid (RNA) content (Figure 1 D).^{6, 8}

As described in dogs¹ in our case the neoplasia was composed of many lymphoid cells, which occasionally contained some vacuoles (Russell bodies), and by lower numbers of Mott cells, in the absence of cells with a morphology consistent with well differentiated plasma cells. For this reason, a plasma cell tumor, including a multiple myeloma, was considered less likely, and the neoplasia was considered morphologically consistent with a lymphoma with Mott cell differentiation. Immunohistochemistry was consistent with the morphologic diagnosis. The transcription factor Pax5 is expressed in all pre-B and mature B cell stages.⁹ Pax5 expression is downregulated when B cells undergo plasma cell differentiation.¹⁰ In our case, as in similar previous cases in other species,^{7, 11} the majority of neoplastic cells expressed the transcription factor Pax5, indicating a B cell origin (Figure 3 B). Some plasma cell tumors, including multiple myeloma, can express Pax5, as well as other B cell markers, such as CD20.¹² Therefore, the diagnosis of B cell lymphoma with Mott cell differentiation, in the present case, was based on the morphological appearance of the population of neoplastic cells, supported by demonstration of Pax5 expression. MUM1 is a transcription factor expressed in plasma cells.¹³ In our case, as in previously reported cases,^{7, 14} cells with

Mott cell differentiation expressed nuclear immunohistochemical positivity for MUM1 (Figure 3 C). The presence of scattered CD3 positive T-cells (Figure 3 D) was interpreted to be due to the presence of tumor infiltrating T lymphocytes. Negative control samples from an African pygmy hedgehog without neoplasia provided validation for the immunohistochemical markers in this species (Supplementary Figure 4 – 6).

Analogous B cell lymphomas have been described in dogs,^{1, 11, 14, 15} in a cat,⁷ and in a ferret.¹⁶ Similar to the case reported here, previous cases were characterized by a bi-phasic population of larger, more immature lymphoid B cells that differentiate to Mott cells filled with characteristic Russell bodies. While plasma globulins can be within the reference interval, and no obvious monoclonal peak may be detected using electrophoresis,¹⁶ immunofixation demonstrated the presence of monoclonal bands in the IgM and IgA proteins in two dogs.¹⁵ Circulating neoplastic cells have been described in two dogs with B cell lymphomas with Mott cell differentiation.¹ Unfortunately the overall low number of reported cases is insufficient to reach any definitive conclusions on the prognosis for this neoplasia.

In conclusion, this is the first case report of B cell lymphoma with Mott cell differentiation in a hedgehog, with validation of Pax5, CD3, and MUM1 immunohistochemical markers in this species.

Acknowledgments

The authors would like to thank Neil Macintyre from Easter Bush Pathology, The University of Edinburgh and The Roslin Institute, for the assistance with the special stains and immunohistochemistry.

Figures

Figure 1. Fine needle aspirate of a splenic mass of an African pygmy hedgehog. (A and B) Note the presence of a pleomorphic population of round cells that often have clear cytoplasmic vacuoles; (B) mitotic figures are also present. May-Grünwald-Giemsa stain. (A) 60x objective. (B) 100x objective. (C) Cytoplasmic vacuoles are PAS positive, supporting their identification as Russell bodies. Periodic acid-Schiff stain, 60x objective. (D) The presence of cytoplasmic red staining suggests a high RNA content. Methyl green pyronin stain, 100x objective.

Figure 2. Gross findings at post-mortem examination of a hedgehog. The spleen is enlarged, mottled, and covered with a fibrin exudate. The liver is pale and has rounded borders. There is a fluid effusion in the pleural cavity.

Figure 3. Histologic sections from an abdominal mass in a hedgehog. (A) Histological appearance of the spleen showing diffuse infiltrates of neoplastic cells with plasmacytoid morphology and several Mott cells containing Russell bodies (arrows). 40x objective. Scale bar = 50 μ m. (B) Immunohistochemistry for Pax5, (C) MUM1, and (D) CD3 in the spleen. 40x objective. Scale bars = 50 μ m.

Supplementary figure 4. Immunohistochemistry for Pax5 in lymphoid tissue in the white pulp of a normal spleen in a control African pygmy hedgehog. 40x objective. Scale bars = 50 μ m.

Supplementary figure 5. Immunohistochemistry for MUM in lymphoid tissue in the white pulp of a normal spleen in a control African pygmy hedgehog. 40x objective. Scale bars = 50 μ m.

Supplementary figure 6. Immunohistochemistry for CD3 in lymphoid tissue in the white pulp of a normal spleen in a control African pygmy hedgehog. 40x objective. Scale bars = 50 μ m.

References

1. Stacy NI, Nabity MB, Hackendahl N, Buote M, Ward J, Ginn PE, Vernau W, Clapp WL and Harvey JW. B-cell lymphoma with Mott cell differentiation in two young adult dogs. *Vet Clin Pathol.* 2009; **38**(1): 113-20.

- 245 2. Raymond JT and Garner MM. Spontaneous tumours in captive African hedgehogs
246 (*Atelerix albiventris*): a retrospective study. *J Comp Pathol*. 2001; **124**(2-3): 128-33.
- 247 3. Raymond JT, Clarke KA and Schafer KA. Intestinal lymphosarcoma in captive
248 African hedgehogs. *J Wildl Dis*. 1998; **34**(4): 801-6.
- 249 4. Raskin R. Lymphoid system. In: *Canine and feline cytology : a color atlas and*
250 *interpretation guide*, 2nd edn., St. Louis, Mo., Saunders/Elsevier, 2010: 77-122.
- 251 5. Cazzini P, Watson VE and Brown HM. The many faces of Mott cells. *Vet Clin*
252 *Pathol*. 2013; **42**(2): 125-6.
- 253 6. Ioachim HL and Medeiros LJ. Cytochemistry. In: *Ioachim's lymph node pathology*,
254 edn., Lippincott Williams & Wilkins, 2009: 35-7.
- 255 7. Kanehara T, Matsui N, Murakami M, Maruo K, Mori T, Hirata A, Yanai T and Sakai
256 H. B-cell lymphoma with Mott cell differentiation in a cat. *Vet Clin Pathol*. 2016; **45**(2):
257 356-60.
- 258 8. D'Ablaing G, Rogers ER, Parker JW and Lukes RJ. Laboratory suggestion: a
259 simplified and modified methyl green pyronin Stain. *Am J Clin Pathol*. 1970; **54**(4): 667-9.
- 260 9. Nutt SL, Heavey B, Rolink AG and Busslinger M. Commitment to the B-lymphoid
261 lineage depends on the transcription factor Pax5. *Nature*. 1999; **401**(6753): 556-62.
- 262 10. Nera KP and Lassila O. Pax5 - a critical inhibitor of plasma cell fate. *Scand J*
263 *Immunol*. 2006; **64**(3): 190-9.
- 264 11. Kodama A, Sakai H, Kobayashi K, Mori T, Maruo K, Kudo T, Yanai T and Masegi
265 T. B-cell intestinal lymphoma with Mott cell differentiation in a 1-year-old miniature
266 Dachshund. *Vet Clin Pathol*. 2008; **37**(4): 409-15.
- 267 12. Lin P, Mahdavy M, Zhan F, Zhang H-Z, Katz RL and Shaughnessy JD. Expression
268 of PAX5 in CD20-positive multiple myeloma assessed by immunohistochemistry and
269 oligonucleotide microarray. *Modern pathology*. 2004; **17**(10): 1217.

- 270 13. Falini B, Fizzotti M, Pucciarini A, Bigerna B, Marafioti T, Gambacorta M, Pacini R,
271 Alunni C, Natali-Tanci L, Ugolini B, Sebastiani C, Cattoretti G, Pileri S, Dalla-Favera R and
272 Stein H. A monoclonal antibody (MUM1p) detects expression of the MUM1/IRF4 protein
273 in a subset of germinal center B cells, plasma cells, and activated T cells. *Blood*. 2000; **95**(6):
274 2084-92.
- 275 14. Snyman HN, Fromstein JM and Vince AR. A rare variant of multicentric large B-
276 cell lymphoma with plasmacytoid and mott cell differentiation in a dog. *J Comp Pathol*.
277 2013; **148**(4): 329-34.
- 278 15. Seelig DM, Perry JA, Zaks K, Avery AC and Avery PR. Monoclonal
279 immunoglobulin protein production in two dogs with secretory B-cell lymphoma with mott
280 cell differentiation. *J Am Vet Med Assoc*. 2011; **239**(11): 1477-82.
- 281 16. Gupta A, Gumber S, Schnellbacher R, Bauer RW and Gaunt SD. Malignant B-cell
282 lymphoma with Mott cell differentiation in a ferret (*Mustela putorius furo*). *J Vet Diagn*
283 *Invest*. 2010; **22**(3): 469-73.
- 284