



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Does free schooling affect pathways from adverse childhood experiences via mental health distress to HIV risk among adolescent girls in South Africa**

A longitudinal moderated pathway model

**Citation for published version:**

Meinck, F, Orkin, FM & Cluver, L 2019, 'Does free schooling affect pathways from adverse childhood experiences via mental health distress to HIV risk among adolescent girls in South Africa: A longitudinal moderated pathway model', *Journal of the International AIDS Society*, vol. 22, no. 3, e25262, pp. 1-9. <https://doi.org/10.1002/jia2.25262>

**Digital Object Identifier (DOI):**

[10.1002/jia2.25262](https://doi.org/10.1002/jia2.25262)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Journal of the International AIDS Society

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



RESEARCH ARTICLE

# Does free schooling affect pathways from adverse childhood experiences via mental health distress to HIV risk among adolescent girls in South Africa: a longitudinal moderated pathway model

Franziska Meinck<sup>1,2§</sup> , FM Orkin<sup>3</sup> and Lucie Cluver<sup>1,4</sup> 

§Corresponding author: Franziska Meinck, Barnett House, 32 Wellington Square, Oxford OX1 2ER. Tel: +44 (0) 1865 270327. ([franziska.meinck@spi.ox.ac.uk](mailto:franziska.meinck@spi.ox.ac.uk))

## Abstract

**Introduction:** Adolescent girls are at high risk of HIV infection in sub-Saharan Africa. Mental health distress, driven by adverse childhood experiences (ACEs) such as abuse, poverty and family HIV, may be an important driver of HIV risk behaviour among adolescent girls, while education may mitigate these risks. This study aimed to develop an empirically based theoretical model between ACEs, mental health distress and HIV risk behaviour among adolescent girls in South Africa and to investigate the potential moderating effects of free schooling provision.

**Methods:** Self-report questionnaires using validated scales were completed by adolescent girls aged 12 to 17 at baseline in two provinces in South Africa in 2011, with a 99% one-year follow-up in 2012 (n = 1498). Sampling included every household in randomly selected census enumeration areas of four deprived health districts. Confirmatory factor analysis was employed to identify measurement models and a structural equation model was developed to test pathways of risk and protection.

**Results:** Internalizing and externalizing mental health distress fully mediated the positive relationship between ACEs at baseline and HIV risk behaviour at follow-up among adolescent girls. Internalizing mental health distress was associated with increased sexual risk at follow-up via higher externalizing problems. Free schooling provision at baseline and follow-up eliminated the pathway from internalizing to externalizing mental health distress by moderating the pathway between ACEs and internalizing mental health distress. It also weakened the pathway from externalizing mental health distress to HIV risk behaviour at follow-up through a direct negative effect on externalizing mental health distress.

**Conclusions:** Reducing ACEs and adolescent mental health distress is essential for reducing HIV risk behaviour among girls in South Africa. Free schooling provision may be an important tool for reducing these problems and mitigating negative pathways to HIV risk among vulnerable adolescent girls.

**Keywords:** sexual risk behaviour; adolescent girls; mental health; adverse childhood experiences; social protection

Additional Supporting Information may be found online in the Supporting information tab for this article.

Received 28 July 2018; Accepted 20 February 2019

Copyright © 2019 The Authors. *Journal of the International AIDS Society* published by John Wiley & Sons Ltd on behalf of the International AIDS Society.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

## 1 | INTRODUCTION

Human immunodeficiency virus (HIV) remains a major public health concern in sub-Saharan Africa. Despite impressive global achievements in access to treatment and declines in overall AIDS-related mortality and morbidity, AIDS remains the leading cause of death among adolescents in Africa [1]. New infections are unequally distributed among genders, with young women 44% more affected than young men [2].

There is increasing evidence that adolescent HIV risk behaviours are associated with social, psychological and economic vulnerability. Previous studies in the US showed

clear cumulative associations between adverse childhood experiences (ACEs) such as physical, emotional and sexual abuse, exposure to domestic violence, or parental absence and increased risk for HIV risk behaviour among women [3]. A recent global meta-analysis found a combined odds ratio of 5.92 for exposure to four or more ACEs and sexually transmitted diseases in adulthood [4]. In sub-Saharan Africa, factors associated with increased HIV risk behaviour include ACEs such as being AIDS-affected, poverty, abuse and psychological distress [5,6]. However, actual pathways between ACEs and HIV risk behaviours are not well understood.

A potential pathway by which ACEs may increase HIV risk behaviours is through mental health distress. It is well established that exposure to ACEs is associated with an increased risk for depression, suicide ideation, anxiety, delinquency and behaviour problems [4], with internalizing mental health distress preceding externalizing distress [7]. Associations between poor mental health and HIV risk behaviour have been shown in multiple studies in sub-Saharan Africa [8–10], however, the role of mental health in pathways from ACEs to HIV risk behaviours is under-examined.

In recent years, there has been an increasing interest in the association between education and HIV risk behaviour as a potential mechanism for intervention, particularly among adolescent girls. For children in low-resource settings and affected by HIV, paying school fees can be a substantial strain on households, causing school dropout and uncertainty regarding education. Qualitative work identifies that this is a source of distress for children [11]. Recent research evidence suggests that school non-attendance is linked to higher HIV risk behaviour in multiple contexts [12,13] while school attendance and increases in education levels have been shown to significantly reduce HIV acquisition and prevalence [14–17]. Recent randomized controlled trials (RCTs) show that free-schooling social protection, including school feeding, school uniform and tuition fees, decreases rates of school-dropout and HIV risk behaviour among orphaned children [18–20]. Furthermore, experimental and quasi-experimental studies have shown that “cash,” “care” and “classroom” (free schooling) social protection interventions in combination or by themselves can be effective in reducing HIV risk among adolescent girls [21–23]. A previous study has shown lower HIV risk behaviour and suicidality among adolescents with high school connectedness [24] but thus far no study examines whether classroom interventions reduce adolescent mental health distress. Questions also remain about whether these social protection interventions, and in particular free schooling, work by affecting the pathways to HIV risk behaviour among adolescent girls in areas with high HIV prevalence. An understanding of such mechanisms is important to develop effective interventions. This study presents a unique opportunity to examine the hypothesized beneficial effects of the national free schooling roll-out in South Africa on pathways to HIV risk behaviour among adolescent girls in HIV endemic settings.

The current paper therefore has two aims: (1) To develop a theoretical pathway model by investigating hypothesized associations from ACEs to HIV-risk behaviours via internalizing and externalizing mental health distress in adolescent girls in South Africa (Figure 1). (2) To further develop intervention theory by investigating whether the established pathways between ACEs, mental health distress and HIV risk behaviours obtain when moderated by classroom social protection in the form of free schooling.

## 2 | METHODS

### 2.1 | Participants and procedure

Adolescents ( $n = 3516$ ) aged 10 to 17 (56.7% female) were recruited from urban and rural areas in two provinces in South Africa and interviewed at baseline (2010 to 2011) and follow-up (2011 to 2012). Retention rate at one-year follow-up was 96.8%,


refusal rate at baseline was <2.5%. Adolescents were recruited through door-to-door sampling in randomly selected census enumeration areas in four health districts with antenatal HIV-prevalence >30%. One randomly selected adolescent was interviewed per household, for 70 minutes, by interviewers trained in working with vulnerable youth. No exclusion criteria were applied unless the adolescent was deemed incapable of giving consent or understanding the questionnaire. The sample thus included children with chronic illnesses, physical disabilities and severe mental health problems. Informed consent was sought from both the primary caregiver and the adolescent. Consent forms and questionnaires were translated into Zulu, Xhosa, Sotho, Sepedi, Swati and Tsonga and checked with back translation.

Ethical clearance was provided by the Universities of Oxford (SSD/CUREC2/09-52), Cape Town (389/209) and KwaZulu-Natal (HSS/0254/09) and by the provincial government departments of Health and Education. Participants were interviewed in private spaces (i.e. gardens) to ensure privacy and confidentiality. Confidentiality was maintained throughout the study except where participants requested help or were at risk of significant harm. Then referrals were made to health or child protective services with follow-up support. There were no monetary incentives, but all participants received a certificate and refreshments.

### 2.2 | Measures

The *adverse childhood experiences latent construct* was measured as follows: Lifetime *child abuse* was measured using five physical and emotional abuse items from the UNICEF Scales for National-Level Monitoring of Orphans and Other Vulnerable Children [25] and two sexual abuse items designed by local social workers. Items were summed to create a total child abuse score ( $\alpha = 0.66$ ). *Poverty* was measured using one item on the number of days of food insufficiency in the household in the past week. *Family AIDS* was measured using Verbal Autopsy methods validated in previous studies of adult mortality in South Africa [26]. Determination of caregiver HIV/AIDS-illness required three or more AIDS-defining illnesses (i.e. tuberculosis, shingles or Kaposi's sarcoma) or reported HIV positive status. *Witnessing domestic violence* was measured using one item querying the number of days where adults were hitting each other in the household in the past week. The *internalizing mental health latent construct* was measured as follows: *Depression* in the past two weeks was measured using the Children's Depression Inventory which has been previously used in South Africa and has acceptable internal consistency in this sample ( $\alpha = 0.73$ ). *Anxiety* was measured using the Revised Children's Manifest Anxiety Scale [27] which has been validated in South Africa [28] and showed good internal consistency in this sample ( $\alpha = 0.84$ ). *Suicidal ideation* during the past month was measured using the Mini International Psychiatric Interview for Children and Adolescents (Mini-Kid) [29] which has been previously used in South Africa and showed acceptable internal consistency in this sample ( $\alpha = 0.73$ ). The *externalizing mental health latent construct* was measured using the following: *Alcohol and drug use* in the past month was measured using two items from the National Survey of HIV and Risk Behaviors amongst Young South Africans [30]. *Behaviour problems* in the past six months were measured using the delinquency subscale of the Child Behaviour Checklist ( $\alpha = 0.58$ ) [31]. *Peer relationship problems* in

## What are the pathways from adverse childhood experiences to HIV-risk behaviours in adolescent girls in South Africa – the hypothesised model



**Figure 1.** Hypothesized path model from adverse childhood experiences to HIV risk behaviour among adolescent girls.

the past six months were measured using the Peer Problems subscale of the Strengths and Difficulties Questionnaire ( $\alpha = 0.60$ ) [32]. All have been previously used in South Africa. The *HIV risk behaviour latent construct* was measured using three binary items from the National Survey of HIV and Sexual Behavior among Young South Africans [30] measuring past year *infrequent condom use*, *lifetime sex whilst drunk or on drugs* and *multiple partners* (>2 in the past year). *Free schooling social protection* was measured using three items on whether or not children went to a *no-fees school*, received a *free school meal* and *free text books* and defined as receiving all three provisions at baseline and follow-up versus not receiving all three at baseline and/or follow-up. *Pre-selected covariates urban/rural location* and *age* were also collected.

### 2.3 | Analysis

Analyses were conducted in four stages with Stata 14.2 and MPlus 8 using the longitudinal sample of adolescent girls ( $n = 1919$ ). Few adolescents younger than 12 years reported being sexually active ( $n = 9$ ) and thus analyses were limited to the sample of adolescent girls aged 12 to 17 years at baseline (retention 99.3%;  $n = 1498$  at follow-up). There were no differences between those retained and lost to follow-up at baseline. First, descriptive statistics for all outcomes, potential mediators, school social protection variables and covariates were calculated (Table 1). Second, measurement models were assessed for all latent constructs, which comprised sum scores, dichotomous variables or both. Third, latent constructs were included in a model with hypothesized pathways from ACEs via adolescent mental health to HIV-risk behaviour, controlling for baseline HIV risk behaviour, age and urban/rural location. Non-significant pathways were dropped and small modifications made to improve model fit. These modifications included re-specification of covariance between error terms [33]. Finally, hypothesized moderation of pathways through the free schooling variable was tested in three sequential models, each testing the moderation on a separate pathway of the model [34]. Analyses were conducted in MPlus 8, using weighted least squares and variance adjusted estimation due to categorical data of the outcome construct indicators. As

some variables were non-normally distributed, all parameters were also estimated using the bootstrapping procedure with 1000 bootstrapped samples. Model fit was assessed using  $\chi^2$ /d.f., comparative fit index (CFI), root mean square error of approximation (RMSEA), standard root mean residual (SRMR) and weighted root mean square residual (WRMR). By convention, the maximum acceptable value for  $\chi^2$ /d.f. is 5. For CFI, a value >0.95 indicates good fit and >0.90 adequate fit. For RMSEA and SRMR a value of 0.05 indicates good model fit, and for WRMR a value close to 1 indicates good model fit. Bayesian Information Criterion (BIC) and Akaike Information Criterion (AIC) were applied for testing the model fit improvement of moderation against the specified baseline model as other model fit criteria were not available for interactions of latent variables in MPlus. For AIC and BIC smaller values indicate better model fit [35]. Missing data were less than 1% on all variables and thus no imputation was conducted.

## 3 | RESULTS

At baseline adolescent girls had a mean age of 14.3 years (range 12 to 17). 34.3% had a family member ill with AIDS. They reported a mean of 1.9 days without sufficient food in the household and 2.9 abusive events in their lifetime. Mean scores for depression, anxiety, suicidal ideation, peer problems, alcohol and drug use and delinquency can be found in Table 1. At follow-up 11.7% reported inconsistent condom use, 2.4% reported having sex drunk or on drugs and 8.0% had multiple sexual partners over the past year. 71.0% of participants received free schooling provision (Table 1). Sample proportions of adolescent girls' risk exposure by HIV-risk category can be found in Table S1. Proportions and means for the free schooling provision by different risk exposures can be found in Table S2 and show no systematic differences.

### 3.1 | Measurement models

Confirmatory factor analysis was exploited to examine hypothesized latent constructs for the predictors, mediators and outcome. A simultaneous confirmatory factor analysis was carried

**Table 1. Socio-demographic characteristics at baseline and follow-up among vulnerable South African girls (n = 1498)**

	Baseline % (n)	Follow-up % (n)
Mean age	14.3 (SE 0.043) SD 1.66	15.54 (SE 0.05) SD 1.78
Rural Area	52.1 (781)	52.3 (784)
AIDS-ill caregiver	34.3 (514)	19.3 (289)
Poverty (days without food/week, Scale 0 to 7)	0.94 (SE 0.04) SD 1.59	0.98 (SE 0.04) SD 1.63
Any abuse (lifetime)	54.2 (812)	65.4 (979)
Past week domestic violence (past week)	6.6 (99)	5.3 (79)
Anxiety (Scale 0 to 14)	5.02 (SE 0.09) SD 3.55	3.99 (SE 0.09) SD 3.33
Suicidal ideation (past month, Scale 0 to 5)	0.49 (SE 0.03) SD 1.09	0.49 (SE 0.03) SD 1.11
Depression (past two weeks, Scale 0 to 16)	2.16 (SE 0.07) SD 2.72	1.50 (SE 0.06) SD 2.18
Delinquency (past six months, Scale 0 to 14)	1.86 (SE 0.05) SD 2.12	2.03 (SE 0.05) SD 2.13
Alcohol and drug use (past month)	6.8 (102)	25.2 (377)
Peer problems (past six months, Scale 0 to 18)	4.17 (SE 0.09) SD 3.30	4.02 (SE 0.08) SD 3.21
Inconsistent condom use (past year)	8.5 (127)	11.7 (176)
Multiple sex partners (>2) (past year)	4.3 (64)	8.0 (120)
Sex drunk or on drugs (ever)	1.4 (21)	2.4 (36)
Free schooling (receipt at both time points)	71.0 (1063)	

**Table 2. Mediation model from ACEs to HIV risk behaviour among adolescent girls (n = 1498)**

Predictor	Adverse experiences	Internalizing	Externalizing	Sexual Risk
Adverse experiences	–	0.689***	0.251*	ns
Internalizing	–	–	0.196*	ns
Externalizing	–	–	–	0.370***
Age	0.100*	0.128***	0.075*	0.361***
Baseline sexual risk	0.141***	0.068*	0.207***	0.291***
Urban location	ns	ns	0.064*	ns
Correlated item residuals	Peer problems with depression: 0.245***; Alcohol with delinquency: –0.429***			
Goodness of fit	$\chi^2$ (d.f.): 2552.965 (73), CFI 0.940; TFI 0.917, RMSEA 0.037 (0.031 to 0.042),			
$r^2$	0.039	0.544	0.288	0.552

– Refers to non-applicable pathways, ns refers to pathways that were removed in the mediation model. ACEs, adverse childhood experiences; CFI, comparative fit index; RMSEA, root mean square error of approximation. All effects reported are standardized, \*\*\* $p < 0.001$ , \* $p < 0.05$ .

out to examine model fit of the individual constructs (Table S3). The ACEs latent construct was identified by the total score for the poverty measure, the total score for child abuse victimization, and familial AIDS at baseline, which all loaded >0.4. Domestic Violence was removed from the construct as its factor loading was <0.2. The indicators for the latent construct of internalizing mental health distress were the respective averages of depression, anxiety and suicide ideation sum scores at baseline and follow-up [36]. They loaded onto the construct >0.4. The latent construct for externalizing mental health distress was identified by the respective averages of child delinquency, peer problems, and drug and alcohol abuse sum scores at baseline and follow-up [36]. They loaded onto the construct >0.3. The HIV risk behaviour latent construct was identified by the dichotomous variables of frequent condom use, sex drunk or on drugs, and having multiple sexual partners at follow-up, with all indicators loading onto the construct >0.6. All indicator variables within each of the four constructs had equal weight

(Table S3). Model fit for the simultaneous CFA was good on all criteria except CFI, which was adequate ( $\chi^2 = 223.23$ , d.f. = 48,  $p < 0.001$ ,  $\chi^2$ /d.f. = 4.65, CFI 0.916, WRMR 1.428, RMSEA 0.049 (0.043 to 0.056)).

### 3.2 | Path model

The four latent constructs were therefore included in the hypothesized model with the following pathways (Figure 1): ACEs associated with internalizing and externalizing mental health distress and HIV risk behaviour, internalizing mental health distress associated with externalizing mental health distress and each associated with HIV risk behaviour (Table 2). The hypothesized model had poor model fit ( $\chi^2 = 342.931$ , d.f. = 72,  $p < 0.001$ ,  $\chi^2$ /d.f. = 4.76, CFI 0.889, RMSEA 0.050, WRMR 1.561). All non-significant pathways were then removed from the model one pathway at a time: non-significant covariates, the direct effect of ACEs on HIV risk behaviour, and the direct effect of internalizing mental health

distress on HIV risk behaviour. This resulted in the following adequate model fit: ( $\chi^2 = 352.117$ , d.f. = 77,  $p < 0.001$ ,  $\chi^2$ /d.f. = 4.57, CFI 0.900, RMSEA 0.049, WRMR 1.622). Further modifications were carried out based on modification indices as specified in Table S4. The final model shows direct paths from ACEs to internalizing ( $\beta = 0.69$ ,  $p < 0.001$ ) and externalizing mental health distress ( $\beta = 0.25$ ,  $p = 0.005$ ), from internalizing to externalizing mental health distress ( $\beta = 0.20$ ,  $p = 0.028$ ) and from externalizing mental health distress to HIV-risk behaviour ( $\beta = 0.37$ ,  $p < 0.001$ ) (Figure 2).

Three indirect effects were established: (1) from ACEs via internalizing to externalizing mental health distress ( $\beta = 0.14$ ,  $p = 0.023$ ); (2) from adverse experiences via externalizing mental health distress to HIV risk behaviour ( $\beta = 0.09$ ,  $p = 0.011$ ); and (3) from internalizing via externalizing mental health distress to HIV risk behaviour ( $\beta = 0.07$ ,  $p = 0.046$ , Figure 2). The final model accounted for 55% of the variance in HIV risk behaviour. Model fit was good on all criteria set out above except CFI, which was adequate: ( $\chi^2 = 219.284$ , d.f. = 73,  $p < 0.001$ ,  $\chi^2$ /d.f. = 3.00, CFI 0.940, RMSEA 0.032, WRMR 1.279). AIC (58654.4) and BIC (58888.12) were also obtained for this model to allow comparison with the final model.

Direct effects of free schooling provision on the latent constructs, as well as interactions between free schooling social protection and the latent constructs, were examined one at a time because of the computationally heavy demands of categorical latent constructs [37, p. 471]. AIC and BIC were examined in each case and compared to the mediation model in Figure 2. In the first instance, no significant direct effect of free schooling provision on internalizing mental health distress was found (Table S5, Model 1). There was a significant interaction between free schooling provision and ACEs ( $\beta = -0.11$ ,  $p = 0.008$ ) moderating the pathway from ACEs to internalizing mental health distress. A direct effect of free schooling provision was found on externalizing mental health distress, but no interaction effect between ACEs and free schooling provision was found to moderate the pathway between ACEs and externalizing mental health distress. Furthermore, the introduction of free schooling provision renders the pathway from internalizing to externalizing mental health distress insignificant. In the second instance, a

significant direct effect of free schooling provision could be observed on externalizing mental health distress ( $\beta = -0.10$ ,  $p = 0.008$ ), but no interaction effect between internalizing behaviour and free schooling social protection was found (see Table S5, Model 2). In the third instance, neither a direct effect of free schooling social protection on HIV risk behaviour nor an interaction effect between free schooling social protection and externalizing mental health distress was found (see Table S5, Model 3). A final model was then run which included the significant interaction effect and the direct effect of free schooling provision as well as the significant mediation pathway (Figure 3, Table 3).

## 4 | DISCUSSION

Reducing HIV risk among adolescent girls is a huge challenge, particularly in HIV-endemic countries. The empirical model tested in this study shows a set of pathways of vulnerability leading to increased HIV risk behaviour. It is noteworthy that no significant direct pathway was found between ACEs and HIV risk behaviour. The relationship between ACEs and HIV risk behaviour was fully mediated by internalizing and externalizing mental health distress. Moreover, there was no significant pathway from internalizing mental health distress to HIV risk behaviour. This supports the conception of internalizing symptoms preceding externalizing symptoms [7]. Earlier studies, which tested correlations and not pathways found strong associations between ACEs and HIV risk behaviour as well as between internalizing and HIV risk behaviour [38–40]. However, these did not test theoretical pathway models for HIV risk behaviour and therefore had not tested for mediation. This study demonstrates pathways through mental health distress and thereby provides new evidence for opportunities to interrupt cycles of increased risk.

This study also provided evidence that classroom social protection in the form of free schooling for adolescents (i.e. free schooling, school-books and school meals) is associated with lower externalizing mental health distress and modifies the risk pathway from ACEs to internalizing mental health distress. This





Figure 2. Empirical pathway model from adverse childhood experiences to HIV risk behaviour among adolescent girls.



**Figure 3. Final model with moderation: direct effect and moderating effect of free schooling social protection on pathways from adverse childhood experiences to HIV-risk behaviour among adolescent girls.**

**Table 3. Final model including all significant mediation pathways and the significant direct and interaction effects of free schooling social protection on HIV risk among South African adolescent girls (N = 1498)**

	Final model			
	ACEs	Internalizing	Externalizing	Sexual risk
ACEs	–	0.808***	0.344***	ns
Internalizing	–	–	ns	ns
Externalizing	–	–	–	0.306
Free schooling social protection	–	ns	–0.092*	ns
ACE × free schooling	–	–0.099*	–	–
Internalizing × free schooling	–	–	ns	–
Externalizing × free schooling	–	–	ns	–
Sexual risk × free schooling	–	–	–	ns
Age	0.100*	0.125***	0.086**	0.448***
Baseline sexual risk	0.145**	ns	0.184***	0.189***
Rural location	ns	ns	0.094**	ns
Measurement models				
AIDS-ill caregiver	0.295***			
Child abuse	0.606***			
Poverty	0.349***			
Suicide ideation		0.635***		
Anxiety		0.743***		
Depression		0.664***		
Peer problems			0.191***	
Delinquency			0.748***	
Drug and alcohol use			0.580***	
Inconsistent condom use				0.802***
Multiple partners				0.814***
Sex drunk or on drugs				0.721***
Goodness of fit	BIC: 49059.763; AIC: 48804.793			

– Refers to non-applicable pathways, ns refers to non-significant pathways that were removed in the mediation model. AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion.

All effects reported are standardized, \*\*\* $p < 0.001$ , \*\* $p < 0.005$ , \* $p < 0.05$ .

could operate through a number of mechanisms which need to be further explored: for instance, adolescent girls who receive free schooling may experience lower mental health distress

because they do not have the additional worry of being able to pay the school fees [11]. In addition, by receiving free schooling social protection, adolescent girls receive one hot meal a day

which may prevent hunger and thus reduce risk for mental health distress [41]. Further school social protection may increase school attendance [42], giving girls the chance to make friendships and build support networks. This study suggests that free education is an important tool to promote mental health among adolescent girls in South Africa, and thereby mitigate pathways to HIV risk behaviour. However, HIV risk behaviour and mental health distress vary across age group and thus may require age-sensitive interventions. Therefore, further research could valuably investigate which age groups benefit most from classroom social protection for the prevention of HIV risk behaviours. Future qualitative research should also examine why free schooling is directly associated with reducing externalizing mental health distress and also modifies the risk pathway to internalizing mental health distress.

This study has notable limitations. First, data are available only from two time points, and although mean factor scores of baseline and follow-up mental health data were used for the mediator variables to establish temporal sequence, causality cannot be asserted. While reverse causality might be unlikely for many of the pathways, for some pathways – particularly between internalizing and externalizing mental health distress – bi-directionality is possible. However, the cautious attribution of causal order with temporal precedence, and based on plausible theoretical grounds, is well established in structural equation modelling [33]. Future research using rigorous randomized designs is needed to examine the effectiveness of free-schooling social protection on HIV risk behaviour among adolescent girls. Such studies can complement the existing evidence on orphaned children only [19] and of cash social protection on HIV risk behaviour [14] and school attendance [43].

Second, data were collected in 2011 and 2012 and results and potential intervention requirements need to be interpreted with this in mind. Non-fee schools and the school feeding programme were started in 2006/2007 with schools in lowest income areas and have since been extended across the country. They reach nine million children in schools in deprived areas and thus effects of these programmes may be different now that they have been scaled up [44]. However, testing intervention effects at the start of rollout when an intervention is not universally available does allow determination of differences between those that are in receipt and not in receipt of it and thus this can be considered a real strength of the presented data. Third, receipt of social protection was only measured at baseline and follow-up and it is therefore impossible to distinguish between children who have been in long-term receipt of the intervention or those who are recent recipients. Fourth, all items were measured using adolescent self-report only. As child abuse and HIV risk behaviours are highly stigmatized, some underreporting may have occurred. However, other types of respondents, that is, caregivers have been shown to be even more unreliable [45] and thus adolescent self-report is preferred over caregiver report. Fifth, HIV risk taking behaviours are low in this sample, possibly due to the young age of the participants (mean age 15.5 years at follow-up), and this will have implications for programming. Sixth, this study did not use randomization for the intervention and thus some selection bias cannot be ruled out as students receiving the interventions were in the most highly deprived areas of South Africa and thus

may have experienced higher levels of poverty, mental health distress and adversity. Seventh, this study focused on pathways which increase HIV risk behaviour between latent constructs, as moderated by free schooling provision. Future studies should examine pathways of additional resilience, protective factors and the contributions of each of the exposures towards HIV risk behaviour. Eighth, this study focused on adolescent girls, as the group within the region at highest risk of HIV infection. Future research should investigate pathways to HIV-risk behaviours among boys and examine the effects of free schooling and other potential moderators on these pathways. Finally, the study only sampled adolescents in highly deprived areas of South Africa and is thus not representative of the overall population. However, the study benefited from in-sample variation that included participants from six different language groups, and is therefore likely to be indicative for vulnerable adolescent girls in deprived areas in South Africa.

This study also has multiple strengths. It is the first longitudinal study to test multiple pathways associated with increased HIV risk behaviour among girls in South Africa. It moves the analysis beyond direct associations by indicating the moderating effect of a feasible intervention, free schooling. The sample size was large, participation high, attrition at follow-up low and model fit good.

## 5 | CONCLUSIONS

Free access to education, school feeding and books at primary and secondary school level may be an important part of an HIV prevention package for adolescent girls in South Africa. Previous research has focused on single provisions such as grants to alleviate poverty, access to education to reduce risk behaviour, or parenting to reduce abuse and improve adolescent mental health [14,22,46]. In recent years, research focus has shifted to combination interventions of “cash and care”, in the form of grants combined with parenting. Cash and care provisions together have been shown to reduce transactional and age disparate sex [47]. This study shows that “classroom” intervention, in the form of free schooling, may be an additional means to reduce internalizing and externalizing mental health distress and thereby interrupt pathways to HIV-risk behaviours among adolescent girls in South Africa.

### AUTHORS' AFFILIATIONS

<sup>1</sup>Centre for Evidence-Based Intervention, Department of Social Policy and Intervention, University of Oxford, Oxford, United Kingdom; <sup>2</sup>OPTENTIA, Faculty of Health Sciences, North-West University, Vanderbijlpark, South Africa; <sup>3</sup>MRC Developmental Pathways for Health Research Unit, School of Clinical Medicine, University of the Witwatersrand, Johannesburg, South Africa; <sup>4</sup>Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa

### COMPETING INTERESTS

The authors declare that they have no competing interests.

### AUTHORS' CONTRIBUTIONS

FM and MO conceptualized the analysis and the paper and also led the analyses. LC, FM and MO conceptualized the overall study and oversaw data



collection. LC contributed to the analyses and interpretation of the findings. All authors reviewed and approved the final version.

## ACKNOWLEDGEMENTS

The authors thank the adolescents who participated in the study, their families, all the fieldworkers working tirelessly to interview as many adolescents as possible, the Rural AIDS Development Action Research programme (RADAR) at the University of the Witwatersrand, Cape Town Child Welfare, Jenny Doubt & Sarah Hoeksma. We thank Jennifer Rabedeau for editing and proof reading.

## FUNDING

This study was funded by the Economic and Social Research Council (UK) and the National Research Foundation (RES-062-23-2068), the National Department of Social Development, the Claude Leon Foundation, the Nuffield Foundation (OPD/31598), the Health Economics and HIV/AIDS Research Division at the University of KwaZulu-Natal (R14304/AA002), the John Fell Fund (103/757), the University of Oxford Impact Acceleration Account (1602-KEA-189, 1311-KEA-004, 1069-GCRF-227) and the Leverhulme Trust (PLP-2014-095). FM is funded by an ESRC Future Research Leader Award (ES/N01747/1). LC is funded by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013, grant agreement 313421).

## REFERENCES

1. WHO. Global burden of disease 2015. Geneva: WHO; 2016.
2. UNAIDS. AIDSinfo [Internet]. 2017[cited 2018 Jan 19]. Available from: [aidsinfo.unaids.org](http://aidsinfo.unaids.org)
3. Hillis SD, Anda RF, Felitti VJ, Marchbanks PA. Adverse childhood experiences and sexual risk behaviors in women: a retrospective cohort study. *Fam Plann Perspect*. 2001;33(5):206–11.
4. Hughes K, Bellis MA, Hardcastle KA, Sethi D, Butchart A, Mikton C, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health*. 2017;2(8):e356–66.
5. Gibbs A, Dunkle K, Washington L, Willan S, Shai N, Jewkes R. Childhood traumas as a risk factor for HIV-risk behaviours amongst young women and men living in urban informal settlements in South Africa: a cross-sectional study. *PLoS One*. 2018;13(4):e0195369.
6. Operario D, Underhill K, Chuong C, Cluver L. HIV infection and sexual risk behaviour among youth who have experienced orphanhood: systematic review and meta-analysis. *J Int AIDS Soc*. 2011;14:25.
7. Ritakallio M, Koivisto A-M, von der Pahlen B, Pelkonen M, Marttunen M, Kaltiala-Heino R. Continuity, comorbidity and longitudinal associations between depression and antisocial behaviour in middle adolescence: a 2-year prospective follow-up study. *J Adolesc*. 2008;31(3):355–70.
8. Cluver L, Orkin F, Meinck F, Boyes M, Sherr L. Structural drivers and social protection: mechanisms of HIV risk and HIV prevention for South African adolescents. *J Int AIDS Soc*. 2016;19(1):20646.
9. Lundberg P, Rukundo G, Ashaba S, Thorson A, Allebeck P, Östergren P-O, et al. Poor mental health and sexual risk behaviours in Uganda: a cross-sectional population-based study. *BMC Public Health*. 2011;11(1):125.
10. De Beaudrap P, Mac-Seing M, Pasquier E. Disability and HIV: a systematic review and a meta-analysis of the risk of HIV infection among adults with disabilities in Sub-Saharan Africa. *AIDS Care*. 2014;26(12):1467–76.
11. Watkins JA, Sello OM, Cluver L, Kaplan L, Boyes M. 'At school I got myself a certificate': HIV/AIDS Orphanhood and Secondary Education: a Qualitative Study of Risk and Protective Factors. *Glob Soc Welf*. 2014;1(3):111–21.
12. Pufall EL, Eaton JW, Robertson L, Mushati P, Nyamukapa C, Gregson S. Education, substance use, and HIV risk among orphaned adolescents in Eastern Zimbabwe. *Vulnerable Child Youth Stud*. 2017;12(4):360–74.
13. Pettifor A, Levandowski BA, MacPhail C, Padian NS, Cohen MS, Rees HV. Keep them in school: the importance of education as a protective factor against HIV infection among young South African women. *Int J Epidemiol*. 2008;37(6):1266–73.
14. Pettifor A, MacPhail C, Hughes JP, Selin A, Wang J, Gómez-Olivé FX, et al. The effect of a conditional cash transfer on HIV incidence in young women in rural South Africa (HPTN 068): a phase 3, randomised controlled trial. *Lancet Glob Health*. 2016;4(12):e978–88.
15. De Neve J-W, Fink G, Subramanian SV, Moyo S, Bor J. Length of secondary schooling and risk of HIV infection in Botswana: evidence from a natural experiment. *Lancet Glob Health*. 2015;3(8):e470–7.

16. Baird S, Chirwa E, McIntosh C, Özler B. The short-term impacts of a schooling conditional cash transfer program on the sexual behavior of young women. *Health Econ*. 2010;19 Suppl:55–68.
17. Stoner MCD, Pettifor A, Edwards JK, Aiello AE, Halpern CT, Julien A, et al. The effect of school attendance and school dropout on incident HIV and HSV-2 among young women in rural South Africa enrolled in HPTN 068. *AIDS*. 2017;31(15):2127–34.
18. Hallfors D, Cho H, Rusakaniko S, Iritani B, Mapfumo J, Halpern C. Supporting adolescent orphan girls to stay in school as HIV-risk prevention: evidence from a randomized controlled trial in Zimbabwe. *Am J Public Health*. 2011;101:1082–91.
19. Cho H, Mbai I, Luseno WK, Hobbs M, Halpern C, Hallfors DD. School Support as Structural HIV Prevention for Adolescent Orphans in Western Kenya. *J Adolesc Health*. 2018;62(1):44–51.
20. Cho H, Catherine Ryberg R, Hwang K, Pearce LD, Iritani BJ. A school support intervention and educational outcomes among orphaned adolescents: results of a cluster randomized controlled trial in Kenya. *Prev Sci*. 2017;18(8):943–54.
21. Cluver L, Hodes R, Sherr L, Orkin F, Meinck F, Lim Ah Ken P, et al. Social protection: potential for improving HIV outcomes among adolescents. *J Int AIDS Soc*. 2015;18 Suppl 6:20260.
22. Baird S, Garfein RS, McIntosh CT, Ozler B. Effect of a cash transfer programme for schooling on prevalence of HIV and herpes simplex type 2 in Malawi: a cluster randomised trial. *Lancet*. 2012;379(9823):1320–9.
23. Pettifor A, MacPhail C, Nguyen N, Rosenberg M. Can money prevent the spread of HIV? A review of cash payments for HIV prevention. *AIDS Behav*. 2012;16(7):1729–38.
24. Govender K, Naicker SN, Meyer-Weitz A, Fanner J, Naidoo A, Penfold WL. Associations between perceptions of school connectedness and adolescent health risk behaviors in South African high school learners. *J Sch Health*. 2013;83(9):614–22.
25. Snider L, Dawes A. Psychosocial vulnerability and resilience measures for national-level monitoring of orphans and other vulnerable children: recommendations for revision of the UNICEF psychological indicator. Cape Town: UNICEF; 2006.
26. Lopman B, Barnabas R, Boerma T, Chawira J, Gaitskell K, Harrop T, et al. Creating and validating an algorithm to measure AIDS mortality in the adult population using verbal autopsy. *PLoS Med*. 2006;3:e312.
27. Reynolds CR, Richmond BO. What I think and feel: A revised measure of children's manifest anxiety. *J. Abnorm. Child Psychol*. 1978;6(2):271–80.
28. Boyes M, Cluver L. Performance of the revised children's manifest anxiety scale in a sample of children and adolescents from poor urban communities in Cape Town. *Eur J Psychol Assess*. 2013;29:113–20.
29. Sheehan D, Shytle D, Milo K. MINI KID: mini international neuropsychiatric interview for children and adolescents. English Version 4.0. Paris: University of South Florida, Tampa and Hospital de la Salpêtrière; 2004.
30. Reproductive Health Research Unit, Lovelife. The National Survey of HIV and sexual behaviour among young South Africans. Johannesburg: University of Witwatersrand; 2005.
31. Achenbach T. Child Behaviour Checklists (CBCL/2-3 and CBCL/4-18), Teacher Report Form (TRF) and Youth Self-Report (YSR). In: Rush J, First M, Blacker D, editors. The handbook of psychiatric measures. 1st ed. Arlington, VA: The American Psychiatric Association; 2000.
32. Goodman R. Psychometric properties of the Strengths and Difficulties Questionnaire (SDQ). *J Am Acad Child Adolesc Psychiatry*. 2001;40:1337–45.
33. Byrne B. Structural equation modeling with MPlus. New York: Routledge; 2012.
34. Muthen B, Muthen L, Asparouhov T. Regression and mediation analysis using MPlus. Los Angeles, CA: Muthen & Muthen; 2016. 519 p.
35. Hox J, Bechger T. An introduction to structural equation modeling. *Fam Sci Rev*. 2007;11:354–73.
36. Twisk J. Applied longitudinal data analysis for epidemiology – a practical guide. Cambridge: Cambridge University Press; 2003.
37. Muthen B, Muthen L. MPlus User Guide Version 7. Los Angeles: Statmodel; 2012.
38. Cluver L, Orkin M, Boyes M, Gardner F, Meinck F. Transactional sex amongst AIDS-orphaned and AIDS-affected adolescents predicted by abuse and extreme poverty. *J Acquir Immune Defic Syndr*. 2011;58:336–43.
39. Peltzer K, Pengpid S. Sexual abuse, violence AND HIV risk among adolescents in South Africa. *Gend Behav*. 2008;6:1462–78.
40. Richter L, Komárek A, Desmond C, Celentano D, Morin S, Sweat M, et al. Reported physical and sexual abuse in childhood and adult HIV risk behaviour in

three African countries: findings from project accept (HPTN-043). *AIDS Behav.* **2014**;18(2):381–9.

41. Sorsdahl K, Slopen N, Siefert K, Seedat S, Stein DJ, Williams DR. Household food insufficiency and mental health in South Africa. *J Epidemiol Community Health.* **2011**;65(5):426–31.

42. Hall K. Children's access to education. *South African Child Gauge* 2015. Cape Town: University of Cape Town; **2015**.

43. Baird S, Ferreira F, Ozler B, Woolcock M. Relative effectiveness of conditional and unconditional cash transfers for schooling outcomes in developing countries: a systematic review. *Campbell Collab Libr.* **2013**;9(8):4–124.

44. Department of Basic Education. National School Nutrition Programme (NSNP): annual report 2013/2014. Pretoria: Department of Basic Education; **2014**.

45. Johnsona R, Kotch J, Catellier D, Winsor J, Dufort V, Hunter W, et al. Adverse behavioural and emotional outcomes from child abuse and witnessed violence. *Child Maltreat.* **2002**;7(3):179–86.

46. Cluver L, Meinck F, Steinert J, Shenderovich Y, Doubt J, Lombard C, et al. Parenting for lifelong health: a pragmatic cluster randomised controlled trial of a non-commercialised parenting programme for adolescents and their families in South Africa. *BMJ Glob Health.* **2018**;3:e000539.

47. Cluver L, Orkin F, Boyes M, Sherr L. Cash plus care: social protection cumulatively mitigates HIV-risk behaviour among adolescents in South Africa. *AIDS.* **2014**;28 Suppl 3:S389–97.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

**Table S1.** Proportions of adolescent girls (n = 1498) engaging in HIV risk behaviour based on their risk exposure

**Table S2.** Proportions and mean differences of adolescent girls (n = 1498) exposed to risk based on receipt of free schooling

**Table S3.** Standardized factor loadings for adverse childhood experiences, internalizing and externalizing mental health distress and sexual risk behaviour among South African adolescent girls using simultaneous confirmatory factor analysis (n = 1498)

**Table S4.** Modification steps of the model to achieve final model fit

**Table S5.** Interaction effects of free schooling social protection on pathways from adverse childhood experiences to HIV risk behaviour among adolescent girls (n = 1498)