



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Genome-wide transposon mutagenesis identifies a role for host neuroendocrine stress hormones in regulating the expression of virulence genes in *Salmonella*

Citation for published version:

Spencer, H, Karavolos, MH, Bulmer, DM, Aldridge, P, Chhabra, SR, Winzer, K, Williams, P & Khan, CMA 2010, 'Genome-wide transposon mutagenesis identifies a role for host neuroendocrine stress hormones in regulating the expression of virulence genes in *Salmonella*', *Journal of Bacteriology*, vol. 192, no. 3, pp. 714-24. <https://doi.org/10.1128/JB.01329-09>

Digital Object Identifier (DOI):

[10.1128/JB.01329-09](https://doi.org/10.1128/JB.01329-09)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Bacteriology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Genome-Wide Transposon Mutagenesis Identifies a Role for Host Neuroendocrine Stress Hormones in Regulating the Expression of Virulence Genes in *Salmonella*[∇]

H. Spencer,¹ M. H. Karavolos,¹ D. M. Bulmer,¹ P. Aldridge,¹ S. R. Chhabra,² K. Winzer,²
P. Williams,² and C. M. A. Khan^{1*}

Institute for Cell and Molecular Biosciences, The Medical School, Newcastle University, Newcastle NE2 4HH,¹ and School of Molecular Medical Sciences, Centre for Biomolecular Sciences, University Park, University of Nottingham, Nottingham NG7 2RD,² United Kingdom

Received 8 October 2009/Accepted 10 November 2009

Bacterial sensing of environmental signals plays a key role in regulating virulence and mediating bacterium-host interactions. The sensing of the neuroendocrine stress hormones epinephrine (adrenaline) and norepinephrine (noradrenaline) plays an important role in modulating bacterial virulence. We used MudJ transposon mutagenesis to globally screen for genes regulated by neuroendocrine stress hormones in *Salmonella enterica* serovar Typhimurium. We identified eight hormone-regulated genes, including *yhaK*, *iroC*, *nrdF*, *accC*, *yedP*, STM3081, and the virulence-related genes *virK* and *mig14*. The mammalian α -adrenergic receptor antagonist phentolamine reversed the hormone-mediated effects on *yhaK*, *virK*, and *mig14* but did not affect the other genes. The β -adrenergic receptor antagonist propranolol had no activity in these assays. The *virK* and *mig14* genes are involved in antimicrobial peptide resistance, and phenotypic screens revealed that exposure to neuroendocrine hormones increased the sensitivity of *S. Typhimurium* to the antimicrobial peptide LL-37. A *virK* mutant and a *virK mig14* double mutant also displayed increased sensitivity to LL-37. In contrast to enterohemorrhagic *Escherichia coli* (EHEC), we have found no role for the two-component systems QseBC and QseEF in the adrenergic regulation of any of the identified genes. Furthermore, hormone-regulated gene expression could not be blocked by the QseC inhibitor LED209, suggesting that sensing of hormones is mediated through alternative signaling pathways in *S. Typhimurium*. This study has identified a role for host-derived neuroendocrine stress hormones in downregulating *S. Typhimurium* virulence gene expression to the benefit of the host, thus providing further insights into the field of host-pathogen communication.

Bacterial sensing of environmental signals plays a key role in regulating virulence gene expression and bacterium-host interactions. It is increasingly recognized that detection of host-derived molecules, such as the neuroendocrine stress hormones (catecholamines) epinephrine (adrenaline) and norepinephrine (noradrenaline), plays an important role in modulating bacterial virulence (29, 42).

Physical and psychological stress has been linked to increased severity and susceptibility to infection in humans and other animals (23, 42), and epinephrine/norepinephrine levels are an important factor in this. Stress triggers an increase in plasma epinephrine levels (31), and plasma levels of epinephrine and norepinephrine have been reported to increase with patients suffering from postoperative sepsis compared to patients with no complications (32). Administration of norepinephrine and epinephrine to otherwise healthy subjects increases the severity of bacterial infections, including *Clostridium perfringens* in humans and enterohemorrhagic *Escherichia coli* (EHEC) in calves (42, 63, 65). Treatment with norepinephrine also increases the virulence of *Salmonella enterica* serovar Enteritidis in chicks and

Salmonella enterica serovar Typhimurium in mice, with a substantial increase in bacterial numbers recovered from the cecum and liver in both cases (47, 65).

Norepinephrine is found in large concentrations in the gut due to release by gastrointestinal neurones; indeed up to half the norepinephrine in the body may be produced in the enteric nervous system (ENS) (3). Epinephrine, while not normally found in the gut, is present in the bloodstream and is also produced by macrophages in response to bacteria-derived lipopolysaccharide (LPS) (12, 26). *S. Typhimurium* is an enteropathogen, can also cross the epithelial barrier to cause systemic infection, and will therefore encounter both these molecules in the normal infection cycle.

Phenotypes induced by stress hormones in bacteria include increased adherence of EHEC to bovine intestinal mucosa (63), upregulation of type III secretion and Shiga toxin production in EHEC (22, 60), upregulation of type III secretion in *Vibrio parahaemolyticus* (51), increase in invasion of epithelial cells and breakdown of epithelial tight junctions by *Campylobacter jejuni* (15), affected motility and expression of iron uptake genes in *S. Typhimurium* (8, 9, 36), and modulated virulence in *Borrelia burgdorferi* (59). Epinephrine and norepinephrine can overcome the growth inhibition of many bacteria, including *Salmonella*, in serum-containing media (13, 43), due to the ability to act as a siderophore to facilitate iron uptake (13, 28, 47).

Norepinephrine and epinephrine also interact with bacterial

* Corresponding author. Mailing address: Institute for Cell and Molecular Biosciences and School of Biomedical Sciences, The Medical School, Newcastle University, Framlington Place, Newcastle upon Tyne, NE2 4HH, United Kingdom. Phone and fax: 44 191 2227066. E-mail: anjam.khan@ncl.ac.uk.

[∇] Published ahead of print on 20 November 2009.

quorum-sensing (QS) systems. QS is a process of bacterial cell-cell communication in which each cell produces small signal molecules termed "autoinducers" (AIs), which regulate gene expression when a critical threshold concentration and therefore population density have been reached. QS affects diverse processes, including motility, virulence, biofilm formation, type III secretion, and luminescence (6, 64).

The EHEC AI-3 QS system is important for motility and expression of the type III secretion system encoded by the locus of enterocyte effacement (LEE) (60). AI-3 sensing and signal transduction are mediated via the QseBC and QseEF two-component systems, respectively. Epinephrine and norepinephrine can substitute for AI-3, causing cross talk between the two signaling systems and induction of type III secretion and motility (57, 60). The sensor kinase QseC is autophosphorylated upon binding either epinephrine or norepinephrine (14), demonstrating the presence of adrenergic receptors in bacteria. These adrenergic phenotypes can also be blocked by the mammalian α - and β -adrenergic antagonists phentolamine and propranolol, although it should be noted that QseC is blocked only by the former (14, 60). This suggests the occurrence of cross talk between bacterial and mammalian cell signaling systems and the existence of multiple bacterial adrenergic sensors.

To elucidate the role of host-derived stress hormones in the physiology and pathogenicity of *S. Typhimurium*, we used *MudJ* transposon mutagenesis to screen globally for epinephrine- and norepinephrine-regulated genes in *S. Typhimurium*.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions. *Salmonella enterica* serovar Typhimurium strains, plasmids, and primers are shown in Table 1. Bacteria were grown overnight in 5 ml LB broth at 37°C and 200 rpm. A total of 25 μ l of overnight broth was used to inoculate 25 ml fresh LB, or 250 μ l to inoculate 25 ml M9 minimal media (48) supplemented with 0.3% Casamino Acids and 40 μ g/ml histidine, and the bacteria were grown in 250-ml conical flasks at 37°C and 200 rpm. LB medium is commonly used to mimic intestinal conditions (54, 62). We also used minimal media, as much previous work investigating catecholamines has been done in SAPI-serum minimal media (13, 28, 43). As supplementation with catecholamines drastically affects growth rate in this media, we used M9 to negate the possibility of growth rate indirectly affecting the phenotypes seen.

General physiological and molecular biological manipulations were performed according to standard laboratory protocols (58). Antibiotics were added at the following concentrations: kanamycin, 50 μ g/ml, and ampicillin, 100 μ g/ml.

Epinephrine/norepinephrine hydrochloride (Sigma) was added to a final concentration of 50 μ M or 500 μ M in distilled water (dH₂O). Propranolol and phentolamine (Sigma) were added to a final concentration of 500 μ M in dH₂O.

Chemical synthesis of LED209. *N*-phenyl-4-[(phenylamino)thioxomethyl]-amino-benzenesulfonamide (LED209) was synthesized in three steps essentially by the procedure described by Rasko et al. (55). Reaction of aniline with 4-acetamidobenzenesulfonyl chloride in ethanol in the presence of sodium acetate at room temperature afforded 4-acetamido-*N*-phenyl-benzenesulfonamide in 80% yield as a white solid (melting point, 207 to 209°C). The acid hydrolysis of this product with 6 M HCl in ethanol at reflux gave 4-amino-*N*-phenyl-benzenesulfonamide as a white solid in 98% yield. In the final step, the 4-amino-*N*-phenyl-benzenesulfonamide product was condensed with phenylisothiocyanate in dry tetrahydrofuran in the presence of triethylamine at reflux for 48 h. After concentration, the crude oily product was purified by flash chromatography on silica gel in acetone-hexane (1:3) to deliver LED209 as a white crystalline solid (melting point of 161 to 163°C) in 25% yield. The structure and purity of the product was unequivocally confirmed by ¹H NMR, which showed the expected resonances at δ 7.00 to 7.47 (9H, 5 \times m, ArH), 7.70 (5H, s, Ph), 10.07 (2H, br s, NHCSNH), and 10.23 (1H, br s, SO₂NH). The structure was further confirmed by electrospray mass spectrometry (ES-MS) (negative-ion mode), which displayed the correct molecular ion at an *m/z* of 382 (M-H, C₁₉H₁₇N₃O₂S₂ requires

an *m/z* of 383) and expected fragmentation ions at *m/z* values of 348 (M-H-H₂S), 289 (M-H-PhNH₂), and 247 (M-H-Ph-N=C=S). LED209 was dissolved in dimethyl sulfoxide (DMSO) at 50 mM and then diluted in dH₂O as stated.

***MudJ* mutagenesis.** *MudJ* mutagenesis was done as described previously (35). Briefly, a phage lysate of strain TPA356 was prepared according to standard laboratory protocol (58) and transduced into SL1344. Transductants were plated onto no-carbon E salts (NCE) plates (19) with 100 μ l 50 mM epinephrine overlaid on the agar and incubated at 37°C for 48 h. Resulting colonies were patched to MacConkey agar plates (Sigma), with or without an overlay of 100 μ l 50 mM epinephrine, to give a final concentration of approximately 250 μ M.

Genomic sequencing. *MudJ* gene insertions were identified using inverse PCR as previously described (2). Briefly, genomic DNA was isolated using a DNeasy blood and tissue kit (Qiagen), digested with *TaqI* and self ligated with T4 DNA ligase. One μ l was used as a template for PCR using Phusion polymerase (New England Biolabs) and primers MudOut and MudTaq (Table 1). PCR products were purified using PCR cleanup columns (Invitrogen) and ligated into the *SmaI* site of pPCR-Script (Stratagene). Plasmid DNA was purified using an Invitrogen miniprep kit and sequenced with primer Universal-PCR. Sequencing was performed at Pinnacle, Newcastle University, Newcastle upon Tyne, United Kingdom. Sequence homologies were identified using BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

β -Galactosidase assays. Strains were grown in 25-ml volumes in M9 minimal medium for 2.5 h or LB for 3.5 h (for *yhaK::MudJ*), 50 μ M epinephrine/norepinephrine was added, and samples were removed after 30 min. β -Galactosidase assays were performed using the chloroform-sodium dodecyl sulfate (SDS) permeabilization method as previously described (48). Experiments were carried out at least three times.

Construction of luciferase reporters and luciferase assays. Promoter regions were identified using the Berkeley Drosophila Genome Project (http://www.fruitfly.org/seq_tools/promoter.html) and cloned into the EcoRI and BamHI restriction sites of plasmid pMK1*lux* (36), using primers listed in Table 1.

Promoter-luciferase expression assays were carried out in 200 μ l LB or M9 in 96-well plates at 37°C, with hormones, adrenergic antagonists, and LED209 added immediately after inoculation. Optical density (OD) and luminescence were measured in a Tecan Infinite200 spectrophotometer every 15 min for 16 h, and the maximum luminescence/OD at 600 nm (OD₆₀₀) of this was recorded.

Construction of gene knockouts. Genes were deleted using lambda Red mutagenesis (18). Briefly, primers P1 and P4 (Table 1) were used to amplify the kanamycin cassette from plasmid pKD13, which was transformed into the parent strain containing plasmid pKD46. The kanamycin cassette was removed using the FLP recombinase expressed from the temperature-sensitive plasmid pCP20, and mutations were confirmed by PCR.

Antimicrobial peptide resistance assays. Strains were grown in M9 for 2.5 h, epinephrine or norepinephrine was added, and incubation was continued for 30 min. Assays were done as previously described (36). Briefly, cultures were diluted to $\sim 10^6$ CFU/ml, and 50 μ l was inoculated into a 96-well plate and incubated with 100 μ l LL-37 (5 μ g/ml) (Phoenix Pharmaceuticals) for 1 h at 37°C and 200 rpm. Survival rates were determined by serial dilutions and viable counts on LB plates. Percent survival was calculated from the numbers of surviving bacteria divided by the numbers for the initial inoculum.

RESULTS

A genetic screen to identify genes regulated by epinephrine in *S. Typhimurium*. The detection of host molecules, such as epinephrine and norepinephrine, has an increasingly recognized role in bacterial virulence and host infection (14, 51, 59, 65). We used a global screen to identify epinephrine-regulated pathways in *S. Typhimurium*, by using *MudJ* transposon mutagenesis to produce random chromosomal *lacZ* gene insertions which were screened for differential regulation with epinephrine. A transposon library of 10,000 *S. Typhimurium* mutants was patched onto MacConkey agar plates with or without epinephrine, and mutants that showed a color change indicative of differential regulation of the inserted *lacZ* gene by epinephrine were isolated.

Eight mutants were isolated; seven mutants carried genes that were downregulated by epinephrine, and one carried a gene that was upregulated by epinephrine.

Inverse PCR was used to amplify and clone the DNA flank-

TABLE 1. *Salmonella enterica* serovar Typhimurium strains, plasmids, and primers used in this study^a

Strain, plasmid, or primer	Description, genotype, or sequence	Reference, source, and/or comment
Strains		
SL1344	Wild-type parent strain	34
LT2 TPA356	MudJ donor strain, <i>hisD9953::MudJ</i> , <i>hisA9944::MudI</i>	35
SL1344-YK	<i>yhaK::MudJ</i>	This study
SL1344-VK	<i>virK::MudJ</i>	This study
SL1344-MIG	<i>mig14::MudJ</i>	This study
SL1344-IC	<i>iroC::MudJ</i>	This study
SL1344-YP	<i>yedP::MudJ</i>	This study
SL1344-3	STM3081::MudJ	This study
SL1344-AC	<i>accC::MudJ</i>	This study
SL1344-NF	<i>nrdF::MudJ</i>	This study
SL1344-pYK	SL1344 pMK1 <i>lux</i> -P _{yhaK}	This study
SL1344-pVK	SL1344 pMK1 <i>lux</i> -P _{virK}	This study
SL1344-pMIG	SL1344 pMK1 <i>lux</i> -P _{mig14}	This study
SL1344-pACC	SL1344 pMK1 <i>lux</i> -P _{accB}	This study
SL1344-pYP	SL1344 pMK1 <i>lux</i> -P _{yedP}	This study
SL1344-p3	SL1344 pMK1 <i>lux</i> -P _{STM3083}	This study
SL1344-pIRO	SL1344 pMK1 <i>lux</i> -P _{iroB}	This study
SL1344-pNRD	SL1344 pMK1 <i>lux</i> -P _{nrdH}	This study
SL1344 <i>basS</i>	SL1344 Δ <i>basS</i>	36
SL1344 <i>qseBC</i>	SL1344 Δ <i>qseBC</i>	This study
SL1344 <i>qseF</i>	SL1344 Δ <i>qseF</i>	This study
SL1344 <i>qseBC qseF</i>	SL1344 Δ <i>qseBC \Delta</i> <i>qseF</i>	This study
SL1344 <i>virK</i>	SL1344 Δ <i>virK</i>	This study
SL1344 <i>mig14</i>	SL1344 Δ <i>mig14</i>	This study
SL1344 <i>virK mig14</i>	SL1344 Δ <i>virK \Delta</i> <i>mig14</i>	This study
SL1344 <i>tonB</i>	SL1344 Δ <i>tonB</i>	This study
Plasmids		
pPCR-Script	Sequencing vector	Stratagene
pMK1 <i>lux</i>	pBR322 with <i>luxCDABE</i> operon and MCS	36
pMK1 <i>lux</i> -P _{yhaK}	pMK1 <i>lux</i> with P _{yhaK} cloned as 5'EcoRI-3'BamHI fragment	This study
pMK1 <i>lux</i> -P _{virK}	pMK1 <i>lux</i> with P _{virK} cloned as 5'EcoRI-3'BamHI fragment	This study
pMK1 <i>lux</i> -P _{mig14}	pMK1 <i>lux</i> with P _{mig14} cloned as 5'EcoRI-3'BamHI fragment	This study
pMK1 <i>lux</i> -P _{accB}	pMK1 <i>lux</i> with P _{accB} cloned as 5'EcoRI-3'BamHI fragment	This study
pMK1 <i>lux</i> -P _{yedP}	pMK1 <i>lux</i> with P _{yedP} cloned as 5'EcoRI-3'BamHI fragment	This study
pMK1 <i>lux</i> -P _{STM3083}	pMK1 <i>lux</i> with P _{STM3083} cloned as 5'EcoRI-3'BamHI fragment	This study
pMK1 <i>lux</i> -P _{iroB}	pMK1 <i>lux</i> with P _{iroB} cloned as 5'EcoRI-3'BamHI fragment	This study
pMK1 <i>lux</i> -P _{nrdH}	pMK1 <i>lux</i> with P _{nrdH} cloned as 5'EcoRI-3'BamHI fragment	This study
Primers		
MudOut	CCGAATAATCCAATGTCCTCCCGGT	Inverse PCR (2)
MudTaq	AGTGCGCAATAACTTGCTCTCGTTC	Inverse PCR (2)
Universal-PCR	GTAAAACGACGGCCAGTGAGCGCGC	Inverse PCR (2)
<i>yhaK</i> 5	GCG GAATTC TCCAGCAATACCCGCCGCG	Cloning
<i>yhaK</i> 3	GCG GGATCC CGTTATTAACCTCTCTTTAC	Cloning
<i>virK</i> 5	GCG GAATTC CCTCACCCAGTCA	Cloning
<i>virK</i> 3	GCG GGATCC CTACTACTCGATCG	Cloning
<i>mig14</i> 5	GCG GAATTC ATACGCGGATGTGC	Cloning
<i>mig14</i> 3	GCG GGATCC TGACTTAGCTTACT	Cloning
<i>accB</i> 5	GCG GAATTC GTGAACTGTGCGGTCGATTG	Cloning
<i>accB</i> 3	GCG GGATCC GAGTGGGTTCCGACTCTTTG	Cloning
<i>yedP</i> 5	GCG GAATTC GCGGGTGAGCGTAGACGGACG	Cloning
<i>yedP</i> 3	GCG GGATTC GGGTCATGGATTGAGAGC	Cloning
STM3083 5	GTG GAATTC CCGTAACTGATGTTCTTAGG	Cloning
STM3083 3	GTG GGATCC ACTGGCGAGCTATGGTGTC	Cloning
<i>iroB</i> 5	GCG GAATTC TCGAACGGTGACGTTAAAGC	Cloning
<i>iroB</i> 3	GCG GGATCC AATACGCATGAGAAATCCTC	Cloning
<i>nrdH</i> 5	GCG GAATTC AATCAGCGATCTGACCTTCG	Cloning
<i>nrdH</i> 3	GCG GGATCC TGCTCATGATTCGTATTTC	Cloning
<i>qseBC</i> P1	GTTAACTGACGGCAACGCGGATTACCGCAAGGAAGAACAGGT GTAGGCTGGAGCTGCTTC	λ Red
<i>qseBC</i> P4	AAATGTGCAAAGTCTTTTGCGTTTTTGCAAAAAGTCTCTGATT CCGGGGATCCGTCGACC	λ Red
<i>qseBC</i> -5	ACATCGCTGCGGCGACAAG	λ Red
<i>qseBC</i> -3	GCGGTGCGGTGAAATTAGCA	λ Red

Continued on following page

TABLE 1—Continued

Strain, plasmid, or primer	Description, genotype, or sequence	Reference, source, and/or comment
<i>qseF</i> P1	GGCGCCGTCGCCGTCACAAGATGAGGTAACGCCATGATAAGT GTAGGCTGGAGCTGCTTC	λ Red
<i>qseF</i> P4	TTAAACGTAACATATTTTCGCGCTACTTTACGGCATGAAAAATT CCGGGGATCCGTCGACC	λ Red
<i>qseF</i> -5	CAAACCCGCGACGTCTGAAG	λ Red
<i>qseF</i> -3	GTCGCCTGTGTTTTGATCGG	λ Red
<i>virK</i> P1	CCGTCTCGGTTATAAGTCTATCGAGTAGTAGAGCCGTAGTGTG TAGGCTGGAGCTGCTTC	λ Red
<i>virK</i> P4	GAACGGTAGCAGGATCTGCACATCCGCGTATGAAATAGTTAT TCCGGGGATCCGTCGACC	λ Red
<i>virK</i> -5	CGTTAACTCAATCAGGCTA	λ Red
<i>virK</i> -3	CAGATTTCAAAGATCCGCT	λ Red
<i>mig14</i> P1	TCTTATCCCTTACCATAACGTCATCGATTAGCATGTTAGTGT AGGCTGGAGCTGCTTC	λ Red
<i>mig14</i> P4	TTCGAGTAATTGTCATTATGATTATATGGTGGTTAACTTAATTC CGGGGATCCGTCGACC	λ Red
<i>mig14</i> -5	CATGTTATGATCAACATCTT	λ Red
<i>mig14</i> -3	CCGAGTATAGTGTAAGTGAA	λ Red
<i>tonB</i> P1	ATTCAGCTCTGGTTTTTCAACTGAAACGATTATGACTTCAGTG TAGGCTGGAGCTGCTTC	λ Red
<i>tonB</i> P4	CAGCAGGTGATGGTATATTCTGGCTGGCGGCCAGAGAAT TCCGGGGATCCGTCGACC	λ Red
<i>tonB</i> -5	GATTGCTATTTGCATTTAAA	λ Red
<i>tonB</i> -3	AGTATACCCGCTTACGCCGC	λ Red

^a Restriction sites are underlined.

ing each *MudJ* insertion, which was then sequenced. The identities of the genes are listed in Table 2, and the chromosomal organization of the genes is shown in Fig. 1. A single gene was upregulated by epinephrine and was identified as *yhaK*, whose product is a putative cytoplasmic protein of unknown function. Downregulated genes included *virK* and *mig14*, which are important for virulence and survival in a mouse model, survival in macrophages, and resistance to the antimicrobial peptides polymyxin B and CRAMP (cathelicidin-related antimicrobial peptide) (10, 11, 20).

A fatty acid biosynthesis gene, *accC*, which encodes the biotin carboxylase subunit of acetyl-coenzyme A (CoA) car-

boxylase and catalyzes the first committed step of fatty acid biosynthesis, was among the downregulated genes (16).

A siderophore transport gene, *iroC*, part of the gene locus involved in the synthesis, transport, and utilization of the siderophore salmochelin, a glucosylated form of enterobactin (17, 33), was also downregulated.

TABLE 2. The identities of *MudJ* gene insertions affected by epinephrine in *S. Typhimurium*

Locus tag	Gene identity	Function
Epinephrine-upregulated gene STM3236	<i>yhaK</i>	Putative cytoplasmic protein
Epinephrine-downregulated genes		
STM2781	<i>virK</i>	Virulence protein
STM2782	<i>mig14</i>	Virulence protein
STM2774	<i>iroC</i>	ABC transporter
STM3380	<i>accC</i>	Acetyl-CoA carboxylase subunit
STM2808	<i>nrdF</i>	Ribonucleotide diphosphate reductase subunit
STM1986	<i>yedP</i>	Putative mannosyl-3-phosphoglycerate phosphatase
STM3081		Putative L-lactate/malate dehydrogenase




FIG. 1. Chromosomal organization of epinephrine-regulated genes. *MudJ* insertions are indicated by arrows. Upregulated genes and downregulated genes are shown with vertical stripes and horizontal stripes, respectively.




FIG. 2. Expression of isolated MudJ insertions. Cells were grown to mid-exponential phase, 50 μ M epinephrine (EPI) or 50 μ M norepinephrine (NE) was added, and β -galactosidase activity (in Miller units) was determined after 30 min. Asterisks indicate significant difference using Student's *t* test ($P < 0.01$). Error bars are shown. Experiments were performed at least three times. ND, result not determined.

A ribonucleotide diphosphate reductase (RRase) subunit, *nrdF*, which catalyzes conversion of ribonucleotides to deoxyribonucleotides for DNA synthesis, was also downregulated (45, 50, 52). The gene *STM3081*, which encodes a putative L-lactate/malate dehydrogenase, and *yedP*, which encodes a putative mannosyl-3-phosphoglycerate phosphatase, both of uncharacterized function, were also downregulated by epinephrine.

Epinephrine and norepinephrine regulate gene expression in similar manners. β -Galactosidase assays of isolated mutants grown to mid-exponential phase showed the same regulatory phenotype upon addition of 50 μ M epinephrine as seen in the initial screen using MacConkey agar plates. The addition of norepinephrine also resulted in the same phenotype, suggesting that both hormones may act in similar manners to affect *S. Typhimurium* gene expression (Fig. 2).

To further confirm the epinephrine- and norepinephrine-regulated phenotypes, promoter-luciferase reporter constructs for each gene or gene operon shown in Fig. 1 were made and introduced into the wild-type background. The activity of each promoter was measured by luminescence produced by the luciferase. The promoters of *STM3083*, *virK*, *mig14*, *nrdH*, *accB*, *iroB*, and *yedP* were all downregulated by epinephrine and norepinephrine, and expression of *yhaK* was upregulated by epinephrine and norepinephrine (Fig. 3). These results are in agreement with the β -galactosidase assays, confirming the initial observations. Each hormone was added to 500 μ M for luciferase assays to induce the same phenotype seen in the β -galactosidase assays. This may be explained by differences in growth conditions between the two assays; β -galactosidase assays were done in 25-ml volumes in shaking flasks with good aeration, whereas luciferase assays were conducted in 200- μ l volumes in 96-well plates with occasional agitation.

Epinephrine/norepinephrine-mediated gene expression is blocked by the α -adrenergic antagonist phentolamine. Some neuroendocrine stress hormone-mediated phenotypes, such as upregulation of type III secretion and motility in EHEC, are blocked by mammalian adrenergic antagonists, such as phentolamine (PE) (α -blocker) and propranolol (PO) (β -blocker) (14, 60). The neuroendocrine stress hormone-associated growth promotion of *S. Typhimurium* in SAPI-serum minimal media is also blocked by PE but not PO (27), and it was reported that the motility of *S. Typhimurium* is reduced by PE (8).

We investigated whether PE or PO could block the epineph-

rine- and norepinephrine-regulated phenotypes seen with *S. Typhimurium* by measuring the luminescence of promoter-luciferase constructs for each identified gene. The β -blocker PO did not reverse the epinephrine- and norepinephrine-regulated phenotype for any gene (data not shown). On the other hand, addition of the α -blocker PE could reverse the epinephrine- and norepinephrine-mediated regulation of *yhaK*, *virK*, and *mig14*, while having no significant effect on the other epinephrine/norepinephrine-regulated genes (see Fig. 3).

Epinephrine and norepinephrine decrease antimicrobial peptide resistance. *virK* and *mig14* are involved in resistance to the murine antimicrobial peptide CRAMP, both showing decreased survival rates in relation to the wild type (11). As expression of both genes is decreased by epinephrine and norepinephrine, we investigated whether these hormones affect resistance to cathelicidin LL-37, a human antimicrobial peptide with 67% identity to CRAMP and produced by macrophages and intestinal epithelial cells (4, 53).

Exposure to 500 μ M epinephrine or norepinephrine resulted in a significant decrease in resistance to LL-37 to approximately 20% of the water control (Fig. 4A). We constructed precise deletion mutants of *virK* and *mig14*, along with a *virK mig14* double mutant. The $\Delta virK$ mutant was significantly more sensitive to LL-37 than the wild type, and the $\Delta mig14$ mutant showed a slight, nonsignificant decrease in resistance. The $\Delta virK \Delta mig14$ double mutant was also more sensitive to LL-37, suggesting that the epinephrine/norepinephrine-induced LL-37 sensitivity is mediated through these genes (Fig. 4B).

Recently, we have shown that epinephrine increases sensitivity to polymyxin B in a manner dependent on the two-component system BasSR (36). We saw no change in expression of *virK* or *mig14* in the $\Delta basS$ mutant compared to the wild type, either with or without epinephrine/norepinephrine (data not shown), suggesting that *virK* and *mig14* are regulated by epinephrine/norepinephrine through a mechanism independent of BasSR.

The two-component systems QseBC and QseEF do not mediate adrenergic signaling responses. The QseBC two-component system has been identified as an adrenergic receptor in EHEC, regulating the epinephrine-, norepinephrine-, and autoinducer-3-mediated induction of type III secretion and motility (14, 60). The two-component system QseEF is also reported to be involved in adrenergic regulation of type III




FIG. 3. Maximum expression of promoter-luciferase fusions in the presence of 500 μ M EPI, NE, or phentolamine (PE). Luminescence is expressed as relative light units (RLU) per OD of culture standardized to water control. Asterisks indicate significant difference using Student's *t* test ($P < 0.01$). Error bars are shown. Experiments were performed at least three times for each fusion.

secretion in EHEC (56, 57). To determine whether QseBC or QseEF are involved in the adrenergic regulation of the identified genes, we constructed a precise *qseBC* mutant and a mutant of the response regulator *qseF*, which negates the sig-

naling effect of QseEF in EHEC (56). We introduced the promoter-luciferase reporter constructs for each identified gene into the Δ *qseBC* and Δ *qseF* mutants and found no difference in the expression levels of any genes in the Δ *qseBC* or




FIG. 4. Sensitivity to the antimicrobial peptide cathelicidin LL-37 (5 μ g/ml). Results are expressed as percent survival normalized to wild-type (WT)-H₂O. (A) Survival of SL1344 when grown in M9 supplemented with H₂O or 500 μ M EPI or NE. (B) Survival of SL1344, SL1344 *virK* mutant, SL1344 *mig-14* mutant, or SL1344 *virK mig-14* mutant when grown in M9 without supplementation. Asterisks indicate significant difference to wild-type-H₂O using Student's *t* test ($P < 0.01$). Error bars are shown. Experiments were performed at least three times.




FIG. 5. Maximum expression of promoter-luciferase fusions of stated genes in the WT or $\Delta qseBC \Delta qseF$ mutant (*qse*). EPI or NE were added at 500 μ M. Luminescence is expressed as RLU per OD of culture standardized to water control. Error bars are shown. Experiments were performed at least three times for each fusion.

$\Delta qseF$ mutant compared to the wild type upon addition of water, epinephrine, or norepinephrine (data not shown). As many phenotypes attributed to QseBC have been analyzed in a *qseC* single mutant (8, 14, 55, 60), we constructed and analyzed a *qseC* single mutant, which also displayed no change in phenotype compared to the wild type (data not shown). Furthermore, we constructed a $\Delta qseBC \Delta qseF$ triple mutant and introduced each promoter-luciferase construct to investigate the possibility of functional redundancy between the two signaling systems. We found no change in expression in the $\Delta qseBC \Delta qseF$ mutant compared to the wild type (Fig. 5). This suggests that these genes are regulated through a mechanism independent of known bacterial adrenergic signaling pathways in *S. Typhimurium*.

N-phenyl-4-[(phenylamino)thioxomethyl]amino-benzene-sulfonamide (LED209) was recently identified as inhibiting adrenergic signaling in EHEC by preventing the binding of norepinephrine to QseC and thus blocking QseC-mediated cell signaling at a concentration of 5 pM (55). Addition of LED209 at 5 pM, 5 nM, or 5 μ M did not affect expression of any of the identified genes when supplemented with water, epinephrine, or norepinephrine (Fig. 6). This further supports the argument that QseC is not involved in adrenergic gene regulation, at least of these genes, in *S. Typhimurium*.

Role of TonB-dependent siderophore uptake in mediating adrenergic sensing. Some adrenergic phenotypes in bacteria are associated with altered iron uptake via the siderophore enterobactin (13, 28, 47). To determine whether the same mechanism is responsible for the adrenergic regulation of the identified genes, we constructed a *tonB* mutant defective in siderophore uptake and introduced the promoter-luciferase reporter constructs for each gene. We found that each fusion showed differential regulation upon addition of each hormone, suggesting that the adrenergic regulation is mediated through a mechanism independent of TonB (Fig. 7). We found that expression of *mig14* and *nrdH* were significantly upregulated in the $\Delta tonB$ mutant compared to the wild type, suggesting that these genes may be affected by the altered iron status of the

cell in the $\Delta tonB$ mutant. Upon addition of either hormone, we saw that expression was reduced to approximately the same level in both the wild type and the $\Delta tonB$ mutant. These results suggest that in contrast to other adrenergic phenotypes, such as growth promotion in SAPI-serum (13, 28, 47), the adrenergic regulation of these genes is mediated through a mechanism independent of TonB-dependent siderophore transport and iron uptake.

DISCUSSION

This work has identified several genes in *S. Typhimurium* which are regulated by epinephrine and norepinephrine. We have shown adrenergic downregulation of *virK* and *mig14*, which are important virulence genes affecting survival and long-term persistence in the host. A mutation in either gene reduces the virulence of *S. Typhimurium* in a mouse infection model and reduces survival in macrophages. *S. Typhimurium mig14* mutants colonize mouse liver/spleen as wild type but fail to replicate after 5 days postinfection, suggesting a role in persistence and the late stages of infection (11, 20). The production of epinephrine recently observed with macrophages (12, 26) may therefore act as a host defense system to reduce the pathogenic potential of *Salmonella* by downregulating expression of these virulence genes. The downregulation of *mig14* may reduce levels of persistent infection and promote clearance of bacteria.

Both *virK* and *mig14* are reported to affect antimicrobial peptide resistance, mutants showing decreased resistance to polymyxin B, and the mouse-derived peptide CRAMP (10, 11). Brodsky et al. (11) showed increased levels of fluorescein isothiocyanate (FITC)-labeled CRAMP associated with *virK* and *mig14* mutant cells compared to the wild type, suggesting that these genes may act to prevent association of antimicrobial peptides with the bacterial cell, although the mechanism is currently unknown.

We have shown a stress hormone-mediated decrease in resistance to the peptide cathelicidin LL-37, a human antimicro-




FIG. 6. Maximum expression of promoter-luciferase fusions of stated genes. LED209 was added at 5 μ M, and EPI and NE were added at 500 μ M. Luminescence is expressed as RLU per OD of culture standardized to water control. Error bars are shown. Experiments were performed at least three times for each fusion.

bial peptide displaying 67% identity to CRAMP, which is produced by diverse tissues, including the gastrointestinal tract, bone marrow, and macrophages, and has antimicrobial activity against many Gram-positive and Gram-negative bacteria.




FIG. 7. Maximum expression of promoter-luciferase fusions of stated genes in the WT or Δ *tonB* mutant. EPI or NE were added at 500 μ M. Luminescence is expressed as RLU per OD of culture standardized to water control. Error bars are shown. Asterisks indicate significant difference between Δ *tonB* mutant and the respective wild-type values ($P < 0.001$). Experiments were performed at least three times for each fusion.

LL-37 plays an important role in innate immunity, serving to protect epithelia from bacterial pathogens (4, 53). Levels of host stress hormones increase in response to infection (23, 32), and an epinephrine/norepinephrine-mediated increase in sensitivity to LL-37 may act as a host defense system to combat infection, suggesting that bacterial sensing of stress hormones may be a double-edged sword; although bacteria can sense and exploit these molecules, the host can use the same signals to manipulate the bacteria. The effects of epinephrine and norepinephrine are likely to be complex, involving multiple effects on both bacteria and host to influence the outcome of infection. It is possible that the adrenergic downregulation of these genes may confer an advantage to the bacteria under certain in vivo conditions but an unavoidable disadvantage in others; for example, it was recently shown that the downregulation of the lipopolysaccharide (LPS)-modifying enzymes PmrF and PagL causes an increase in sensitivity to polymyxin B but also concomitantly reduces activation of the TLR-4 Toll-like receptors, reducing the host inflammatory response to infection (37, 38, 49).

Pretreatment of mice with norepinephrine enhances recovery of *S. Typhimurium* from the liver and cecum of infected BALB/c mice by up to 2 logs (47, 65). These mice are genetically highly susceptible to *Salmonella* infection and usually die a few days postinfection. Mig14 is involved in the later stages of infection; when infecting the less susceptible mouse strain 129Xi/SvJ, mutants initially show no virulence defect but are cleared from the host more effectively after 28 days of infection (11). Speculatively, epinephrine and norepinephrine may initially act to increase bacterial virulence during acute infection, as observed with mice, chicks, and cattle, (44, 47, 65), but the hormone-mediated downregulation of genes such as *mig14* may then help clear infection and reduce long-term persistence of bacteria in the host.

Epinephrine has been shown to decrease resistance to polymyxin B through downregulation of the *pmr* operon, in a man-

ner dependent on the two-component system BasSR (36). We have shown that expression of *virK* and *mig14* is independent of BasSR, suggesting that stress hormones affect LL-37 and polymyxin B resistance by two independent mechanisms. Supporting this theory, the epinephrine-mediated downregulation of the *pmr* operon and polymyxin sensitivity are blocked by the β -adrenergic antagonist propranolol (36), whereas *mig14* and *virK* expression is blocked by the α -adrenergic antagonist phentolamine.

The two-component system QseBC has been identified as an adrenergic receptor in EHEC, regulating virulence-related processes in a manner blockable by the adrenergic antagonist phentolamine (14, 60). We found that deletion of *qseBC* had no impact on the regulation of any hormone-regulated gene in this study. Despite this, we have observed that the adrenergic regulation of *virK*, *mig14*, and *yhaK* is blockable by phentolamine. This suggests that a different receptor may be responsible for mediating the adrenergic phenotypes seen with *S. Typhimurium*. In support of this hypothesis, it has recently been published that QseBC does not affect any of the identified epinephrine/norepinephrine-regulated phenotypes in *S. Typhimurium* as determined by microarray analysis (46). Furthermore, we found no effect of LED209 on gene expression, although this compound inhibits QseC-mediated adrenergic signaling in EHEC (55), further supporting the theory that adrenergic cell signaling is mediated through different pathways in *S. Typhimurium* and EHEC. Furthermore, as the remaining identified genes were unaffected by phentolamine or propranolol, this suggests that at least two signaling pathways are present, one blockable by phentolamine and one not.

Epinephrine and norepinephrine have been shown to act as siderophores to increase bacterial iron uptake (13, 28, 36, 47). This increase in intracellular iron may cause the downregulation of genes involved in siderophore synthesis and transport, normally expressed under iron-limiting conditions. We have observed this with the epinephrine/norepinephrine-mediated downregulation of the *iro* genes, which synthesize and transport the siderophore salmochelin (17, 33). The downregulation of the *nrd* genes may also be due to changes in iron status, as their expression has been reported to increase during iron-limited conditions (45).

Enterobactin is a major siderophore in *Salmonella*. A glucosylated form of enterobactin, salmochelin displays lower iron binding affinity but is advantageous under certain conditions (33). Enterobactin is bound by host proteins, such as serum albumin and neutrophil lipocalin NGAL, which is produced by neutrophils and epithelial cells (30, 40), which reduces efficiency of enterobactin-mediated iron acquisition. The glucosylation present on salmochelin inhibits the binding of lipocalin (25), making it more effective for iron acquisition in some environments, such as inside macrophages, but it has been reported that the *Salmonella*-containing vacuole (SCV) of macrophages is an iron-rich environment (24), so expression of iron uptake genes would be limited and of no significance in modulating pathogenesis.

The range of epinephrine and norepinephrine concentrations used in this study reflect those widely used in other studies (28, 36, 43, 60). Up to half the norepinephrine in the body may be produced in the enteric nervous system (ENS) (3), but the actual concentrations found are very difficult to

determine accurately. Furthermore, the levels vary according to, for example, the tissue location, stress levels, and circadian rhythms and can potentially reach millimolar concentrations (12), adding physiological relevance to the concentrations used in this study.

The genes identified in this work show no overlap with genes previously identified as being regulated by epinephrine in *S. Typhimurium* (36). This may be due to following differences in the screening methods utilized: MudJ transcriptional fusions compared to mRNA levels, differences in growth conditions affecting bacterial physiology, and agar plates compared to LB liquid culture. The two methods may highlight important differences in the *S. Typhimurium* responses to epinephrine/norepinephrine under different growth conditions. It must be noted that concomitant with our data, Bearson et al. (9) identified the gene *iroC* as being downregulated by norepinephrine in SAPI-serum minimal media.

The bacterial phenotypes associated with epinephrine/norepinephrine have been linked to effects on iron uptake or interaction with quorum-sensing (QS) systems (13, 43, 60). We have found that the adrenergic regulation of the identified genes is independent of TonB-dependent siderophore transport. Interaction of stress hormones with quorum-sensing systems is thought to be distinct from iron uptake, but the divide between iron uptake and quorum sensing is becoming increasingly blurred. For example, the *Pseudomonas aeruginosa* siderophore pyoverdine also has a separate role in cell signaling (41), and the *Pseudomonas* signaling molecule PQS along with other quinolones can chelate iron and facilitate iron uptake (21). A mutant in *tonB*, essential for iron-bound siderophore uptake, has been reported to be defective in the production of the QS molecule 3-oxo-C12-AHL in *Pseudomonas* (1). Stress hormone-regulated effects on bacteria are likely to be complex, involving multiple factors, including quorum sensing and iron uptake.

It is interesting to note that three of the identified genes, *virK*, *mig14*, and *iroC*, are all part of a 45-kb *S. enterica*-specific genomic region, thought to have been acquired by horizontal transfer after divergence from *E. coli* (7). Both *mig14* and *virK* are found in some (but not all) *S. enterica* serovars, but neither is found in *E. coli*. The *iro* locus is found in *S. enterica* and some uropathogenic *E. coli* strains (33, 61). The adrenergic regulation of these genes may therefore form part of a unique *Salmonella*-specific or even *Salmonella* serovar-specific response to host-derived hormones.

None of the remaining genes that we identified that are present in *E. coli* are regulated by epinephrine or norepinephrine in *E. coli*, to the best of our knowledge, although several microarray studies have been conducted to investigate the global effects of these hormones in *E. coli* (5, 22, 39). This may be due to the differences in pathogenesis of the two; *S. Typhimurium* is an invasive pathogen infecting macrophages and epithelial cells, whereas *E. coli* is mainly noninvasive, remaining in the host intestine. The two very different niches occupied by each pathogen require different sets of genes to be expressed to gain the maximum advantage, so epinephrine and norepinephrine may regulate different sets of genes to the advantage or detriment of the pathogen.

In conclusion, we have identified epinephrine/norepinephrine-mediated downregulation of two important virulence-

related genes, *virK* and *mig14*, and increased sensitivity to the antimicrobial peptide LL-37, potentially highlighting a mechanism for neuroendocrine stress hormone-mediated down-regulation of bacterial virulence.

This study has further elucidated the *S. Typhimurium* response to the host-derived hormones epinephrine and norepinephrine, providing further insights into the area of bacterium-host communication.

ACKNOWLEDGMENTS

We thank Judith Hall and Claire Townes for providing LL-37 and Joe Gray (Pinnacle) for DNA sequencing.

This work was supported by research grants from the Medical Research Council (MRC) and the Biotechnology and Biological Sciences Research Council (BBSRC) to C.M.A.K. and P.W. In addition, H.S. was a recipient of a BBSRC DTA Ph.D. studentship with C.M.A.K.

REFERENCES

1. Abbas, A., C. Adams, N. Scully, J. Glennon, and F. O'Gara. 2007. A role for TonB1 in biofilm formation and quorum sensing in *Pseudomonas aeruginosa*. *FEMS Microbiol. Lett.* **274**:269–278.
2. Ahmer, B. M., J. van Reeuwijk, C. D. Timmers, P. J. Valentine, and F. Heffron. 1998. *Salmonella typhimurium* encodes an SdiA homolog, a putative quorum sensor of the LuxR family, that regulates genes on the virulence plasmid. *J. Bacteriol.* **180**:1185–1193.
3. Aneman, A., G. Eisenhofer, L. Olbe, J. Dalenback, P. Nitescu, L. Fandriks, and P. Friberg. 1996. Sympathetic discharge to mesenteric organs and the liver. Evidence for substantial mesenteric organ norepinephrine spillover. *J. Clin. Invest.* **97**:1640–1646.
4. Bals, R., X. Wang, M. Zasloff, and J. M. Wilson. 1998. The peptide antibiotic LL-37/hCAP-18 is expressed in epithelia of the human lung where it has broad antimicrobial activity at the airway surface. *Proc. Natl. Acad. Sci. U. S. A.* **95**:9541–9546.
5. Bansal, T., D. Englert, J. Lee, M. Hegde, T. K. Wood, and A. Jayaraman. 2007. Differential effects of epinephrine, norepinephrine, and indole on *Escherichia coli* O157:H7 chemotaxis, colonization, and gene expression. *Infect. Immun.* **75**:4597–4607.
6. Bassler, B. L., and R. Losick. 2006. Bacterially speaking. *Cell* **125**:237–246.
7. Bäuml, A. J., and F. Heffron. 1998. Mosaic structure of the *smgB-nrdE* intergenic region of *Salmonella enterica*. *J. Bacteriol.* **180**:2220–2223.
8. Bearson, B. L., and S. M. D. Bearson. 2008. The role of the QseC quorum-sensing sensor kinase in colonization and norepinephrine-enhanced motility of *Salmonella enterica* serovar Typhimurium. *Microb. Pathog.* **44**:271–278.
9. Bearson, B. L., S. M. D. Bearson, J. J. Uthe, S. E. Dowd, J. O. Houghton, I. Lee, M. J. Toscano, and D. C. Lay, Jr. 2008. Iron regulated genes of *Salmonella enterica* serovar Typhimurium in response to norepinephrine and the requirement of *jepDGC* for norepinephrine-enhanced growth. *Microbes Infect.* **10**:807–816.
10. Brodsky, I. E., R. K. Ernst, S. I. Miller, and S. Falkow. 2002. *mig-14* is a *Salmonella* gene that plays a role in bacterial resistance to antimicrobial peptides. *J. Bacteriol.* **184**:3203–3213.
11. Brodsky, I. E., N. Ghoris, S. Falkow, and D. Monack. 2005. Mig-14 is an inner membrane-associated protein that promotes *Salmonella typhimurium* resistance to CRAMP, survival within activated macrophages and persistent infection. *Mol. Microbiol.* **55**:954–972.
12. Brown, S. W., R. T. Meyers, K. M. Brennan, J. M. Rumble, N. Narasimhachari, E. F. Perozzi, J. J. Ryan, J. K. Stewart, and K. Fischer-Stenger. 2003. Catecholamines in a macrophage cell line. *J. Neuroimmunol.* **135**:47–55.
13. Burton, C. L., S. R. Chhabra, S. Swift, T. J. Baldwin, H. Withers, S. J. Hill, and P. Williams. 2002. The growth response of *Escherichia coli* to neurotransmitters and related catecholamine drugs requires a functional enterobactin biosynthesis and uptake system. *Infect. Immun.* **70**:5913–5923.
14. Clarke, M. B., D. T. Hughes, C. Zhu, E. C. Boedeker, and V. Sperandio. 2006. The QseC sensor kinase: a bacterial adrenergic receptor. *Proc. Natl. Acad. Sci. U. S. A.* **103**:10420–10425.
15. Cogan, T. A., A. O. Thomas, L. E. N. Rees, A. H. Taylor, M. A. Jepson, P. H. Williams, J. Ketley, and T. J. Humphrey. 2007. Norepinephrine increases the pathogenic potential of *Campylobacter jejuni*. *Gut* **56**:1060–1065.
16. Cronan, J. E., Jr., and G. L. Waldrop. 2002. Multi-subunit acetyl-CoA carboxylases. *Prog. Lipid Res.* **41**:407–435.
17. Crouch, M.-L. V., M. Castor, J. E. Karlinsey, T. Kalthorn, and F. C. Fang. 2008. Biosynthesis and IroC-dependent export of the siderophore salmochelin are essential for virulence of *Salmonella enterica* serovar Typhimurium. *Mol. Microbiol.* **67**:971–983.
18. Datsenko, K. A., and B. L. Wanner. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97**:6640–6645.
19. Davis, R. W., D. Botstein, and J. R. Roth. 1980. Advanced bacterial genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
20. Detweiler, C. S., D. M. Monack, I. E. Brodsky, H. Mathew, and S. Falkow. 2003. *virK*, *somA* and *rscC* are important for systemic *Salmonella enterica* serovar Typhimurium infection and cationic peptide resistance. *Mol. Microbiol.* **48**:385–400.
21. Diggle, S. P., S. Matthijs, V. J. Wright, M. P. Fletcher, S. R. Chhabra, I. L. Lamont, X. Kong, R. C. Hider, P. Cornelis, M. Camara, and P. Williams. 2007. The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS play multifunctional roles in quorum sensing and iron entrapment. *Chem. Biol.* **14**:87–96.
22. Dowd, S. E. 2007. *Escherichia coli* O157:H7 gene expression in the presence of catecholamine norepinephrine. *FEMS Microbiol. Lett.* **273**:214–223.
23. Dreu, D., G. Sonnenfeld, N. Fowler, D. S. Morton, and M. Lyte. 1999. Effects of social conflict on immune responses and *E. coli* growth within closed chambers in mice. *Physiol. Behav.* **67**:133–140.
24. Eriksson, S., S. Lucchini, A. Thompson, M. Rhen, and J. C. D. Hinton. 2003. Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*. *Mol. Microbiol.* **47**:103–118.
25. Fischbach, M. A., H. Lin, L. Zhou, Y. Yu, R. J. Abergel, D. R. Liu, K. N. Raymond, B. L. Wanner, R. K. Strong, C. T. Walsh, A. Aderem, and K. D. Smith. 2006. The pathogen-associated *iroA* gene cluster mediates bacterial evasion of lipocalin 2. *Proc. Natl. Acad. Sci. U. S. A.* **103**:16502–16507.
26. Flierl, M. A., D. Rittirsch, B. A. Nadeau, A. J. Chen, J. V. Sarma, F. S. Zetoune, S. R. McGuire, R. P. List, D. E. Day, L. M. Hoesel, H. Gao, N. Van Rooijen, M. S. Huber-Lang, R. R. Neubig, and P. A. Ward. 2007. Phagocyte-derived catecholamines enhance acute inflammatory injury. *Nature* **449**:721–725.
27. Freestone, P. P. E., R. D. Haigh, and M. Lyte. 2007. Blockade of catecholamine-induced growth by adrenergic and dopaminergic receptor antagonists in *Escherichia coli* O157:H7, *Salmonella enterica* and *Yersinia enterocolitica*. *BMC Microbiol.* **7**:8.
28. Freestone, P. P. E., R. D. Haigh, P. H. Williams, and M. Lyte. 2003. Involvement of enterobactin in norepinephrine-mediated iron supply from transferrin to enterohaemorrhagic *Escherichia coli*. *FEMS Microbiol. Lett.* **222**:39–43.
29. Freestone, P. P. E., and M. Lyte. 2008. Microbial endocrinology: experimental design issues in the study of interkingdom signalling in infectious disease. *Adv. Appl. Microbiol.* **64**:75–105.
30. Goetz, D. H., M. A. Holmes, N. Borregaard, M. E. Bluhm, K. N. Raymond, and R. K. Strong. 2002. The neutrophil lipocalin NGAL is a bacteriostatic agent that interferes with siderophore-mediated iron acquisition. *Mol. Cell* **10**:1033–1043.
31. Goldberg, A. D., L. C. Becker, R. Bonsall, J. D. Cohen, M. W. Ketterer, P. G. Kaufman, D. S. Krantz, K. C. Light, R. P. McMahon, T. Noreuil, C. J. Pepine, J. Raczynski, P. H. Stone, D. Strother, H. Taylor, and D. S. Sheps. 1996. Ischemic, hemodynamic, and neurohormonal responses to mental and exercise stress. Experience from the Psychophysiological Investigations of Myocardial Ischemia Study (PIMI). *Circulation* **94**:2402–2409.
32. Groves, A. C., J. Griffiths, F. Leung, and R. N. Meek. 1973. Plasma catecholamines in patients with serious postoperative infection. *Ann. Surg.* **178**:102–107.
33. Hantke, K., G. Nicholson, W. Rabsch, and G. Winkelmann. 2003. Salmochelins, siderophores of *Salmonella enterica* and uropathogenic *Escherichia coli* strains, are recognized by the outer membrane receptor IroN. *Proc. Natl. Acad. Sci. U. S. A.* **100**:3677–3682.
34. Hoiseit, S. K., and B. A. Stocker. 1981. Aromatic-dependent *Salmonella typhimurium* are non-virulent and effective as live vaccines. *Nature* **291**:238–239.
35. Hughes, K. T., and J. R. Roth. 1988. Transitory cis complementation: a method for providing transposition functions to defective transposons. *Genetics* **119**:9–12.
36. Karavolos, M. H., H. Spencer, D. M. Bulmer, A. Thompson, K. Winzer, P. Williams, J. C. Hinton, and C. M. Khan. 2008. Adrenaline modulates the global transcriptional profile of *Salmonella* revealing a role in the antimicrobial peptide and oxidative stress resistance responses. *BMC Genomics* **6**:458–471.
37. Kawasaki, K., R. K. Ernst, and S. I. Miller. 2004. 3-O-deacylation of lipid A by PagL, a PhoP/PhoQ-regulated deacylase of *Salmonella typhimurium*, modulates signaling through Toll-like receptor 4. *J. Biol. Chem.* **279**:20044–20048.
38. Kawasaki, K., R. K. Ernst, and S. I. Miller. 2005. Inhibition of *Salmonella enterica* serovar Typhimurium lipopolysaccharide deacylation by aminoarabinose membrane modification. *J. Bacteriol.* **187**:2448–2457.
39. Kendall, M. M., D. A. Rasko, and V. Sperandio. 2007. Global effects of the cell-to-cell signaling molecules autoinducer-2, autoinducer-3, and epinephrine in a *luxS* mutant of enterohemorrhagic *Escherichia coli*. *Infect. Immun.* **75**:4875–4884. [Author's correction, **76**:1319, 2008.]
40. Konopka, K., and J. B. Neilands. 1984. Effect of serum albumin on siderophore-mediated utilization of transferrin iron. *Biochemistry* **23**:2122–2127.
41. Lamont, I. L., P. A. Beare, U. Ochsner, A. I. Vasil, and M. L. Vasil. 2002.

- Siderophore-mediated signaling regulates virulence factor production in *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. U. S. A. **99**:7072–7077.
42. Lyte, M. 2004. Microbial endocrinology and infectious disease in the 21st century. Trends Microbiol. **12**:14–20.
 43. Lyte, M., and S. Ernst. 1992. Catecholamine induced growth of gram negative bacteria. Life Sci. **50**:203–212.
 44. McCuddin, Z. P., S. A. Carlson, and V. K. Sharma. 2008. Experimental reproduction of bovine *Salmonella* encephalopathy using a norepinephrine-based stress model. Vet. J. **175**:82–88.
 45. McHugh, J. P., F. Rodriguez-Quinones, H. Abdul-Tehrani, D. A. Svislunenko, R. K. Poole, C. E. Cooper, and S. C. Andrews. 2003. Global iron-dependent gene regulation in *Escherichia coli*. A new mechanism for iron homeostasis. J. Biol. Chem. **278**:29478–29486.
 46. Merighi, M., A. N. Septer, A. Carroll-Portillo, A. Bhatiya, S. Porwollik, M. McClelland, and J. S. Gunn. 2009. Genome-wide analysis of the PreA/PreB (QseB/QseC) regulon of *Salmonella enterica* serovar Typhimurium. BMC Microbiol. **9**:42.
 47. Methner, U., W. Rabsch, R. Reissbrodt, and P. H. Williams. 2008. Effect of norepinephrine on colonisation and systemic spread of *Salmonella enterica* in infected animals: role of catecholate siderophore precursors and degradation products. Int. J. Med. Microbiol. **298**:429–439.
 48. Miller, J. H. 1992. A short course in bacterial genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 49. Miller, S. I., R. K. Ernst, and M. W. Bader. 2005. LPS, TLR4 and infectious disease diversity. Nat. Rev. Microbiol. **3**:36–46.
 50. Monje-Casas, F., J. Jurado, M. J. Prieto-Alamo, A. Holmgren, and C. Pueyo. 2001. Expression analysis of the *nrdHIEF* operon from *Escherichia coli*. Conditions that trigger the transcript level in vivo. J. Biol. Chem. **276**:18031–18037.
 51. Nakano, M., A. Takahashi, Y. Sakai, and Y. Nakaya. 2007. Modulation of pathogenicity with norepinephrine related to the type III secretion system of *Vibrio parahaemolyticus*. J. Infect. Dis. **195**:1353–1360.
 52. Nordlund, P., and P. Reichard. 2006. Ribonucleotide reductases. Annu. Rev. Biochem. **75**:681–706.
 53. Overhage, J., A. Campisano, M. Bains, E. C. W. Torfs, B. H. A. Rehm, and R. E. W. Hancock. 2008. Human host defense peptide LL-37 prevents bacterial biofilm formation. Infect. Immun. **76**:4176–4182.
 54. Papezova, K., D. Gregorova, J. Jonuschies, and I. Rychlik. 2007. Ordered expression of virulence genes in *Salmonella enterica* serovar Typhimurium. Folia Microbiol. **52**:107–114.
 55. Rasko, D. A., C. G. Moreira, D. R. Li, N. C. Reading, J. M. Ritchie, M. K. Waldor, N. Williams, R. Taussig, S. Wei, M. Roth, D. T. Hughes, J. F. Huntley, M. W. Fina, J. R. Falck, and V. Sperandio. 2008. Targeting QseC signaling and virulence for antibiotic development. Science **321**:1078–1080.
 56. Reading, N. C., D. A. Rasko, A. G. Torres, and V. Sperandio. 2009. The two-component system QseEF and the membrane protein QseG link adrenergic and stress sensing to bacterial pathogenesis. Proc. Natl. Acad. Sci. U. S. A. **106**:5889–5894.
 57. Reading, N. C., A. G. Torres, M. M. Kendall, D. T. Hughes, K. Yamamoto, and V. Sperandio. 2007. A novel two-component signaling system that activates transcription of an enterohemorrhagic *Escherichia coli* effector involved in remodeling of host actin. J. Bacteriol. **189**:2468–2476.
 58. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 59. Scheckelhoff, M. R., S. R. Telford, M. Wesley, and L. T. Hu. 2007. *Borrelia burgdorferi* intercepts host hormonal signals to regulate expression of outer surface protein A. Proc. Natl. Acad. Sci. U. S. A. **104**:7247–7252.
 60. Sperandio, V., A. G. Torres, B. Jarvis, J. P. Nataro, and J. B. Kaper. 2003. Bacteria-host communication: the language of hormones. Proc. Natl. Acad. Sci. U. S. A. **100**:8951–8956.
 61. Valdivia, R. H., D. M. Cirillo, A. K. Lee, D. M. Bouley, and S. Falkow. 2000. *mig-14* is a horizontally acquired, host-induced gene required for *Salmonella enterica* lethal infection in the murine model of typhoid fever. Infect. Immun. **68**:7126–7131.
 62. Van Immerseel, F., J. De Buck, F. Pasmans, P. Velge, E. Bottreau, V. Fievez, F. Haesebrouck, and R. Ducatelle. 2003. Invasion of *Salmonella enteritidis* in avian intestinal epithelial cells *in vitro* is influenced by short-chain fatty acids. Int. J. Food Microbiol. **85**:237–248.
 63. Vlisidou, I., M. Lyte, P. M. van Diemen, P. Hawes, P. Monaghan, T. S. Wallis, and M. P. Stevens. 2004. The neuroendocrine stress hormone norepinephrine augments *Escherichia coli* O157:H7-induced enteritis and adherence in a bovine ligated ileal loop model of infection. Infect. Immun. **72**:5446–5451.
 64. Williams, P. 2007. Quorum sensing, communication and cross-kingdom signalling in the bacterial world. Microbiology **153**:3923–3938.
 65. Williams, P. H., W. Rabsch, U. Methner, W. Voigt, H. Tschape, and R. Reissbrodt. 2006. Catecholate receptor proteins in *Salmonella enterica*: role in virulence and implications for vaccine development. Vaccine **24**:3840–3844.