



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Direct measurements of low-energy resonance strengths of the $^{23}\text{Na}(p,)^{24}\text{Mg}$ reaction for astrophysics

Citation for published version:

Boeltzig, A, Best, A, Pantaleo, FR, Imbriani, G, Junker, M, Aliotta, M, Balibrea-Correa, J, Bemmerer, D, Broggini, C, Bruno, CG, Buompane, R, Caciolli, A, Cavanna, F, Chillery, T, Ciani, GF, Corvisiero, P, Csedreki, L, Davinson, T, deBoer, RJ, Depalo, R, Di Leva, A, Elekes, Z, Ferraro, F, Fiore, EM, Formicola, A, Fülöp, Z, Gervino, G, Guglielmetti, A, Gustavino, C, Gyürky, G, Kochanek, I, Lugaro, M, Marigo, P, Menegazzo, R, Mossa, V, Munnik, F, Patocchio, V, Perrino, R, Piatti, D, Prati, P, Schiavulli, L, Stöckel, K, Straniero, O, Strieder, F, Szucs, T, Takács, MP, Trezzi, D, Wiescher, M & Zavatarelli, S 2019, 'Direct measurements of low-energy resonance strengths of the $^{23}\text{Na}(p,)^{24}\text{Mg}$ reaction for astrophysics', *Physics Letters B*, vol. 795, pp. 122-128. <https://doi.org/10.1016/j.physletb.2019.05.044>

Digital Object Identifier (DOI):

[10.1016/j.physletb.2019.05.044](https://doi.org/10.1016/j.physletb.2019.05.044)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Physics Letters B

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Direct Measurements of Low-Energy Resonance Strengths of the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ Reaction for Astrophysics

A. Boeltzig^{a,b,c,d,*}, A. Best^{e,f}, F. R. Pantaleo^{g,h}, G. Imbriani^{e,f}, M. Junker^{d,a}, M. Aliottaⁱ, J. Balibrea-Correa^{e,f}, D. Bemmerer^j, C. Broggini^l, C. G. Brunoⁱ, R. Buompane^{f,k}, A. Cacioli^{m,l}, F. Cavannaⁿ, T. Chilleryⁱ, G. F. Ciani^{a,d}, P. Corvisiero^{o,n}, L. Csedreki^{d,a}, T. Davinsonⁱ, R. J. deBoer^{b,c}, R. Depalo^{m,l}, A. Di Leva^{e,f}, Z. Elekes^p, F. Ferraro^{o,n}, E. M. Fiore^{g,h}, A. Formicola^d, Zs. Fülöp^p, G. Gervino^{q,r}, A. Guglielmetti^{s,t}, C. Gustavino^u, Gy. Gyürky^p, I. Kochanek^d, M. Lugaro^v, P. Marigo^{l,m}, R. Menegazzo^l, V. Mossa^{g,h}, F. Munnik^j, V. Patricchio^h, R. Perrino^{h,l}, D. Piatti^{m,l}, P. Prati^{o,n}, L. Schiavulli^{g,h}, K. Stöckel^{l,w}, O. Straniero^{x,d}, F. Strieder^y, T. Szücs^j, M. P. Takács^{j,w,2}, D. Trezzi^{s,t}, M. Wiescher^{b,c}, S. Zavatarelli^a

^aGran Sasso Science Institute, Viale F. Crispi 7, 67100 L'Aquila, Italy

^bThe Joint Institute for Nuclear Astrophysics, University of Notre Dame, Notre Dame, Indiana 46556, USA

^cDepartment of Physics, University of Notre Dame, Nieuwland Science Hall 225, Notre Dame, Indiana 46556-5670, USA

^dIstituto Nazionale di Fisica Nucleare Laboratori Nazionali del Gran Sasso (LNGS), Via G. Acitelli 22, 67100 Assergi, Italy

^eUniversità degli Studi di Napoli "Federico II", Dipartimento di Fisica "E. Pancini", Via Cinthia 21, 80126 Napoli, Italy

^fIstituto Nazionale di Fisica Nucleare, Sezione di Napoli, Via Cintia 21, 80126 Napoli, Italy

^gUniversità degli Studi di Bari, Dipartimento Interateneo di Fisica, Via G. Amendola 173, 70126 Bari, Italy

^hIstituto Nazionale di Fisica Nucleare, Sezione di Bari, Via E. Orabona 4, 70125 Bari, Italy

ⁱSchool of Physics and Astronomy, University of Edinburgh, Peter Guthrie Tait Road, EH9 3FD Edinburgh, United Kingdom

^jHelmholtz-Zentrum Dresden-Rossendorf, Bautzner Landstraße 400, 01328 Dresden, Germany

^kUniversità degli Studi della Campania "L. Vanvitelli", Dipartimento di Matematica e Fisica, Viale Lincoln 5, 81100 Caserta, Italy

^lIstituto Nazionale di Fisica Nucleare, Sezione di Padova, Via F. Marzolo 8, 35131 Padova, Italy

^mUniversità degli Studi di Padova, Via F. Marzolo 8, 35131 Padova, Italy

ⁿIstituto Nazionale di Fisica Nucleare, Sezione di Genova, Via Dodecaneso 33, 16146 Genova, Italy

^oUniversità degli Studi di Genova, Via Dodecaneso 33, 16146 Genova, Italy

^pInstitute for Nuclear Research of the Hungarian Academy of Sciences (MTA Atomki), PO Box 51, 4001 Debrecen, Hungary

^qUniversità degli Studi di Torino, Via P. Giuria 1, 10125 Torino, Italy

^rIstituto Nazionale di Fisica Nucleare, Sezione di Torino, Via P. Giuria 1, 10125 Torino, Italy

^sUniversità degli Studi di Milano, Via G. Celoria 16, 20133 Milano, Italy

^tIstituto Nazionale di Fisica Nucleare, Sezione di Milano, Via G. Celoria 16, 20133 Milano, Italy

^uIstituto Nazionale di Fisica Nucleare, Sezione di Roma, Piazzale A. Moro 2, 00185 Roma, Italy

^vKonkoly Observatory, Research Centre for Astronomy and Earth Sciences, Hungarian Academy of Sciences, 1121 Budapest, Hungary

^wTechnische Universität Dresden, Institut für Kern- und Teilchenphysik, Zellescher Weg 19, 01069 Dresden, Germany

^xINAF – Osservatorio Astronomico d'Abruzzo, Via Mentore Maggini, 64100 Teramo, Italy

^yDepartment of Physics, South Dakota School of Mines and Technology, 501E St Joseph Street, Rapid City, South Dakota 57701, USA

Abstract

The NeNa and the MgAl cycles play a fundamental role in the nucleosynthesis of asymptotic giant branch stars undergoing hot bottom burning. The $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction links these two cycles and a precise determination of its rate is required to correctly estimate the contribution of these stars to the chemical evolution of various isotopes of Na, Mg and Al. At temperatures of $50 \lesssim T \lesssim 110$ MK, narrow resonances at $E_p = 140$ and 251 keV are the main contributors to the reaction rate, in addition to the direct capture that dominates in the lower part of the temperature range. We present new measurements of the strengths of these resonances at the Laboratory for Underground Nuclear Astrophysics (LUNA). We have used two complementary detection approaches: high efficiency with a 4π BGO detector for the 140 keV resonance, and high resolution with a HPGe detector for the 251 keV resonance. Thanks to the reduced cosmic ray background of LUNA, we were able to determine the resonance strength of the 251 keV resonance as $\omega\gamma = 482(82) \mu\text{eV}$ and observed new gamma ray transitions for the decay of the corresponding state in ^{24}Mg at $E_x = 11931$ keV. With the highly efficient BGO detector, we observed a signal for the 140 keV resonance for the first time in a direct measurement, resulting in a strength of $\omega\gamma_{140} = 1.46^{+0.58}_{-0.53}$ neV (68% CL). Our measurement reduces the uncertainty of the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction rate in the temperature range from 0.05 to 0.1 GK to at most $^{+50\%}_{-35\%}$ at 0.07 GK. Accordingly, our results imply a significant reduction of the uncertainties in the nucleosynthesis calculations.

Keywords: Experimental Nuclear Astrophysics, Underground Nuclear Physics, Hydrogen Burning, Stellar Evolution

1. Introduction

Asymptotic Giant Branch (AGB) stars provide a major contribution to the synthesis of the elements in the cosmos and specifically to the chemical evolution of stellar clusters and galaxies. The more massive AGB stars ($M \gtrsim 4 M_{\odot}$) undergo proton captures at the base of the convective envelope (hot bottom burning, HBB) [1, 2]. When the convective envelope extends down to the H-burning shell, the ashes of the CNO, NeNa and MgAl cycles appear at the stellar surface. The coupling between convection and nuclear burning is responsible for the synthesis of various isotopes of Na, Mg and Al.

Combined, the uncertainties of the $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$ and the $^{22}\text{Ne}(p, \gamma)^{23}\text{Na}$ reaction rates account for the major nuclear contribution to the uncertainty of predicted ^{23}Na and ^{24}Mg yields from these massive AGB stars [3]. The $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$ reaction, in particular, is the bridge between the NeNa and MgAl cycles. The $^{22}\text{Ne}(p, \gamma)^{23}\text{Na}$ reaction has been subject of recent experimental studies [4, 5, 6, 7, 8]; prior studies of $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$ are described below.




Figure 1: Fractional contributions of the narrow resonances and the non-resonant cross section to the reaction rate at different stellar temperatures. The bands result from sampling the resonance parameters according to the parameters and procedures in [9, 10, 11].

During HBB, the temperatures at the bottom of the convective envelopes range between $50 \lesssim T \lesssim 110$ MK, increasing with the stellar mass and decreasing with the metallicity. The $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$ reaction rate in this temperature range is dominated by two resonances at $E_p = 140$ and 251 keV, where E_p is the proton beam energy in the laboratory system throughout this paper. At lower energies, the non-resonant capture contribution dominates the reaction rate, whereas another narrow resonance at

$E_p = 309$ keV is the principal component of the reaction rate at higher temperatures, up to $T = 1$ GK. The relative importance of each contribution for the reaction rate at different temperatures is illustrated in Fig. 1. The reaction rate calculations shown in this figure use the current best upper limit for the strength of the 140 keV resonance quoted in the literature [9] and treated as in [10]. Strengths of the 251 keV and 309 keV resonances have been measured by [12, 13, 14, 15]; and the calculations presented in Fig. 1 use the values as compiled in [16]. Information on the direct capture cross section was used as obtained indirectly in [17]. The nuclear properties of the resonances are summarized in Table 1.

Table 1: Nuclear resonance parameters for the $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$ reaction rate calculation as used in [9] and shown in Fig. 1. The upper limit for $\omega\gamma_{140\text{keV}}$ corresponds to 95% CL.

	140 keV	251 keV	309 keV
E_x (keV)	[18, 19], [17] 11827(4), 11830.7(15)	[18, 19] 11931.2(6)	[18, 19] 11987.72(10)
E_p (keV)	[18, 19], [17, 9] 140.2(42), 144.0(16)	[16] 250.9(2)	[16] 308.75(6)
Γ_{tot} (keV)		[16, 20] < 0.02	[16, 21] < 0.002
$\omega\gamma$ (eV)	[9] < 5.17×10^{-9}	[12, 16] $5.25(175) \times 10^{-4}$	[12, 16] $1.05(19) \times 10^{-1}$

The upper limit on the strength of the 140 keV resonance [9] does not exclude this resonance from being the dominant contribution to the reaction rate at $T \sim 80 - 90$ MK. In fact, the uncertainty of the $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$ reaction rate in the discussed temperature range, associated with the upper limit of the 140 keV resonance strength and the 33% relative uncertainty of the 251 keV resonance strength, provide a strong motivation for an experimental study of these resonances.

2. Methodology

2.1. Setup

The measurements were performed at the Laboratory for Underground Nuclear Astrophysics (LUNA), located at the Gran Sasso National Laboratory. Proton beams with energies of 130 to 400 keV were provided by the LUNA400 accelerator [22] and used to bombard water-cooled solid targets. Typical beam intensities of 100-250 μA on target were obtained.

Two complementary setups were used to study the cross section: one employed a High-Purity Germanium (HPGe) detector, the other a segmented Bismuth Germanium Oxide (BGO) summing detector. The use of two setups is advisable, as the efficiency of the HPGe detector is too

*Corresponding author

Email address: axel.boeltzig@gssi.infn.it (A. Boeltzig)

¹Permanent address: Istituto Nazionale di Fisica Nucleare, Sezione di Lecce, Via Arnesano, 73100 Lecce, Italy

²Current address: Physikalisch-Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany

low to study the weak 140 keV resonance. The energy res-
 olution of the HPGe detector, however, significantly facili-
 tates the study the gamma-ray branchings of the stronger
 resonances at 251 keV and 309 keV. Both setups benefit
 from the underground location of LUNA at the Gran Sasso
 National Laboratory, and target and detectors were sur-
 rounded by a massive lead shielding to reduce background
 from environmental radiation [23, 24, 25].

2.2. BGO Detector Phase: 140 keV Resonance

The segmented BGO summing detector was used to
 search for the 140 keV resonance with a large detection
 efficiency, utilizing a digital data acquisition system to ac-
 quire independent spectra of the six individual crystals
 and generating a sum energy spectrum offline [25, 26,
 and references therein]. The response of the BGO de-
 tector was estimated through a Monte Carlo simulation
 using Geant4 [27, 26], which was validated with calibra-
 tion measurements. The efficiency for the detection of
 the full gamma-ray energy depends on the gamma-ray
 branching ratios and the low-energy threshold of the in-
 dividual crystal segment. In the case of the 309 keV re-
 sonance, for example, the detection efficiency above 10 MeV
 is about 50% [25]; the efficiency of this detector is fur-
 ther discussed in subsection 3.2. In the search for the
 $E_p = 140$ keV resonance, measurements were taken at
 beam energies around 130, 140, 145 and 147 keV, in view of
 the uncertainty of the resonance energy (see Table 1). Due
 to a narrow resonance in $^{18}\text{O}(p, \gamma)^{19}\text{F}$ ($Q = 7994$ keV) at
 $E_p = 151$ keV, the background from random coincidence
 summing (pile-up) of gamma rays from this reaction pro-
 hibited the $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$ resonance search for beam en-
 ergies above 151 keV. At energies of 140 keV and below
 we observed yields compatible with zero within the sta-
 tistical uncertainty. For the runs at 147 keV ($147 \text{ keV} \lesssim$
 $E_p \lesssim 149$ keV) we observe a statistically significant excess
 of counts over the beam-induced background. This beam
 energy is compatible with the resonance energy in litera-
 ture, and we analyzed this data set considering these runs
 as on-resonance, taking into account the target thickness
 (described in sec. 2.4).

2.3. HPGe Detector Phase: 251 keV and 309 keV Resonances

The HPGe detector was employed in close geometry
 at an angle of 55° to study the resonance strengths and
 gamma-ray branchings of the resonances at 251 keV and
 309 keV (cf. [28, and references therein]). The efficiency
 calibration was based on radioactive sources (^{137}Cs , ^{60}Co)
 and gamma rays from narrow resonances at $E_p = 326$ keV
 in $^{27}\text{Al}(p, \gamma)^{28}\text{Si}$ and at $E_p = 278$ keV in $^{14}\text{N}(p, \gamma)^{15}\text{O}$. An
 empirical calibration curve was fitted to the total and full
 energy peak efficiency, using the parametrization in [29].
 Summing effects that arise from the coincident detection of
 multiple gamma rays of the same cascade (enhanced by the
 close detector geometry) were taken into account, based

on the parametrized efficiency curve; angular correlations
 were neglected for the calculation of the summing effects.
 The parameters of the efficiency model were determined in
 a global maximum likelihood fit. We considered an uncer-
 tainty of 8% for the efficiency calibration curve, taking into
 account the scatter of the calibration data points, the re-
 peatability of the (p, γ) calibration measurements, and the
 few available sources to calibrate the efficiency at gamma
 energies above 8 MeV.

2.4. Targets

All targets were produced by evaporation of sodium
 compounds onto a high purity tantalum backing at the
 Institute for Nuclear Research, Hungarian Academy of Sci-
 ences (MTA Atomki). The backing was chemically cleaned
 and heated in vacuum before the evaporation. We tested
 several sodium compounds (NaCl , Na_2SiO_3 , Na_2WO_4),
 and obtained the best results (i. e., reproducible target
 properties and stability under beam) with Na_2WO_4 (as
 in [9]). In a test evaporation of Na_2WO_4 onto a carbon
 backing we observed no major deviations in the stoichiom-
 etry of the sodium compound with respect to the nominal
 composition in a Rutherford backscattering measurement.
 Materials supplied by different vendors were tested in an
 effort to reduce beam-induced backgrounds from light con-
 taminants, and the cleanest material was chosen for target
 production.

Typical target thicknesses for the Na_2WO_4 targets were
 $50 \mu\text{g}/\text{cm}^2$ (beam energy loss of about 14 keV at 150 keV
 proton energy). Between experimental runs we regularly
 performed measurements of the target thickness by means
 of nuclear resonance analysis, scanning over the 309 keV
 resonance in $^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$. Targets were replaced when
 the target thickness (width of the plateau in a 309 keV
 resonance scan) had decreased strongly, or a drop in on-
 resonance yield was observed. An example for the devel-
 opment of the target scan under bombardment is shown
 in Fig. 2. Typical accumulated charges on a single target
 ranged between 10 and 30 C.

Elastic Recoil Detection Analysis (ERDA) was per-
 formed on selected targets at the Ion Beam Center at the
 Helmholtz-Center Dresden-Rossendorf, using a 43 MeV
 Cl^{7+} beam and detecting the recoils at an angle of 31°
 (cf. [30] for a description of the setup). Analysis of the
 data with the program NDF [31] resulted in depth pro-
 file information and allowed for the study of the chemi-
 cal composition of various targets during the initial target
 development phase. Furthermore, two targets were pro-
 duced under identical conditions: one was used to measure
 the 251 keV and 309 keV resonances with the HPGe de-
 tector setup at LUNA, the other was not bombarded. Both
 of these targets were also analyzed with ERDA to study
 the chemical composition for the analysis of the HPGe
 measurements. Depth profiles were obtained and the sto-
 ichiometry of Na, W, O and H in the main target layer
 resulted in effective stopping powers that were on the or-
 der of 10% larger compared to the nominal composition




Figure 2: Yield curves showing scans of the narrow resonance at 309 keV for the same target after different amounts of charge accumulated on this target.

of pure Na_2WO_4 . Traces of carbon were detected in the bulk of the samples, but the amount was irrelevant in the context of our measurement.

3. Analysis and Results

3.1. HPGe Detector Phase: 251 keV and 309 keV Resonances

With the effective stopping power of the material at the resonance energy, ε_{eff} , the absolute strength $\omega\gamma$ of the narrow resonances can be determined through the well-known relation $Y = \frac{\lambda^2}{2} \frac{\omega\gamma}{\varepsilon_{\text{eff}}}$ [32], where Y is the reaction yield (number of reactions per incident projectile), and λ the de Broglie wavelength of the projectile. Stopping power tables were generated using SRIM 2013.00 [33].

Branching ratios and resonance strengths were determined solving the system of equations: $N_i = \sum_j (B_{ij} \cdot P_j) \cdot Y \cdot \frac{Q}{q_p}$, where the vector elements N_i are the number of counts in the primary gamma-ray peaks, and the matrix elements B_{ij} correspond to the probability that a decay through the primary branch i is registered in the primary peak j (including detection efficiency and summing effects). The vector elements P_i are the primary branching ratios that fulfill the normalization condition $\sum P_i = 1$ and Q and q_p are the accumulated charge on target and the charge of a single projectile (proton), respectively.

When calculating the summing effects, gamma-ray branching ratios for secondary transitions were taken from literature [18]. The uncertainties of Y and the P_i were determined by a Monte Carlo approach, repeating the analysis after varying the N_i within the respective experimental uncertainties and the secondary branching ratios within their uncertainties given in literature. The uncertainty of ε_{eff} , derived from the uncertainty of the chemical composition of the target as determined with ERDA, typically amounted to 15% and was the dominant contribution to

the resonance strength uncertainty. A systematic uncertainty of 3% was assumed for the measurement of the accumulated charge on target.

Among the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ resonances in Table 1, the strength of that at $E_p = 309$ keV is the best known in literature [16]. To verify our results, we redetermined the resonance strength and gamma-ray branching ratios as described above. The branching ratios of the 12 primary transitions in literature [18] were found to be in agreement with our measurement within the experimental uncertainties. The height of the plateau of two resonance scans on target and the composition of the not bombarded target as determined by ERDA resulted in a resonance strength of $\omega\gamma_{309} = 108(19)$ meV, in good agreement with the literature value of $105(19)$ meV [16].

The strength of the resonance at $E_p = 251$ keV is known with a larger uncertainty of about 33% (Table 1). Additionally, the gamma-ray branchings of the corresponding level in ^{24}Mg are not given in current compilations (such as [19]), and the only publication of these branching ratios known to the authors indicates only two strong transitions from this level [34]. A scan of the 251 keV resonance, followed by a long run ($Q \approx 11$ C) at an energy on the resonance, was used to determine the strength of this resonance and the gamma-ray branching ratios of the corresponding level in ^{24}Mg . The most intense transitions are marked in the spectrum in Fig. 3 and the resulting branching ratios are given in Table 2. In addition to the two major transitions described in [34], we identified multiple minor transitions and determined their branching ratios. The intensities of several minor peaks in the spectrum, most prominently the transition to the ground state of ^{24}Mg , were consistent with the expectation from true coincidence summing (summing-in) alone and are thus not included in Table 2.

Table 2: Branching ratios (in percent) for the state at $E_x = 11931$ keV in ^{24}Mg , populated through $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ at $E_p = 251$ keV, to the levels E_f .

E_f (keV)	9300	8864	7812	7349	6011	5235	4238	4123	1369
[34]	-	-	-	-	-	-	-	11	89
this	0.14	0.25	1.05	1.21	3.21	0.95	0.38	18.2	74.6
work	± 0.05	0.05	0.12	0.14	0.34	0.16	0.14	1.8	7.5

With the yield of the long run on the resonance plateau and using the same target stoichiometry as in the analysis of the 309 keV resonance, we determined a resonance strength of $\omega\gamma_{251} = 482(82)$ μeV , in agreement with the literature value but with an improved relative uncertainty of 17% (previously 33% [16]) that is now comparable to that of the 309 keV resonance.

3.2. BGO Detector Phase: 140 keV Resonance

During the search for the 140 keV resonance with the BGO detector, the 309 keV resonance was used as a reference to account for the influence of target composition




Figure 3: HPGe spectrum acquired at $E_p = 255$ keV, $Q = 11.4$ C, indicating the major primary transitions from the 11931 keV state in ^{24}Mg . Dotted lines mark gamma rays and escape peaks from secondary transitions. The line at an energy corresponding to the ground state transition has been identified as caused purely by summing-in (see text for details).




Figure 4: Comparison of sum energy spectra for a target made from unfavorable raw material (large beam-induced background, blue) and an example of an analyzed spectrum taken with a cleaner target (red, 147 keV, 9.0 C). The modeled background contributions are shown to demonstrate the influence of $^7\text{Li}(p,\gamma)^8\text{Be}$ and $^{11}\text{B}(p,\gamma)^{12}\text{C}$ on the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ ROI. The inset shows the result (best fit) of a sum energy spectrum for one of the possible branching ratios of $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$, with the binning used for the Bayesian analysis. See text for details.

variations on the yield. The yields Y and the strengths $\omega\gamma$ of the two resonances are related through $\frac{\omega\gamma_{140}}{\omega\gamma_{309}} = \frac{Y_{140}}{Y_{309}} \cdot \frac{\lambda_{309}^2}{\lambda_{140}^2} \cdot \frac{\varepsilon_{\text{eff},140}}{\varepsilon_{\text{eff},309}}$. Whilst the effective stopping power can vary considerably with changes in stoichiometry, the ratio of effective stopping powers at the two resonance energies is much more robust (with $< 1\%$ variation in this case).

We found two beam-induced backgrounds that were critical for the observation of the 140 keV resonance in our measurements: $^7\text{Li}(p,\gamma)^8\text{Be}$ with $Q = 17.25$ MeV, and $^{11}\text{B}(p,\gamma)^{12}\text{C}$ with $Q = 15.96$ MeV. With the Q -values of both background reactions larger than that of $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ ($Q = 11.69$ MeV), these reactions can contribute to the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ region of interest (ROI) in the sum energy spectrum. The contribution of these backgrounds has been modeled as shown in Fig. 4. To illustrate the potential influence of these impurities, we compare the expected yields: for a proton beam with an energy of 147 keV impinging on a $60 \mu\text{g}/\text{cm}^2$ Na_2WO_4 target, the yield of a narrow $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ resonance with an $\omega\gamma$ at the current upper limit [9] is 15 reactions/C. The same reaction yield is obtained for $^{11}\text{B}(p,\gamma)^{12}\text{C}$ by a contamination of about 2500 ppm of boron, or for $^7\text{Li}(p,\gamma)^8\text{Be}$ by about 700 ppm of lithium.

Before and after each run in search of the 140 keV resonance, we scanned the 309 keV resonance to obtain a target profile and used the known gamma rays of the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction to constrain the energy calibration of the individual detector segments. Slightly non-linear energy responses from the detector segments (especially at the higher gamma ray energies produced by the beam-induced backgrounds) were allowed by introducing a quadratic term in the energy calibration function. Using the energy calibrations of the individual crystals, we created the sum spectrum.

Using the sum spectrum at energies above the total gamma-ray energy expected for the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction, we obtained constraints on the background-contributions from ^7Li and ^{11}B to the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ ROI. A Bayesian approach using Poisson-likelihood for the bin contents in this energy region (1 MeV wide bins between 10 MeV and 18 MeV) and uniform prior distributions for the yields of the background reactions results in a posterior probability distribution for the resonance strength, based on the excess of the number of counts in the ROI for the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction over the contribution from the two beam-induced backgrounds (cf. inset of Fig. 4). The Bayesian Analysis Toolkit (BAT) [35] was used to calculate the posterior probability density function.

The relation between an excess over background in the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction ROI and the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction resonance strength requires knowledge of the detection efficiency, i.e., the probability of an $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ event to cause a detected energy deposition in the corresponding ROI. This efficiency depends on the gamma-ray branchings, and spin and parity of the resonant level, which are unknown. We considered levels in ^{24}Mg with known branching ratios and E_x above 9 MeV, and simulated the corresponding detection efficiencies in the ROI around the respective E_x . This procedure corresponds to the approach in [9], except that all levels above 9 MeV were considered in this work, covering a wider range of possible branching ratios than the states selected in [9] and resulting in a larger detection efficiency uncertainty by comparison. Random sampling from the efficiencies corresponding to the branching ratios of different states was applied to include the effect of this uncertainty on the results for the resonance strength.

Different runs were performed within each chosen beam

energy group. The posterior distributions for the resonance strength of the individual runs were combined to obtain a final posterior distribution. The posterior distribution obtained in the analysis for runs with $147 \text{ keV} \lesssim E_p \lesssim 149 \text{ keV}$ is shown in Fig. 5. The posterior distribution has a mode at 1.46 neV , and the shortest interval to contain 68% probability ranges from 0.94 and 2.05 neV (95% interval from 0.51 to 2.74 neV).




Figure 5: Posterior probability density function for the strength of the 140 keV resonance.

Table 3: Resonance strengths for $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ (in eV) of this work, compared to previous literature values. All uncertainties are given at 68% CL.

140 keV	251 keV	309 keV
[9]	[12, 16]	[12, 16]
$2.15(129) \times 10^{-9} *$	$5.25(175) \times 10^{-4}$	$1.05(19) \times 10^{-1}$
this work	this work	this work
$1.46^{+0.58}_{-0.53} \times 10^{-9}$	$4.82(82) \times 10^{-4}$	$1.08(19) \times 10^{-1}$

* when interpreting result of [9] as a positive detection; corresponds to an upper limit of $< 5.17 \text{ neV}$ at 95% CL otherwise.

4. Reaction Rate and Astrophysical Implications

To summarize our results, the obtained strengths for the 309, 251 and 140 keV resonances are shown in Table 3. The value for $\omega\gamma_{309}$ is in good agreement with the current literature value and has a comparable uncertainty. The $\omega\gamma_{251}$ has a reduced relative uncertainty of 17% (previously: 33% [16]). Moreover, we have found previously unobserved decay branches for the corresponding state in ^{24}Mg . We have directly observed the 140 keV resonance with an excess over zero of more than 2σ for the first time. The new experimentally determined value for the strength of the 140 keV resonance is compatible with the result of [9] when interpreted as a positive detection, but we established it as a value different from zero with a much

larger statistical significance and a relative uncertainty of 38%.

With these results we calculated the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction rate and determined its uncertainty by a Monte Carlo approach. Resonance strengths for $\omega\gamma_{309}$ and $\omega\gamma_{251}$ were sampled from Gaussian distributions, $\omega\gamma_{140}$ was sampled from the posterior distribution obtained in this measurement as discussed above. The direct capture strength and resonance energies were sampled as described in [11, 10]. The resulting reaction rate, compared to the rates in [9], is shown in Fig. 6. The different treatment of the uncertainty of $\omega\gamma_{140}$ in this work and the upper limit in [9] should be noted for this comparison.




Figure 6: Reaction rate and uncertainty (68% CL) based on the results presented in this work, compared to the reaction rate in [9]. Refer to the text for details on the different treatment of the uncertainty of the 140 keV resonance.

The relative uncertainty of the reaction rate is decreased in the temperature window $T = 0.05 - 0.1 \text{ GK}$, thanks to the improved knowledge on the 140 keV resonance and the 251 keV resonance, with uncertainties between $+50\%$ at 0.07 GK and -14% at 0.1 GK . At $T = 0.06 - 0.09 \text{ GK}$ the reaction rate based on the results of this work is slightly higher than previously determined, because of the larger median value of $\omega\gamma_{140}$. Below $T = 0.05 \text{ GK}$ and above $T = 0.2 \text{ GK}$, the reaction rate is dominated by the direct capture contribution and the 309 keV resonance (plus higher resonances), respectively, and is therefore unaffected by this work.

The main limiting factors for the determination of $\omega\gamma_{140}$ were the beam-induced backgrounds from traces of ^7Li and ^{11}B , the uncertainty of the resonance energy, and the missing branching ratios for the resonant level.




Figure 7: Network calculations to follow ^{20}Ne , ^{23}Na and ^{24}Mg mass fractions, using the reaction rates of [9] (top) and this work (bottom) at $T = 80$ MK. See text for details.

To understand the potential impact of our results, we performed a network calculation for a constant $T = 80$ MK and $\rho = 10$ g/cm³. The outcome of this calculation is shown in Fig. 7. The network includes all the isotopes of the NeNa cycle plus ^{24}Mg . The H and He mass fractions are kept constant, at $X(\text{H}) = 0.5$ and $X(^4\text{He}) = 0.499$ to simulate continuous replenishment via convection. The initial mass fractions of Ne and Na isotopes are scaled to solar abundances, with $X(\text{Ne}) + X(\text{Na}) = 0.001$, and initially $X(^{24}\text{Mg}) = 0$. The calculations cover 3×10^3 yr, the typical time elapsed between two subsequent thermal pulses (instabilities of the He-burning shell) in an AGB star with initial mass $M = 7 M_{\odot}$. Except for $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$, reaction rates were taken from StarLib (version 6) [36], which includes the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ rates of [9]. For $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$ we used the most recent reaction rate from [8] (which is similar to the rate of [7]). The results of these calculations are shown in the upper panel of Fig. 7. The lower panel shows the results of the same calculation when using the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ rate based on this work.

The Ne and Na isotopes almost instantaneously reach their equilibrium abundances, and their abundance ratios, e.g. the $^{23}\text{Na}/^{20}\text{Ne}$ ratio, remain constant as they only depend on the temperature and the choice of the reaction rates. The continuous depletion of the total abundance of the Ne and Na isotopes is driven by the leakage of the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction. As a consequence the ^{24}Mg piles up. The shaded areas in Fig. 7 represent the influence of the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction rate on the three most abundant Ne, Na, and Mg isotopes during this nucleosynthesis episode. The reduction of the uncertainties when passing from the previous [9] and the new rate is evident. This

result will improve future calculations of the yields of massive AGB stars, providing, in particular, a more accurate understanding of the evolution of the Na and Mg isotopes in galaxies and stellar clusters.

Acknowledgements

This work was supported by INFN. The authors acknowledge the support at the Gran Sasso National Laboratory, and would like to thank D. Ciccotti and L. Roscilli as well as to the mechanical workshops of the INFN sections of LNGS and Naples for their technical support. Financial support of this work was granted by the Helmholtz Association (VH-VI 417 and ERC-RA-0016), by the NK-FIH (grant K120666), by the Deutsche Forschungsgemeinschaft (BE 4100/4-1), by the Science and Technology Facilities Council (STFC), by the National Science Foundation (PHY-1713857, and PHY-1430152: JINA Center for the Evolution of the Elements), and through the STAR program of the University of Naples “Federico II” and the Compagnia di San Paolo.

- [1] M. Forestini, C. Charbonnel, Nucleosynthesis of light elements inside thermally pulsing AGB stars: I. The case of intermediate-mass stars, *Astronomy and Astrophysics Supplement Series* 123 (1997) 241–272. arXiv:astro-ph/9608153, doi:10.1051/aas:1997348.
- [2] J. C. Lattanzio, C. A. Frost, R. C. Cannon, P. R. Wood, Hot Bottom Burning Nucleosynthesis in $6 M_{\text{Solar}}$ Stellar Models, *Nuclear Physics A* 621 (1997) 435–438. doi:10.1016/S0375-9474(97)00286-8.
- [3] R. G. Izzard, M. Lugaro, A. I. Karakas, C. Iliadis, M. van Raaij, Reaction rate uncertainties and the operation of the NeNa and MgAl chains during HBB in intermediate-mass AGB stars, *Astronomy and Astrophysics* 466 (2) (2007) 641–648. doi:10.1051/0004-6361:20066903.
- [4] F. Cavanna, R. Depalo, M. Aliotta, M. Anders, D. Bemmerer, A. Best, A. Boeltzig, C. Brogгинi, C. G. Bruno, A. Cacioli, P. Corvisiero, T. Davinson, A. di Leva, Z. Elekes, F. Ferraro, A. Formicola, Z. Fülöp, G. Gervino, A. Guglielmetti, C. Gustavino, G. Gyürky, G. Imbriani, M. Junker, R. Menegazzo, V. Mossa, F. R. Pantaleo, P. Prati, D. A. Scott, E. Somorjai, O. Straniero, F. Strieder, T. Szücs, M. P. Takács, D. Trezzi, The LUNA Collaboration, Three New Low-Energy Resonances in the $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$ Reaction, *Physical Review Letters* 115 (25). doi:10.1103/PhysRevLett.115.252501.
- [5] F. Cavanna, R. Depalo, M. Aliotta, M. Anders, D. Bemmerer, A. Best, A. Boeltzig, C. Broggini, C. G. Bruno, A. Cacioli, P. Corvisiero, T. Davinson, A. di Leva, Z. Elekes, F. Ferraro, A. Formicola, Z. Fülöp, G. Gervino, A. Guglielmetti, C. Gustavino, G. Gyürky, G. Imbriani, M. Junker, R. Menegazzo, V. Mossa, F. R. Pantaleo, P. Prati, D. A. Scott, E. Somorjai, O. Straniero, F. Strieder, T. Szücs, M. P. Takács, D. Trezzi, The LUNA Collaboration, Erratum: Three New Low-Energy Resonances in the $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$ Reaction [Phys. Rev. Lett. 115, 252501 (2015)], *Physical Review Letters* 120 (23) (2018) 239901. doi:10.1103/PhysRevLett.120.239901.
- [6] R. Depalo, F. Cavanna, M. Aliotta, M. Anders, D. Bemmerer, A. Best, A. Boeltzig, C. Broggini, C. G. Bruno, A. Cacioli, G. F. Ciani, P. Corvisiero, T. Davinson, A. Di Leva, Z. Elekes, F. Ferraro, A. Formicola, Z. Fülöp, G. Gervino, A. Guglielmetti, C. Gustavino, G. Gyürky, G. Imbriani, M. Junker, R. Menegazzo, V. Mossa, F. R. Pantaleo, D. Piatti, P. Prati, O. Straniero, T. Szücs, M. P. Takács, D. Trezzi, LUNA Collaboration, Direct measurement of low-energy $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$ resonances, *Physical Review C* 94 (5). doi:10.1103/PhysRevC.94.055804.

- [7] K. J. Kelly, A. E. Champagne, L. N. Downen, J. R. Dermigny,⁵⁴⁶ S. Hunt, C. Iliadis, A. L. Cooper, New measurements of low-⁵⁴⁷ energy resonances in the $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$ reaction, *Physical Review C* 95 (1) (2017) 015806. doi:10.1103/PhysRevC.95.015806.⁵⁴⁹
- [8] F. Ferraro, M. P. Takács, D. Piatti, F. Cavanna, R. Depalo,⁵⁵⁰ M. Aliotta, D. Bemmerer, A. Best, A. Boeltzig, C. Broggini,⁵⁵¹ C. G. Bruno, A. Cacioli, T. Chillery, G. F. Ciani, P. Corvisiero,⁵⁵² T. Davinson, G. D'Erasmio, A. Di Leva, Z. Elekes, E. M.⁵⁵³ Fiore, A. Formicola, Z. Fülöp, G. Gervino, A. Guglielmetti,⁵⁵⁴ C. Gustavino, G. Gyürky, G. Imbriani, M. Junker, A. Karakas,⁵⁵⁵ I. Kochanek, M. Lugaro, P. Marigo, R. Menegazzo, V. Mossa,⁵⁵⁶ F. R. Pantaleo, V. Paticchio, R. Perrino, P. Prati, L. Schi-⁵⁵⁷avulli, K. Stöckel, O. Straniero, T. Szücs, D. Trezzi, S. Za-⁵⁵⁸vatarelli, LUNA Collaboration, Direct Capture Cross Section⁵⁵⁹ and the $E_p = 71$ and 105 keV Resonances in the $^{22}\text{Ne}(p,\gamma)^{23}\text{Na}$ ⁵⁶⁰ Reaction, *Physical Review Letters* 121 (17) (2018) 172701. doi:10.1103/PhysRevLett.121.172701.⁵⁶¹
- [9] J. M. Cesaratto, A. E. Champagne, M. Q. Buckner, T. B.⁵⁶² Clegg, S. Daigle, C. Howard, C. Iliadis, R. Longland, J. R.⁵⁶⁴ Newton, B. M. Oginni, Measurement of the $E_r^{\text{c.m.}} = 138$ keV⁵⁶⁵ resonance in the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ reaction and the abundance of⁵⁶⁶ sodium in AGB stars, *Physical Review C* 88 (6) (2013) 065806. doi:10.1103/PhysRevC.88.065806.⁵⁶⁷
- [10] C. Iliadis, R. Longland, A. Champagne, A. Coc, Charged-⁵⁶⁹ particle thermonuclear reaction rates: III. Nuclear physics⁵⁷⁰ input, *Nuclear Physics A* 841 (1-4) (2010) 251–322. doi:10.1016/j.nuclphysa.2010.04.010.⁵⁷¹
- [11] R. Longland, C. Iliadis, A. Champagne, J. Newton, C. Ugalde,⁵⁷³ A. Coc, R. Fitzgerald, Charged-particle thermonuclear re-⁵⁷⁴ action rates: I. Monte Carlo method and statistical dis-⁵⁷⁵ tributions, *Nuclear Physics A* 841 (1-4) (2010) 1–30. doi:10.1016/j.nuclphysa.2010.04.008.⁵⁷⁶
- [12] Z. Switkowski, R. O'Brien, A. Smith, D. Sargood, Total Yield⁵⁷⁸ Measurements in $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$, *Australian Journal of Physics*⁵⁷⁹ 28 (2) (1975) 141. doi:10.1071/PH750141.⁵⁸⁰
- [13] B. M. Paine, S. R. Kennett, D. G. Sargood, (p, γ) resonance⁵⁸¹ strengths in the s-d shell, *Physical Review C* 17 (5) (1978) 1550–⁵⁸² 1554. doi:10.1103/PhysRevC.17.1550.⁵⁸³
- [14] B. Paine, D. Sargood, (p, γ) Resonance strengths in the s-d shell,⁵⁸⁴ *Nuclear Physics A* 331 (2) (1979) 389–400. doi:10.1016/0375-⁵⁸⁵ 9474(79)90349-X.⁵⁸⁶
- [15] J. Görres, M. Wiescher, C. Rolfs, Hydrogen burning of ^{23}Na ⁵⁸⁷ in the NeNa cycle, *The Astrophysical Journal* 343 (1989) 365. doi:10.1086/167710.⁵⁸⁸
- [16] P. Endt, Energy levels of $A = 21$ -44 nuclei (VII), *Nuclear*⁵⁹⁰ *Physics A* 521 (1990) 1–400. doi:10.1016/0375-9474(90)90598-⁵⁹¹ G.⁵⁹²
- [17] S. Hale, A. Champagne, C. Iliadis, V. Hansper, D. Pow-⁵⁹³ ell, J. Blackmon, Investigation of the $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ and⁵⁹⁴ $^{23}\text{Na}(p,\alpha)^{20}\text{Ne}$ reactions via ($^3\text{He},d$) spectroscopy, *Physical Re-⁵⁹⁵ view C* 70 (4) (2004) 045802. doi:10.1103/PhysRevC.70.045802.⁵⁹⁶
- [18] National Nuclear Data Center, information extracted from the⁵⁹⁷ NuDat 2 database, <http://www.nndc.bnl.gov/nudat2/> (2017).⁵⁹⁸
- [19] R. Firestone, *Nuclear Data Sheets for $A = 24$,*⁵⁹⁹ *Nuclear Data Sheets* 108 (11) (2007) 2319–2392. doi:10.1016/j.nds.2007.10.001.⁶⁰⁰
- [20] S. Wagner, M. Heitzmann, Resonanzenergien für den Pro-⁶⁰² toneneinfang II. Natrium, *Zeitschrift für Naturforschung A*⁶⁰³ 15 (1). doi:10.1515/zna-1960-0111.⁶⁰⁴
- [21] S. Wüstenbecker, H. Ebbing, W. Schulte, H. Baumeister,⁶⁰⁵ H. Becker, B. Cleff, C. Rolfs, H. Trautvetter, G. Mitchell,⁶⁰⁶ J. Schweitzer, C. Peterson, Improvements in targetry and high⁶⁰⁷ voltage stability for high resolution ion beam experiments, *Nu-⁶⁰⁸ clear Instruments and Methods in Physics Research Section A:*⁶⁰⁹ *Accelerators, Spectrometers, Detectors and Associated Equip-⁶¹⁰ ment* 279 (3) (1989) 448–466. doi:10.1016/0168-9002(89)91292-⁶¹¹ 8.⁶¹²
- [22] A. Formicola, G. Imbriani, M. Junker, D. Bemmerer,⁶¹³ R. Bonetti, C. Broggini, C. Casella, P. Corvisiero, H. Costan-⁶¹⁴ tini, G. Gervino, C. Gustavino, A. Lemut, P. Prati, V. Roca,⁶¹⁵ C. Rolfs, M. Romano, D. Schürmann, F. Strieder, F. Terrasi,⁶¹⁶ H.-P. Trautvetter, S. Zavatarelli, The LUNA II 400 kV accel-
erator, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* 507 (3) (2003) 609–616.
- [23] H. Costantini, A. Formicola, G. Imbriani, M. Junker, C. Rolfs,
F. Strieder, LUNA: a laboratory for underground nuclear astro-
physics, *Reports on Progress in Physics* 72 (8) (2009) 086301. doi:10.1088/0034-4885/72/8/086301.
- [24] D. Bemmerer, F. Confortola, A. Lemut, R. Bonetti, C. Broggini,
P. Corvisiero, H. Costantini, J. Cruz, A. Formicola, Z. Fülöp,
G. Gervino, A. Guglielmetti, C. Gustavino, G. Gyürky, G. Im-
briani, A. P. Jesus, M. Junker, B. Limata, R. Menegazzo,
P. Prati, V. Roca, D. Rogalla, C. Rolfs, M. Romano,
C. Rossi Alvarez, F. Schümann, E. Somorjai, O. Straniero,
F. Strieder, F. Terrasi, H. P. Trautvetter, A. Vomiero, Feasibil-
ity of low-energy radiative-capture experiments at the LUNA
underground accelerator facility, *The European Physical Jour-
nal A* 24 (2) (2005) 313–319. doi:10.1140/epja/i2004-10135-4.
- [25] A. Boeltzig, A. Best, G. Imbriani, M. Junker, M. Aliotta, D. Bem-
merer, C. Broggini, C. G. Bruno, R. Buompane, A. Cacioli,
F. Cavanna, T. Chillery, G. F. Ciani, P. Corvisiero, L. Csedreki,
T. Davinson, R. J. deBoer, R. Depalo, A. Di Leva, Z. Elekes,
F. Ferraro, E. M. Fiore, A. Formicola, Z. Fülöp, G. Gervino,
A. Guglielmetti, C. Gustavino, G. Gyürky, I. Kochanek,
R. Menegazzo, V. Mossa, F. R. Pantaleo, V. Paticchio, R. Per-
rino, D. Piatti, P. Prati, L. Schiavulli, K. Stöckel, O. Straniero,
F. Strieder, T. Szücs, M. P. Takács, D. Trezzi, M. Wiescher,
S. Zavatarelli, Improved background suppression for radiative
capture reactions at LUNA with HPGe and BGO detectors,
Journal of Physics G: Nuclear and Particle Physics 45 (2) (2018)
025203. doi:10.1088/1361-6471/aaa163.
- [26] F. Strieder, B. Limata, A. Formicola, G. Imbriani, M. Junker,
D. Bemmerer, A. Best, C. Broggini, A. Cacioli, P. Corvisiero,
H. Costantini, A. DiLeva, Z. Elekes, Z. Fülöp, G. Gervino,
A. Guglielmetti, C. Gustavino, G. Gyürky, A. Lemut, M. Marta,
C. Mazzocchi, R. Menegazzo, P. Prati, V. Roca, C. Rolfs,
C. Rossi Alvarez, E. Somorjai, O. Straniero, F. Terrasi,
H. Trautvetter, The $^{25}\text{Mg}(p,\gamma)^{26}\text{Al}$ reaction at low astro-
physical energies, *Physics Letters B* 707 (1) (2012) 60–65. doi:10.1016/j.physletb.2011.12.029.
- [27] J. Allison, K. Amako, J. Apostolakis, P. Arce, M. Asai, T. Aso,
E. Bagli, A. Bagulya, S. Banerjee, G. Barrand, B. Beck, A. Bog-
danov, D. Brandt, J. Brown, H. Burkhardt, P. Canal, D. Cano-
Ott, S. Chauvie, K. Cho, G. Cirrone, G. Cooperman, M. Corts-
Giraldo, G. Cosmo, G. Cuttone, G. Depaola, L. Desorgher,
X. Dong, A. Dotti, V. Elvira, G. Folger, Z. Francis, A. Gal-
loyan, L. Garnier, M. Gayer, K. Genser, V. Grichine, S. Guatelli,
P. Guye, P. Gumplinger, A. Howard, I. Hivnov, S. Hwang,
S. Incerti, A. Ivanchenko, V. Ivanchenko, F. Jones, S. Jun,
P. Kaitaniemi, N. Karakatsanis, M. Karamitros, M. Kelsey,
A. Kimura, T. Koi, H. Kurashige, A. Lechner, S. Lee, F. Longo,
M. Maire, D. Mancusi, A. Mantero, E. Mendoza, B. Morgan,
K. Murakami, T. Nikitina, L. Pandola, P. Paprocki, J. Perl,
I. Petrovi, M. Pia, W. Pokorski, J. Quesada, M. Raine, M. Reis,
A. Ribon, A. Risti Fira, F. Romano, G. Russo, G. Santin,
T. Sasaki, D. Sawkey, J. Shin, I. Strakovsky, A. Taborda,
S. Tanaka, B. Tom, T. Toshito, H. Tran, P. Truscott, L. Ur-
ban, V. Uzhinsky, J. Verbeke, M. Verderi, B. Wendt, H. Wenzel,
D. Wright, D. Wright, T. Yamashita, J. Yarba, H. Yoshida, Re-
cent developments in Geant 4, *Nuclear Instruments and Meth-
ods in Physics Research Section A: Accelerators, Spectrome-
ters, Detectors and Associated Equipment* 835 (2016) 186–225. doi:10.1016/j.nima.2016.06.125.
- [28] A. Di Leva, D. A. Scott, A. Cacioli, A. Formicola,
F. Strieder, M. Aliotta, M. Anders, D. Bemmerer, C. Broggini,
P. Corvisiero, Z. Elekes, Z. Fülöp, G. Gervino, A. Guglielmetti,
C. Gustavino, G. Gyürky, G. Imbriani, J. José, M. Junker,
M. Laubenstein, R. Menegazzo, E. Napolitani, P. Prati,
V. Rigato, V. Roca, E. Somorjai, C. Salvo, O. Straniero,
T. Szücs, F. Terrasi, D. Trezzi, Underground study of the
 $^{17}\text{O}(p,\gamma)^{18}\text{F}$ reaction relevant for explosive hydrogen burning,

- 617 Physical Review C 89 (1). doi:10.1103/PhysRevC.89.015803.
- 618 [29] G. Imbriani, H. Costantini, A. Formicola, A. Vomiero, C. Angulo,
619 D. Bemmerer, R. Bonetti, C. Brogini, F. Confortola,
620 P. Corvisiero, J. Cruz, P. Descouvemont, Z. Fülöp, G. Gervino,
621 A. Guglielmetti, C. Gustavino, G. Gyürky, A. P. Jesus,
622 M. Junker, J. N. Klug, A. Lemut, R. Menegazzo, P. Prati,
623 V. Roca, C. Rolfs, M. Romano, C. Rossi-Alvarez, F. Schümann,
624 D. Schürmann, E. Somorjai, O. Straniero, F. Strieder, F. Ter-
625 rasi, H. P. Trautvetter, S-factor of $^{14}\text{N}(p,\gamma)^{15}\text{O}$ at astrophysical
626 energies, *The European Physical Journal A* 25 (3) (2005) 455–
627 466. doi:10.1140/epja/i2005-10138-7.
- 628 [30] U. Kreissig, S. Grigull, K. Lange, P. Nitzsche, B. Schmidt,
629 In situ ERDA studies of ion drift processes during anodic
630 bonding of alkali-borosilicate glass to metal, *Nuclear Instru-
631 ments and Methods in Physics Research Section B: Beam Inter-
632 actions with Materials and Atoms* 136-138 (1998) 674–679.
633 doi:10.1016/S0168-583X(97)00778-7.
- 634 [31] N. P. Barradas, C. Jeynes, R. P. Webb, Simulated annealing
635 analysis of Rutherford backscattering data, *Applied Physics
636 Letters* 71 (2) (1997) 291–293. doi:10.1063/1.119524.
- 637 [32] C. E. Rolfs, W. S. Rodney, *Cauldrons in the cosmos: nuclear
638 astrophysics*, Theoretical astrophysics, University of Chicago
639 Press, Chicago, 1988.
- 640 [33] J. F. Ziegler, SRIM 2013, <http://srim.org>.
- 641 [34] J. L’Ecuyer, R. J. A. Lévesque, La réaction $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ pour
642 des protons d’énergie inférieure à 500 keV, *Canadian Journal of
643 Physics* 44 (10) (1966) 2201–2210. doi:10.1139/p66-178.
- 644 [35] A. Caldwell, D. Kollár, K. Kröninger, BAT The Bayesian anal-
645 ysis toolkit, *Computer Physics Communications* 180 (11) (2009)
646 2197–2209. doi:10.1016/j.cpc.2009.06.026.
- 647 [36] A. L. Sallaska, C. Iliadis, A. E. Champagne, S. Goriely, S. Star-
648 rfield, F. X. Timmes, STARLIB: A Next-Generation Reaction-
649 Rate Library for Nuclear Astrophysics, *The Astrophysical Jour-
650 nal Supplement Series* 207 (1) (2013) 18. doi:10.1088/0067-
651 0049/207/1/18.