



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Evidence of thalattosuchian crocodylomorphs in the Portland Stone Formation (Late Jurassic) of England, and a discussion on Cretaceous teleosauroids

Citation for published version:

Young, M & Sachs, S 2020, 'Evidence of thalattosuchian crocodylomorphs in the Portland Stone Formation (Late Jurassic) of England, and a discussion on Cretaceous teleosauroids', *Historical Biology*.
<https://doi.org/10.1080/08912963.2019.1709453>

Digital Object Identifier (DOI):

[10.1080/08912963.2019.1709453](https://doi.org/10.1080/08912963.2019.1709453)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Historical Biology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Evidence of thalattosuchian crocodylomorphs in the Portland Stone Formation (Late Jurassic) of England, and a discussion on Cretaceous teleosauroids

Journal:	<i>Historical Biology</i>
Manuscript ID	GHBI-2019-0150.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Young, Mark; University of Edinburgh, School of GeoSciences Sachs, Sven; Naturkunde-Museum Bielefeld
Keywords:	Crocodylomorpha, England, Portland Stone Formation, Tithonian, Teleosauroidea, Machimosaurus

SCHOLARONE™
Manuscripts

1
2
3 **Evidence of thalattosuchian crocodylomorphs in the Portland Stone Formation (Late**
4 **Jurassic) of England, and a discussion on Cretaceous teleosauroids**
5
6
7
8
9

10 MARK T. YOUNG^{1*}, ~~and SVEN SACHS², and PASCAL ABEL³~~
11
12
13

14 ¹ School of GeoSciences, Grant Institute, The King's Buildings, University of Edinburgh, James
15 Hutton Road, Edinburgh, EH9 3FE, United Kingdom
16
17

18 ² Naturkunde-Museum Bielefeld, Abteilung Geowissenschaften, Adenauerplatz 2, 33602 Bielefeld,
19 Germany
20
21

22 ³ ~~Senckenberg Centre for Human Evolution and Palaeoenvironment, Eberhard-Karls-University~~
23 ~~Tübingen, Sigwartstraße 10, 72076 Tübingen, Germany~~
24
25
26
27
28
29

30 We report the first definite specimen of a thalattosuchian crocodylomorph from the Portland
31 Stone Formation of England. This specimen (an isolated tooth crown) can be referred to the
32 teleosauroid genus *Machimosaurus* based on its conical shape, distinctive enamel
33 ornamentation and lack of carinae. Understanding the faunal composition of the Portland
34 Stone Formation is key to elucidating the distinct shift in crocodylomorph taxa that occurred
35 during the Tithonian-to-Berriasian in Europe. One of the most striking aspects of this faunal
36 shift is the hypothesised extinction of Teleosauroidea in Europe. The presence of
37 *Machimosaurus* in the Portland Stone Formation supports the hypothesis that the localised
38 marine regression in Europe at the Jurassic–Cretaceous boundary, and the resultant habitat
39 loss, contributed to the absence of teleosauroids in Europe during the Berriasian. However,
40 the fossil record of thalattosuchians during the Cretaceous is notorious scarce. We review the
41 purported Cretaceous record of teleosauroids, and agree that closer to the equator this clade
42 survived for at least 20 million years after the Jurassic–Cretaceous boundary.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Keywords:** Crocodylomorpha; England; Portland Stone Formation; Tithonian;
4
5 Teleosauroidea
6
7
8
9

10 * **Corresponding author.** Email: marktyoung1984@gmail.com
11
12
13

14 **1. Introduction**

15
16 In southern England there is a distinct faunal shift in crocodylomorph taxa from the
17 Tithonian-to-Berriasian (i.e. crossing the Jurassic–Cretaceous boundary). During the
18 Tithonian, pelagic metriorhynchid thalattosuchians are overwhelmingly the most abundant
19 component of the crocodylomorph fauna, with teleosauroid thalattosuchians being a rare
20 component (see Young and Steel in press for a recent overview). By the Berriasian, the
21 crocodylomorph fauna of southern England is significantly different, being composed of the
22 neosuchian genera *Theriosuchus*, *Goniopholis* and *Pholidosaurus* (e.g. Benton and Spencer
23 1995; Salisbury 2002; Andrade et al. 2011; Tennant et al. 2016). However, the falling sea
24 levels in Europe across the Tithonian-Berriasian (Hallam, 1988, 2001) is most likely the
25 cause of this shift, especially as *Theriosuchus* and *Goniopholis* are known from freshwater
26 Jurassic strata in Europe (e.g. Schwarz 2002; Young et al. 2016), with pholidosaurids also
27 known from the Jurassic (Fortier et al. 2011). Moreover, metriorhynchid biodiversity peaks
28 coincide with high sea levels and the greater number of shallow marine formations, and
29 biodiversity troughs coincide with low sea levels and lower numbers of shallow marine
30 formations (Young, 2009; Young et al. 2010).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50
51 However, when this regionalised shift occurred is still not understood. This is because
52 the late Tithonian thalattosuchian fossil record is exceptionally poor, restricted to the
53 exceptional fossils from Argentina (e.g. Gasparini et al. 2006; [Fernández et al. 2019](#)). In
54 England, thalattosuchian fossils from the Tithonian are exclusively come-known from the
55
56
57
58
59
60

1
2
3 Kimmeridge Clay Formation (e.g. Benton and Spencer 1995; Young and Steel in press). The
4
5 crocodylomorph fauna from geological younger strata, such as the Portland Stone Formation,
6
7 are unknown. Here we begin to rectify this issue by describing an incomplete tooth crown
8
9 from the Portland Stone Formation of the Isle of Portland, which we assign to the
10
11 teleosauroid genus *Machimosaurus*.
12
13
14
15
16

17 **1.1 Institutional Abbreviation**

18
19 **MB**, Museum für Naturkunde der Humboldt Universität, Berlin, Germany; [MNHN](#),
20
21 [Muséum national d'Histoire naturelle, Paris, France](#).
22
23
24
25

26 **2. Systematic Palaeontology**

27
28
29
30
31 Crocodylomorpha Hay, 1930 (*sensu* Nesbitt, 2011)

32
33 Thalattosuchia Fraas, 1901 (*sensu* Young and Andrade, 2009)

34
35 Teleosauroidea Geoffroy Saint-Hilaire, 1831 (*sensu* Young and Andrade, 2009)

36
37 *Machimosaurus* von Meyer, 1837 (emendation von Meyer, 1838)

38 39 40 41 42 **Type species**

43
44 *Machimosaurus hugii* von Meyer, 1837

45
46
47
48
49 *Machimosaurus* sp.

50
51 (Figure 1)

52 53 54 55 56 **Specimen**

57
58 MB.R.4059, an incomplete isolated tooth crown.
59
60

Locality

Isle of Portland, Dorset, England, United Kingdom.

Horizon

Portland Stone Formation, Portland Group. Tithonian, Upper Jurassic.

3. Description

The tooth appears to have been broken in three places, first near the root-crown junction, second along its apicobasal axis, and finally in the apical region. Thus, the tooth does not preserve the root, at least one third of the crown in the basal-and-mid regions, and the entire apical region (Fig. 1). From what is preserved, the tooth crown would have had a conical shape, with slight mediolateral compression. The preserved external surface shows a slight curvature towards the apex. In *Machimosaurus* teeth this slight curvature occurs on the labial surface (Krebs 1967; Hua 1999; Lepage et al. 2008; Young et al. 2014a, 2014b). As such, the lingual surface is largely missing. Unfortunately, we cannot orientate the tooth mesially and distally.

Mesial and distal carinae are absent in MB.R.4059. Although the tooth crown is incomplete (Fig. 1), the carinae should still be visible as most of the crown is preserved. The presence, and morphology, of carinae is variable in *Machimosaurus*, ranging from being readily identifiable, being hard to distinguish from the superficial enamel ornamentation, to being completely absent (see Krebs 1967; Young et al. 2014b).

The preserved external surfaces of the enamel are covered by superficial ornamentation. It is composed of numerous, tightly packed apicobasally aligned ridges that are (sub)-parallel to one another. Most are continuous from the basal region to the preserved

1
2
3 apical-most region, although shorter ridges are also present. As the apex is missing, the
4
5 anticipated shift in enamel ornamentation to an anastomosed pattern cannot be seen.
6
7
8
9

10 **4. Discussion**

11 **4.1 Referral of MB.R.4059 to *Machimosaurus***

12
13 The isolated tooth crown described herein has a suite of morphological characteristics that are
14
15 only known in two genera of thalattosuchians, the teleosauroid *Machimosaurus* and the
16
17 metriorhynchid *Torvoneustes* (von Meyer 1837; Krebs 1967; Hua 1999; Lepage et al. 2008;
18
19 Andrade et al. 2010; Barrientos-Lara et al. 2016; Young et al. 2013, 2014a, 2014b):
20
21
22

- 23
24 1. Tooth crowns that are largely conical (poor mediolateral compression) - although this
25
26 is variable between species in these genera, and across the tooth-row (for
27
28 *Torvoneustes* see Barrientos-Lara et al. 2016).
- 29
30 2. Basal third to four-fifths of the tooth crown have numerous and tightly-packed
31
32 apicobasally aligned enamel ridges that are (sub)parallel to one another, with most
33
34 being continuous from the base to the apical region.
35
36
37
38
39

40
41 Based on this character suite we can safely refer MB.R.4059 to either *Machimosaurus*
42
43 or *Torvoneustes*. We can exclude this tooth from *Torvoneustes* due to the lack of carinae. The
44
45 dentition of *Torvoneustes* have strongly developed mesiodistal carinae (Andrade et al. 2010;
46
47 Young et al. 2013; Barrientos-Lara et al. 2016), whereas in *Machimosaurus* the presence of
48
49 carinae is variable across the tooth-row, including tooth crowns lacking carinae (Young et al.
50
51 2014a, 2014b).
52

53
54 These morphological features are characteristic of the ‘crunch guild’ of Massare
55
56 (1987). As shown by Foffa et al. (2018), other Late Jurassic marine reptile groups from the
57
58 Sub-Boreal seaway did not convergently evolve this character suite. Morphologically, the
59
60

1
2
3 closest to these ‘crunch guild’ teeth, are those of the ‘smash guild’, occupied by ichthyosaurs.
4
5 However, plicidentine enamel is considered an apomorphy of Ichthyosauria (Maxwell et al.,
6
7 2011), a unique enamel ornamentation arrangement composed of apicobasal ridges
8
9 interspaced with furrows.
10
11

12 13 14 **4.2 Cretaceous teleosauroids**

15
16 The thalattosuchian fossil record in the Cretaceous is notoriously poor, especially for
17
18 teleosauroids. Young et al. (2014a) revised the genus *Machimosaurus* and found there to be
19
20 no conclusive evidence of *Machimosaurus*, and thus teleosauroids, to be present in the
21
22 Cretaceous. Fanti et al. (2016) described a new species from Tunisia, *Machimosaurus rex*
23
24 Fanti et al. 2016, which they considered to be from the Early Cretaceous. While Fanti et al.
25
26 (2016) considered *M. rex* to be from the Hauterivian, the age of *M. rex* is far from certain (see
27
28 Martin et al. 2018; Cortes et al. in press). The possibility that the *M. rex* holotype is Tithonian
29
30 in age cannot be discounted (M. Johnson pers. comm. 2019). Thus, *M. rex* cannot be taken as
31
32 definitive evidence for the survival of teleosauroids into the Cretaceous.
33
34
35

36
37 _____ An incomplete fossil from late Barremian strata of Colombia was recently referred to
38
39 Teleosauroidea (Cortes et al. in press). Unfortunately, no apomorphic characters were used to
40
41 make this determination, with mesoeucrocodylian symplesiomorphies being used to justify
42
43 the identification (e.g. hourglass-shaped amphiplatyan dorsal vertebral centra and osteoderms
44
45 that are noticeably rectangular in shape). This is understandable, given that the specimen
46
47 consists of dorsal vertebrae and ribs, dorsal and ventral trunk osteoderms, and epipodial
48
49 elements preserved in matrix. Alas, the possibility of a pholidosaurid identification was not
50
51 discussed. This is unfortunate as the large-bodied pholidosaurid *Sarcosuchus hartti* (Marsh,
52
53 1869) is known from the late Hauterivian-early Barremian of Brazil (Souza et al. 2019).
54
55 Moreover, Cortes et al. (in press) estimated the body length of the Colombian specimen to be
56
57
58
59
60

1
2
3 9.6m (with 95% confidence interval of 7.6m – 13.4m), this size is more consistent with
4 *Sarcosuchus* Broin & Taquet 1966 (O'Brien et al. 2019) than a teleosauroid (Young et al.
5 2016). However, the measurements of the dorsal osteoderms and their dorsal ornamentation
6 are more consistent with a teleosauroid than a pholidosaurid. The shape of the dorsal
7 vertebrae are also more reminiscent of a teleosauroid than *Sarcosuchus* (examining the
8 *Sarcosuchus imperator* Broin & Taquet 1966 display skeleton in the MNHN), given their
9 proportionally long centra.

10
11
12
13
14
15
16
17
18
19 Despite the shortcomings of the description and comparisons in Cortes *et al.* (in
20 press), the taxonomic identification does appear to be correct. It yields an approximate 20
21 million-year gap between the Colombian indeterminate specimen and the late Tithonian
22 *Machimosaurus* tooth described herein (MB.R.4059). Prior to the description of the
23 MB.R.4059, the youngest definitive teleosauroids were from the early Tithonian of Europe,
24 including *Aeolodon priscus* (von Sömmerring, 1814) and *Machimosaurus hugii* von Meyer
25 (1837) (see Young et al. 2014a; Foffa et al. 2019; Young & Steel in press).

26
27
28
29
30
31
32
33
34
35 More diagnostic remains, and remains that have better age constraint, are needed
36 before any definitive statements can be made regarding Cretaceous teleosauroids. However,
37 closer to the equator it does appear that teleosauroids survived far longer than previously
38 supposed, and grew to lengths rivalling even the largest Cretaceous neosuchians.

4.23 *Late Tithonian thalattosuchians in Europe*

46
47
48
49 With the description of MB.R.4059, we have the first late Tithonian thalattosuchian known
50 from Europe. Given the incredible diversity of European thalattosuchians in the early
51 Tithonian (in particular metriorhynchids), it is highly likely that other fossils attributable to
52 Thalattosuchia are currently in museum collections waiting to be described. We hope this
53 paper will spur further investigation into these collections, as until more fossils are
54
55
56
57
58
59
60

1
2
3 discovered and described, we cannot elucidate the dramatic shift in the crocodylomorph biota
4
5 across the Jurassic-Cretaceous boundary.
6
7
8
9

10 **Acknowledgements**

11
12 We would like to thank Daniela Schwarz for access to the MB collection. Furthermore, we
13
14 would like to thank Florian Witzmann (MB) for photography. We would like to thank the two
15
16 reviewers for their constructive comments. MT Young is supported by a Leverhulme Trust
17
18 Research Project grant (number RPG-2017-167). MT Young received support to visit the MB
19
20 and MNHN collections from the SYNTHESYS Project (<http://www.synthesys.info/>) which is
21
22 financially supported by European Community Research Infrastructure Action under the FP7
23
24 ‘Capacities’ Programme.
25
26
27
28
29

30 **References**

- 31
32
33 Andrade MB, Young MT, Desojo JB, Brusatte SL. 2010. The evolution of extreme
34
35 hypercarnivory in Metriorhynchidae (Mesoeucrocodylia: Thalattosuchia): evidence
36
37 from microscopic denticle morphology and a new tri-faceted Kimmeridgian tooth
38
39 from Germany. *Journal of Vertebrate Paleontology* 30:1451–1465.
40
41
42 Andrade MB, Edmonds R, Benton MJ, Schouten R. 2011. A new Berriasian species of
43
44 *Goniopholis* (Mesoeucrocodylia, Neosuchia) from England, and a review of the
45
46 genus. *Zoological Journal of the Linnean Society* 163:S66–S108.
47
48
49 Barrientos-Lara JJ, Herrera Y, Fernández MS, Alvarado-Ortega J. 2016. Occurrence of
50
51 *Torvoneustes* (Crocodylomorpha, Metriorhynchidae) in marine Jurassic deposits of
52
53 Oaxaca, Mexico. *Revista Brasileira de Paleontologia* 19:415–424.
54
55
56 Benton MJ, Spencer PS. 1995. Fossil reptiles of Great Britain. Geological Conservation
57
58 Review Series, No. 10. London: Chapman and Hall. 386 pp.
59
60

1
2
3 Broin FP de, Taquet P. 1966. Découverte d'un crocodylien nouveau dans le Crétacé Inférieur
4 du Sahara. Comptes Rendus de l'Académie des Sciences Paris 262:2326–2329.
5

6
7 Cortes D, Larsson HCE, Maxwell EE, Parra Ruge ML, Patarroyo P, Wilson JA. In press. An
8 Early Cretaceous teleosauroid (Crocodylomorpha: Thalattosuchia) from Colombia.
9 Ameghiniana. doi: 10.5710/AMGH.26.09.2019.3269
10

11
12 Fanti F, Miyashita T, Cantelli L, Mnasri F, Dridi J, Contessi M, Cau A. 2016. The largest
13 thalattosuchian (Crocodylomorpha) supports teleosaurid survival across the Jurassic-
14 Cretaceous boundary. Cretaceous Research 61:263–274.
15

16
17 Fernández MS, Herrera Y, Vennari VV, Campos L, de la Fuente M, Talevi M, Aguirre-
18 Urreta B. 2019. Marine reptiles from the Jurassic/Cretaceous transition at the High
19 Andes, Mendoza, Argentina. Journal of South American Earth Sciences 92:658–673.
20

21
22 Foffa D, Young MT, Stubbs TL, Dexter KG, Brusatte BL. 2018. The long-term ecology and
23 evolution of marine reptiles in a Jurassic seaway. Nature Ecology & Evolution
24 2:1548–1555.
25

26
27 Foffa D, Johnson MM, Young MT, Steel L, Brusatte SL. 2019. A revision of the deep-water
28 teleosauroid crocodylomorph *Teleosaurus megarhinus* Hulke, 1871 from the
29 Kimmeridge Clay Formation (Late Jurassic) of England, UK. PeerJ 7:e6646.
30

31
32 Fortier D, Perea D, Schultz, C. 2011. Redescription and phylogenetic relationships of
33 *Meridiosaurus vallisparadisi*, a pholidosaurid from the Late Jurassic of Uruguay.
34 Zoological Journal of the Linnean Society 163:S257–S272.
35

36
37 Fraas E. 1901. Die Meerkrokodile (Thalattosuchia n. g.) eine neue sauriergruppe der
38 Juraformation. Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg
39 57:409–418.
40

41
42 Gasparini ZB, Pol D, Spalletti LA. 2006. An unusual marine crocodyliiform from the
43 Jurassic-Cretaceous boundary of Patagonia. Science 311:70–73.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Geoffroy Saint-Hilaire É. 1831. Recherches sur de grands sauriens trouvés à l'état fossile aux
4 confins maritimes de la Basse-Normandie, attribués d'abord au crocodile, puis
5 déterminés sous les noms de *Teleosaurus* et *Steneosaurus*. Mémoires de l'Académie
6 des sciences 12:1–138.
7
8
9
10

11
12 Hallam A. 1988. A re-evaluation of Jurassic eustasy in the light of new data and the revised
13 Exxon curve. In: Wilgus CK, Hastings BS, Kendall CGC, Posamentier HW, Ross
14 CA, Van Wagoner JC. (eds.) *Sea-level changes – an integrated approach*, SEPM
15 *SPec. Publ* 42:261–273.
16
17
18
19
20

21 Hallam A. 2001. A review of the broad pattern of Jurassic sea-level changes and their
22 possible causes in the light of current knowledge. *Palaeogeography*
23 *Palaeoclimatology Palaeoecology* 167:23–37.
24
25
26
27

28 Hay OP. 1930. Second bibliography and catalogue of the fossil vertebrata of North America,
29 volume 2. Washington, DC: Carnegie Institute Washington. 1094 pp.
30
31
32

33 Hua S. 1999. Le crocodilien *Machimosaurus mosae* (Thalattosuchia, Teleosauridae) du
34 Kimmeridgien du Boulonnais (Pas de Calais, France). *Palaeontographica Abteilung A*
35 252:141–170.
36
37
38
39

40 Krebs B. 1967. Der Jura-Krokodilier *Machimosaurus* H. v. Meyer. *Paläontologische*
41 *Zeitschrift* 41:46–55.
42
43
44

45 Lepage Y, Buffetaut E, Hua S, Martin JE, Tabouelle, J. 2008. Catalogue descriptif,
46 anatomique, géologique et historique des fossiles présentés à l'exposition « Les
47 Crocodiliens fossiles de Normandie » (6 novembre - 14 décembre 2008). *Bulletin de*
48 *la Société Géologique de Normandie et des Amis du Muséum du Havre* 95:5–152.
49
50
51
52

53 [Marsh OC. 1869. Notice of some new reptilian remains from the Cretaceous of Brazil.](#)
54 [American Journal of Science 47:390–392.](#)
55
56
57
58
59
60

1
2
3 [Martin JE, Suteethorn S, Lauprasert K, Tong H, Buffetaut E, Liard R, Salaviale C, Deesri U,](#)
4 [Suteethorn V, Claude J. 2018. A new freshwater teleosaurid from 636 the Jurassic of](#)
5 [northeastern Thailand. Journal of Vertebrate Paleontology 38 \(6\):e1549059.](#)
6
7

8
9
10 [Massare JA. 1987. Tooth morphology and prey preference of Mesozoic marine reptiles.](#)
11 [Journal of Vertebrate Paleontology 7:121–137.](#)
12

13
14 [Maxwell EE, Caldwell M, Lamoureux DO. 2011. The structure and phylogenetic distribution](#)
15 [of amniote plicidentine. Journal of Vertebrate Paleontology 31:553–561.](#)
16

17
18
19 Meyer CEH von. 1837. Mitteilungen, an Professor Bronn gerichtet. Neues Jahrbuch für
20
21 Mineralogie, Geologie, Geognosie und Petrefaktenkunde 4:413–418.
22

23
24 Meyer CEH von. 1838. Briefliche Mitteilungen an Professor Bronn. Neues Jahrbuch für
25
26 Mineralogie, Geologie, Geognosie und Petrefaktenkunde 1838:668.
27

28
29 Nesbitt SJ. 2011. The early evolution of archosaurs: relationships and the origin of major
30
31 clades. Bulletin of the American Museum of Natural History 352:1–292.
32

33 [O'Brien HD, Lynch LM, Vliet KA, Brueggen J, Erickson GM, Gignac PM. 2019.](#)
34 [Crocodylian head width allometry and phylogenetic prediction of body size in extinct](#)
35 [crocodyliforms. Integrative Organismal Biology 1 \(1\):obz006.](#)
36
37

38
39
40 Salisbury SW. 2002. Crocodylians from the Lower Cretaceous (Berriasian) Purbeck
41
42 Limestone Group of Dorset, Southern England. In: Milner AR, Batten DJ, eds. *Life*
43 [and environments in Purbeck times](#). Special Papers in Palaeontology 68:121–144.
44
45

46
47 Schwarz D. 2002. A new species of *Goniopholis* from the Upper Jurassic of Portugal.
48
49 Palaeontology 45:185–208.
50

51 [Souza RG, Figueiredo RG, Azevedo SAK, Riff D, Kellner AWA. 2019. Systematic revision](#)
52 [of *Sarcosuchus hartti* \(Crocodyliformes\) from the Recôncavo Basin \(Early](#)
53 [Cretaceous\) of Bahia, north-eastern Brazil. Zoological Journal of the Linnean Society:](#)
54 [zlz057.](#)
55
56
57
58
59
60

Sömmerring ST. von. 1814. Über den *Crocodylus priscus*, oder ein in Baiern versteint
gefundenes schmalkie-feriges Krokodil, Gavial der Vorwelt. Denkschriften der
Königlichen Akademie der Wissenschaften zu München. Classe der Mathematik und
Naturwissenschaften 5:9–82.

- Tennant JP, Mannion PD, Upchurch P. 2016. Evolutionary relationships and systematics of Atoposauridae (Crocodylomorpha: Neosuchia): implications for the rise of Eusuchia. *Zoological Journal of the Linnean Society* 177:854–936.
- Young MT 2009. Quantifying macroevolutionary patterns in highly specialised clades of archosaurs. Unpublished PhD thesis, University of Bristol. 350 pp.
- Young MT, Andrade MB. 2009. What is *Geosaurus*? Redescription of *Geosaurus giganteus* (Thalattosuchia, Metriorhynchidae) from the Upper Jurassic of Bayern, Germany. *Zoological Journal of the Linnean Society* 157:551–585.
- Young MT, Andrade MB, Etches S, Beatty BL. 2013. A new metriorhynchid crocodylomorph from the Lower Kimmeridge Clay Formation (Late Jurassic) of England, with implications for the evolution of dermatocranium ornamentation in Geosaurini. *Zoological Journal of the Linnean Society* 169:820–848.
- Young MT, Hua S, Steel L, Foffa D, Brusatte SL, Thüring S, Mateus O, Ruiz-Omeñaca JJ, Havlik P, Lepage Y, Andrade MB. 2014a. Revision of the Late Jurassic teleosaurid genus *Machimosaurus* (Crocodylomorpha, Thalattosuchia). *Royal Society Open Science* 1 (2):140222. <http://dx.doi.org/10.1098/rsos.140222>
- Young MT, Steel L, Brusatte SL, Foffa D, Lepage Y. 2014b. Tooth serration morphologies in the genus *Machimosaurus* (Crocodylomorpha, Thalattosuchia) from the Late Jurassic of Europe. *Royal Society Open Science* 1 (3):140269. <http://dx.doi.org/10.1098/rsos.140269>

1
2
3 Young MT, Tennant JP, Brusatte SL, Challands TJ, Fraser NC, Clark NDL, Ross DA. 2016.

4
5 The first definitive Middle Jurassic atoposaurid (Crocodylomorpha, Neosuchia), and a
6
7 discussion on the genus *Theriosuchus*. *Zoological Journal of the Linnean Society*
8
9
10 176:443–462.

11
12 Young MT, Steel L. In press. Crocodiles: Thalattosuchia. In: Martill D. (ed.) *Field Guide to*
13
14 *fossils of the Kimmeridge Clay Formation*. Palaeontological Association Field Guide
15
16
17 16.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

FIGURE CAPTION

Figure 1. *Machimosaurus* sp. MB.R.4059. Isolated tooth crown. A, lingual view, B, lingual view rotated towards basal cracked region, C, preserved labial view, and D, lingual view rotated away from basal cracked region. Scale bar is two centimetres.

For Peer Review Only




Figure 1. *Machimosaurus* sp. MB.R.4059. Isolated tooth crown. A, lingual view, B, lingual view rotated towards basal cracked region, C, preserved labial view, and D, lingual view rotated away from basal cracked region. Scale bar is two centimetres.

184x79mm (600 x 600 DPI)