



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **The challenge of using routinely collected data to compare hospital admission rates by ethnic group**

**Citation for published version:**

Knox, S, Bhopal, RS, Thomson, CS, Millard, A, Fraser, A, Gruer, L & Buchanan, D 2019, 'The challenge of using routinely collected data to compare hospital admission rates by ethnic group: a demonstration project in Scotland', *Journal of Public Health*. <https://doi.org/10.1093/pubmed/fdz175>

**Digital Object Identifier (DOI):**

[10.1093/pubmed/fdz175](https://doi.org/10.1093/pubmed/fdz175)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Journal of Public Health

**Publisher Rights Statement:**

This is the author's peer-reviewed manuscript as accepted for publication.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



<http://mc.manuscriptcentral.com/jph>

**The challenge of using routinely collected data to compare hospital admission rates by ethnic group: a demonstration project in Scotland**

Journal:	<i>Journal of Public Health</i>
Manuscript ID	JPH-19-0553.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Knox, Stuart; NHS National Services Scotland, ISD Bhopal, Raj; University of Edinburgh, Public health sciences Thomson, Catherine; NHS National Services Scotland, ISD Millard, Andrew; Scottish Public Health Observatory, NHS Health Scotland, Public Health Science Fraser, Andrew; NHS Health Scotland, Public Health Science Gruer, Laurence; University of Edinburgh, Usher Institute of Population Health Sciences and Informatics Buchanan, Duncan; ISD Scotland,
Keywords:	Ethnicity, Secondary and tertiary services, Methods

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6 **The challenge of using routinely collected data to compare hospital**  
7 **admission rates by ethnic group: a demonstration project in**  
8 **Scotland**  
9  
10  
11  
12  
13

14 Knox S<sup>1</sup>, Bhopal RS<sup>2</sup>, Thomson CS<sup>1</sup>, Millard A<sup>3</sup>, Fraser A<sup>4</sup>, Gruer L<sup>2</sup>, Buchanan D<sup>1</sup>

16  
17 Knox S BSc, Senior Information Analyst

18  
19 Bhopal RS Professor, Emeritus Professor of Public Health

20  
21 Thomson CS MSc, BSc, Information Lead for Population Health

22  
23 Andrew Millard, Ph.D., M.P.H., M.Sc., B.A., Research Fellow

24  
25 Dr Andrew Fraser MPH, Director of Public Health Science

26  
27 Gruer L BSc MB ChB MD MPH, Honorary Professor of Public Health

28  
29 Buchanan D BSc, C Stat, Head of Consultancy Services

- 30  
31  
32  
33  
34  
35 1. Information Services Division, NHS National Services Scotland, Gyle Square, 1  
36 South Gyle Crescent, Edinburgh, EH12 9EB, UK  
37  
38 2. Usher Institute of Population Health Sciences and Informatics, University of  
39 Edinburgh, Edinburgh, UK  
40  
41 3. MRC/CSO Social and Public Health Sciences Unit, University of Glasgow,  
42 Glasgow, UK.  
43  
44 4. NHS Health Scotland, Gyle Square, 1 South Gyle Crescent, Edinburgh, EH12  
45 9EB, UK  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 **Corresponding Author:** D Buchanan  
56  
57  
58  
59  
60

## ABSTRACT

### Background

Recording patients' ethnic group supports efforts to achieve equity in health care provision. Before the Equality Act (2010), recording ethnic group at hospital admission was poor in Scotland but has improved subsequently. We describe the first analysis of the utility of such data nationally for monitoring ethnic variation.

### Methods

We analysed all in-patient or day case hospital admissions in 2013. We imputed missing data using the most recent ethnic group recorded for a patient from 2009-2015. For episodes lacking an ethnic code we attributed known ethnic codes proportionately. Using the 2011 Census population, we calculated rates and rate ratios for all-cause admissions and ischaemic heart diseases (IHD) directly standardised for age.

### Results

Imputation reduced missing ethnic group codes from 24% to 15%, and proportionate redistribution to zero. While some rates for both all-cause and IHD admissions appeared plausible, unexpectedly low or high rates were observed for several ethnic groups particularly among White groups and newly coded groups.

### Conclusions

Completeness of ethnicity recoding on hospital admission records has improved markedly since 2010. However the validity of admission rates based on these data

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

is variable across ethnic groups and further improvements are required to support monitoring of inequality.

For Peer Review

## INTRODUCTION

The quest for equity in health and health-care in modern, multicultural societies requires accurate, recent, quantitative information on similarities and differences in health status and health care utilisation by important characteristics including ethnicity (1-3). A practical way to achieve this for ethnicity is by including an ethnic group code in individual health records. In 1991, the Department of Health recommended that general practitioners in England should include patients' ethnic group in letters referring them to hospital, but without clear guidance on how to do so (4). After this approach failed, mandatory recording in secondary care was implemented in England in 1995, backed up by detailed guidance (5). By 2005, recording ethnic group was recommended but not mandatory in secondary care in Scotland (6,7). Perhaps because of this, ethnicity was only 42% complete for hospital admissions in 2010, with four of the 15 Scottish health Boards achieving less than 10% (8). The Equality Act 2010 made an explicit legal requirement of the NHS across the UK to monitor equity and equality (9). In the light of the Act, senior NHS managers in Scotland prioritised the recording of ethnic group. Rates for hospital admissions rose rapidly to 75% in 2012, reaching 82% in 2016, and remaining at 81% in 2018 (10,11).

The Scottish Health and Ethnicity Linkage Study (SHELS) (12) linked variables from the 2001 Scotland Census including self-reported ethnic group to NHS records at an individual level, enabling comparisons to be made between the larger ethnic groups for hospital admission rates for all-causes, ischaemic heart disease and a wide range of other health conditions (13-21). Such academic research linkage projects have many strengths in terms of data accuracy and robustness, but due to the

1  
2  
3 extensive resources required, the long time lag before results become available and  
4  
5 the need for stringent information governance covering research projects, such  
6  
7 studies cannot provide routine ethnic monitoring as required in health policy.  
8  
9

10  
11  
12 Building on the efforts to improve completeness, this project aimed to investigate the  
13  
14 utility of ethnicity recording in routine hospital admission data for monitoring equity of  
15  
16 health care provision. Analyses based on incomplete data would be potentially  
17  
18 biased if completeness varied between ethnic groups or if some ethnic groups were  
19  
20 more likely to be inaccurately recorded than others. We thus initially focused on  
21  
22 methods to maximise completeness. We produced the first analysis of hospital  
23  
24 admission rates by ethnic group at national level, focussing on all-cause and  
25  
26 ischaemic heart disease (IHD) admissions to permit comparisons with previous  
27  
28 findings. While no 'gold standard' ethnicity data were available to directly assess the  
29  
30 validity of the rates, we used published results from the SHELS cohort as a general  
31  
32 benchmark across ethnic groups, taking account of methodological differences.  
33  
34  
35  
36  
37  
38  
39

## 40 **Methods**

41  
42 In-patients and/or day cases in Scotland in 2013 from the Scottish hospital episodes  
43  
44 database (SMR01) formed our study population. Ethnic group codes were available  
45  
46 in 76% of these records, with the remainder recorded as missing, unknown or patient  
47  
48 refused. Because ethnic group may be recorded whenever a patient is admitted,  
49  
50 different ethnic group codes could be given for the same patient. We extracted all  
51  
52 SMR01 episodes in 2009-2015 for the patients in our 2013 study population,  
53  
54 grouping multiple episodes and ethnic codes by patient. Imputation of an individual's  
55  
56 ethnic code for 2013 was based on the most recent recorded ethnic group during the  
57  
58  
59  
60

1  
2  
3 period 2019-15, excluding the codes for Other Ethnic Group, Unknown or Not  
4 Provided. For the remaining patients with an Unknown or Not Provided ethnic group  
5 code after imputation, these admissions were redistributed in the same proportions  
6 as those with a known ethnic group within each five-year age band and sex  
7 category. There was thus complete ethnic group assignment across the patient  
8 cohort.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

19 Due to small numbers and to ensure confidentiality, the two African and three Black/  
20 Caribbean ethnic groups were each combined into one group. Pre-2011 episodes  
21 used the 2001 census ethnic groups, and therefore could not include the new 2011  
22 codes for White Polish, White Gypsy/Traveller and Arab. Table A1 in the online  
23 supplementary information details the 2001 and 2011 census codes, the mappings  
24 between them, and the shortened text used to label the ethnic groups throughout this  
25 paper. Population data for each ethnic group were obtained from the 2011 Scottish  
26 Census. Three sets of hospitalisation admission rates were calculated and reported  
27 by sex for 2013, directly age-standardised using the 2013 European Standard  
28 Population structure (22). These were for the original recorded patient ethnicity in  
29 2013 episodes, the imputed codes using the 2009-2015 records, and the records  
30 with complete ethnic codes following proportionate redistribution of the Unknown or  
31 Not Provided ethnic group cases (labelled as 'original ethnic codes', 'imputed codes'  
32 and 'imputed and redistributed codes' respectively).  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 Confidence intervals were calculated for directly age-standardised rates assuming a  
55 Poisson distribution (23) and rate ratios (RR) relative to the White Scottish  
56  
57  
58  
59  
60



1  
2  
3 population produced for our 'imputed and redistributed codes' method for the full  
4  
5 2013 cohort.  
6  
7

8 We used International Classification of Disease (ICD10) codes I20, I21, I22, I23, I24,  
9  
10 I25 to select all diagnoses of Ischaemic Heart Diseases. Ethics approval was  
11  
12 unnecessary as this was an internal quality audit of the routinely-collected hospital  
13  
14 data submitted and used by the Information Services Division of the NHS in Scotland  
15  
16 (ISD).  
17  
18  
19  
20  
21  
22

### 23 **Results**

24 Among more than 1.5 million hospital admission episodes in 2013, 76% had an  
25  
26 ethnic group recorded. Imputation increased completeness to 85% (Table A2 online).  
27  
28 Within ethnic groups the increase varied from 7% for White Other and Mixed groups  
29  
30 to 26% for the Caribbean group. The number of Other Ethnic Group episodes  
31  
32 decreased by 4% since some were replaced by more specific codes.  
33  
34  
35  
36  
37  
38

39 Figures 1 and 2 show the age-standardised all-cause hospitalisation rates per 1,000  
40  
41 population by ethnic group in 2013 for the three methods for males and females  
42  
43 respectively. As expected, rates were higher after imputation and redistribution, with  
44  
45 more episodes assigned to a known, specific code, although the observed variation  
46  
47 and relative rankings of the ethnic groups remained similar. Very low admission  
48  
49 rates were observed for both males and females in the White Gypsy, Arab and White  
50  
51 Irish ethnic groups, being only 14-36% of those in White Scottish group. On the other  
52  
53 hand, the Other Ethnic Group rates were extremely high, using all three methods.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Age-standardised rate ratios (RR) for all-cause hospital admissions are shown for  
4 the 'imputed and redistributed codes' method in Figures 3 (males) and 4 (females).  
5  
6 For males, the highest RR was for White Other (1.88, 95% CI 1.84 – 1.93) followed  
7  
8 by White Other British (1.39, 95% CI 1.38 – 1.41). The lowest RRs were for White  
9  
10 Irish (0.36, 95% CI 0.34 – 0.38) followed by Chinese (0.64, 95% CI 0.59 – 0.68) and  
11  
12 White Polish (0.78, 95% CI 0.73 – 0.83) ethnic groups. The RRs for the remaining  
13  
14 ethnic groups ranged from 0.81 to 1.17. Very similar results were observed for the  
15  
16 females.  
17  
18  
19  
20  
21  
22  
23

24 Comparison with the incident rate ratios derived from the SHELS cohort (13,14) are  
25 shown in supplementary Tables A3 (all causes) and A4 (IHD). For all causes, rate  
26 ratios for both males and females were markedly lower in the current study for the  
27  
28 White Irish and much higher for the White Other British and White Other (Table A3).  
29  
30 The differences in the ratio ratios for the other groups were much smaller, being less  
31  
32 than 10% for Mixed, Pakistani, Indian and Chinese males and Mixed Females. For  
33  
34 IHD, in comparison with SHELS results, the rate ratios were markedly lower for  
35  
36 White Irish males and females, and markedly higher for White Other British, White  
37  
38 Other, Mixed, Pakistani and Indian males and for White Other British, White Other,  
39  
40 Mixed and Pakistani females. Only for Chinese and Asian Other males and Indian  
41  
42 females were the corresponding rate ratios within 10% of each other (Table A4).  
43  
44  
45  
46  
47  
48  
49  
50  
51

52 The detailed data behind the 2013 rate ratios are provided in the supplementary  
53  
54 online Tables A3 – A7.  
55  
56  
57  
58  
59  
60

## DISCUSSION

### *Main findings of this study*

Using routinely collected hospital admission data for Scotland in 2013, we found 76% of hospitalisation records had an ethnic group recorded. This was increased to 85% by imputation and to 100% by proportionate redistribution of the Unknown/Not Provided category. We subsequently calculated admission rates and rate ratios for 14 ethnic groups, both for all-cause hospital admissions and for IHD. There was considerable variation in admission rates among the White groups, less so among the other non-White groups. The White Irish, Arab and White Gypsy groups were implausibly low, suggesting they had been under-recorded in the Scottish hospital admission records. Apart from these, the lowest admission rates were found among the Chinese. Comparison with previously published rates based on self-reported census ethnicity suggested that, despite relatively high levels of data completeness, there was variable validity of admission rates across ethnic groups based on these routine data, particularly for IHD.

### *What is already known on this topic*

Very few countries have included ethnic coding in their hospital admission records. In a 2009 review, Rafnsson and Bhopal found only England, Scotland and four other countries in the European Union had population or hospital admission registers with coding for migrants or ethnic minorities for cardiovascular diseases and diabetes (24). Of these, only England, Scotland, Sweden and Italy were at a national level. England and Scotland identified a number of different ethnic groups, but the rate of recording ethnic group in Scotland at that time was too low to allow any analysis. Sweden coded for country of birth and Italy for citizenship: neither specified

1  
2  
3 individual ethnic groups. Where ethnicity is routinely recorded, studies have found  
4  
5 completeness to be sub-optimal and approaches have been proposed to maximise  
6  
7 the usability of the data.  
8  
9

10 The English National Cancer Intelligence Network compared cancer rates in different  
11  
12 ethnic groups by linking cancer registry and Hospital Episode Statistics (HES)  
13  
14 datasets for 2002-2006. They found that among almost 1.2 million patients, 24% had  
15  
16 a missing ethnic group code, and the proportion missing varied by cancer type (25).  
17  
18 They used three different proportionality methods to assign the unknowns to ethnic  
19  
20 groups to allow age-standardised rates to be calculated but recognised they could  
21  
22 not exclude the possibility of bias. Three studies of cancer patients in England  
23  
24 addressed incompleteness by assuming that the proportions of the missing values  
25  
26 for each ethnic group were the same for those with a known ethnic group (26,27), or  
27  
28 have focussed on rate ratios by ethnic group as in the current study (28). Another  
29  
30 highlighted the benefits of linking patient data across multiple administrative  
31  
32 datasets to maximise completeness (29). Aspinall and Jacobson suggested a  
33  
34 range of approaches that might be used to address incompleteness and encourage  
35  
36 greater use of the data (30).  
37  
38  
39  
40  
41  
42

43 While much of the literature has focussed on data completeness, few studies have  
44  
45 been able to adequately measure and benchmark the validity and accuracy of  
46  
47 routine ethnic data. Comparing self-reported ethnic group in a survey of over  
48  
49 58,000 cancer patients in England with hospital records for the same patients,  
50  
51 Saunders et al found that in the hospital records, ethnic group was correct in 98%  
52  
53 of White British but varied between 80% and only 10% in 15 minority groups (31).  
54  
55 In the absence of a similar gold standard comparator, others have relied on linking  
56  
57 records between routine datasets (29, 32) A review of ethnicity data in the health  
58  
59  
60

1  
2  
3 service in New Zealand found that recording in hospitals and other health sectors  
4 was very variable (33). For example, one audit found that only 65% of Maori were  
5 correctly classified on the hospital computer system.  
6  
7  
8  
9

### 10 11 12 **What this study adds**

13  
14  
15 After a decision by the Scottish Government in 2009 to prioritise ethnic coding for  
16 hospital admissions and out-patient attendances, coding completeness increased  
17 rapidly from below 10% to around 75% overall and around 90% in some regions.  
18 This was achieved through focused academic, professional and managerial actions  
19 (34). However, until now the impact of these improvements on the validity and utility  
20 of the data has not been formally reviewed. In this study, we were able to improve  
21 the completeness of the data through a simple imputation method. We were also  
22 able to compare results based on hospital admission records with a reliable  
23 benchmark provided by a large scale census-based cohort (13,14). While imputation  
24 increased data completeness in this cohort from 76 to 85%, we found that  
25 considerable unexplained variation remained among some ethnic groups, particularly  
26 the White groups. Some more recently introduced ethnicity codes were still clearly  
27 under-recorded suggesting delays in full implementation. Presenting commonly  
28 used health service measures such as admission rates and rate ratios encourages  
29 more focus on the utility and value of these data, not just the completeness, and  
30 emphasises the need to improve data quality in line with other routinely collected  
31 service data.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

56 Taken together, the previous published results and the present paper indicate that  
57 no country has yet found a way of guaranteeing accurate recording of ethnic group in  
58  
59  
60

1  
2  
3 routine health data systems. This study provides a valuable evidence base to aid  
4  
5 future improvements to ethnicity monitoring data in Scotland. Our work suggests low  
6  
7 cost imputation and linkage methods can help increase the validity of the routine  
8  
9 data. Census- or population register-based databases are clearly more robust but  
10  
11 are expensive to set up and maintain and cannot provide data in real time. If policy-  
12  
13 makers are committed to demonstrating ethnic equality in health care through  
14  
15 monitoring, as in Scotland, then continued efforts will be needed to improve and  
16  
17 monitor the accuracy of data recorded on administrative systems, in addition to their  
18  
19 completeness.  
20  
21  
22  
23  
24  
25

### 26 ***Limitations of the study***

27  
28 This demonstration project is restricted to two outcomes for patients hospitalised in a  
29  
30 single year. As the first analysis of such routine data for the whole of Scotland, it is  
31  
32 open to refinement. We could have used other methods of imputing missing ethnic  
33  
34 codes, for example the most frequently used code rather than the most recent.  
35  
36 However, on balance, we decided the most recent was likely to be the most  
37  
38 accurate. We assumed that all the records with unknown values after imputation  
39  
40 were missing at random and could be assigned according to the known distribution  
41  
42 of ethnic codes in each 5-year age group by sex. This assumption would be flawed if  
43  
44 the records with missing values were seriously biased towards particular ethnic  
45  
46 groups. We had no way of assessing this although comparison of demographic and  
47  
48 other more complete admission variables did not indicate a more general bias.  
49  
50  
51  
52  
53  
54  
55

56 We restricted imputation to a single available dataset of hospital admissions.

57  
58 Including additional routinely available datasets could have improved ethnicity  
59  
60

1  
2  
3 completeness further though this would have increased the complexity of linkage;  
4 ethnicity recording in non-hospital-based datasets remains low. Other methods of  
5 imputation could have been considered such as using geographically and  
6 demographically close donor records though this is likely to be problematic given the  
7 relatively low ethnic minority population in Scotland and others have found this to  
8 perform poorly (30, 32).  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

19 The year 2013 was chosen to benefit from the improvements in completeness of  
20 ethnicity seen in the preceding years but yet remain close to the 2011 census as the  
21 source of population denominator data for rates. However, some degree of  
22 numerator-denominator bias was still likely in the rates. We made no attempt to  
23 extrapolate the Census populations by ethnic group to provide denominators for later  
24 years in this study. While desirable, getting accurate estimates of ethnic group  
25 populations between decennial census years remains a significant challenge.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 There was no gold standard benchmark for the comparison of the validity of ethnicity  
39 recording in hospital records. However we aimed to advance the debate around  
40 ethnicity monitoring by demonstrating the potential utility of the data now available  
41 through much effort in Scotland. Therefore we drew general comparison with the  
42 published results from the SHELS programme. That cohort was based on the 2001  
43 census followed up for a number of years and thus not fully contemporaneous with  
44 the 2013 data set we used. There were some differences in statistical methods it  
45 used to calculate the standardised rates. It also did not include White Polish, White  
46 Gypsy and Arab groups, as these were only introduced from at the 2011 census.  
47 While the 2013 dataset included all ICD10 codes for IHD, SHELS only included  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 diagnoses of acute myocardial infarction. However, given the broad similarities of the  
4  
5 epidemiology of different IHD diagnoses, this is unlikely to have affected the  
6  
7 comparisons. We adjusted rates for age and sex, which vary in distribution across  
8  
9 ethnic groups, but not for socio-economic variables. However these have been  
10  
11 shown to have little effect on variation in admission rates among ethnic groups (13).  
12  
13  
14  
15  
16

## 17 **CONCLUSIONS**

19 In this first national analysis of routinely collected hospital admission data analysed  
20  
21 by ethnic group in Scotland, we have confirmed relatively high rates of ethnic group  
22  
23 recording and demonstrated simple methods to increase the utility of the data for  
24  
25 analytical purposes. Analysis of routine hospital records by ethnicity brings many  
26  
27 benefits. However, the validity of admission rates based on these data is variable  
28  
29 across ethnic groups. As in other countries, there remains the continuing challenge  
30  
31 of achieving and sustaining high levels of accurate ethnic coding in routine health  
32  
33 systems consistently across all ethnic groups.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## References

1 Bhopal RS. Migration, Ethnicity, Race and Health in Multicultural Societies. 2nd ed. Oxford: Oxford University Press; 2014.

2 World Health Organization. Health of Migrants - The Way Forward: Report of a global consultation. Madrid, Spain, 3-5 March 2010: World Health Organization, 2010.  
[https://apps.who.int/iris/bitstream/handle/10665/44336/9789241599504\\_eng.pdf?sequence=1&isAllowed=y](https://apps.who.int/iris/bitstream/handle/10665/44336/9789241599504_eng.pdf?sequence=1&isAllowed=y) (last accessed 17 April 2019)

3 Bhopal RS. The quest for culturally sensitive health-care systems in Scotland: insights for a multi-ethnic Europe. Journal of Public Health 2012;34:5-11  
<https://doi.org/10.1093/pubmed/fdr094> (last accessed 17 April 2019)

4 Heath I. The role of ethnic monitoring in general practice. British Journal of General Practice. 1991;41:310-11.

5 NHS Executive. Collecting Ethnic Group Data for Admitted Patient Care : Implementation Guidance and Training Material. Leeds: NHS Executive; 1994.

6 ISD Scotland. Communications Guidelines for the Introduction of Ethnic monitoring in Health Boards in Scotland; Edinburgh: ISD Scotland; 2006.  
<http://www.equalitiesinhealth.org/publications/COMMUNICATIONSGUIDELINES.pdf> (last accessed 17 April 2019)

7 ISD Scotland. Ethnic Monitoring Tool. Edinburgh: ISD Scotland; 2005.  
[http://www.equalitiesinhealth.org/public\\_html/publications/ETHNIC\\_MONITORING\\_TOOL.pdf](http://www.equalitiesinhealth.org/public_html/publications/ETHNIC_MONITORING_TOOL.pdf) (last accessed 17 April 2019)

8 ISD Scotland. Improving data collection for equality and diversity monitoring All Scotland: Ethnicity Completeness in SMR01 and SMR00; Edinburgh: ISD Scotland;.

1  
2  
3 2011 [http://www.isdscotland.org/Health-Topics/Equality-and-](http://www.isdscotland.org/Health-Topics/Equality-and-Diversity/Publications/Briefing%20Paper0211-Feb11%20_JJ.pdf)  
4 [Diversity/Publications/Briefing%20Paper0211-Feb11%20\\_JJ.pdf](http://www.isdscotland.org/Health-Topics/Equality-and-Diversity/Publications/Briefing%20Paper0211-Feb11%20_JJ.pdf) (last accessed 17  
5  
6 [April 2019](http://www.isdscotland.org/Health-Topics/Equality-and-Diversity/Publications/Briefing%20Paper0211-Feb11%20_JJ.pdf))  
7  
8

9 UK Government. Equality Act 2010  
10  
11 [http://www.legislation.gov.uk/ukpga/2010/15/pdfs/ukpga\\_20100015\\_en.pdf](http://www.legislation.gov.uk/ukpga/2010/15/pdfs/ukpga_20100015_en.pdf)  
12  
13 (Last accessed 17 April 2019)  
14  
15

16  
17  
18  
19 10 ISD Scotland. Improving ethnic data collection for equality and diversity  
20 monitoring October 2010 – September 2012; Edinburgh: ISD Scotland; 2013  
21  
22 [https://www.isdscotland.org/Health-Topics/Equality-and-Diversity/Publications/2013-](https://www.isdscotland.org/Health-Topics/Equality-and-Diversity/Publications/2013-02-26/2013-02-26-EDIP-Report.pdf?34324282408)  
23 [02-26/2013-02-26-EDIP-Report.pdf?34324282408](https://www.isdscotland.org/Health-Topics/Equality-and-Diversity/Publications/2013-02-26/2013-02-26-EDIP-Report.pdf?34324282408) (last accessed 10 April 2019).  
24  
25  
26

27  
28  
29  
30 11 ISD Scotland. Improving ethnic data collection for equality and diversity  
31 monitoring July – September 2018; Edinburgh: ISD Scotland; 2019  
32  
33 [https://www.isdscotland.org/products-and-Services/Data-Support-and-](https://www.isdscotland.org/products-and-Services/Data-Support-and-Monitoring/SMR-Ethnic-Group-Recording/)  
34 [Monitoring/SMR-Ethnic-Group-Recording/](https://www.isdscotland.org/products-and-Services/Data-Support-and-Monitoring/SMR-Ethnic-Group-Recording/) (last accessed 17 April 2019)  
35  
36  
37

38  
39  
40  
41 12 Bhopal R, Fischbacher C, Povey C, Chalmers J, Mueller G, Steiner M, et al.  
42 Cohort profile: Scottish Health and Ethnicity Linkage Study of 4.65 million people  
43 exploring ethnic variations in disease in Scotland. *Int J Epidemiol.* 2011;40:1168-75.  
44  
45  
46

47  
48  
49  
50 13 Gruer L, Millard A, Williams L, Bhopal R, Katikireddi S V, Cezard G, et al.  
51 Differences in all-cause hospitalisation by ethnic group: a data linkage cohort study  
52 of 4.62 million people in Scotland, 2001-2013. *Public Health* 2018;161:5-11.  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 14 Bansal N, Fischbacher CM, Bhopal RS, Brown H, Steiner MF, Capewell S, et al.  
4  
5 Myocardial infarction incidence and survival by ethnic group: Scottish Health and  
6  
7 Ethnicity Linkage retrospective cohort study. *BMJ Open* 2013;3:  
8  
9 e003415.doi:10.1136/bmjopen-2013-003415.  
10  
11  
12  
13  
14  
15  
16 15 Bhopal R, Steiner MF, Cezard G, Bansal N, Fischbacher C, Simpson CR, et al.  
17  
18 Risk of respiratory hospitalization and death, readmission and subsequent mortality:  
19  
20 Scottish health and ethnicity linkage study. *EurJPublicHealth* 2015;25:769-74.  
21  
22  
23  
24  
25 16 Cezard GI, Bhopal RS, Ward HJ, Bansal N, Bhala N. Ethnic variations in upper  
26  
27 gastrointestinal hospitalizations and deaths: the Scottish Health and Ethnicity  
28  
29 Linkage Study. *EurJPublic Health*. 2015;26:254-60.  
30  
31  
32  
33  
34 17 Gruer L, Cezard G, Clark E, Douglas A, Steiner M, Millard A, et al. Life  
35  
36 expectancy of different ethnic groups using death records linked to population  
37  
38 census data for 4.62 million people in Scotland. *Journal of Epidemiology and*  
39  
40 *Community Health*. 2016;70:1251-54.  
41  
42  
43  
44  
45  
46 18 Millard A, Guthrie C, Fischbacher C, Jamieson J. Pilot ethnic analysis of routine  
47  
48 hospital admissions data and comparison with census linked data: IHD rates remain  
49  
50 high in Pakistanis. *Ethnicity and Inequalities in Health and Social Care*. 2012;5:98-  
51  
52 107.  
53  
54  
55  
56  
57 19 Fischbacher CM, Bhopal R, Povey C, Steiner M, Chalmers J, Mueller G, et al.  
58  
59 Record linked retrospective cohort study of 4.6 million people exploring ethnic  
60

1  
2  
3 variations in disease: myocardial infarction in South Asians. *BMCPublic Health*.  
4  
5 2007;7:142.  
6  
7  
8  
9

10 20 Bhala N, Cezard G, Ward HJ, Bansal N, Bhopal R. Ethnic Variations in Liver-  
11 and Alcohol-Related Disease Hospitalisations and Mortality: The Scottish Health and  
12 Ethnicity Linkage Study. *Alcohol and Alcoholism*, 2016;51:593-601.  
13  
14  
15  
16  
17  
18

19 21 Katikireddi, SV, Cezard, G, Bhopal, RS, Williams, L, Douglas, A, Millard, A et al.  
20 Assessment of health care, hospital admissions, and mortality by ethnicity:  
21 population-based cohort study of health-system performance in Scotland. *Lancet*.  
22 *Public Health* 2018;3:e226-e236. doi: 10.1016/S2468-2667(18)30068-9  
23  
24  
25  
26  
27

28 22 2013 European Standard Population  
29 ([http://ec.europa.eu/eurostat/en/web/products-manuals-and-guidelines/-/KS-RA-13-](http://ec.europa.eu/eurostat/en/web/products-manuals-and-guidelines/-/KS-RA-13-028)  
30 [028](http://ec.europa.eu/eurostat/en/web/products-manuals-and-guidelines/-/KS-RA-13-028)). (last accessed 17 April 2019).  
31  
32  
33  
34  
35  
36

37 23 Rothman KJ, Greenland S, Lash TL. *Modern Epidemiology* 3<sup>rd</sup> Ed. Philadelphia:  
38 LWW.com; 2008.  
39  
40  
41  
42  
43

44 24 Rafnsson SB, Bhopal RS. Large-scale epidemiological data on cardiovascular  
45 diseases and diabetes in migrant and ethnic minority groups in Europe. *Eur J Public*  
46 *Health*. 2009;19(5):484-91.  
47  
48  
49  
50  
51  
52

53 25 National Cancer Intelligence Network & Cancer Research UK. *Cancer Incidence*  
54 *and Survival by Major Ethnic Group, England, 2002 - 2006*. London: Cancer  
55 *Research UK*; 2009.  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 26 Jack RH, Davies ES, Moller H. Breast cancer incidence, stage, treatment and  
7 survival in ethnic groups in South East England. *Br J Cancer* 2009;100:545–50.  
8  
9

10  
11  
12 27 Jack RH, Davies EA, Moller H. Prostate cancer incidence, stage at diagnosis,  
13 treatment and survival in ethnic groups in South-East England. *BJUInt*.  
14  
15 2010;105(9):1226-30.  
16  
17  
18

19  
20  
21 28 Coupland VH, Lagergren J, Konfortion J, Allum W, Mendall MA, Hardwick RH, et  
22 al. Ethnicity in relation to incidence of oesophageal and gastric cancer in England.  
23  
24 *BrJCancer*. 2012;107:1908-14.  
25  
26  
27

28  
29  
30 29 Mathur R, Bhaskaran K, Chaturvedi N, Leon DA, vanStaa T, Grundy E, Smeeth  
31 L. Completeness and usability of ethnicity data in UK-based primary care and  
32 hospital database. *Journal of Public Health*. 2014; 36(4): 684-692.  
33  
34  
35  
36

37  
38  
39 30 Aspinall PJ, Jacobson B. Why poor quality of ethnicity data should not preclude  
40 its use for identifying disparities in health and healthcare. *QualSaf Health Care*.  
41  
42 2007;16:176-80.  
43  
44  
45

46  
47  
48 31 Saunders CL, Abel GA, El Turabi A, Ahmed F, Lyratzopoulos G. Accuracy of  
49 routinely recorded ethnic group information compared with self-reported ethnicity:  
50 evidence from the English Cancer Patient Experience Survey. *BMJ Open*. 2013; 3:  
51  
52 e002882.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 32 Ryan R, Vernon S, Lawrence G, Wilson S. Use of name recognition software  
4 census data, and multiple imputation to predict missing data on ethnicity: application  
5 to cancer registry records. BMC Medical Informatics and Decision Making. 2012:  
6 12:3.  
7  
8  
9

10  
11  
12  
13  
14 33 Cormack D & McLeod M. Improving and maintaining quality in ethnicity data  
15 collections in the health and disability sector. Wellington, New Zealand, 2010.  
16  
17  
18

19  
20  
21 34 Douglas A, Glover J, Bhopal R. Lothian Ethnic Coding Task Force, March 2009-  
22 March 2012, Final Report. Edinburgh: NHS Lothian 2012.  
23  
24  
25  
26  
27  
28  
29  
30  
31

### 32 **Acknowledgements**

33  
34 We would like to thank Maighread Simpson for her contribution to the project in the  
35 initial planning stages.  
36  
37  
38  
39  
40

### 41 **Conflicts of Interest:**

42  
43  
44 None  
45  
46  
47  
48

### 49 **Funding:**

50  
51 A proportion of the analysis for the study was supported by a small grant from the  
52 University of Edinburgh.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

For Peer Review

1  
2  
3  
4  
5  
6 **The challenge of using routinely collected data to compare hospital**  
7 **admission rates by ethnic group: a demonstration project in**  
8 **Scotland**  
9  
10  
11  
12  
13

14 Knox S<sup>1</sup>, Bhopal RS<sup>2</sup>, Thomson CS<sup>1</sup>, Millard A<sup>3</sup>, Fraser A<sup>4</sup>, Gruer L<sup>2</sup>, Buchanan D<sup>1</sup>

16  
17 Knox S BSc, Senior Information Analyst

18  
19 Bhopal RS Professor, Emeritus Professor of Public Health

20  
21 Thomson CS MSc, BSc, Information Lead for Population Health

22  
23 Andrew Millard, Ph.D., M.P.H., M.Sc., B.A., Research Fellow

24  
25 Dr Andrew Fraser MPH, Director of Public Health Science

26  
27 Gruer L BSc MB ChB MD MPH, Honorary Professor of Public Health

28  
29 Buchanan D BSc, C Stat, Head of Consultancy Services  
30  
31  
32  
33

- 34  
35 1. Information Services Division, NHS National Services Scotland, Gyle Square, 1  
36 South Gyle Crescent, Edinburgh, EH12 9EB, UK  
37  
38 2. Usher Institute of Population Health Sciences and Informatics, University of  
39 Edinburgh, Edinburgh, UK  
40  
41 3. MRC/CSO Social and Public Health Sciences Unit, University of Glasgow,  
42 Glasgow, UK.  
43  
44 4. NHS Health Scotland, Gyle Square, 1 South Gyle Crescent, Edinburgh, EH12  
45 9EB, UK  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 **Corresponding Author:** D Buchanan  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

## Figures Legends


Figure 1 – All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (males)

Figure 2 - All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (females)

Figure 3 Hospital Admissions, All causes in Scotland, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; males

Figure 4 Hospital Admissions, All causes in Scotland for All causes, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; females

**Figure 1 All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (males)**



See Table A1 for mappings and shortened text

1  
2  
3  
4  
5  
6 **The challenge of using routinely collected data to compare hospital**  
7 **admission rates by ethnic group: a demonstration project in**  
8 **Scotland**  
9  
10  
11  
12  
13

14 Knox S<sup>1</sup>, Bhopal RS<sup>2</sup>, Thomson CS<sup>1</sup>, Millard A<sup>3</sup>, Fraser A<sup>4</sup>, Gruer L<sup>2</sup>, Buchanan D<sup>1</sup>

16  
17 Knox S BSc, Senior Information Analyst

18  
19 Bhopal RS Professor, Emeritus Professor of Public Health

20  
21 Thomson CS MSc, BSc, Information Lead for Population Health

22  
23 Andrew Millard, Ph.D., M.P.H., M.Sc., B.A., Research Fellow

24  
25 Dr Andrew Fraser MPH, Director of Public Health Science

26  
27 Gruer L BSc MB ChB MD MPH, Honorary Professor of Public Health

28  
29 Buchanan D BSc, C Stat, Head of Consultancy Services  
30  
31  
32  
33

- 34  
35 1. Information Services Division, NHS National Services Scotland, Gyle Square, 1  
36 South Gyle Crescent, Edinburgh, EH12 9EB, UK  
37  
38 2. Usher Institute of Population Health Sciences and Informatics, University of  
39 Edinburgh, Edinburgh, UK  
40  
41 3. MRC/CSO Social and Public Health Sciences Unit, University of Glasgow,  
42 Glasgow, UK.  
43  
44 4. NHS Health Scotland, Gyle Square, 1 South Gyle Crescent, Edinburgh, EH12  
45 9EB, UK  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 **Corresponding Author:** D Buchanan  
56  
57  
58  
59  
60

## Figures Legends


Figure 1 – All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (males)

Figure 2 - All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (females)

Figure 3 Hospital Admissions, All causes in Scotland, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; males

Figure 4 Hospital Admissions, All causes in Scotland for All causes, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; females

Figure 2 All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (females)



See Table A1 for mappings and shortened text

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

For Peer Review

1  
2  
3  
4  
5  
6 **The challenge of using routinely collected data to compare hospital**  
7 **admission rates by ethnic group: a demonstration project in**  
8 **Scotland**  
9  
10  
11  
12  
13

14 Knox S<sup>1</sup>, Bhopal RS<sup>2</sup>, Thomson CS<sup>1</sup>, Millard A<sup>3</sup>, Fraser A<sup>4</sup>, Gruer L<sup>2</sup>, Buchanan D<sup>1</sup>

16  
17 Knox S BSc, Senior Information Analyst

18  
19 Bhopal RS Professor, Emeritus Professor of Public Health

20  
21 Thomson CS MSc, BSc, Information Lead for Population Health

22  
23 Andrew Millard, Ph.D., M.P.H., M.Sc., B.A., Research Fellow

24  
25 Dr Andrew Fraser MPH, Director of Public Health Science

26  
27 Gruer L BSc MB ChB MD MPH, Honorary Professor of Public Health

28  
29 Buchanan D BSc, C Stat, Head of Consultancy Services  
30  
31  
32  
33

- 34  
35 1. Information Services Division, NHS National Services Scotland, Gyle Square, 1  
36 South Gyle Crescent, Edinburgh, EH12 9EB, UK  
37  
38 2. Usher Institute of Population Health Sciences and Informatics, University of  
39 Edinburgh, Edinburgh, UK  
40  
41 3. MRC/CSO Social and Public Health Sciences Unit, University of Glasgow,  
42 Glasgow, UK.  
43  
44 4. NHS Health Scotland, Gyle Square, 1 South Gyle Crescent, Edinburgh, EH12  
45 9EB, UK  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 **Corresponding Author:** D Buchanan  
56  
57  
58  
59  
60



## Figures Legends


Figure 1 – All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (males)

Figure 2 - All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (females)

Figure 3 Hospital Admissions, All causes in Scotland, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; males

Figure 4 Hospital Admissions, All causes in Scotland for All causes, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; females

**Figure 3 Hospital Admissions, All causes in Scotland, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; males**



See Table A1 for mappings and shortened text

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

For Peer Review

1  
2  
3  
4  
5  
6 **The challenge of using routinely collected data to compare hospital**  
7 **admission rates by ethnic group: a demonstration project in**  
8 **Scotland**  
9  
10  
11  
12  
13

14 Knox S<sup>1</sup>, Bhopal RS<sup>2</sup>, Thomson CS<sup>1</sup>, Millard A<sup>3</sup>, Fraser A<sup>4</sup>, Gruer L<sup>2</sup>, Buchanan D<sup>1</sup>

16  
17 Knox S BSc, Senior Information Analyst

18  
19 Bhopal RS Professor, Emeritus Professor of Public Health

20  
21 Thomson CS MSc, BSc, Information Lead for Population Health

22  
23 Andrew Millard, Ph.D., M.P.H., M.Sc., B.A., Research Fellow

24  
25 Dr Andrew Fraser MPH, Director of Public Health Science

26  
27 Gruer L BSc MB ChB MD MPH, Honorary Professor of Public Health

28  
29 Buchanan D BSc, C Stat, Head of Consultancy Services

- 30  
31  
32  
33  
34  
35 1. Information Services Division, NHS National Services Scotland, Gyle Square, 1  
36 South Gyle Crescent, Edinburgh, EH12 9EB, UK  
37  
38 2. Usher Institute of Population Health Sciences and Informatics, University of  
39 Edinburgh, Edinburgh, UK  
40  
41 3. MRC/CSO Social and Public Health Sciences Unit, University of Glasgow,  
42 Glasgow, UK.  
43  
44 4. NHS Health Scotland, Gyle Square, 1 South Gyle Crescent, Edinburgh, EH12  
45 9EB, UK  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 **Corresponding Author:** D Buchanan  
56  
57  
58  
59  
60

## Figures Legends


Figure 1 – All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (males)

Figure 2 - All-cause Hospital Admissions, Scotland, 2013; European Age-Standardised Rates per 1,000 population (females)

Figure 3 Hospital Admissions, All causes in Scotland, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; males

Figure 4 Hospital Admissions, All causes in Scotland for All causes, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; females

**Figure 4 Hospital Admissions, All causes in Scotland, 2013: European Age Standardised Rate Ratios relative to White Scottish with 95% Confidence intervals after imputation and redistribution; females**



See Table A1 for mappings and shortened text

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1  
2  
3  
4  
5  
6 **The challenge of using routinely collected data to compare hospital**  
7 **admission rates by ethnic group: a demonstration project in**  
8 **Scotland**  
9  
10  
11  
12  
13

14 Knox S<sup>1</sup>, Bhopal RS<sup>2</sup>, Thomson CS<sup>1</sup>, Millard A<sup>3</sup>, Fraser A<sup>4</sup>, Gruer L<sup>2</sup>, Buchanan D<sup>1</sup>

16  
17 Knox S BSc, Senior Information Analyst

18  
19 Bhopal RS Professor, Emeritus Professor of Public Health

20  
21 Thomson CS MSc, BSc, Information Lead for Population Health

22  
23 Andrew Millard, Ph.D., M.P.H., M.Sc., B.A., Research Fellow

24  
25 Dr Andrew Fraser MPH, Director of Public Health Science

26  
27 Gruer L BSc MB ChB MD MPH, Honorary Professor of Public Health

28  
29 Buchanan D BSc, C Stat, Head of Consultancy Services  
30  
31  
32  
33

- 34  
35 1. Information Services Division, NHS National Services Scotland, Gyle Square, 1  
36 South Gyle Crescent, Edinburgh, EH12 9EB, UK  
37  
38 2. Usher Institute of Population Health Sciences and Informatics, University of  
39 Edinburgh, Edinburgh, UK  
40  
41 3. MRC/CSO Social and Public Health Sciences Unit, University of Glasgow,  
42 Glasgow, UK.  
43  
44 4. NHS Health Scotland, Gyle Square, 1 South Gyle Crescent, Edinburgh, EH12  
45 9EB, UK  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 **Corresponding Author:** D Buchanan  
56  
57  
58  
59  
60



## Online supplementary data

### List of Tables

Table A1 – List of the codes from the 2001 Census and 2011 Census recorded in SMR01 records (along with dates they could be used within SMR01)

Table A2 – Impact of method of imputing ethnicity for patient episodes in 2013

Table A3 Rate ratios for ISD and SHELS (Reference 13) analyses for All-cause admissions and the differences between them. White Scottish is the reference group (RR=1.0)

Table A4 Rate ratios for ISD and SHELS (Reference 13) analyses for IHD admissions for Ischaemic Heart Disease (IHD) and the differences between them. White Scottish is the reference group (RR=1.0)

Table A5a: European Age-Standardised Rates (EASR) per 1,000 population for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; males: Original, imputed and imputed and redistributed rates.

Table A5b: EASR per 1,000 population for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; females: Original, imputed and imputed and redistributed rates.

Table A6a: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; males.

Table A6b: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; females.

Table A7a: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for Ischaemic Heart Disease Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; males.

Table A7b: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for Ischaemic Heart Disease Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; females.

**Table A1 – List of the codes from the 2001 Census and 2011 Census recorded in SMR01 records (along with dates they were could be used within SMR01)**

<b>2001 Census Codes</b> (Last acceptable date is 31/03/2012)	<b>Final 2011 Census Codes</b> (Available from 01/10/2011)	<b>Text used within publication</b>
<b>A – White</b>	<b>A – White</b>	<b>A – White</b>
1A Scottish	1A Scottish	White Scottish
1B Other British	1B Other British	White Other British
1C Irish	1C Irish	White Irish
	1K Gypsy/ Traveller	White Gypsy
	1L Polish	White Polish
1D Any other white background	1Z Other white ethnic group	White Other
<b>B – Mixed</b>	<b>B - Mixed or multiple ethnic groups</b>	<b>B - Mixed or multiple ethnic groups</b>
2A Any mixed background	2A Any mixed or multiple ethnic groups	Mixed
<b>C - Asian, Asian Scottish or Asian British</b>	<b>C - Asian, Asian Scottish or Asian British</b>	<b>C - Asian, Asian Scottish or Asian British</b>
3A Indian	3G Indian, Indian Scottish or Indian British	Indian
3B Pakistani	3F Pakistani, Pakistani Scottish or Pakistani British	Pakistani
3C Bangladeshi	3H Bangladeshi, Bangladeshi Scottish or Bangladeshi British	Bangladeshi
3D Chinese	3J Chinese, Chinese Scottish or Chinese British	Chinese
3E Any other Asian background	3Z Other Asian, Asian Scottish or Asian British	Asian Other
<b>D - African, Caribbean or Black</b>	<b>D – African</b>	<b>D – African</b>
4B African	4D African, African Scottish or African British	African
	4Y Other African	African Other
		<b>(these two codes merged into one group known as African)</b>
	<b>E - Caribbean or Black</b>	<b>E - Caribbean or Black</b>
4A Caribbean	5C Caribbean, Caribbean Scottish or Caribbean British	Caribbean
	5D Black, Black Scottish or Black British	Black

4C Any other black background	5Y Other Caribbean or Black	Caribbean or Black Other
		<b>(these three codes merged into one group known as Caribbean/Black)</b>
<b><i>E - Other Ethnic Background</i></b>	<b><i>F - Other ethnic group</i></b>	<b><i>F - Other ethnic group</i></b>
	6A Arab, Arab Scottish or Arab British	Arab
5A Any other ethnic background	6Z Other ethnic group	Other Ethnic Group
<b><i>F - Refused/Not provided by patient</i></b>	<b><i>G - Refused/Not provided by patient</i></b>	<b><i>G - Refused/Not provided by patient</i></b>
98 Refused/Not provided by patient	98 Refused/Not provided by patient	Refused/Not provided by patient
<b><i>G - Not Known</i></b>	<b><i>H - Not Known</i></b>	<b><i>H - Not Known</i></b>
99 Not Known	99 Not Known	Not Known

**Table A2 – Impact of method of imputing ethnicity for patient episodes in 2013**

Ethnic Group	Original data	After imputation	
	No. Of episodes	No. Of episodes	Increase %
White Scottish	982,618	1,084,589	10%
White Other British	119,311	142,545	19%
White Irish	3,998	4,758	19%
White Gypsy	107	125	17%
White Polish	5,038	5,516	9%
White Other	20,094	21,504	7%
Mixed	2,608	2,801	7%
Pakistani	8,139	9,247	14%
Indian	3,692	4,164	13%
Bangladeshi	433	497	15%
Chinese	2,200	2,459	12%
Asian Other	2,251	2,577	14%
African	2,442	2,710	11%
African Other	358	424	18%
Black	433	471	9%
Caribbean	219	277	26%
Caribbean or Black	447	482	8%
Other			
Arab	399	447	12%
Other Ethnic Group	2,149	2,056	-4%
Unknown\Not Provided	362,349	231,636	-36%
<b>Total</b>	<b>1,519,285</b>	<b>1,519,285</b>	
Percentage of Episodes with ethnicity unknown\ not provided	23.8%	15.2%	

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

**Table A3**

**Comparison of rate ratios (RRs) for the current study and SHELS study (Reference 13) for All-cause admissions and the differences between them. White Scottish is the reference group for the current study (RR=1.0)**

Ethnic Group	Male (RR)			Female (RR)				
	Current study	SHELS cohort	Difference	Current study	SHELS cohort*	Difference		
White Scottish	1.00	1.0		1.0	1.0			
White Other British	1.40	0.81	0.59	1.39	0.85	0.54		
White Irish	0.36	1.0	-0.64	0.34	0.95	-0.61		
White Polish	0.78	0.81	-0.03	0.71	0.78	-0.07		
White Other	1.88	0.81	1.07	1.48	0.78	0.70		
Mixed	0.98	0.92	0.06	0.94	0.87	0.07		
Pakistani	1.11	1.19	-0.08	1.28	1.14	0.14		
Indian	0.84	0.95	-0.11	1.05	0.91	0.14		
Bangladeshi	1.05	0.92	0.13	1.26	0.82	0.44		
Chinese	0.64	0.62	0.02	0.55	0.66	-0.11		
Asian Other	1.17	0.92	0.25	1.02	0.91	0.11		
African	0.91	0.79	0.12	0.76	1.07	-0.31		
Caribbean/Black	0.81	0.89	-0.07	1.05	1.16	-0.11		
		Black	0.81	1.18	-0.27	1.05	0.86	0.19

**(\*) Note the rate ratios for the SHELS cohort are based on a different statistical method that means they are not directly comparable. These rates were based on poisson regression models of the risk of hospitalisation for a population-based cohort followed up from the 2001 census to 2013. The SHELS cohort use the 2001 census ethnicity groupings which differ from the 2011 census groupings used in the current study.**

**Table A4**

**Comparison of the rate ratios (RRs) for current study and SHELS study (Reference 14) analyses for Ischaemic Heart Disease (IHD) admissions and the differences between them. White Scottish is the reference group for the current study (RR=1.0)**

Ethnic Group	Males (RR)				Females (RR)			
	Current study	SHELS cohort	Current study	SHELS cohort*	Difference	Current study	SHELS cohort*	Difference
White Scottish			1.0	1.0	-	1.0	1.0	-
White Other British			1.3	0.77	0.53	1.32	0.72	0.6
White Irish			0.39	0.93	-0.54	0.30	0.95	-0.65
White Polish		White Other	0.81	0.82	-0.01	1.10	0.81	0.29
White Other		White Other	1.96	0.82	1.14	1.59	0.81	0.78
Mixed			1.78	1.29	0.49	1.55	1.2	0.35
Pakistani			2.4	1.42	0.98	2.5	1.29	1.21
Indian			1.45	1.21	0.24	1.16	1.24	-0.08
Bangladeshi		Asian Other	2.66	1.33	1.33	3.46	1.53	1.93
Asian Other		Asian Other	1.27	1.33	-0.06	2.04	1.53	0.51
Chinese			0.48	0.45	0.03	0.47	0.71	-0.24
African		African	1.11	1.0	0.11	0.86	1.21	-0.35
Caribbean/Black		African	0.83	1.0	-0.17	1.0	1.21	-0.21

**(\*) Note the rate ratios for the SHELS cohort are based on a different statistical method and disease group that means they are not directly comparable. These rates were based directly standardised rates on of the risk of Acute Myocardial Infarction (ICD codes I21, I22) for a population-based cohort followed up from the 2001 census to 2008. The reference group was the White Scottish population. The SHELS cohort use the 2001 census ethnicity groupings which differ from the 2011 census groupings used in the current study.**

**Table A5a: European Age-Standardised Rates (EASR) per 1,000 population after imputation and redistribution for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; males**

<b>Ethnicity</b>	<b>Original EASR per 1,000 Population</b>	<b>Imputed EASR per 1,000 Population</b>	<b>Imputed &amp; redistributed EASR per 1,000 Population</b>	<b>Lower Confidence Limit</b>	<b>Upper Confidence Limit</b>
White Scottish	175.0	193.3	230.8	230.1	231.5
White Other British	226.0	270.5	322.0	319.4	324.5
White Irish	60.2	70.0	83.2	79.4	86.9
White Gypsy	16.8	26.6	31.7	21.4	41.9
White Polish	130.3	149.8	179.2	167.2	191.1
White Other	337.4	367.7	434.7	424.7	444.7
Mixed	165.1	189.5	226.1	205.7	246.5
Pakistani	184.5	214.0	255.4	245.8	265.0
Indian	139.4	162.3	193.6	182.7	204.5
Bangladeshi	192.1	204.7	242.9	206.1	279.6
Chinese	112.7	123.6	146.9	136.3	157.4
Asian Other	188.9	227.4	270.1	245.2	295.0
African	152.1	176.7	210.0	188.2	231.8
Caribbean/Black	155.7	155.6	187.8	167.3	208.2
Arab	54.0	56.0	66.9	53.0	80.9
Other Ethnic Group	574.7	530.6	579.2	531.5	627.0

See Table A1 for mappings and shortened text

**Table A5b: European Age-Standardised Rates (EASR) per 1,000 population after imputation and redistribution for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; females**

<b>Ethnicity</b>	<b>Original EASR per 1,000 Population</b>	<b>Imputed EASR per 1,000 Population</b>	<b>Imputed &amp; Redistributed EASR per 1,000 Population</b>	<b>Lower Confidence Limit</b>	<b>Upper Confidence Limit</b>
White Scottish	175.1	192.8	229.2	228.5	229.8
White Other British	227.4	268.6	318.7	316.2	321.1
White Irish	54.0	64.7	76.7	73.4	80.1
White Gypsy	35.3	37.6	44.6	32.1	57.1
White Polish	127.2	136.5	162.0	149.5	174.6
White Other	271.4	288.0	340.0	332.6	347.5
Mixed	158.1	181.1	215.2	196.3	234.0
Pakistani	215.8	246.1	292.2	281.1	303.3
Indian	176.1	202.3	239.7	226.5	252.9
Bangladeshi	180.1	244.7	289.5	237.2	341.7
Chinese	90.1	105.8	125.6	116.4	134.8
Asian Other	178.2	197.5	233.3	212.7	253.9
African	120.7	145.7	173.2	155.1	191.3
Caribbean/Black	170.5	200.9	239.7	216.5	263.0
Arab	33.4	39.9	47.6	35.9	59.4
Other Ethnic Group	552.2	583.4	691.8	636.7	747.0

See Table A1 for mappings and shortened text



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

**Table A6a: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; males**

<b>Ethnicity</b>	<b>Rate Ratio</b>	<b>Lower Confidence Limit</b>	<b>Upper Confidence Limit</b>
White Scottish	1.000	0.000	0.000
White Other British	1.395	1.383	1.407
White Irish	0.360	0.344	0.377
White Polish	0.776	0.726	0.830
White Other	1.884	1.841	1.928
Mixed	0.980	0.895	1.072
Pakistani	1.107	1.066	1.149
Indian	0.839	0.793	0.888
Bangladeshi	1.052	0.905	1.224
Chinese	0.636	0.592	0.684
Asian Other	1.170	1.067	1.284
African	0.910	0.820	1.009
Caribbean/Black	0.814	0.730	0.907

See Table A1 for mappings and shortened text

**Table A6b: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for All-cause Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; females**

<b>Ethnicity</b>	<b>Rate Ratio</b>	<b>Lower Confidence Limit</b>	<b>Upper Confidence Limit</b>
White Scottish	1.000	0.000	0.000
White Other British	1.391	1.379	1.402
White Irish	0.335	0.321	0.350
White Polish	0.707	0.654	0.764
White Other	1.484	1.452	1.517
Mixed	0.939	0.860	1.025
Pakistani	1.275	1.227	1.324
Indian	1.046	0.990	1.105
Bangladeshi	1.263	1.054	1.513
Chinese	0.548	0.509	0.590
Asian Other	1.018	0.932	1.112
African	0.756	0.681	0.839
Caribbean/Black	1.046	0.949	1.153

See Table A1 for mappings and shortened text

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

**Table A7a: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for Ischaemic Heart Disease Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; males**

<b>Ethnicity</b>	<b>Rate Ratio</b>	<b>Lower Confidence Limit</b>	<b>Upper Confidence Limit</b>
White Scottish	1.000	0.000	0.000
White Other British	1.296	1.260	1.334
White Irish	0.394	0.344	0.452
White Polish	0.812	0.648	1.017
White Other	1.958	1.808	2.121
Mixed	1.775	1.374	2.293
Pakistani	2.402	2.186	2.640
Indian	1.450	1.237	1.699
Bangladeshi	2.657	1.878	3.759
Chinese	0.476	0.352	0.645
Asian Other	1.274	0.925	1.755
African	1.105	0.749	1.630
Caribbean/Black	0.831	0.512	1.347

See Table A1 for mappings and shortened text

**Table A7b: European Age-Standardised Rate Ratios (RRs) after imputation and redistribution for Ischaemic Heart Disease Hospital Admissions in Scotland 2013 with 95% Confidence Intervals; females**

<b>Ethnicity</b>	<b>Rate Ratio</b>	<b>Lower Confidence Limit</b>	<b>Upper Confidence Limit</b>
White Scottish	1.000	0.000	0.000
White Other British	1.319	1.268	1.372
White Irish	0.301	0.246	0.370
White Polish	1.095	0.769	1.560
White Other	1.593	1.425	1.780
Mixed	1.550	1.040	2.310
Pakistani	2.501	2.150	2.910
Indian	1.161	0.879	1.534
Bangladeshi	3.464	1.851	6.483
Chinese	0.468	0.291	0.752
Asian Other	2.041	1.428	2.918
African	0.858	0.445	1.655
Caribbean/Black	0.999	0.530	1.884

See Table A1 for mappings and shortened text