



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Chronological and dietary aspects of the human burials from Ajdovska Cave, Slovenia

Citation for published version:

Bonsall, C, Horvat, M, McSweeney, K, Masson, M, Higham, T, Pickard, C & Cook, G 2007, 'Chronological and dietary aspects of the human burials from Ajdovska Cave, Slovenia', *Radiocarbon*, vol. 49, no. 2, pp. 727-740. <<https://journals.uair.arizona.edu/index.php/radiocarbon/article/view/2968>>

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Radiocarbon

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



CHRONOLOGICAL AND DIETARY ASPECTS OF THE HUMAN BURIALS FROM AJDOVSKA CAVE, SLOVENIA

C Bonsall^{1,2} • M Horvat³ • K McSweeney¹ • M Masson¹ • T F G Higham⁴ • C Pickard¹ • G T Cook⁵

ABSTRACT. Ajdovska Jama (The Pagan's Cave) in southeast Slovenia lies within the catchment of the River Sava, a major tributary of the Danube. The site is well known for its Neolithic burials and has been excavated to a high standard on various occasions since 1884. The human remains at the site occurred as distinct clusters of mainly disarticulated bones belonging to at least 31 individuals. Hitherto, dating of the burials has been based on the associated archaeological finds, including a few low-precision radiometric radiocarbon measurements on charred plant material. In the present study, bones from 15 individuals were subsampled for accelerator mass spectrometry (AMS) and stable isotope analyses. These comprised adults and children from 3 of the clusters. The results of the study indicate that the burials all belong to a relatively short time interval, while the stable isotope data indicate a mixed diet based on C₃ plant and animal food sources. These interpretations differ somewhat from those of previous researchers. The AMS ¹⁴C and stable isotope analyses form part of a wider investigation of dietary and demographic change from the Mesolithic to the Iron Age in the Danube Basin.

INTRODUCTION

This paper forms part of a wider study of prehistoric human populations in the Danube Basin, which focuses particularly on mortality profiles, variations in health and diet, and the incidence of violent trauma. Our sample populations are drawn from different archaeological periods and environments spanning the period from Mesolithic to Iron Age (~10,000–2000 BP) (Figure 1).

Our initial research concentrated on the Iron Gates section of the Danube Valley along the Romania/Serbia border (Bonsall et al. 1997, 2000, 2004). The Iron Gates has the largest concentration of Mesolithic open-air settlements in southeast Europe. More than 30 sites have been investigated along this 230-km stretch of the Danube and human remains have been recovered from 11 of these, with 4 sites—Lepenski Vir, Padina, and Vlasac in Serbia; and Schela Cladovei in Romania—each producing more than a hundred burials. Many of these sites were also occupied during later periods. Radiocarbon dating demonstrates that the burials in the Iron Gates sites range in age from early Mesolithic (~10,000 BP) to post-Medieval (~400 BP) (Bonsall et al. 2004).

Our research in the Iron Gates, involving paired accelerator mass spectrometry (AMS) ¹⁴C and stable (C and N) isotope measurements on human bone collagen, has indicated a heavy dietary emphasis on freshwater resources throughout the Mesolithic (Bonsall et al. 1997, 2000, 2004). It also demonstrated a ¹⁴C reservoir effect, of a similar magnitude to the marine reservoir effect, due to consumption of these resources. This significantly influences the dating of human remains, which are important for developing archaeological chronologies in the Iron Gates region. However, it was possible to correct for this effect using the $\delta^{15}\text{N}$ value of the human bone collagen as an indicator of the extent of freshwater resource consumption and the offset between paired ungulate and human bones as an indicator of the magnitude of the effect (Cook et al. 2001, 2002).

¹Archaeology, University of Edinburgh, Infirmary Street, Edinburgh EH1 1LT, United Kingdom.

²Corresponding author. Email: C.Bonsall@ed.ac.uk.

³Archaeology, University of Ljubljana, Slovenia.

⁴Oxford Radiocarbon Accelerator Unit, Research Laboratory for Archaeology and the History of Art, Oxford University, Dyson Perrins Building, Oxford OX1 3QY, United Kingdom.

⁵SUERC, Scottish Enterprise Technology Park, East Kilbride G75 0QF, United Kingdom.



Figure 1 Map of the Danube Basin showing the locations of Ajdovska Cave and other sample populations included in the wider research program.

There is a marked change in the stable isotope values around 6000 cal BC. The median $\delta^{15}\text{N}$ for 12 burials from Lepenski Vir dated between ~6300–6000 cal BC is +15.7‰, while the median value for 5 burials dated between 6000 and 5600 cal BC is +10.8‰. This suggests a substantial increase in the consumption of terrestrial foods, which has been interpreted as reflecting the transition to farming in this region (Bonsall et al. 1997, 2000). If so, the transition occurred no later in the Iron Gates than elsewhere in neighboring areas of the northern Balkans. This evidence contradicts previous ideas of a lengthy “survival” of the Mesolithic hunter-gatherer way of life in the Iron Gates. Nevertheless, the first farmers within the Iron Gates continued to use freshwater resources, as evidenced by i) $\delta^{15}\text{N}$ values for human bone collagen that are still enriched relative to those typical of a totally terrestrial-based diet, and ii) the occurrence of large numbers of fish bones in the archaeological record of this period in the Iron Gates (Bartosiewicz et al. 2001).

The continued use of aquatic resources in the earliest Neolithic is not confined to the Iron Gates. Stable isotope data suggest significant use of freshwater resources at some Early Neolithic (Körös culture) sites on the Hungarian Plain (Figure 2). This is perhaps not surprising given that the rivers in this region are some of the largest in Europe and between 6000–5500 cal BC the Danube Basin was effectively the agricultural “frontier.”




Figure 2 Carbon and nitrogen isotope ratios for human bone collagen from Early Neolithic sites on the Hungarian Plain plotted against the typical ranges for domestic and wild herbivores (data from Whittle et al. 2002). In this environment, a $\delta^{13}\text{C}$ value lighter than approximately -21‰ combined with a $\delta^{15}\text{N}$ value heavier than approximately $+10\text{‰}$ suggests a diet that included significant amounts of protein from freshwater fish. Whittle et al. (2002) interpreted the skeleton with the lightest $\delta^{13}\text{C}$ value as “Mesolithic”; however, the contracted body position is characteristic of the Early Neolithic, and the ^{14}C age of 7765 ± 55 BP may include a significant reservoir effect owing to consumption of river fish (Bonsall et al. 2004).

Some later burials at Lepenski Vir that have $\delta^{15}\text{N}$ values within the Early Neolithic range exhibit significantly heavier $\delta^{13}\text{C}$ values. These are thought to be due to the inclusion of millet (*Panicum miliaceum* or *Setaria italica*), an introduced C_4 crop plant, in the diet either directly or indirectly through its use as fodder for livestock (Bonsall et al. 2000). Judging from the ^{14}C evidence this change occurred in the Iron Gates sometime between the Chalcolithic (~ 4000 cal BC) and the Roman period (\sim AD 300), possibly in the latter part of this time range. Elsewhere in the Danube Basin, elevated $\delta^{13}\text{C}$ values consistent with direct or indirect consumption of millet are clearly evident in skeletons belonging to the pre-Roman Iron Age (Murray and Schoeninger 1988; Whittle et al. 2002). Tracing the rise of millet cultivation in the Danube Basin is one of our research priorities. Broomcorn millet (*Panicum miliaceum*) occurs sporadically in the archaeobotanical record of southeast Europe as early as 5500–6000 cal BC (Zohary and Hopf 1994). Throughout the region as a whole, it appears to have been a minor crop for several millennia after its introduction, though locally it may have been important. Conceivably, millet consumption accounts for the slightly heavier $\delta^{13}\text{C}$ values exhibited by Late Neolithic/Bronze Age skeletons from the Hungarian Plain compared to earlier Neolithic populations (Figure 2), although more data are needed to test this hypothesis.




Figure 3 Plan of Ajdovska Jama (Slovenia) showing the locations of the Neolithic burials.

AJDOVSKA JAMA

Ajdovska Jama (The Pagan's Cave), which is the focus of this paper, is situated in southeast Slovenia near the city of Krško ($45^{\circ}58'N$, $15^{\circ}29'E$). Occupying an east-facing position in limestone hills 5 km from the River Sava, the cave has 2 entrances at 240 and 244 m asl, respectively, which lead along passages to a central chamber ~ 17 m in diameter (Figure 3). Archaeological investigations, conducted on various occasions since 1884, have demonstrated periodic human use of the cave from the Paleolithic to the Middle Ages.

The site is well known for its burial remains, the majority of which were recovered in excavations by Milena Horvat between 1982 and 1987 (Horvat 1989). The human bones occurred in a horizon ~ 10 cm thick, which was divided into 2 stratigraphic units (43 and 44), on the basis of their colors and the differing amounts of charcoal present. Five distinct clusters of bones were identified, 4 in the south passage and 1 toward the rear of the main chamber (Figure 3). Bone preservation was generally poor and the crania were missing. Therefore, aging and sexing of the burials were based on surviving long bones and mandibles (Corrain and Capitanio 1991). The entire assemblage comprises a minimum of 31 individuals—15 adults (7 males and 8 females) and 16 children. On the basis of associated pottery and other archaeological finds, the burials were assigned to the Late Neolithic Alpine Lengyel culture, and this is broadly confirmed by radiometric ^{14}C ages on charred plant material from the same archaeological horizon, which range from 5620 ± 130 (Z-1044) to 5120 ± 130 BP (Z-1042), with outliers of 4700 ± 200 BP (Z-1179) and 2900 ± 120 BP (Z-1603) (Srdoč et al. 1984, 1987, 1989, 1992; Obelić et al. 1994).

The cave has been interpreted as a “necropolis” used by a farming community that practiced excarnation. It has been suggested that the dead were first left exposed on the cave floor (possibly in the north passage) and once the skeletons were defleshed, some of the bones were collected up and deposited in the various locations shown in Figure 3. In some cases, stones were placed around the bone clusters, defining the burial area. Personal items were placed next to the bodies, including items of jewelry with the women and weapons with the men. Food items were also placed next to the bodies. These were represented by bones of cattle, sheep, and deer, and pots containing carbonized remains of cereals and pulses (Horvat 1989).

The human remains from Ajdovska Jama were the subject of a previous paleodietary study by Ogrinc (1999; Ogrinc and Budja 2005) in which the carbon and nitrogen isotopes of the bone collagen were analyzed, although no direct ¹⁴C dating of the bones was undertaken. Ogrinc analyzed bones from both stratigraphic units in 3 of the clusters (Figure 3 A–C). She assumed that the 2 stratigraphic units related to distinct periods: the lower unit (44) she dated to ~6000 BP (revised to 6400 BP by Ogrinc and Budja [2005]) and the upper unit (43) to ~5300 BP. In total, Ogrinc undertook stable isotope analysis of bones from 25 individuals (adults and children) (Figure 4a); she also analyzed animal bones and plant remains to obtain comparative information on the local food web.




Figure 4 Carbon and nitrogen stable isotope ratios for human bone collagen from Ajdovska Jama obtained by Ogrinc (1999): a) all samples analyzed (adults + children); b) infants (0–2 yr) and adult females; c) adults, apparently showing 2 groups with differing $\delta^{15}\text{N}$ signatures.

Ogrinc (1999) interpreted the stable isotope results for the human bones as indicative of a population with a relatively homogeneous diet that consisted mainly of meat from domestic and wild herbivores, but in which individual food preferences could be observed. She was also able to show a “nursing effect” where the bone collagen $\delta^{15}\text{N}$ values of infants (0–2 yr) were on average $\sim 2\text{‰}$ heavier than the adult females (Figure 4b), whereas older children did not consistently show this effect. Ogrinc and Budja (2005) applied the concentration-weighted linear mixing model (IsoConc) proposed by Phillips and Koch (2002) and re-interpreted the stable isotope data as representing a diet based mainly on meat from domestic and wild herbivores (average of 44%) and cereals (average of 39%).

The present study undertook paired AMS ^{14}C and stable C and N isotope analyses of bone collagen from 15 individuals that had previously been analyzed by Ogrinc. These bones came from the 3 richest clusters (A, B, and C) and both stratigraphic units (43 and 44). The objectives were to: i) obtain direct ^{14}C age measurements on the human bones and compare the results with the radiometric ages previously obtained on associated materials; ii) investigate any age and stable isotope differences between the different bone clusters and stratigraphic units and compare the stable isotope data with Ogrinc’s previous study; and iii) assess the contributions of freshwater resources and C_4 plants to the human diet.

MATERIALS AND METHODS

The bone fragments were decalcified and gelatinized using standard techniques, and transferred to an ultrafilter (VivaspinTM 15, 30 kD MWCO) and centrifuged. The ultrafilter retains the $>30\text{-kD}$ molecular weight fraction, which will include un-degraded collagen. The $<30\text{-kD}$ fraction, which contains low molecular weight components such as salts, degraded collagen fragments, and sometimes soil-derived contaminants, was discarded. The $>30\text{-kD}$ fraction was recovered and freeze-dried in preparation for analysis. The analysis system comprised a Europa Scientific ANCA-MS system consisting of a 20–20 isotope ratio mass spectrometer interfaced to a Roboprep CHN sample converter unit, operating in continuous-flow mode. This enables the measurement of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$, nitrogen and carbon content, and C/N ratios. Graphite was prepared from the CO_2 prior to AMS ^{14}C measurement using published techniques (Bronk Ramsey and Hedges 1997; Bronk Ramsey et al. 2000).

The ultrafiltrated gelatin usually produces collagen of an improved quality, as shown by the C/N ratios (Bronk Ramsey et al. 2004). Collagen quality and chemical integrity were assessed using the atomic ratio of carbon to nitrogen (C/N atomic ratio), the percentage of collagen extracted compared with the starting weight of bone (wt% collagen) and the carbon yield of the collagen on combustion. Problem bones may be screened on the basis of these parameters. Bone is considered acceptable if measured C/N ratios of collagen fall between 2.9 and 3.5. In addition, bone that is composed of less than 1 wt% collagen is not dated.

RESULTS AND DISCUSSION

The AMS ^{14}C and stable isotope results from the present study are presented in Table 1 along with the corresponding stable isotope data obtained by Ogrinc (1999). Of the 15 samples analyzed, only 10 produced collagen suitable for analysis. The ^{14}C results all fall within a tightly constrained age range of 5485 ± 50 to 5340 ± 36 BP, confirming the archaeological dating to the Late Neolithic. There is no evidence of a population dating to 6400 BP as proposed by Ogrinc and Budja (2005).

Table 1 AMS ^{14}C dates and stable carbon and nitrogen values for Late Neolithic burials from Ajdovska Jama, Slovenia. Ogrinc's (1999) stable isotope data are included for comparison. In the majority of cases, the same bone was analyzed in both studies; the exceptions are samples Aj03, Aj06, and Aj14 (denoted with an asterisk), although these were from the same skeletons. Dates have been calibrated using atmospheric data from Reimer et al. (2004) and OxCal v. 4.0 beta (Bronk Ramsey 1995, 2001, 2006), and are shown at 95.4% probability. N.B. Ogrinc and Budja (2005) give different $\delta^{15}\text{N}$ values for samples Aj08 and Aj13 (in parentheses) compared to Ogrinc (1999). The data from Ogrinc (1999) are shown in Figure 4.

Sample ID	Catalog #	Age at death	Sex	Sample description	Bone cluster	Strat unit	Lab ID	^{14}C age BP	cal BC age range (2 σ)	Present study					Ogrinc (1999)		
										C/N	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N	$\delta^{13}\text{C}$
Aj01	7	Adult	M	R ulna	B	44	OxA-15041	5485 \pm 50	4448–4243	3.2	-20.3	8.1	3.3	-20.6	7.9		
Aj02	19	Adult	M	R talus	A	44	Insufficient collagen						3.2	-22.2	4.9		
Aj03*	20	Adult	M	L clavicle	A	43	Insufficient collagen						3.5	-20.5	9.0		
Aj04	21	Adult	F	Ulna	A	43	OxA-15072	5365 \pm 31	4328–4055	3.2	-20.4	9.0	3.3	-20.5	8.7		
Aj05	22	Adult	M	Proximal phalanx of hand	A	43	Insufficient collagen						3.4	-21.2	5.6		
Aj06*	12	Child (~1 yr)		L ulna	A	44	Insufficient collagen						3.5	-20.7	10.8		
Aj07	14	Child (1–2 yr)		Metatarsal	A	44	Insufficient collagen						3.5	-20.4	11.5		
Aj08	10	Child (~5 yr)		Radius	A	44	OxA-15073	5369 \pm 31	4328–4061	3.2	-20.9	10.2	3.5	-20.3	10.7 (8.7)		
Aj09	1	Adult	F	L clavicle	B	44	OxA-15119	5340 \pm 36	4317–4049	3.2	-20.1	8.1	3.3	-20.2	7.3		
Aj10	2	Adult	M	L clavicle	B	44	OxA-15074	5416 \pm 35	4345–4173	3.2	-20.6	8.3	3.2	-20.9	7.7		
Aj11	3	Adult	F	L fibula	B	44	OxA-15091	5421 \pm 30	4340–4235	3.2	-20.6	8.1	3.2	-20.9	8.5		
Aj12	6	Child (1–2 yr)		Tibia	B	44	OxA-15092	5436 \pm 30	4343–4243	3.2	-20.8	9.0	2.9	-20.7	10.2		
Aj13	15	Child (~2 yr)		Long bone diaphysis frag.	C	43	OxA-15093	5389 \pm 30	4335–4075	3.2	-20.5	8.7	3.3	-20.7	10.0 (10.7)		
Aj14*	16	Child (~10 yr)		L rib	C	43	OxA-15094	5405 \pm 31	4339–4172	3.2	-20.6	8.6	2.9	-20.8	7.5		
Aj15	17	Child (~6 yr)		Long bone diaphysis frag.	C	43	OxA-15095	5471 \pm 31	4363–4257	3.2	-20.5	8.8	3.0	-20.6	9.1		




Figure 5 Probability distributions of Bayesian modeled age ranges of human bone samples dated from Ajdovska Cave. Plot generated using OxCal v 4.0 (Bronk Ramsey 1995, 2001, 2006). The overall agreement index for this model was 82.9%. ^{14}C likelihoods are shown with lighter shading, while posterior probability distributions are shown in bold shades. See Table 2 for actual posterior range data.

The AMS dates have been calibrated against the IntCal04 data set (Reimer et al. 2004) using OxCal v 4.0 (Bronk Ramsey 1995, 2001, 2006). There is a dearth of prior archaeological information associated with the selected skeletons from the site, particularly in terms of any relative sequencing, since the total thickness of the bone-bearing horizon is only about 10 cm. The dated samples were modeled as 2 phases (44 and 43), in accordance with Horvat's original stratigraphic interpretation. The results of this Bayesian analysis are shown in Figure 5 and Table 2. The overall agreement index for the model was 82.9%. Subsequent runs of the model showed consistent agreement indices. One of the posterior distributions yielded low agreement indices, suggesting strongly that it did not fit the inferred phasing (OxA-15119), and this was therefore questioned (Table 2). The modeling indicates that burial deposition began after 4340–4290 BC (68.2% probability) and that the burials represent a brief period of activity; total span is equivalent to 5–120 yr (95.4% probability) with the highest probability associated with 10–20 yr (Figure 6). The terrestrial stable isotope signals (see below) suggest a lack of a reservoir effect, and this strengthens the confidence in the reliability of these human bone determinations.

The underlying principle of dietary tracing using stable isotopes is that the ratios of carbon and nitrogen isotopes in bone collagen reflect those in the average diet—more specifically, the protein portion of the diet. With each step along the food chain, there is fractionation of 1 isotope relative to another, resulting in a change in ratio (“trophic enrichment”). Most workers assume a slight enrichment in $\delta^{13}\text{C}$ (up to 1‰) and an average enrichment in $\delta^{15}\text{N}$ of 3‰ between food source and consumer.

Table 2 Posterior distributions for the Ajdovska Jama Bayesian model. Ranges for start and end boundaries as well as overall span are shown in cal yr BC, as well as for each posterior distribution. The results have been rounded to the nearest 5 yr. These data are shown in Figure 5.

	Posterior distributions (cal age range BC)				Agreement index
	68.2%		95.4%		
	from	to	from	to	
Phase 44					
Boundary Start 44	4340	4290	4355	4265	
OxA-15041	4330	4270	4340	4260	88.5
OxA-15092	4330	4280	4335	4265	103.9
OxA-15091	4330	4280	4335	4265	111.1
OxA-15074	4329	4285	4335	4265	116
OxA-15073	4330	4290	4330	4260	88.2
OxA-15119	4245	4070	4320	4050	questioned
Boundary End 44/Start 43	4320	4265	4330	4260	
Phase 43					
OxA-15095	4300	4260	4325	4250	53.3
OxA-15094	4300	4250	4320	4245	112.1
OxA-15093	4310	4250	4320	4240	113.4
OxA-15072	4310	4250	4325	4240	98.5
Boundary End 43	4295	4235	4325	4220	
Span	5	60	5	120	




Figure 6 Bayesian span for the modeled data presented in Figure 5. The distribution represents the span of both archaeological phases and suggests that the highest probability is associated with a very brief period.

In the studies of Ogrinc (1999) and Ogrinc and Budja (2005), the $\delta^{15}\text{N}$ values for the bone collagen of the adults ranged from +4.9 to +9.3‰, which is unusually large for a single population (Figure 4c). These data were interpreted as indicating a dietary regime with wide variation in individual food preferences.

The $\delta^{15}\text{N}$ values for adults obtained during the present study show a much narrower range of <1‰, from +8.1 to +9.0‰ (Figure 7). None of the adult samples for which Ogrinc obtained low (<6‰) $\delta^{15}\text{N}$ values produced a quality of collagen suitable for analysis, based on the parameters discussed above. It could be argued that this is why the range in our study is smaller and there are no low $\delta^{15}\text{N}$ values. However, it may be that the low $\delta^{15}\text{N}$ samples analyzed by Ogrinc contained non-collagenous contaminants, which have influenced her results. There has been little direct comparison of stable isotope bone collagen values obtained with and without the ultrafiltration step, and so at present we have no empirical data to support this argument. However, an effect has been demonstrated on ^{14}C ages (Bronk Ramsey et al. 2004; Higham et al. 2006), which implies that in some circumstances (e.g. low collagen content) stable isotope values could also be influenced.




Figure 7 Collagen carbon and nitrogen stable isotope ratios for human adults and children from Ajdovska Jama obtained in the present study, plotted against Ogrinc's (1999) food source data.

Of the 4 children for which results were obtained in the present study, only 1 shows a significantly higher $\delta^{15}\text{N}$ value compared to the adults, which may be indicative of a "nursing effect" (Figure 7). Interestingly, this is not one of the youngest children in the group, having an age at death of ~5 yr (Table 1, sample Aj08).

From a comparison of her human and food source stable isotope data, Ogrinc (1999) concluded that the human diet consisted predominantly of meat from domestic and wild herbivores, with very little of the protein being derived from plant sources. The re-interpretation (Ogrinc and Budja 2005) proposed a diet based mainly on meat from domestic and wild herbivores (average of 44%) and cereals (average of 39%).

The present study favors a somewhat different interpretation. In Figure 7, our human collagen stable isotope results are compared with Ogrinc's data for animal and plant food sources. The difference in average $\delta^{15}\text{N}$ between the human adults and herbivores is 2.3‰. This enrichment would be consis-

tent with a largely meat-based diet if one assumes a 3‰ trophic shift. However, judging from ethnohistorical evidence, such a diet would be highly unusual for a peasant farming society in south-eastern Europe where cultivated plants (cereals, legumes, and fruits) typically provide the bulk of the diet, with a modest contribution from dairy products, and meat is regarded as a luxury.

Since animal products generally contain more protein per unit weight than do most plant foods, particularly in a cooked state, which is how they would normally be consumed by humans (Table 3), the meat portion of the diet may be disproportionately represented in bone collagen (Bonsall et al. 1997). Moreover, while most isotopic studies of human diet employ a trophic enrichment of 3‰ for $\delta^{15}\text{N}$, recent research suggests that the enrichment in humans can often be significantly greater than this. For example, our work in the Iron Gates indicates an enrichment of 5.4–6.7‰ in Mesolithic humans over the riverine fish on which their diet was based (Bonsall, unpublished data), while average collagen $\delta^{15}\text{N}$ values for Middle and Early Upper Paleolithic populations in western Europe are typically 5‰ higher than those of their main prey animals such as reindeer, bovinds, and horse (estimate based on data presented by Drucker and Bocherens 2004).

Table 3 Nutritional value of selected (modern) animal and plant foods (data from Paul and Southgate 1978). In cooked form, the protein content of meat on average is approximately 4 times greater than that of cereals and pulses.

Food	CH (g)	Fat (g)	Protein (g)	Kcal/100 g
Beef (sirloin, raw)	0	22.8	16.6	272
Beef (sirloin, roast)	0	21.1	23.6	284
Beef (rump, grilled)	0	12.1	27.3	218
Lamb (leg, raw)	0	18.7	17.9	240
Lamb (leg, roast)	0	17.9	26.1	266
Pork (leg, raw)	0	22.5	16.6	269
Pork (leg, roast)	0	19.8	26.9	286
Venison (roast)	0	6.4	35.0	198
Cheese (Camembert type)	Trace	23.2	22.8	300
Milk (cow's)	4.7	3.8	3.3	65
Wheat (wholemeal flour)	65.8	2.0	13.2	318
Barley (raw)	83.6	1.7	7.9	360
Barley (boiled)	27.6	0.6	2.7	120
Beans (haricot, raw)	45.5	1.6	21.4	271
Beans (haricot, boiled)	16.6	0.5	6.6	93
Beans (broad, boiled)	7.1	0.6	4.1	48
Peas (raw)	10.6	0.4	5.8	67
Peas (boiled)	7.7	0.4	5.0	52
Lentils (raw)	53.2	1.0	23.8	304
Lentils (boiled)	17.0	0.5	7.6	99

Therefore, we would suggest that the average $\delta^{15}\text{N}$ value of +8.3‰ for the adult humans from Ajdovska Jama does not necessarily indicate heavy reliance on the meat of herbivores as suggested by Ogrinc (1999) and Ogrinc and Budja (2005), but is entirely consistent with a mixed diet of plant and animal products. Furthermore, the tight clustering of the adult $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in the present study does not support Ogrinc's (1999) suggestion of significant variation in individual food preferences among the Late Neolithic population.

It is instructive to compare our data with those for modern people who are vegans and vegetarians. In a study by Macko et al. (1999), hair samples from 2 vegans gave $\delta^{15}\text{N}$ values of +7.2‰, while hair samples from 8 ovo-lacto-vegetarians (who consumed a combination of plants, eggs, and milk products) yielded $\delta^{15}\text{N}$ values in the range +8.5 to +10.9‰.

The suggestion that plant foods contributed significantly to the diet of the Ajdovska Jama population also receives support from dental evidence. Table 3 shows that cereals and pulses are rich sources of carbohydrates when compared to meat from terrestrial herbivores. Heavier reliance on carbohydrates may be reflected in a greater prevalence of dental caries (“tooth decay”: Larsen 1997). Although adult dental remains at Ajdovska Cave were under-represented and the caries rate cannot be calculated reliably, the presence of 2 out of 4 mandibles exhibiting carious lesions and/or *ante mortem* tooth loss and 3 decayed loose teeth out of a total of 19 implies that it would have been high. This contrasts with the situation among Mesolithic populations in the Iron Gates, where the diet was high in animal (mainly fish) protein and low in carbohydrate and caries rates were correspondingly very low (Bonsall et al. 1997).

The average $\delta^{13}\text{C}$ value for the adults analyzed in the present study is -20.5‰, which suggests a diet based on C_3 food sources. There is no evidence for significant consumption of C_4 crop plants or meat from animals fed regularly on C_4 plants, and this is highlighted by a comparison of the $\delta^{13}\text{C}$ values for the Late Neolithic skeletons from Ajdovska Jama with the much heavier values found in early Iron Age burials from Magdalenska Gora also in the Sava catchment, 60 km to the west of Ajdovska Jama, which according to Murray and Schoeninger (1988) reflect high levels of millet consumption (Figure 8).




Figure 8 Collagen carbon and nitrogen stable isotope ratios for human adults from Ajdovska Jama (Late Neolithic) and Magdalenska Gora (early Iron Age). Both sites are in the Sava river catchment, ~60 km apart (Magdalenska Gora data from Murray and Schoeninger 1988).

Similarly, $\delta^{13}\text{C}$ values of -20.1‰ to -20.9‰ coupled with the relatively low $\delta^{15}\text{N}$ values argue against a significant contribution of fish to the diet, and no fish bones were present among the faunal remains from Late Neolithic contexts at Ajdovska Jama.

CONCLUSIONS

The AMS ^{14}C ages on human bone collagen presented in this paper confirm the archaeological dating of the burials in Ajdovska Jama, Slovenia, to the Late Neolithic, ~5400 BP (4300 cal BC), with no evidence of burial activity around 6400 BP as suggested by Ogrinc and Budja (2005). Bayesian analysis of the ^{14}C data suggests the people were buried during a period of no more than 120 yr (95.4% probability). The associated carbon and nitrogen isotope ratios indicate a population with a “terrestrial” diet in which the protein was derived very largely or exclusively from C_3 plant and animal food sources. Taken together with the archaeobotanical, physical anthropological, and zooarchaeological evidence from the site, the data are suggestive of a mixed farming economy. Meat from wild herbivores was probably included in the diet, but there is no archaeological or isotopic evidence for the regular consumption of fish, in spite of the relative proximity of the River Sava. Nor is there evidence for the use of C_4 plants as human food or animal fodder. The Ajdovska Jama burials thus belong to a period when farming had already been established in the region for over a millennium, but before millet (the only C_4 crop plant grown widely in prehistoric Europe) became economically important. From the perspective of stable isotope research, Ajdovska Jama constitutes an important *reference* sample of an established Neolithic farming community from the Danube Basin.

ACKNOWLEDGMENTS

Clive Bonsall’s work on the human bone material from Ajdovska Jama was undertaken as part of his *Northern Adriatic Project*, supported by research grants from the British Academy, the Carnegie Trust for the Universities of Scotland, and the Munro Fund, Hayter Fund, and Development Trust Research Fund of the University of Edinburgh. The AMS ^{14}C measurements and stable isotope analyses were funded by an AHRC–NERC ORADS award to CB. The authors wish to thank László Bartosiewicz and an anonymous referee for their comments on a preliminary draft of the paper. We are also grateful to Nives Kokeza and Matija Črešnar for Slovene–English translations, and to Professor Mihael Budja for his invaluable assistance at an early stage of the research.

REFERENCES

- Bartosiewicz L, Bonsall C, Boroneanț V, Stallibrass S. 2001. New data on the prehistoric fauna of the Iron Gates: a case study from Schela Cladovei, Romania. In Kertés R, Makkay J, editors. *From the Mesolithic to the Neolithic*. Budapest: Archaeolingua (Main Series). p 15–21.
- Bonsall C, Lennon R, McSweeney K, Stewart C, Harkness D, Boroneanț V, Payton R, Bartosiewicz L, Chapman JC. 1997. Mesolithic and Early Neolithic in the Iron Gates: a palaeodietary perspective. *Journal of European Archaeology* 5(1):50–92.
- Bonsall C, Cook G, Lennon R, Harkness D, Scott M, Bartosiewicz L, McSweeney K. 2000. Stable isotopes, radiocarbon and the Mesolithic–Neolithic transition in the Iron Gates. *Documenta Praehistorica* 27:119–32.
- Bonsall C, Cook GT, Hedges REM, Higham TFG, Pickard C, Radovanović I. 2004. Radiocarbon and stable isotope evidence of dietary change from the Mesolithic to the Middle Ages in the Iron Gates: new results from Lepenski Vir. *Radiocarbon* 46(1):293–300.
- Bronk Ramsey C. 1995. Radiocarbon calibration and analysis of stratigraphy: the OxCal program. *Radiocarbon* 37(2):425–30.
- Bronk Ramsey C. 2001. Development of the radiocarbon calibration program. *Radiocarbon* 43(2A):355–63.
- Bronk Ramsey C. 2006. OxCal program v 4.0 [software and documentation]. URL: http://c14.arch.ox.ac.uk/oxcal/hlp_contents.html.
- Bronk Ramsey C, Hedges REM. 1997. Hybrid ion sources: radiocarbon measurements from microgram to milligram. *Nuclear Instruments and Methods in Physics Research B* 123(1–4):539–45.
- Bronk Ramsey C, Pettitt PB, Hedges REM, Hodgins GWL, Owen DC. 2000. Radiocarbon dates from the Oxford AMS system: Archaeometry Datelist 30. *Archaeometry* 42:459–79.
- Bronk Ramsey C, Higham TFG, Bowles A, Hedges REM. 2004. Improvements to the pretreatment of bone at Oxford. *Radiocarbon* 46(1):155–63.
- Cook GT, Bonsall C, Hedges REM, McSweeney K, Boroneanț V, Pettitt PB. 2001. A freshwater diet-derived ^{14}C reservoir effect at the Stone Age sites in the Iron Gates Gorge. *Radiocarbon* 43(2A):453–60.
- Cook GT, Bonsall C, Hedges REM, McSweeney K, Boroneanț V, Bartosiewicz L, Pettitt PB. 2002. Problems of dating human bones from the Iron Gates. *Antiquity*

- 76(291):77–85.
- Corrain C, Capitanio M. 1991. La necropoli di Adjovska jama (Slovenia). *Poročilo o raziskovanju paleolita, neolita in eneolita v Sloveniji* 29:207–47. In Italian.
- Drucker D, Bocherens H. 2004. Carbon and nitrogen stable isotopes as tracers of change in diet breadth during Middle and Upper Palaeolithic Europe. *International Journal of Osteoarchaeology* 14(3–4):162–77.
- Higham T, Bronk Ramsey C, Karavanić I, Smith FH, Trinkaus E. 2006. Revised direct radiocarbon dating of the Vindija G₁ Upper Paleolithic Neandertals. *Proceedings of the National Academy of Sciences of the USA* 103(3):553–7.
- Horvat M. 1989. *Ajdovska jama pri Nemški vasi*. Ljubljana: Znanstveni inštitut Filozofske fakultete. In Slovene.
- Larsen CS. 1997. *Bioarchaeology: Interpreting Behaviour from the Human Skeleton*. Cambridge: Cambridge University Press. 474 p.
- Macko SA, Lubec G, Teschler-Nicola M, Andrusevich V, Engel MH. 1999. The Ice Man's diet as reflected by the stable nitrogen and carbon isotopic composition of his hair. *The FASEB Journal* 13:559–62.
- Murray M, Schoeninger M. 1988. Diet, status and complex social structure in Iron Age Central Europe: some contributions of bone chemistry. In: Gibson DB, Geselowitz MN, editors. *Tribe and Polity in Late Prehistoric Europe: Demography, Production and Exchange in the Evolution of Complex Social Systems*. New York: Plenum. p 155–76.
- Obelić B, Horvatinčić N, Srdoč D, Krajcar Bronić I, Sliepčević A, Grgić S. 1994. Rudjer Bošković Institute radiocarbon measurements XIII. *Radiocarbon* 36(2):303–24.
- Ogrinc N. 1999. Stable isotope evidence of the diet of the Neolithic population in Slovenia—a case study: Ajdovska jama. *Documenta Praehistorica* 26:193–200.
- Ogrinc N, Budja M. 2005. Paleodietary reconstruction of a Neolithic population in Slovenia: a stable isotope approach. *Chemical Geology* 218(1–2):103–16.
- Paul AA, Southgate DAT. 1978. *McCance and Widdowson's the Composition of Foods*. 4th edition. London: Her Majesty's Stationery Office.
- Phillips DL, Koch PL. 2002. Incorporating concentration dependence in stable isotope mixing models. *Oecologia* 130(1):114–25.
- Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Bertrand CJH, Blackwell PG, Buck CE, Burr GS, Cutler KB, Damon PE, Edwards RL, Fairbanks RG, Friedrich M, Guilderson TP, Hogg AG, Hughen KA, Kromer B, McCormac G, Manning S, Bronk Ramsey C, Reimer RW, Remmele S, Southon JR, Stuiver M, Talamo S, Taylor FW, van der Plicht J, Weyhenmeyer CE. 2004. IntCal04 terrestrial radiocarbon age calibration, 0–26 cal kyr BP. *Radiocarbon* 46(3):1029–58.
- Srdoč D, Obelić B, Horvatinčić N, Krajcar I, Sliepčević A. 1984. Rudjer Bošković Institute radiocarbon measurements VIII. *Radiocarbon* 26(3):449–60.
- Srdoč D, Horvatinčić N, Obelić B, Krajcar Bronić I, Sliepčević A. 1987. Rudjer Bošković Institute radiocarbon measurements IX. *Radiocarbon* 29(1):115–34.
- Srdoč D, Obelić B, Horvatinčić N, Krajcar Bronić I, Sliepčević A. 1989. Rudjer Bošković Institute radiocarbon measurements XI. *Radiocarbon* 31(1):85–98.
- Srdoč D, Horvatinčić N, Krajcar Bronić I, Obelić B, Sliepčević A. 1992. Rudjer Bošković Institute radiocarbon measurements XII. *Radiocarbon* 34(1):155–75.
- Whittle A, Bartosiewicz L, Borić D, Pettitt P, Richards M. 2002. In the beginning: new radiocarbon dates for the Early Neolithic in northern Serbia and southeast Hungary. *Antaeus* 25:64–117.
- Zohary D, Hopf M. 1994. *Domestication of Plants in the Old World: the Origin and Spread of Cultivated Plants in West Asia, Europe, and the Nile Valley*. 2nd edition. Oxford: Clarendon Press. 264 p.