



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Alcohol intake and cognitive abilities in old age: The Lothian Birth Cohort 1936 study.

### Citation for published version:

Corley, J, Jia, X, Brett, CE, Gow, AJ, Starr, J, Kyle, JAM, Mcneill, G & Deary, IJ 2011, 'Alcohol intake and cognitive abilities in old age: The Lothian Birth Cohort 1936 study.', *Cognitive Neuropsychology*, vol. 25, no. 2, pp. 166-175. <https://doi.org/10.1037/a0021571>

### Digital Object Identifier (DOI):

[10.1037/a0021571](https://doi.org/10.1037/a0021571)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Peer reviewed version

### Published In:

Cognitive Neuropsychology

### Publisher Rights Statement:

© Corley, J., Jia, X., Brett, C. E., Gow, A. J., Starr, J., Kyle, J. A. M., Mcneill, G., & Deary, I. J. (2011). Alcohol intake and cognitive abilities in old age: The Lothian Birth Cohort 1936 study. *Cognitive Neuropsychology*, 25(2), 166-175. [10.1037/a0021571](https://doi.org/10.1037/a0021571)

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



2 **Alcohol intake and cognitive abilities in old age: the Lothian Birth Cohort**

3 **1936 study**

4

5 Janie Corley

6 University of Edinburgh, UK

7

8 Xueli Jia,

9 University of Aberdeen, UK

10

11 Caroline E. Brett, Alan J. Gow, and John M. Starr

12 University of Edinburgh, UK

13

14 Janet A.M. Kyle, and Geraldine McNeill

15 University of Aberdeen, UK

16

17 Ian J. Deary

18 University of Edinburgh, UK

19

20 **Author note**

21 Janie Corley, Department of Psychology, University of Edinburgh, UK; Xueli Jia, Health

22 Services Research Unit, University of Aberdeen, UK; Caroline E. Brett, Department of

23 Psychology, University of Edinburgh, UK; Alan J. Gow, Department of Psychology, Centre

24 for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, UK; John M.

25 Starr, Geriatric Medicine Unit, Centre for Cognitive Ageing and Cognitive Epidemiology,

1 University of Edinburgh, UK; Janet A. M. Kyle, Population Health, Division of Applied  
2 Health Sciences, University of Aberdeen, UK; Geraldine McNeill, Population Health,  
3 Division of Applied Health Sciences, University of Aberdeen, UK; Ian J. Deary, Department  
4 of Psychology, Centre for Cognitive Ageing and Cognitive Epidemiology, University of  
5 Edinburgh, UK.

6  
7 Janie Corley and Alan J. Gow are funded by the Sidney De Haan Award, part of the  
8 Disconnected Mind project funding from Help The Aged to Ian Deary and John Starr. A  
9 Research Into Ageing programme grant to Ian Deary and John Starr supported Alan Gow,  
10 Janie Corley and Caroline Brett. We thank the Scottish Council for Research in Education for  
11 allowing access to the SMS1947. We thank the LBC1936 Study participants. The work was  
12 undertaken by The University of Edinburgh Centre for Cognitive Ageing and Cognitive  
13 Epidemiology, part of the cross council Lifelong Health and Wellbeing Initiative  
14 (G0700704/84698). Funding from the Biotechnology and Biological Sciences Research  
15 Council (BBSRC), Engineering and Physical Sciences Research Council (EPSRC), Economic  
16 and Social Research Council (ESRC) and Medical Research Council (MRC) is gratefully  
17 acknowledged.

18  
19 Correspondence should be addressed to Ian J. Deary, Centre for Cognitive Ageing and  
20 Cognitive Epidemiology, Department of Psychology, University of Edinburgh, 7 George  
21 Square, Edinburgh, EH8 9JZ, Scotland, UK. Tel: +44 131 650 3452. Email  
22 I.Deary@ed.ac.uk

23

24 **Tables:** 4

25 **Supplementary tables:** 2

**Abstract**

1  
2 **Objective:** Moderate alcohol consumption has been associated with better cognitive  
3 performance in late adulthood, possibly via improving vascular health. Few studies have  
4 examined the potentially confounding roles of prior cognitive ability and social class in this  
5 relationship. **Method:** Participants were 922 healthy adults aged about 70 years in the  
6 Lothian Birth Cohort 1936 Study, on whom there are IQ data from age 11 years. Alcohol  
7 consumption was obtained by self-report questionnaire. Cognitive outcome measures  
8 included general cognitive ability, speed of information processing, memory and verbal  
9 ability. **Results:** Moderate-substantial drinking ( $> 2$  units/day) was associated with better  
10 performance on cognitive tests than low-level drinking ( $\leq 2$  units/day) or non-drinking in  
11 men and women. After adjusting for childhood IQ and adult social class, most of these  
12 associations were removed or substantially attenuated. After full adjustment, there remained  
13 a small, positive association between overall alcohol intake and memory (in women and  
14 men) and verbal ability (in women only). Womens' overall alcohol intake was derived almost  
15 exclusively from wine. In men, effects differed according to beverage type: wine and  
16 sherry/port consumption was associated with better verbal ability, but beer with a poorer  
17 verbal ability, and; spirits intake was associated with better memory. **Conclusions:** Prior  
18 intelligence and socioeconomic status influence both amount and/or type of alcohol intake,  
19 and may partly explain the link between alcohol intake and improved cognitive performance  
20 at age 70. Alcohol consumption was found to make a small, independent contribution to  
21 memory performance and verbal ability, but the clinical significance of these findings is  
22 uncertain.

23  
24 *Key words:* alcohol, cognitive function, childhood IQ, aging.  
25

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

## Introduction

Moderate alcohol intake in middle-aged and older adults is associated with reduced all-cause mortality and a lower risk of cardiovascular disease (Naimi et al., 2005). Population-based studies have found that moderate drinking, compared to abstinence, can also benefit cognitive function (Espeland et al., 2005; Lang, Wallace, Huppert, & Melzer, 2007; Stampfer, Kang, Chen, Cherry, & Grodstein, 2005). Evidence from longitudinal assessments of older people suggests that light to moderate drinking has a protective effect against cognitive decline over time (Ganguli, Vander Bilt, Saxton, Shen, & Dodge, 2005; Wright, Elkind, Luo, Paik, & Sacco, 2006). Stampfer et al. (2005) reported that those who drank up to 15 mg of alcohol per day (not more than about one drink), had a 20% decreased risk of cognitive impairment at baseline, compared with non-drinkers, and these ‘moderate’ drinkers experienced less cognitive decline over a two-year period. Some studies have reported a positive association between moderate alcohol intake and improved cognitive performance in women, but not men (Dufouil, Ducimetiere, & Alperovitch, 1997; Leroi, Sheppard, & Lyketsos, 2002; McGuire, Ajani, & Ford, 2007; Stott et al., 2008), and others, a significant effect in both sexes but a larger effect size in women (Britton, Singh-Manoux, & Marmot, 2004). A few studies have failed to find any positive cognitive benefit of non-excessive alcohol consumption (Cervilla, Prince, & Mann, 2000; Elwood et al., 1999). Inconsistencies may be attributable to methodological issues; a meta-analysis found that definitions of ‘moderate’ drinking vary widely across studies (Peters, Peters, Warner, Beckett, & Bulpitt, 2008).

The ‘beneficial’ cognitive effects of moderate alcohol intake are reportedly heterogenous, concentrated in areas of learning, executive functions and psychomotor speed (Ganguli et al., 2005); verbal knowledge and phonemic fluency (Britton et al., 2004); and in a women-only study, verbal knowledge, phonemic fluency and fine motor speed (Espeland et al., 2006). These, and many other studies, have failed to find strong evidence of an association

1 between alcohol intake and memory. At present, it is unclear how different types of alcoholic  
2 beverage are associated with cognitive function in late life. Few epidemiological studies have  
3 looked further than overall alcohol intake. There is some suggestion that wine, but not spirits or  
4 beer, is protective against cognitive decline (Luchsinger, Tang, Siddiqui, Shea, & Mayeux,  
5 2004) and dementia (Truelsen, Thudium, & Grønbaek, 2002). The proposed explanation is  
6 based on the antioxidant activity of the flavonoids found in wine. However, not all studies  
7 report consistent results. According to The Nurses' Health Study, using data derived from  
8 12,480 elderly women, no significant differences in risk of cognitive impairment or decline  
9 were observed according to type of alcoholic beverage (Stampfer et al., 2005).

10 Typically, the direction of causation that is studied and assumed is from alcohol to  
11 cognitive ability, especially in old age. The most widely accepted mechanism behind the  
12 putatively protective role of alcohol on cognitive aging is an indirect benefit via a reduced risk  
13 of vascular disease. A quantitative review of moderate alcohol intake and biological markers of  
14 CHD risk, reported a causal association via alcohol induced changes in lipids and haemostatic  
15 factors (Rimm, Williams, Fosher, Criqui, & Stampfer, 1999). Further support for an effect on  
16 brain vasculature comes from Magnetic Resonance Imaging studies which reveal a lower  
17 prevalence of white matter abnormalities and infarcts in older persons with moderate alcohol  
18 intake than in non-drinkers (Mukamal, Longstreth, Mittleman, Crum, & Siscovick, 2001).  
19 However, it is also possible that the observed relation between alcohol and cognitive health is  
20 attributable to prior intelligence. Previous studies report that intelligence influences the amount  
21 and/or type of alcohol intake. People with a higher IQ tend to drink regularly but moderately  
22 (British Medical Association, 2008). Higher IQ scores have also been associated with a  
23 preference for wine over other types of alcohol in later life (Mortensen, Sørensen, & Grønbaek,  
24 2005). If moderate drinkers are, on average, more intelligent than non-drinkers, the reported  
25 beneficial effect of moderate drinking on cognitive function in adulthood may be confounded

1 by prior ability. Similarly, there appear to be social gradients in drinking. In the Whitehall  
2 study, some of the association between alcohol consumption and cognitive function could be  
3 explained by social position, as measured by employment grade (Britton et al., 2004).  
4 Variations in both IQ and SES may play important confounding roles in the alcohol-cognition  
5 relationship, yet many studies attribute causal effects of alcohol intake to improved cognition  
6 in the absence of this kind of data.

7 The sample in the present study is unusual in having validated measures of cognitive  
8 ability from childhood and old age as well as a range of sociodemographic and health data. The  
9 objectives of the present study were to examine whether: (1) a pattern of light to moderate  
10 drinking is associated with better cognitive outcomes in old age than non-drinking; (2) the  
11 effects on cognitive outcomes vary by type of alcoholic beverage; and (3) specifically, whether  
12 these relationships could be attributed to confounding by prior cognitive ability (age 11 IQ)  
13 and/or SES.

## 15 **Methods**

### 16 **Participants and general methods**

17 Participants were enrolled on the Lothian Birth Cohort 1936 (LBC1936) Study which  
18 comprises 1091 men and women. These individuals are surviving participants of the Scottish  
19 Mental Survey of 1947 (SMS1947: see Deary, Whalley, & Starr, 2009). Full details of the  
20 recruitment and testing of the LBC1936 are given in a free-access protocol paper (Deary et al.,  
21 2007). At the time of recruitment, LBC1936 members mostly resided in Edinburgh and its  
22 surrounding area (Lothian) in Scotland. They were relatively healthy and lived independently.  
23 Between 2004 and 2007, at a mean age of ~70 years, LBC1936 participants attended the  
24 Wellcome Trust Clinical Research Facility in Edinburgh to undergo cognitive testing, a clinical  
25 assessment and an interview. As a part of their general assessment, participants were asked to

1 complete a Food Frequency Questionnaire. Of the 1091 participants interviewed, 922 (84.5%)  
2 provided both alcohol consumption and cognitive data, and formed the present study sample.  
3 Ethics permission for the Lothian Birth Cohort 1936 (LBC1936) study protocol was obtained  
4 from the Multi-Centre Research Ethics Committee for Scotland (MREC/01/0/56) and from  
5 Lothian Research Ethics Committee (LREC/2003/2/29). The research was carried out in  
6 compliance with the Helsinki Declaration. All participants gave their written, informed  
7 consent.

8

## 9 **Procedure**

10 **Measurement of alcohol intake.** Alcohol intake was assessed using the Scottish  
11 Collaborative Group 165-item Food Frequency Questionnaire (FFQ) version 7.0  
12 (<http://www.foodfrequency.org>). The FFQ has good repeatability (dietary intake in later life is  
13 reasonably stable in the short term) and good validity for most nutrients in community  
14 dwelling older populations (Jia, Craig, Aucott, Milne, & McNeill, 2008; McNeill, Winter, &  
15 Jia, 2009). It is also a valid assessment tool for measuring alcohol intake; there is good  
16 agreement between the FFQ and 4-day weighed diet records in men ( $r = 0.83$ ) and women ( $r =$   
17  $0.70$ ) (Masson et al., 2003). This self-report questionnaire measures alcohol use over the  
18 previous 2-3 month period from nine alcoholic beverages, namely: ‘low-alcohol lager or beer’;  
19 ‘dark beer (export, bitter or stout)’; ‘light beer (lager or continental beer)’; ‘white wine’; ‘red  
20 wine’; ‘sherry, port etc’; ‘spirits or liqueurs’; ‘alcopops (e.g. Bacardi Breezer)’ and ‘cider’.  
21 Each item on the questionnaire refers to a standard measure, e.g., 1 half-pint (beer/lager/cider),  
22 1 pub measure (spirits). Participants mark one of nine responses to indicate frequency of  
23 consumption: rarely or never; 1-3 per month; 1 per week; 2-3 per week; 4-6 per week; 1 per  
24 day; 2-3 per day; 4-6 per day; 7+ per day.



1           In the event of any missing responses, a letter was sent requesting the information.  
2   Using standard FFQ protocol, incomplete questionnaires (with 10 or more missing items from  
3   a total of 165) were excluded from the analyses. Thirty-nine questionnaires were ‘incomplete’,  
4   26 were returned blank and 98 were not returned. A total of 928 (85%) completed FFQs were  
5   returned. One was later excluded where the amount of alcohol consumed was a clear statistical  
6   outlier and information collected at interview revealed a history of problem drinking. Five  
7   further individuals were excluded because they were identified as having potential dementia  
8   based on a score of < 24 on the Mini-Mental State Examination (Folstein, Folstein, &  
9   McHugh, 1975). The final sample for analysis in the present study comprised 922 relatively  
10   healthy participants (445 men, 477 women) aged about 70 ( $M = 69.5$ ,  $SD = 0.8$ ) years at time  
11   of testing. Five of these participants each had one item of missing alcohol data. Therefore, total  
12   alcohol intake could be calculated for 917 participants. We calculated the daily alcohol intake  
13   in units derived from each beverage based on UK government guidelines<sup>1</sup> and a combined total  
14   daily alcohol measure from all 9 sources.

15           **Measurement of cognitive performance at age 70.** Cognitive function was assessed  
16   using a battery of neuropsychological tests. See the free-access LBC1936 protocol article for a  
17   full description of tests (Deary et al., 2007). In the present study, we used three composite  
18   cognitive scores derived from principal components analyses (PCA), to represent three distinct  
19   cognitive domains. Regression scores were calculated for the first unrotated principal  
20   component of the tests in each domain. In each case the scree slopes and eigenvalues suggested  
21   that a single component could be extracted.

22           *g factor (general cognitive ability)*

---

<sup>1</sup> Alcohol unit calculations were based on UK Government guidelines ([http://www.direct.gov.uk/en/HealthandWellbeing/DG\\_10036434](http://www.direct.gov.uk/en/HealthandWellbeing/DG_10036434)) as follows: half pint of lager or beer = 1 unit; half pint of low alcohol lager or beer = 0.5 units; standard glass of wine = 2 units; half pint/1 bottle of cider = 1 unit; standard glass of sherry or port = 1 unit; 1 pub measure of spirits or liqueurs = 1 unit; 1 bottle of alcopops = 1.5 units.

1           A *g* factor score, representing general cognitive ability, was derived from a PCA of  
2 scores on six Wechsler Adult Intelligence Scale-III<sup>UK</sup> (WAIS-III; Wechsler, 1998a) subtests,  
3 namely: *Letter-Number Sequencing* (working memory); *Matrix Reasoning* (non-verbal  
4 reasoning); *Block Design* (constructional ability); *Digit Symbol* (speed of information  
5 processing); *Digit Span Backwards* (working memory); *Symbol Search* (speed of information  
6 processing). The first unrotated principal component explained 53% of the variance and all  
7 subtests had high loadings.

8           *Processing speed factor*

9           A processing speed factor was derived from a PCA of scores on a set of speed of  
10 processing measures, namely: *Symbol Search* (WAIS-III); *Digit Symbol* (WAIS-III);  
11 *Inspection Time* (computer-based task used to assess speed of elementary visual processing  
12 with no requirement for speeded reactions; Deary et al., 2004a); *Simple (SRT) and Choice*  
13 *Reaction Time (CRT)* (speed and variability of simple information processing; Cox, Huppert &  
14 Whichelow, 1993; Deary, Der, & Ford, 2001). The reaction time tasks were administered using  
15 a purpose built portable machine with five response keys (1, 2, 0, 3, 4). In *SRT*, the participant  
16 pressed the 0 key as quickly as possible after each 0 was shown on the LED screen (20 trials).  
17 In *CRT (four-choice)*, the participant pressed the appropriate key (1, 2, 3, or 4) according to the  
18 number which appeared on the LED screen, as quickly as possible (40 trials). Mean SRT and  
19 CRT response time and standard deviation were calculated. The first unrotated principal  
20 component for the speed factor explained 51% of the variance and all tests had high loadings.

21           *Memory factor*

22           A memory factor was derived from a PCA of scores on a set of memory measures from  
23 Wechsler Memory Scale-III<sup>UK</sup> (WMS-III; Wechsler, 1998b), namely: *Logical Memory I*  
24 *Immediate and II Delayed Recall* (verbal declarative memory); *Spatial Span Forwards* and  
25 *Spatial Span Backwards* (non-verbal spatial learning and memory); *Verbal Paired Associates I*

1 *Immediate Recall and II Delayed Recall* (verbal learning and memory; immediate and delayed  
2 recall). The first unrotated principal component explained 43% of the variance and all tests had  
3 high loadings.

4 *Verbal ability*

5 Verbal ability was assessed using the *National Adult Reading Test* (NART: Nelson &  
6 Willison, 1991) and the *Wechsler Test of Adult Reading* (WTAR: Holdnack, 2001). These tests  
7 are widely used to estimate prior cognitive ability and each requires the pronunciation of a list  
8 of 50 irregular words.

9 *Mini-Mental State Examination (MMSE)*

10 The *MMSE* is a standardised brief screening measure for cognitive pathology (Folstein  
11 et al., 1975). Scores range from 0-30, with a score of less than 24 often used to indicate  
12 possible dementia.

13 *The Moray House Test (MHT)*

14 Participants had previously taken part in the nationwide Scottish Mental Survey of  
15 1947 (SMS1947) at age 11. All school children born in 1936 and attending Scottish schools on  
16 the 4<sup>th</sup> of June 1947, took a version of the *Moray House Test (MHT)* No.12 (Scottish Council  
17 for Research in Education; SCRE, 1933; 1949), a group administered mental test comprising  
18 71 items and with a time limit of 45 minutes. Often referred to as a ‘verbal reasoning’ test, it  
19 contains items, including: word classification; arithmetic; spatial items; cypher decoding;  
20 same-opposites; reasoning; proverbs; practical items; analogies; and following directions. The  
21 *MHT* was concurrently validated against the Terman-Merrill Revision of the Binet Scales  
22 (Terman & Merrill, 1937) following the SMS1947. The correlation for both boys and girls was  
23 .81 (SCRE, 1949). The test conducted at age 11 reflects cognitive functioning towards the end  
24 of primary school education and is a valid measure of early-life ability. LBC1936 participants  
25 repeated the same test about 60 years later, at ~age 70, as part of the cognitive assessment.

1 *MHT* scores at age 11 and 70 were corrected for age in days at time of testing and converted to  
2 an IQ scale where  $M = 100$  and  $SD = 15$ . In this sample, the correlation between age 11 and  
3 age 70 IQ is .67 ( $p < .001$ ).

4 **Demographic and control variables.** Marital status, education (years of full time  
5 education) and smoking status (current, ex or never smoker) were ascertained during interview.  
6 A medical history was taken (including diagnoses of diabetes, high blood pressure, high  
7 cholesterol, cardiovascular disease and stroke). Body mass index (BMI) was calculated from  
8 height and weight measurements taken during the physical examination. A physical activity  
9 measure (number of days per month of exercise) was obtained from a self-report questionnaire  
10 booklet comprising various social and lifestyle questions. Adult social class was derived from  
11 participants' highest reported occupation and consisted of 6 categories ranging from I  
12 (professional occupations) to V (unskilled occupations), with III (skilled occupations) divided  
13 into IIIN (non-manual) and IIIM (manual) (Office for Population Censuses and Surveys,  
14 1980). Women were assigned a social class based on the highest occupation of the household.  
15 Due to the small number of participants in class V, classes IV and V were combined.

16

#### 17 **Statistical analyses**

18 Analyses were performed using SPSS v.14.0. Participants were categorised as non-  
19 drinkers, low-level drinkers ( $\leq 2$  units per day) or moderate-substantial drinkers ( $> 2$  units per  
20 day). This classification was used to illustrate any demographic and health differences between  
21 alcohol intake groups, using analysis of variance and Chi-Square tests, as appropriate. The  
22 main analyses examined the associations between alcohol intake (units/day) as a continuous  
23 variable and cognitive outcome scores (separately for men and women) using general linear  
24 models. Cognitive outcomes were: age 70 IQ;  $g$  factor; processing speed factor; memory  
25 factor; NART and WTAR. Four models were fitted to the data, each including adjustment for

1 potential confounding factors. Model 1 tested the unadjusted effects of alcohol on each  
2 outcome measure. Model 2 included age 11 IQ to control for early life ability. Model 3  
3 included occupational social class. The final model adjusted for age 11 IQ and occupational  
4 social class in combination. We present relevant estimates of effect size, reported here as  
5 partial eta-squared ( $\eta_p^2$ ), as well as p-values. Post-hoc, the Sobel Test (Sobel, 1982) was used  
6 to test for mediating effects of occupational social class on the associations between childhood  
7 IQ and alcohol consumption. An online resource was used  
8 (<http://www.danielsoper.com/statcalc/calc31.aspx>) to calculate the Sobel Test Statistic and we  
9 report regression coefficients, standard errors, Sobel statistic ( $z$ ) and p-values.

10

11

## Results

### Descriptive

13

14

15

16

17

**[INSERT TABLE 1 APPROX. HERE]**

19

20

21

22

23

24

25

**Characteristics of non-responders.** Compared to those who completed the FFQ, non-responders ( $n = 163$ ) were significantly more likely to: be older ( $p = .002$ ), male ( $p = .006$ ), belong to a less professional occupational social class, have fewer years of education, have a lower MMSE, have lower age 11 and age 70 IQ (all  $p < .001$ ), and a higher BMI ( $p = .002$ ); and were more likely to have had a diagnosis of diabetes ( $p = .049$ ) or stroke ( $p = .020$ ).

**Characteristics of alcohol intake categories.** The characteristics of alcohol intake categories are shown in table 1. 15% (134) reported no current alcohol intake (non-drinkers), 54% (497) drank 2 units or less per day (low-level drinkers;  $M = 0.8$ ,  $SD = 0.6$ ), and 31% (286) reported a daily intake of greater than 2 units a day (moderate-substantial drinkers; range 2.02 - 20.88,  $M = 4.6$ ,  $SD = 2.7$ ). There were significant gender differences between intake groups. More women than men were non- and low-level drinkers, and more men than women were in the moderate-substantial drinking category. Nearly half the men consumed more than 2 units

1 (equivalent to one pint of beer or one glass of wine) per day compared to one in six women.  
2 Apart from being male, those reporting a higher alcohol intake were significantly more likely  
3 to belong to a professional occupational social class (67% of moderate-substantial drinkers  
4 belonged to social class groups I and II, compared to 48% of the non-drinkers), have more  
5 years of education, and were less likely to be smokers. Those with a higher alcohol intake were  
6 significantly more likely to have a higher childhood IQ and a higher age 70 IQ (both  $p < .001$ ).  
7 Moderate-substantial drinkers had, on average, a 6.8 point higher age 11 IQ than non-drinkers,  
8 and a 6.1 point higher age 70 IQ than non-drinkers. Non-drinkers were significantly more  
9 likely to have had a diagnosis of cardiovascular disease than low or moderate-substantial  
10 drinkers. Alcohol intake was not associated with marital status, MMSE score, BMI, level of  
11 physical activity or history of hypertension, stroke, diabetes or high cholesterol.

12 **[INSERT TABLE 2 APPROX. HERE]**

### 13 **Alcohol intake category and cognitive outcomes at age 11 and age 70**

14 Separate analyses of cognition-alcohol intake associations were conducted for men and  
15 women (see table 2). Mean cognitive scores differed significantly between alcohol intake  
16 categories for all cognitive outcome variables in men; and in women, with the exception of age  
17 70 IQ scores (although the trend was in the same direction). The best cognitive scores for men  
18 and women were among those drinking  $> 2$  units per day; these moderate-substantial drinkers  
19 did better on tests at age 11 and age 70 than both low-level and non-drinkers. The lowest  
20 cognitive scores were associated, almost entirely, with the non-drinkers. In an additional  
21 analysis, we re-classified drinkers into those drinking within current UK guidelines: 21 units  
22 per week for men ( $n = 270$ ); 14 units per week for women ( $n = 303$ ) and those exceeding this  
23 weekly upper limit (men  $n = 132$ , women  $n = 78$ ). This did not appreciably change any of the  
24 associations seen in table 2; the best cognitive scores were still found among those in the  
25 highest consumption category. These results are available from the authors on request.

1           **[INSERT TABLE 3 APPROX. HERE]**

2           **Type of alcohol, and associations with childhood IQ and social class**

3           Table 3 presents mean intake of each type of alcohol for men and women. The women  
4 in our sample were found to derive most of their alcohol units (80%) from wine, whereas men  
5 consumed alcohol from a larger range of sources. The mean daily alcohol intake of men (2.63  
6 units) was more than double that of women (1.14 units;  $p < .001$ ). Compared to women, men  
7 consumed significantly more wine, spirits and beer. Consumption of low-alcohol beer,  
8 alcopops and cider were very low in this older cohort; therefore, no further analyses are  
9 presented using these alcohol types. We examined the relationship between alcohol intake, age  
10 11 IQ and social class. Intake of red wine, white wine, sherry/port and total alcohol intake were  
11 associated with a higher childhood IQ and more professional social class ( $p < .001$ ).

12           **[INSERT TABLE 4 APPROX. HERE]**

13           **General linear models**

14           Four stages of general linear model (GLM) were fitted to the data to examine the  
15 contribution of alcohol (total and by type) and potentially confounding variables to age 70  
16 cognitive function. Table 4 presents the main GLM results, by gender. For women, we present  
17 the effects of total alcohol intake only, as associations were found to reflect the predominant  
18 influence of wine intake on cognitive function (full results can be seen in supplementary table  
19 1). Model 1 (unadjusted) showed a positive association between total alcohol (wine) intake and  
20 performance across all cognitive domains; the largest effect sizes were seen for verbal ability  
21 (NART,  $\eta_p^2 = .050$ ; WTAR,  $\eta_p^2 = .042$ ). When age 11 IQ and social class were added  
22 (independently) to the models, the associations with age 70 IQ,  $g$  factor and processing speed  
23 became non-significant. The positive associations with memory and NART (but not WTAR)  
24 scores remain significant, even after controlling for age 11 IQ and social class in combination.

1 However, effect sizes were markedly reduced. Overall, the attenuating effects of childhood IQ  
2 and social class on these positive associations are very substantial.

3 In men, nearly half (48%) their total alcohol intake comes from wine, and the  
4 remainder, from mainly beer and spirits. Table 4 presents the results for total alcohol intake  
5 and wine intake (full results can be seen in supplementary table 2). In the initial unadjusted  
6 model, total alcohol intake was associated with significantly better test performance in all  
7 cognitive domains. The largest effect sizes were for memory ( $\eta_p^2 = .035$ ) and age 70 IQ ( $\eta_p^2 =$   
8  $.032$ ). After full adjustment, the only remaining statistical effect of overall alcohol intake on  
9 cognitive outcomes in men was a positive association with memory scores, which was also  
10 found in women. Contrary to the results found for overall alcohol intake, wine consumption in  
11 men was associated with significantly better performance on both verbal ability tests (NART,  $p$   
12  $= .004$ ,  $\eta_p^2 = .020$ ; WTAR,  $p = .031$   $\eta_p^2 = .011$ ) and statistical significance held throughout the  
13 models. Again, substantial attenuation from age 11 IQ and social class was observed. The type  
14 of alcohol consumed by men was an important factor in relation to cognitive performance (see  
15 supplementary table 2). Notably, wine and sherry/port consumption was associated with better  
16 cognitive performance, especially on both verbal ability tests, even after full adjustment. The  
17 converse was true of beer consumption which was associated with poorer NART performance  
18 after adjustments ( $p = .041$ ,  $\eta_p^2 = .010$ ). However, spirits intake was associated with a better  
19 memory performance ( $p = .004$ ,  $\eta_p^2 = .021$ ).

20 Interaction terms were added in separate models (not presented in Tables) to examine  
21 whether there were any interaction effects between childhood IQ and alcohol intake, or social class  
22 and alcohol intake, on later cognitive outcomes in either sex. In women, we found no evidence of  
23 any interaction effects of childhood IQ and alcohol. However, there was a social class-alcohol  
24 interaction on age 70 IQ ( $p < .001$ ,  $\eta_p^2 = .050$ ) and processing speed ( $p = .026$ ,  $\eta_p^2 = .025$ ). Data  
25 plots suggested that the only deleterious effects of alcohol consumption on age 70 IQ and



1 processing speed were found in the manual social classes (IIIM, IV, and V). In men, we found no  
2 evidence of any interaction effects of social class on any of the cognitive outcomes. However,  
3 there was a significant childhood IQ-alcohol interaction on age 70 IQ ( $p < .001$ ,  $\eta_p^2 = .044$ ); and  
4 significant childhood IQ-wine interactions on age 70 IQ ( $p < .001$ ;  $\eta_p^2 = .043$ ), NART ( $p < .002$ ;  
5  $\eta_p^2 = .023$ ) and WTAR ( $p < .037$ ;  $\eta_p^2 = .010$ ). Data plots suggested that, in those drinking less  
6 alcohol, there were stronger correlations between age 11 and age 70 IQ. That is, in those  
7 participants drinking little or no alcohol, more of the variance in their age 70 IQ scores could be  
8 accounted for by childhood ability when compared with those drinking more alcohol, among  
9 whom causes other than childhood IQ contributed more to cognitive variation in old age.

10 We also examined the potentially mediating effects of social class on the link between  
11 prior cognitive ability and alcohol consumption. Using the Sobel test, we identified a mediating  
12 effect of social class, on both the association between childhood IQ and total alcohol  
13 consumption ( $A = -.024$ ,  $SE_A = .002$ ;  $z = 2.99$ ,  $p < .001$ ), and between childhood IQ and wine  
14 consumption ( $B = -.303$ ,  $SE_B = .098$ ;  $z = 5.59$ ,  $p < .001$ )<sup>2</sup>. The standardised beta of the direct  
15 path (age 11 IQ-total alcohol) was .020, and .012 after social class was introduced as a  
16 mediator. The amount of the relationship accounted for by social class was .008, representing  
17 40% of the direct effect. For wine intake, the standardised beta of the direct path (age 11 IQ-  
18 wine drinking) was .019, and .008 after social class was introduced as a mediator. Therefore,  
19 the amount of the relationship between age 11 IQ and wine drinking accounted for by social  
20 class was .011, representing 58% of the direct effect.

21

22

## Discussion

---

<sup>2</sup> A = the unstandardized regression coefficient for the relationship between the IV and mediator;  $SE_A$  = standard error of the relationship between the IV and mediator; B = the regression coefficient for the relationship between the mediator and the DV;  $SE_B$  = the standard error of the relationship between the mediator and the DV; z = Sobel Test Statistic

1           In the present study, drinking more alcohol was associated with better cognitive  
2 performance at age 70, by comparison with low-level drinking and no drinking. Moderate-  
3 substantial drinkers had better cognitive scores across all cognitive domains tested, with the  
4 exception of age 70 IQ scores in women (although this association did not reach significance  
5 level, it followed the same positive trend). In line with previous research, abstainers performed  
6 more poorly than light and moderate drinkers. However, after controlling for childhood IQ and  
7 SES, there remained little evidence of a relationship between alcohol intake and current  
8 cognitive function. Where significant effects remained, the reductions in effect sizes were  
9 striking. The apparent ‘benefits’ (after controlling for IQ and SES) of a higher overall alcohol  
10 intake were confined to memory performance in both men and women. Male drinkers of wine  
11 and sherry/port also appeared to have a better verbal ability (crystallized intelligence).  
12 However, these effect sizes were modest. It is plausible that the positive alcohol-crystallized IQ  
13 associations were still significant after controlling for age 11 IQ because crystallized IQ tests  
14 measure ‘peak’ ability and capture variance related to the accumulated intellectual  
15 development, and the associated lifestyle and education choices, that take place between  
16 childhood and adulthood. However, these ‘positive’ effects did not extend to drinking beer.  
17 Beer intake was associated with a poorer crystallized IQ (based on the NART score). Women  
18 with a higher wine intake performed better on tests of crystallized IQ and memory, and those  
19 consuming spirits performed better on one of the tests of crystallized IQ (NART).

20           Many previous studies have concluded that alcohol consumption has a direct protective  
21 effect on cognition, via vascular mechanisms or otherwise. The current study indicates that this  
22 premature conclusion may be erroneous. Prior ability and SES were found to significantly  
23 confound the relationship between alcohol intake and cognitive abilities in old age. These  
24 findings are in keeping with the previous literature where higher intelligence, measured as  
25 early as childhood, was related to a higher alcohol intake in adult life (Batty et al., 2008) but

1 less alcohol induced hangovers (Batty, Deary, & Macintyre, 2006) suggestive of moderate  
2 consumption. One other study has attempted to examine the effects of prior ability on the  
3 cognitive effects of alcohol use in old age (Cooper et al., 2009). However, it used only an  
4 estimate, rather than an actual measure of prior ability. The authors concluded that cognitive  
5 ability was no longer associated with overall alcohol use once estimated ‘premorbid IQ’  
6 (NART) was controlled for. In the present study it was clear that in addition to having NART  
7 scores, it was important to have a measure of early life intelligence, and also to analyse the data  
8 according to type of alcohol-based drink.

9         Moderate drinking has previously been reported to be more prevalent in educated,  
10 affluent classes and non-drinking more concentrated in the least educated, deprived groups  
11 (Jefferis, Manor, & Power, 2007). Our findings support those of a prior study linking alcohol  
12 intake, SES and cognitive function, whereby a pattern of higher SES and cognitive scores was  
13 found in men consuming light-moderate levels of alcohol than in abstainers or heavy drinkers,  
14 and in women drinkers, irrespective of level consumed (Richards, Hardy & Wadsworth, 2005).

15         Those with a higher IQ (even when measured in childhood) and more advantaged adult  
16 SES are more likely to develop a preference for wine and sherry/port drinking. This is  
17 consistent with findings from other studies, suggesting that these effects are not particular to  
18 this cohort, or to Scottish culture. For example, in studies using large Danish samples,  
19 preference for wine over beer and other alcoholic beverages was linked to a higher IQ  
20 (Mortensen, Sørensen, & Grønbaek, 2005) and a higher social class (Nielsen, Schnor, Jensen &  
21 Grønbaek, 2004; Osler, Godtfredsen & Prescott, 2008). In France, social and environment  
22 factors have been linked to alcohol preference; wine drinking was associated with a more  
23 favourable social environment whereas the converse was true for beer drinking (Ruidavets et  
24 al., 2004). In the USA, wine preference was associated with significantly higher level of  
25 education (Paschall & Lipton, 2005). We also found in women that the association between

1 overall alcohol intake and cognitive function was not the same for all social classes; there was  
2 some evidence of relative cognitive disadvantage (in age 70 IQ and processing speed) with  
3 alcohol, but only in those women drinking greater amounts and who belonged to the manual  
4 occupational social class groups. In men, the effects of alcohol were less likely to be moderated  
5 by social class but were, to some extent, by prior ability. There was a particularly ‘beneficial’  
6 (modifying) effect of wine consumption on cognitive performance (age 70 IQ and verbal  
7 ability) in those with a higher childhood IQ. Correspondingly, the association between  
8 childhood IQ and cognitive ability in later life was significantly lower among people who  
9 drank alcohol when compared with those who drank little or none.

10 Moderate alcohol consumption and a preference for wine, in those who are more  
11 cognitively able in later life, may be due to the influence of prior ability and social  
12 circumstances, on lifestyle factors. Wine drinkers tend to have more favourable health and  
13 lifestyle characteristics (e.g. a healthier diet) than that of predominantly beer and spirits  
14 drinkers (Paschall & Lipton, 2005). Given the literature documenting the association between a  
15 higher IQ and a healthier lifestyle in large, population-representative samples (Batty, Deary,  
16 Schoon, & Gale, 2007), it may be that people with higher cognitive ability engage in a lifestyle  
17 that protects against cognitive decline. Here, we found no differences between alcohol  
18 consumption groups in the markers of lifestyle that were measured, i.e. physical activity and  
19 BMI. Further research, perhaps incorporating dietary measures, could evaluate this further.

## 20 **Study limitations**

21 Information regarding longer-term history of drinking was not available for this dataset.  
22 This allowed for previous drinking in the non-drinking group. Ex-drinkers may have very  
23 different characteristics from ‘never drinkers’ (Wannamethee & Shaper, 2002). Compared to  
24 lifelong teetotalers, ex-drinkers show higher mortality rates (De Dabry et al., 1992, Shaper &  
25 Wannamethee, 2000) and cardiovascular risk factors (Shaper & Wannamethee, 2000). In the

1 LBC1936, non-drinkers have a higher incidence of hypertension, diabetes, high cholesterol,  
2 stroke and cardiovascular disease than moderate-substantial drinkers. Rather than reflecting a  
3 health benefit to moderate drinking, these data may reflect ex-drinkers having given up alcohol  
4 due to poor physical health.

5         The FFQ measures alcohol intake from the most recent 2-3 month period. Although this  
6 short-term data cannot be assumed to reflect habitual patterns of alcohol consumption, there  
7 are a number of reports supporting the validity of this form of measurement, given the  
8 temporal stability of patterns of alcohol intake in later life. For example, Ruitenberg et al.  
9 (2002) found that only 6% of older participants reported a change in drinking pattern in the  
10 previous five years. In the present study, classification of alcohol intake into groups was used  
11 for illustrative purposes, not for the key analyses. While this could lead to misclassification,  
12 self-reports of alcohol consumption are assumed to be valid for the purpose of classifying  
13 drinkers into broad consumption bands (Eren, 1995). Drinkers were classified into  
14 consumption groups based on units of alcohol intake, as in previous literature (e.g., Huang,  
15 Qiu, Winblad, & Fratiglioni, 2002; Stott et al., 2008). Our method was also consistent with  
16 those studies using a three-tiered classification representing: non-drinkers; low-level/minimal  
17 drinkers (equivalent to  $\leq 1$  drink/2 units per day); and moderate drinkers (equivalent to  $>1$   
18 drink/2 units per day (e.g. McGuire, Ajani, & Ford, 2007; Britton, Marmot, & Shipley, 2008;  
19 Lang, Wallace, Huppert, & Melzer, 2007; Espeland et al., 2006).

#### 20         **Generalisability**

21         The LBC1936 are a somewhat self-selected sample. The sample represents a healthier  
22 and more cognitively able subgroup of the original SMS1947 cohort. This healthy survivor  
23 effect may have restricted the range of cognitive outcome scores in the present study.  
24 However, the range of cognitive abilities (and IQ scores) was still large and a restriction in  
25 range of abilities would likely lead to a modest underestimation of the effect sizes.

1 **Study advantages**

2 All of the participants were born in the same year (1936), thereby eliminating cohort  
3 effects and the effects of chronological age. Other studies have been hampered by inadequate  
4 cognitive test batteries, limited alcohol measures and inadequate control for confounders. The  
5 LBC1936 Study employs a comprehensive battery of cognitive tests. Studies using early life  
6 IQ data are rare and given the stability of IQ across the lifespan (Deary, Whalley, Lemmon,  
7 Crawford, & Starr, 2000; Deary, Whiteman, Starr, Whalley, & Fox, 2004b), the availability of  
8 such data offers a unique advantage in the investigation of factors affecting cognition with  
9 aging. The FFQ has advantages in terms of ascertaining detailed information on frequency and  
10 amount of different sources of alcohol intake, which is often lacking in studies which only ask  
11 about overall alcohol intake. Using FFQ data allowed us to look at the associations between  
12 different types of alcohol consumption on cognitive abilities. This proved to be important. The  
13 associations between alcohol consumption and cognitive domains are not uniform, and  
14 highlight the importance of making a distinction between alcoholic beverage types during data  
15 collection. The cross-sectional nature of this study is a potential limitation, but the LBC1936  
16 Study is ongoing. There will be opportunities for follow-ups of this cohort, offering the  
17 potential to investigate alcohol's effects, if any, on cognitive decline.

18 **Conclusions**

19 Our results support the concept that the previously reported 'moderate alcohol  
20 consumption-better cognition' association is, substantially, a consequence of confounding by  
21 higher prior cognitive ability and adult SES. The exceptions were positive associations  
22 between alcohol intake and memory performance, and verbal ability. However, the effects  
23 were small, and the clinical significance of these findings is uncertain. It is not until we  
24 examine mediating factors such as IQ and SES more fully across the lifespan that we can begin  
25 to examine the two-way nature of the alcohol-cognition relationship.

26

1 **References**

- 2 Batty, G. D., Deary, I. J., & Macintyre, S. (2006). Childhood IQ and life course socioeconomic  
3 position in relation to alcohol induced hangovers in adulthood: the Aberdeen children  
4 of the 1950s study. *Journal of Epidemiology and Community Health, 60*, 872-874.
- 5 Batty, G. D., Deary, I. J., Schoon, I., Emslie, C., Hunt, K., & Gale, C. R. (2008). Childhood  
6 mental ability and adult alcohol intake and alcohol problems: the 1970 British Cohort  
7 Study. *American Journal of Public Health, 98*, 2237-2243.
- 8 Batty, G. D., Deary, I. J., Schoon, I., & Gale, C. R. (2007). Childhood mental ability in relation  
9 to food intake and physical activity in adulthood: The 1970 British Cohort Study.  
10 *Pediatrics, 119*, e38-45.
- 11 British Medical Association (2008). Alcohol misuse: tackling the UK epidemic.
- 12 Britton, A., Marmot, M. G., & Shipley, M. (2008). Who benefits the most from the  
13 cardioprotective properties of alcohol consumption – health freaks or couch potatoes?  
14 *Journal of Epidemiology and Community Health, 62*, 905-908.
- 15 Britton, A., Singh-Manoux, A., & Marmot, M. (2004). Alcohol consumption and cognitive  
16 function in the Whitehall II Study. *American Journal of Epidemiology, 160*, 240-247.
- 17 Cervilla, J. A., Prince, M., & Mann, A. (2000). Smoking, drinking and incident cognitive  
18 impairment: a community based study included in the Gospel Oak project. *Journal of*  
19 *Neurology, Neurosurgery & Psychiatry, 68*, 622-626.
- 20 Cooper, C., Bebbington, P., Meltzer, H., Jenkins, R., Brugha, T., Lindesay, J. E. B., &  
21 Livingston, G. (2009). Alcohol in moderation, premorbid intelligence and cognition in  
22 Older Adults: results from the Psychiatric Morbidity Survey. *Journal of Neurology,*  
23 *Neurosurgery & Psychiatry, 80*, 1236-1239.
- 24 Cox, B. D., Huppert, F. A., & Whichelow, M. J. (1993). *The health and lifestyle survey: seven*  
25 *years on*. Aldershot, UK: Dartmouth.

- 1 <http://www.danielsoper.com/statcalc/calc31.aspx> Retrieved 16th June 2009.
- 2 Deary, I. J., Whalley, L. J., Lemmon, H., Crawford, J. R., & Starr, J. M. (2000). The stability  
3 of individual differences in mental ability from childhood to old age: Follow up of the  
4 1932 Scottish Mental Survey. *Intelligence*, 28, 49-55.
- 5 Deary, I. J., Der, G., & Ford, G. (2001). Reaction times and intelligence differences: a  
6 population-based cohort study. *Intelligence*, 29, 389-399.
- 7 Deary, I. J., Simonotto, E., Meyer, M., Marshall, A., Marshall, I., Goddard, N., & Wardlaw, J.  
8 M. (2004a). The Functional Anatomy of Inspection Time: an event-related fMRI study.  
9 *NeuroImage*, 22, 1466-1479.
- 10 Deary, I. J., Whiteman, M. C., Starr, J. M., Whalley, L. J., & Fox, H. C. (2004b). The impact of  
11 childhood intelligence on later life: following up the Scottish Mental Surveys of 1932  
12 and 1947. *Journal of Personality and Social Psychology*, 86, 130-147.
- 13 Deary, I. J., Gow, A. J., Taylor, M. D., Corley, J., Brett, C., Wilson, V., ... Starr, J. M. (2007).  
14 The Lothian Birth Cohort 1936: a study to examine the influences on cognitive ageing  
15 from age 11 to age 70 and beyond. *BMC Geriatrics*, 7, 28.
- 16 Deary, I. J., Whalley, L. J., & Starr, J. M. (2009). *A Lifetime of Intelligence: Follow-up Studies*  
17 *of the Scottish Mental Surveys of 1932 and 1947*. Washington, DC: American  
18 Psychological Association.
- 19 De Dabry, L. O., Glynn, R. J., Levenson, M. R., Hermos, J. A., Locastro, J. S., & Vokanos, P.  
20 S. (1992). Alcohol consumption and mortality in an American male population.  
21 Recovering the U shaped curve – Findings from the Normative Aging Study. *Journal of*  
22 *Studies on Alcohol*, 53, 25-32.
- 23 [http://www.direct.gov.uk/en/HealthAndWellBeing/DG\\_10036434](http://www.direct.gov.uk/en/HealthAndWellBeing/DG_10036434) Retrieved 16th Jan 2009.



- 1 Dufouil, C., Ducimetière, P., & Alperovitch, A. (1997). Sex differences in the association  
2 between alcohol consumption and cognitive performance. *American Journal of*  
3 *Epidemiology*, *146*, 405-412.
- 4 Elwood, P. C., Gallacher, J. E. J., Hopkinson, C. A., Pickering, J., Rabbitt, P., Stollery, B., ...  
5 Bayer, A. (1999). Smoking, drinking and other lifestyle factors and cognitive function  
6 in men in the Caerphilly cohort. *Journal of Epidemiology and Community Health*, *53*,  
7 9-14.
- 8 Eren, B. (1995). Alcohol consumption; in *The Scottish Health Survey*.
- 9 Espeland, M. A., Gu, L., Masaki, K. H., Langer, R. D., Coker, L. H., Stefanick, M. L., ...  
10 Rapp, S. R. (2005). Association between reported alcohol intake and cognition: Results  
11 from the Women's Health Initiative Memory Study. *American Journal of*  
12 *Epidemiology*, *161*, 228-238.
- 13 Espeland, M. A., Coker, L. H., Wallace, R., Rapp, S. R., Resnick, S. M., Limacher, M., ...  
14 Messina, C. R. (2006). Association between alcohol intake and domain-specific  
15 cognitive function in older women. *Neuroepidemiology*, *27*, 1-12.
- 16 Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). Mini-Mental State: A practical  
17 method for grading the cognitive state of patients for the clinician. *Journal of*  
18 *Psychiatric Research*, *12*, 189-198.
- 19 Ganguli, M., Vander Bilt, J., Saxton, J. A., Shen, C., & Dodge, H. H. (2005). Alcohol  
20 consumption and cognitive function in late life. *Neurology*, *65*, 1210-1217.
- 21 Holdnack, J. A. (2001). *WTAR: Wechsler Test of Adult Reading manual*. San Antonio, TX:  
22 Psychological Corporation.
- 23 Huang, W., Qiu, C., Winblad, B., & Fratiglioni, L. (2002). Alcohol consumption and incidence  
24 of dementia in a community sample aged 75 years and older. *Journal of Clinical*  
25 *Epidemiology*, *55*, 959-964.

- 1 Jefferis B. J. M. H., Manor, O., & Power, C. (2007). Social gradients in binge drinking and  
2 abstaining: trends in a cohort of British adults. *Journal of Epidemiology and*  
3 *Community Health, 61*, 150-153.
- 4 Jia, X., Craig, L. C., Aucott, L. S., Milne, A. C., & McNeill, G. (2008). Repeatability and  
5 validity of a food frequency questionnaire in free-living older people in relation to  
6 cognitive function. *Journal of Nutrition, Health and Aging, 12*(10), 735-41.
- 7 Lang, I., Wallace, R. B., Huppert, F. A., & Melzer, D. (2007). Moderate alcohol consumption  
8 in older adults is associated with better cognition and well-being than abstinence. *Age*  
9 *and Ageing, 36*, 256-261.
- 10 Leroi, I., Sheppard, J. M., & Lyketsos, C. G. (2002). Cognitive function after 11.5 years of  
11 alcohol use: Relation to alcohol use. *American Journal of Epidemiology, 156*, 747-752.
- 12 Luchsinger, J. A., Tang, M. X., Siddiqui, M., Shea, S., & Mayeux, R. (2004). Alcohol intake  
13 and risk of dementia. *Journal of the American Geriatrics Society, 52*, 540-546.
- 14 Masson, L. F., McNeill, G., Tomany, J. O., Simpson, J. A., Peace, H. S., Wei, L., ... Bolton-  
15 Smith, C. (2003). Statistical approaches for assessing the relative validity of a food-  
16 frequency questionnaire: use of correlation coefficients and the kappa statistic. *Public*  
17 *Health Nutrition, 6*, 313-321.
- 18 McGuire, L. C., Ajani, U. A., & Ford, E. S. (2007). Cognitive functioning in late life: The  
19 impact of moderate alcohol consumption. *Annals of Epidemiology, 17*, 93-99.
- 20 McNeill, G., Winter, J., & Jia, X. (2009). Diet and cognitive function in later life: a challenge  
21 for nutritional epidemiology. *European Journal of Clinical Nutrition, 63*, S33-37.
- 22 Mortensen, L. H., Sørensen, T. I. A., & Grønbaek, M. (2005). Intelligence in relation to later  
23 beverage preference and alcohol intake. *Addiction, 100*, 1445-1452.

- 1 Mukamal, K. J., Longstreth, W. T. Jr, Mittleman, M. A., Crum, R. M., & Siscovick, D. S.  
2 (2001). Alcohol consumption and subclinical findings on magnetic resonance imaging  
3 of the brain in older adults: the Cardiovascular Health Study. *Stroke*, 32, 1939-46.
- 4 Naimi, T. S., Brown, T. W., Brewer, R. D., Giles, W. H., Mensah, G., Serdula, M. K., ...  
5 Stroup, D. F. (2005). Cardiovascular risk factors and confounders among nondrinking  
6 and moderate-drinking US adults. *American Journal of Preventive Medicine*, 28, 369-  
7 373.
- 8 Nelson, H. E., & Willison, J. R (1991). *National Adult Reading Test (NART) Test Manual*  
9 *(Part II)*. Windsor, UK: NFER-Nelson.
- 10 Nielsen, N. R., Schnohr, P., Jensen, G., & Grønbaek, M. (2004). Is the relationship between  
11 type of alcohol and mortality influenced by socio-economic status? *Journal of Internal*  
12 *Medicine*, 255, 280-288.
- 13 Office of Population Censuses and Surveys (1980). *Classification of occupations 1980*.  
14 London: HMSO.
- 15 Osler, M., Gotfredsen, N. S., & Prescott, E. (2008). Childhood social circumstances and health  
16 behaviour in midlife: the Metropolit 1953 Danish male birth cohort. *International*  
17 *Journal of Epidemiology*, 37, 1367-1374.
- 18 Paschall, M. & Lipton, R. I. (2005). Wine preference and related health determinants in a U.S.  
19 national sample of young adults. *Drug and Alcohol Dependence*, 78, 339-344.
- 20 Peters, R., Peters, J., Warner, J., Beckett, N., & Bulpitt, C. (2008). Alcohol, dementia and  
21 cognitive decline in the elderly: a systematic review. *Age and Ageing*, 37, 505-512.
- 22 Richards, M., Hardy, R., & Wadsworth, M. E. J. (2005). Alcohol consumption and midlife  
23 cognitive change in the British 1946 Birth Cohort. *Alcohol and Alcoholism*, 40, 112-17.

- 1 Rimm, E. B., Williams, P., Fosher, K., Criqui, M., & Stampfer, M. J. (1999). Moderate alcohol  
2 intake and lower risk of coronary heart disease: meta-analysis of effects on lipids and  
3 haemostatic factors. *British Medical Journal*, *319*, 1523-1528.
- 4 Ruidavets, J-B, Bataille, V., Dallongeville, J., Simon, C., Bingham, A., Amouyel, P.,  
5 ...Ferrières, J. (2004). Alcohol intake and diet in France, the prominent role of lifestyle.  
6 *European Heart Journal*, *25*, 1153-1162.
- 7 Ruitenberg, A., Swieten, J., Witteman, J., Mehta, K. M., van Duijn, C. M., Hofman, A., &  
8 Breteler, M. M. B. (2002). Alcohol consumption and risk of dementia: the Rotterdam  
9 Study. *Lancet*, *359*, 281-6.
- 10 Sobel, M. E. (1982). Asymptotic Confidence Interval for Indirect Effects in Structural  
11 Equation Models. In S. Leinhardt (Eds.), *Sociological Methodology* (pp. 290-213). San  
12 Francisco, CA: Jossey-Bass.
- 13 Scottish Collaborative Group SCG, University of Aberdeen . Scottish Collaborative Group food  
14 frequency questionnaire. Retrieved from <http://www.abdn.ac.uk/deom/ffq/index.htm> 9<sup>th</sup> Mar  
15 2010.
- 16 Scottish Council for Research in Education (1933). *The Intelligence of Scottish children: a*  
17 *national survey of an age group*. London, UK: University of London Press.
- 18 Scottish Council for Research in Education (1949). *The trend of Scottish intelligence: A*  
19 *comparison of the 1947 and 1932 surveys of the intelligence of eleven-year-old pupils*.  
20 London, UK: University of London Press.
- 21 Shaper, A. G., & Wannamethee, S. G. (2000). Alcohol intake and mortality in middle aged  
22 men with diagnosed coronary artery disease. *Heart*, *83*, 394-399.
- 23 Stampfer, M. J., Kang, J. H., Chen, J., Cherry, R., & Grodstein, F. (2005). Effects of moderate  
24 alcohol consumption on cognitive function in women. *New England Journal of*  
25 *Medicine*, *352*, 245-253.

- 1 Stott, D. J., Falconer, A., Kerr, G. D., Murray, H. M., Trompet, S., Westendorp, R. G. J., ...  
2 Ford, I. (2008). Does low to moderate alcohol intake protect against cognitive decline  
3 in older people? *Journal of the American Geriatrics Society*, 56(12), 2217-24.
- 4 Terman, L. M. & Merrill, M. A. (1937). *Measuring intelligence*. London: Harrap.
- 5 Truelsen, T., Thudium, D., & Grønbæk, M. (2002). Copenhagen City Heart Study. Amount  
6 and type of alcohol and risk of dementia: the Copenhagen City Heart Study. *Neurology*,  
7 59(9), 1313-1319.
- 8 Wannamethee, S. G., & Shaper, A. G. (2002). Taking up regular drinking in middle age: effect  
9 on major coronary heart disease events and mortality. *Heart*, 87, 32-36.
- 10 Wechsler, D. (1998a). *WAIS-III<sup>UK</sup> administration and scoring manual*. London, UK:  
11 Psychological Corporation.
- 12 Wechsler, D. (1998b). *WMS-III<sup>UK</sup> administration and scoring manual*. London, UK:  
13 Psychological Corporation.
- 14 Wright, C. B., Elkind, M. S. V., Luo, X., Paik, M. C., & Sacco, R. L. (2006). Reported alcohol  
15 consumption and cognitive decline: The Northern Manhattan Study.  
16 *Neuroepidemiology*, 27, 201-207.

17  
18  
19  
20  
21  
22  
23

Table 1

*Characteristics of the population by alcohol intake category, with reported mean and p-values*

	Total sample		Non-drinkers		Low-level drinkers ( $\leq 2$ units/day)		Moderate-substantial drinkers ( $> 2$ units/day)		<i>p</i>
	<i>n</i> = 917*		<i>n</i> = 134		<i>n</i> = 497		<i>n</i> = 286		
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	
Alcohol intake (units/day)	1.9	2.4			0.8	0.6	4.6	2.7	n/a
Age (in years)	69.5	0.8	69.6	0.9	69.5	0.8	69.4	0.9	<b>.002</b>
Sex									<b>&lt;.001</b>
Male (%)	48.3		29.9		39.0		72.7		
Female (%)	51.7		70.1		61.0		27.3		
Marital status									.062
Married (%)	72.7		64.2		71.8		78.3		
Widowed (%)	13.2		15.0		15.1		9.1		
Unmarried/divorced (%)	14.1		21.0		13.1		12.6		
Social class (%)									<b>.001</b>
I	18.9		12.4		17.3		25.1		
II	38.4		35.6		36.9		41.9		
IIIN	23.6		29.5		27.1		14.7		
IIIM	15.9		17.0		16.1		15.1		
IV+V	3.3		5.4		2.6		3.3		
Education (yrs f/t)	10.8	1.1	10.5	0.9	10.7	1.1	11.0	1.2	<b>&lt;.001</b>
MMSE	28.9	1.2	28.9	1.3	28.9	1.3	29.0	1.2	.712
Age 11 IQ	101.4	14.0	97.6	14.8	100.6	14.1	104.4	13.0	<b>&lt;.001</b>
Age 70 IQ	101.7	13.2	99.0	13.3	100.4	14.2	105.1	10.3	<b>&lt;.001</b>
Body Mass Index (kg/m <sup>2</sup> )	27.6	4.3	27.7	4.7	27.6	4.3	27.7	4.1	.926
Physical activity (days/mth)	7.6	8.1	6.9	8.6	8.0	8.4	7.3	7.1	.286
Smoking status									<b>&lt;.001</b>

Running head: ALCOHOL INTAKE AND COGNITIVE ABILITIES

Non-smokers (%)	44.4	45.5	49.5	35.0	
Ex-smokers (%)	43.3	36.5	38.8	54.2	
Current smokers (%)	12.3	18.0	11.7	10.8	
Hypertension, yes (%)	39.5	46.3	37.6	39.5	.193
Diabetes, yes (%)	7.5	11.9	6.4	7.3	.100
Cholesterol, yes (%)	35.1	39.5	32.4	37.8	.155
Stroke, yes (%)	4.0	6.7	3.8	3.1	.210
Cardiovascular disease, yes (%)	23.8	33.6	20.5	24.8	<b>.006</b>

---

*Note.*

\*5 subjects had some missing alcohol data – unable to calculate total alcohol intake.

Table 2

*Alcohol intake category and cognitive outcomes for men and women, with reported mean and p-values (ANOVA)*

Cognitive outcome	Men						Women							
	Non-drinkers		Low-level drinkers ( $\leq 2$ units/day)		Moderate-substantial drinkers ( $> 2$ units/day)		Non-drinkers		Low-level drinkers ( $\leq 2$ units/day)		Moderate-substantial drinkers ( $> 2$ units/day)			
	$n = 40$		$n = 194$		$n = 208$		$n = 94$		$n = 303$		$n = 78$			
	$M$	$(SD)$	$M$	$(SD)$	$M$	$(SD)$	$p$	$M$	$(SD)$	$M$	$(SD)$	$M$	$(SD)$	$p$
Age 11 IQ	96.2	(18.3)	98.3	(15.0)	103.7	(13.1)	<b>&lt;.001</b>	98.2	(13.0)	102.2	(13.3)	106.3	(12.3)	<b>&lt;.001</b>
Age 70 IQ	99.8	(13.6)	100.1	(15.2)	106.0	(9.7)	<b>&lt;.001</b>	98.6	(13.2)	100.7	(13.5)	102.9	(11.6)	.103
g factor	-0.10	(1.00)	0.02	(0.98)	0.40	(0.90)	<b>&lt;.001</b>	-0.12	(0.91)	0.01	(0.92)	0.34	(0.91)	<b>.003</b>
Processing speed	-0.02	(0.97)	-0.03	(1.00)	0.31	(0.88)	<b>.001</b>	-0.22	(0.88)	0.08	(0.88)	0.34	(0.92)	<b>&lt;.001</b>
Memory	-0.11	(1.00)	-0.12	(0.93)	0.24	(0.93)	<b>&lt;.001</b>	-0.02	(0.94)	0.04	(0.98)	0.46	(0.88)	<b>.002</b>
NART	34.7	(9.2)	33.2	(7.7)	36.5	(7.5)	<b>&lt;.001</b>	32.8	(7.1)	35.2	(7.7)	39.0	(6.1)	<b>&lt;.001</b>
WTAR	40.9	(7.8)	40.0	(6.7)	43.2	(6.2)	<b>&lt;.001</b>	40.2	(6.5)	41.5	(6.7)	44.5	(4.7)	<b>&lt;.001</b>



Table 3

*Alcohol intake (units/day) by gender (Mann-Whitney U and p-values) and correlations between alcohol (total sample), age 11 IQ and occupational social class*

	Total		Men		Women		Sex differences		Age 11 IQ		Social class	
	<i>n</i> = 922*		<i>n</i> = 445		<i>n</i> = 477		Mann-Whitney U		Spearman rho		Spearman rho	
Alcohol (units/day)	<i>M</i> (range)	% reporting	<i>M</i> (range)	% reporting	<i>M</i> (range)	% reporting	<i>U</i>	( <i>p</i> )	<i>r<sub>s</sub></i>	( <i>p</i> )	<i>r<sub>s</sub></i>	( <i>p</i> )
Total alcohol	1.86 (0-20.9)		2.63 (0-20.9)		1.14 (0-15.1)		<b>64413.5</b>	<b>(&lt;.001)</b>	<b>.194</b>	<b>(&lt;.001)</b>	<b>-.158</b>	<b>(&lt;.001)</b>
Red wine	0.64	54.8	0.83	60.6	0.47	49.5	<b>88969.0</b>	<b>(&lt;.001)</b>	<b>.249</b>	<b>(&lt;.001)</b>	<b>-.291</b>	<b>(&lt;.001)</b>
White wine	0.44	56.1	0.44	51.7	0.44	60.2	<b>98023.5</b>	<b>(.034)</b>	<b>.169</b>	<b>(&lt;.001)</b>	<b>-.256</b>	<b>(&lt;.001)</b>
Sherry/port etc	0.53	26.3	0.61	25.2	0.04	27.5	104318.5	(.561)	<b>.168</b>	<b>(&lt;.001)</b>	<b>-.168</b>	<b>(&lt;.001)</b>
Spirits/liqueurs	0.42	54.7	0.69	66.3	0.16	43.8	<b>70256.0</b>	<b>(&lt;.001)</b>	.053	(.117)	.015	(.643)
Beer (reg+dark)	0.28	36.8	0.56	65.4	0.01	10.1	<b>43129.50</b>	<b>(&lt;.001)</b>	.018	(.589)	.058	(.082)

*Note.*

\*total sample = 922; total sample with total alcohol units/day = 917

Table 4

*Relationship between alcohol (units/day) and cognitive outcomes at age 70, by gender. General linear models, p values and measure of effect size (partial eta squared values). Model 1: unadjusted; Model 2: age 11 IQ; Model 3: social class; Model 4: age 11 IQ + social class*

Predictor	Model	Age 70 IQ		g factor		processing speed		Memory		NART		WTAR	
		<i>n</i> = 917	$\eta_p^2$	<i>n</i> = 898	$\eta_p^2$	<i>n</i> = 886	$\eta_p^2$	<i>n</i> = 891	$\eta_p^2$	<i>n</i> = 921	$\eta_p^2$	<i>n</i> = 921	$\eta_p^2$
units per day		<i>p</i>		<i>p</i>		<i>p</i>		<i>p</i>		<i>p</i>		<i>p</i>	
Total alcohol													
<i>Women</i>													
	1	<b>.027<sup>a</sup></b>	.010	<b>.002<sup>a</sup></b>	.021	<b>.006<sup>a</sup></b>	.017	<b>&lt;.001<sup>a</sup></b>	.034	<b>&lt;.001<sup>a</sup></b>	.050	<b>&lt;.001<sup>a</sup></b>	.042
	2	.523	.001	.156	.005	.093	.007	<b>.024<sup>a</sup></b>	.012	<b>.006<sup>a</sup></b>	.017	<b>.007<sup>a</sup></b>	.016
	3	.399	.002	.054	.008	.062	.008	<b>.002<sup>a</sup></b>	.021	<b>.000<sup>a</sup></b>	.028	<b>.003<sup>a</sup></b>	.019
	4	.228	.003	.123	.006	.170	.005	<b>.043<sup>a</sup></b>	.010	<b>.048<sup>a</sup></b>	.009	.084	.007
<i>Men</i>													
	1	<b>&lt;.001<sup>a</sup></b>	.032	<b>.001<sup>a</sup></b>	.028	<b>.028<sup>a</sup></b>	.011	<b>&lt;.001<sup>a</sup></b>	.035	<b>.002<sup>a</sup></b>	.021	<b>&lt;.001<sup>a</sup></b>	.030
	2	<b>.020<sup>a</sup></b>	.013	<b>.030<sup>a</sup></b>	.011	.292	.003	<b>.010<sup>a</sup></b>	.016	.138	.005	<b>.020<sup>a</sup></b>	.013
	3	<b>.015<sup>a</sup></b>	.014	<b>.040<sup>a</sup></b>	.010	.360	.002	<b>.006<sup>a</sup></b>	.018	.178	.004	<b>.037<sup>a</sup></b>	.010
	4	.062	.009	.370	.002	.648	.001	<b>.030<sup>a</sup></b>	.012	.537	.001	.125	.006
Total wine													
<i>Men</i>													
	1	<b>&lt;.001<sup>a</sup></b>	.034	<b>&lt;.001<sup>a</sup></b>	.032	.088	.007	<b>&lt;.001<sup>a</sup></b>	.031	<b>&lt;.001<sup>a</sup></b>	.060	<b>&lt;.001<sup>a</sup></b>	.047
	2	<b>.035<sup>a</sup></b>	.011	<b>.022<sup>a</sup></b>	.013	.771	.000	<b>.037<sup>a</sup></b>	.011	<b>&lt;.001<sup>a</sup></b>	.042	<b>.001<sup>a</sup></b>	.028
	3	.101	.006	.188	.004	.616	.001	.076	.008	<b>.008<sup>a</sup></b>	.016	<b>.032<sup>a</sup></b>	.011
	4	.208	.004	.345	.002	.349	.002	.212	.004	<b>.004<sup>a</sup></b>	.020	<b>.031<sup>a</sup></b>	.011

*Note.*

<sup>a</sup> denotes a positive model correlation coefficient; <sup>b</sup> denotes a negative model correlation coefficient

Running head: ALCOHOL INTAKE AND COGNITIVE ABILITIES

Supplementary table 1

*In women, the relationship between alcohol (units/day) and cognitive outcomes at age 70. General linear models, p values and associated partial eta squared values. Model 1: unadjusted; Model 2: age 11 IQ; Model 3: social class; Model 4: age 11 IQ + social class*

Predictor	Model	Age 70 IQ		g factor		processing speed		Memory		NART		WTAR	
		<i>n</i> = 917	$\eta_p^2$	<i>n</i> = 898	$\eta_p^2$	<i>n</i> = 886	$\eta_p^2$	<i>n</i> = 891	$\eta_p^2$	<i>n</i> = 921	$\eta_p^2$	<i>n</i> = 921	$\eta_p^2$
units per day		<i>p</i>		<i>p</i>		<i>P</i>		<i>P</i>		<i>p</i>		<i>p</i>	
Total alcohol	1	<b>.027<sup>a</sup></b>	.010	<b>.002<sup>a</sup></b>	.021	<b>.006<sup>a</sup></b>	.017	<b>&lt;.001<sup>a</sup></b>	.034	<b>&lt;.001<sup>a</sup></b>	.050	<b>&lt;.001<sup>a</sup></b>	.042
	2	.523	.001	.156	.005	.093	.007	<b>.024<sup>a</sup></b>	.012	<b>.006<sup>a</sup></b>	.017	<b>.007<sup>a</sup></b>	.016
	3	.399	.002	.054	.008	.062	.008	<b>.002<sup>a</sup></b>	.021	<b>.000<sup>a</sup></b>	.028	<b>.003<sup>a</sup></b>	.019
	4	.228	.003	.123	.006	.170	.005	<b>.043<sup>a</sup></b>	.010	<b>.048<sup>a</sup></b>	.009	.084	.007
Wine	1	<b>.036<sup>a</sup></b>	.009	<b>.002<sup>a</sup></b>	.021	<b>.019<sup>a</sup></b>	.012	<b>&lt;.001<sup>a</sup></b>	.032	<b>&lt;.001<sup>a</sup></b>	.044	<b>&lt;.001<sup>a</sup></b>	.039
	2	.419	.001	.108	.006	.160	.005	<b>.024<sup>a</sup></b>	.012	<b>.019<sup>a</sup></b>	.012	<b>.010<sup>a</sup></b>	.015
	3	.445	.001	.063	.008	.161	.004	<b>.004<sup>a</sup></b>	.019	<b>.001<sup>a</sup></b>	.022	<b>.005<sup>a</sup></b>	.017
	4	.188	.004	.289	.003	.295	.003	<b>.043<sup>a</sup></b>	.010	.121	.006	.116	.006
Spirits	1	.786	.000	.659	.000	.254	.003	.304	.002	.100	.006	.280	.002
	2	.749	.000	.958	.000	.393	.002	.439	.001	<b>.041<sup>a</sup></b>	.009	.212	.003
	3	.888	.000	.579	.001	.194	.004	.321	.002	.062	.008	.235	.003
	4	.869	.000	.850	.000	.305	.003	.468	.001	<b>.028<sup>a</sup></b>	.011	.198	.004
Sherry/port	1	.300	.002	.201	.003	.088	.006	.611	.001	.079	.007	.194	.004
	2	.538	.001	.737	.000	.616	.001	.450	.001	.570	.001	.814	.000
	3	.772	.000	.578	.001	.242	.003	.991	.000	.310	.002	.601	.001
	4	.298	.003	.452	.001	.858	.000	.395	.002	.954	.000	.440	.001
Beer	1	.607	.001	.321	.002	.148	.005	.257	.003	.376	.002	.305	.002
(reg + dark)	2	.704	.000	.806	.000	.500	.001	.608	.001	.423	.001	.269	.003
	3	.931	.000	.541	.001	.272	.003	.378	.002	.584	.001	.494	.001
	4	.938	.000	.599	.001	.661	.000	.668	.000	.605	.001	.408	.002

*Note.*

<sup>a</sup> denotes a positive model correlation coefficient; <sup>b</sup> denotes a negative model correlation coefficient

Running head: ALCOHOL INTAKE AND COGNITIVE ABILITIES

Supplementary table 2

*In men, the relationship between alcohol (units/day) and cognitive outcomes at age 70. General linear models, p values and associated partial eta squared values. Model 1: unadjusted; Model 2: age 11 IQ; Model 3: social class; Model 4: age 11 IQ + social class*

Predictor	Model	Age 70 IQ		g factor		processing speed		Memory		NART		WTAR	
		<i>n</i> = 917	$\eta_p^2$	<i>n</i> = 898	$\eta_p^2$	<i>n</i> = 886	$\eta_p^2$	<i>n</i> = 891	$\eta_p^2$	<i>n</i> = 921	$\eta_p^2$	<i>n</i> = 921	$\eta_p^2$
		<i>P</i>		<i>p</i>		<i>P</i>		<i>p</i>		<i>P</i>		<i>p</i>	
Total alcohol	1	<b>&lt;.001<sup>a</sup></b>	.032	<b>.001<sup>a</sup></b>	.028	<b>.028<sup>a</sup></b>	.011	<b>&lt;.001<sup>a</sup></b>	.035	<b>.002<sup>a</sup></b>	.021	<b>&lt;.001<sup>a</sup></b>	.030
	2	<b>.020<sup>a</sup></b>	.013	<b>.030<sup>a</sup></b>	.011	.292	.003	<b>.010<sup>a</sup></b>	.016	.138	.005	<b>.020<sup>a</sup></b>	.013
	3	<b>.015<sup>a</sup></b>	.014	<b>.040<sup>a</sup></b>	.010	.360	.002	<b>.006<sup>a</sup></b>	.018	.178	.004	<b>.037<sup>a</sup></b>	.010
	4	.062	.009	.370	.002	.648	.001	<b>.030<sup>a</sup></b>	.012	.537	.001	.125	.006
Wine	1	<b>&lt;.001<sup>a</sup></b>	.034	<b>&lt;.001<sup>a</sup></b>	.032	.088	.007	<b>&lt;.001<sup>a</sup></b>	.031	<b>&lt;.001<sup>a</sup></b>	.060	<b>&lt;.001<sup>a</sup></b>	.047
	2	<b>.035<sup>a</sup></b>	.011	<b>.022<sup>a</sup></b>	.013	.771	.000	<b>.037<sup>a</sup></b>	.011	<b>&lt;.001<sup>a</sup></b>	.042	<b>.001<sup>a</sup></b>	.028
	3	.101	.006	.188	.004	.616	.001	.076	.008	<b>.008<sup>a</sup></b>	.016	<b>.032<sup>a</sup></b>	.011
	4	.208	.004	.345	.002	.349	.002	.212	.004	<b>.004<sup>a</sup></b>	.020	<b>.031<sup>a</sup></b>	.011
Spirits	1	.179	.000	.103	.006	.065	.008	<b>.002<sup>a</sup></b>	.021	.827	.000	.154	.005
	2	.482	.001	.307	.003	.175	.004	<b>.006<sup>a</sup></b>	.018	.095	.038	.573	.001
	3	.399	.002	.190	.004	.096	.007	<b>.004<sup>a</sup></b>	.019	.297	.003	.477	.001
	4	.582	.001	.290	.003	.148	.005	<b>.004<sup>a</sup></b>	.021	<b>.039<sup>b</sup></b>	.010	.855	.000
Sherry/port	1	<b>.013<sup>a</sup></b>	.014	<b>.019<sup>a</sup></b>	.013	.121	.006	<b>.023<sup>a</sup></b>	.012	<b>.004<sup>a</sup></b>	.019	<b>.003<sup>a</sup></b>	.020
	2	.057	.009	.065	.008	.261	.003	.081	.007	<b>.012<sup>a</sup></b>	.015	<b>.010<sup>a</sup></b>	.016
	3	<b>.025<sup>a</sup></b>	.012	<b>.034<sup>a</sup></b>	.011	.210	.004	<b>.033<sup>a</sup></b>	.011	<b>.005<sup>a</sup></b>	.018	<b>.004<sup>a</sup></b>	.019
	4	.062	.009	.071	.008	.311	.003	.073	.008	<b>.009<sup>a</sup></b>	.017	<b>.008<sup>a</sup></b>	.017
Beer (reg + dark)	1	.680	.000	.920	.000	.821	.000	.675	.000	<b>.032<sup>b</sup></b>	.011	.172	.004
	2	.469	.001	.837	.000	.550	.001	.714	.000	<b>.007<sup>b</sup></b>	.003	.098	.006
	3	.101	.006	.261	.003	.233	.003	.755	.000	.269	.018	.866	.000
	4	.184	.004	.327	.002	.233	.004	.943	.000	<b>.041<sup>b</sup></b>	.010	.356	.002

*Note.*

<sup>a</sup> denotes a positive model correlation coefficient; <sup>b</sup> denotes a negative model correlation coefficient