



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A systematic review of interventions addressing limited health literacy to improve asthma self-management

Citation for published version:

on behalf of the NIHR Global Health Research Unit on Respiratory Health (RESPIRE) collaborations, Salim, H, Ramdzan, K, McClatchey, K, Shariff Ghazali, S, Lee, PY, Young, I & Pinnock, H 2020, 'A systematic review of interventions addressing limited health literacy to improve asthma self-management', *Journal of Global Health*, vol. 10, no. 1, pp. 010427. <https://doi.org/10.7189/jogh.10.010428>

Digital Object Identifier (DOI):

[10.7189/jogh.10.010428](https://doi.org/10.7189/jogh.10.010428)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Global Health

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



A systematic review of interventions addressing limited health literacy to improve asthma self-management

Hani Salim^{1,2}, Siti Nurkamilla Ramdzan^{1,3}, Kirstie McClatchey⁴, Sazlina Shariff Ghazali², Lee Ping Yein², Ingrid Young⁵, Hilary Pinnock¹
and on behalf of the NIHR Global Health Research Unit on Respiratory Health (RESPIRE) collaborations.

¹*NIHR Global Health Research Unit on Respiratory Health (RESPIRE), Usher Institute, The University of Edinburgh*

²*Department of Family Medicine, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, Malaysia*

³*Department of Primary Care Medicine, Faculty of Medicine, University of Malaya, Malaysia*

⁴*Asthma UK Centre for Applied Research, Usher Institute, The University of Edinburgh.*

⁵*Centre for Biomedicine, Self and Society, Usher Institute, The University of Edinburgh.*

Address for correspondence:

Professor Hilary Pinnock

Professor of Primary Care Respiratory Medicine,

NIHR Global Health Research Unit on Respiratory Health (RESPIRE),

Usher Institute,

The University of Edinburgh,

Doorway 3, Medical School, Teviot Place,

Edinburgh EH8 9AG

Tel: 0131 650 9474

E-mail: hilary.pinnock@ed.ac.uk

Keywords: Asthma, Supported self-management, Health literacy, Systematic review, Low- and middle-income countries

ABSTRACT

Background: Supported asthma self-management improves health outcomes. However, people with limited health literacy, especially in lower-middle-income countries (LMICs), may need tailored interventions to enable them to realise the benefits. We aimed to assess the clinical effectiveness of asthma self-management interventions targeted at people with limited health literacy and to identify strategies associated with effective programmes.

Methods: Following Cochrane methodology, we searched ten databases (January 1990 - June 2018; updated October 2019), without language restriction. We included controlled experimental studies whose interventions targeted health literacy to improve asthma self-management. Selection of papers, extraction of data and quality assessment were done independently by two reviewers. The primary outcomes were clinical (asthma control) and implementation (adoption/adherence to intervention). Analysis was narrative.

Findings: We screened 4318 titles and abstracts, reviewed 52 full-texts and included five trials. One trial was conducted in a LMIC. Risk of bias was low in one trial and high in the other four studies. Clinical outcomes were reported in two trials, both at high risk of bias: one of which reported a reduction in unscheduled care (number of visits in 6-month (SD); Intervention:0.9 (1.2) vs Control:1.8 (2.4), $p=0.001$); the other showed no effect. None reported uptake or adherence to the intervention. Behavioural change strategies typically focused on improving an individual's psychological and physical capacity to enact behaviour (e.g. targeting asthma-related knowledge or comprehension). Only two interventions also targeted motivation; none sought to improve opportunity. Less than half of the interventions used specific self-management strategies (e.g. written asthma action plan) with tailoring to limited health literacy status. Different approaches (e.g. video-based and pictorial action plans) were used to provide education.

Conclusion: The paucity of studies and diversity of the interventions to support people with limited health literacy to self-manage their asthma meant that the impact on health outcomes remains unclear. Given the proportion of the global population who have limited health literacy skills, this is a research priority.

PROSPERO registration: CRD 42018118974

INTRODUCTION

Asthma self-management support, including written action plans and regular reviews by healthcare professionals, improves health outcomes [1-4]. Systematic reviews and guidelines highlight that cultural or age-related tailoring enables the successful implementation of supported self-management, although rarely specify tailoring for people with limited health literacy. This is a significant oversight, as health literacy is a problem globally [5], and a particular challenge in low and middle-income countries (LMICs). There is thus a need to address the challenges of providing support for people with limited health literacy [6,7].

A review of health literacy definitions by Sørensen et al. (2012), describes health literacy as people's knowledge, motivation and competence to assess, understand, appraise and apply health information to make decisions on healthcare, disease prevention and health promotion throughout the life course (see table 1) [8]. These skills are essential for individuals to respond to the demands of managing a variable condition such as asthma, including adherence to medication, adjusting treatment and/or deciding to seek advice in the event of deterioration. Health literacy is not linearly related to health outcomes but influences three aspects of healthcare behaviour: access and utilisation of health services, patient-provider interactions and self-management [9].

Two previous systematic reviews have looked at self-management interventions for people with limited health literacy in long-term health conditions [10,11]. One review included 38 studies, but only 22 were randomised trials, and none addressed self-management interventions in asthma [11]. The other defined the target population as people from low socio-economic groups, assuming that these populations had limited self-literacy [10]. Neither, therefore, specifically addressed

supported management for people with limited health literacy in asthma. We this aimed to systematically search and synthesise the trial evidence for asthma self-management interventions targeted at people with limited health literacy, in order to assess their clinical effectiveness and to identify the behaviour change strategies that were associated with effective programmes [12].

METHODS

This review is registered with the PROSPERO database (registration number: CRD 42018118974). Details of the systematic review protocol have been published [12] with salient points described here. We followed the procedures described in the Cochrane Handbook for Systematic Reviews of Interventions [13].

Deviations from the published protocol.

To be inclusive of data from LMICs, we intended to search the African Index Medicus, Africa Portal Digital Library; Index Medicus for the Southeast Asia Region; IndMed; Latin American and Caribbean Health Science Literature Database (LILACS). However, we decided to omit these after a scoping exercise revealed a lack of controlled trials in these databases, and we considered it was very unlikely that any publications would fulfil our inclusion criteria.

We intended to use the Grading of Recommendations Assessment Development and Evaluation (GRADE) to assess the weight of evidence of the reported outcomes from the included studies [14]. However, there was too much missing information to use GRADE. We have, therefore not presented the GRADE assessment in the paper (see Supplementary file 1 for further explanation).

Search strategy

We searched 10 electronic databases (see Table 2). The search strategy used medical subject headings (MeSH) and text words related to health literacy, asthma, self-management and controlled trial. The initial search (January 1990 to June 2018) was updated in October 2019. We conducted forward citation on included studies and contacted experts in the field to identify related trials. We did not perform manual searches as no journal(s) emerged as having a particular interest in this topic. There was no language restriction, though we did not find any non-English publications. We searched the databases using PICOS criteria (See Table 2). We used the definitions in Table 1 to confirm eligibility.

Study selection and data extraction

After training and quality control, two authors (HS and SNR) independently screened the de-duplicated titles and abstracts. We obtained the full text of potentially relevant studies, and both reviewers independently assessed for eligibility. Disagreements or uncertainties at any stage were resolved by discussion within the team (HP, IY, SGS or PYL).

Studies which had multiple publications (e.g. a protocol, trial findings, process evaluations, qualitative studies, translations) were treated as one study, and reference made to the different publications.

We piloted a data extraction form adapted from the Effective Practice and Organisation of Care (EPOC) recommendations for describing interventions [15] and the Template for Intervention Description and Replication (TIDieR) checklist [16]. Two reviewers (HS and SNR) independently extracted data. We contacted authors for any information which was not found within the included paper(s).

Quality assessment

We used the Cochrane Risk of Bias tool [13], and the guidance from the EPOC group [15], to assess selection, performance, detection, attrition, reporting and other potential sources of bias [13]. The risk of bias for each domain was classified as 'low', 'high' or 'unclear' based on the information available [13]. We generated 'risk of bias' summary graphs and figures using Review Manager 5.3 [17].

Outcomes

Outcomes are described in Table 2. We were primarily interested in health outcomes (e.g. asthma control; acute attacks) and implementation outcomes (e.g. adoption of intervention). Secondary outcomes included intermediate self-management measures (e.g. knowledge improvement), health literacy outcomes and impact indicators (e.g. cost-effectiveness).

Data synthesis.

We conducted two analyses to answer the two objectives of our systematic review. First, we considered the effectiveness of asthma self-management interventions which addressed health literacy needs compared with the control group. From scoping work, we anticipated that the studies included in this review would vary substantially in design, target populations, outcomes measured and duration of follow-up precluding meta-analysis. We, therefore, conducted a narrative synthesis of the data.

Second, we described and characterised the included interventions using the Behaviour Change Wheel (BCW) framework (figure 1), which provides a systematic way to describe and characterise the techniques used in the interventions in this review [18,19]. The BCW has three layers; its core components consist of the COM-B system (Capability, Opportunity and Motivation); interactions between these

components determine Behaviour [18]. Capability is the individual's psychological and physical capacity to engage in the behaviour. It includes having the required knowledge and skills. Motivation is defined as processes that contribute towards both reflective and automatic mechanisms that activate or inhibit behaviour. Opportunity includes aspects of the physical and social environment that lie outside the individual that prompt or make behaviour possible. The second layer of the BCW describes the nine functions of interventions that are designed to change behaviour. The intervention functions are; education, persuasion, coercion, training, enablement, modelling, environmental restructuring and restrictions. The third layer of the BCW identifies seven types of policies (e.g. legislation, fiscal measures, etc.) that can be applied to deliver these intervention functions [18].

It is proposed that specific intervention functions are likely to influence change in the specific target behaviour. This underpins a matrix (Figure 5), produced through a consensus exercise amongst behaviour change experts [18], that enables gaps in intervention functions required to impact on the three core components (capability, opportunity and motivation) that govern behaviour change [18].

We plotted the components of interventions in this review onto the matrix. In the mapping process, which was completed independently by two reviewers (HS and KM), we first identified the core components of behaviour that were targeted, and also the intervention functions used in each included study. Through a consensus approach (see supplementary file 2 for the exercise), we plotted our findings within the matrix (Figure 5).

RESULTS

The selection process is illustrated in the PRISMA diagram (Figure 2). From 3359 papers, we selected six papers describing five randomised control trials [20-24]

(the sixth paper described the development of the intervention[25]). The studies included a total of 731 participants in the intervention groups and 561 participants in the control groups [20-24].

Characteristics of included studies

The randomised control trials were conducted from 2011 to 2017; four studies were conducted in high-income countries [20-22,24] (three in the United States (US); one in Canada) and one in Turkey (a middle-income country) [23]. Table 3 summarises population characteristics (see supplementary file 3 for further details).

Participants characteristics: The three US studies included majority and minority populations [20,21,24]. Yin et al. (2017) included mainly Latin Americans (Hispanics); Apter et al. (2011) included mainly African-Americans, and the majority of the population in the study by Macy et al. (2011) was White American. The trial conducted in Canada by Poureslami et al. (2012) included participants from minority Chinese and Punjabi ethnic groups [22]. The study conducted in Turkey by Ozyigit et al. (2014) did not specify the ethnicity of the population [23]. Participants' asthma status was described as uncontrolled [20,23]; mild intermittent, persistent or moderate-severe asthma [24]; mild asthma [21]. One study did not describe the participants' level of asthma control [22].

Study setting: Two studies were conducted in primary care settings [20,23]. Three studies were conducted in secondary/tertiary care settings (specialist paediatric [24] or emergency department [21], university-based pulmonary medicine clinic [22]).

Geographical area and socioeconomic status: Four studies were described as set in an urban environment [20-22,24]; three described their population as of low socioeconomic status [20,21,24], the fourth had less than a third in the 'working-class

group' [22]. The non-urban study described the population as living in the most socio-economically under-developed province in the country [23].

Health literacy status of the population: Only three studies measured the level of health literacy of their participants. One study, which used the validated Newest Vital Sign (NVS), estimated that 70% of the study population had limited health literacy level [24]. Two other studies measured the health literacy level of the study population using the Short Test of Functional Health Literacy in Adults (sTOFHLA) (stating that the mean reading comprehension score was 'adequate'[20]) or the Rapid Estimate of Adult Literacy in Medicine (REALM) (reporting that 'two-thirds of the study population had an 'adequate' level of health literacy') [21]. Two studies included 'immigrants' [22] or 'illiterates' [23] as their study population.

Intervention characteristics: Table 3 summarises the interventions (see supplementary file 4 for further details). All studies had one intervention and one control group [20,21,23,24] except Poureslami [22], which had three intervention groups [22].

All interventions included education delivered through various methods; one used a face-to-face personalised problem-solving approach [20], two used video-based education [21,22], and two used education with pictorial asthma action plans [23,24] although only one of these explicitly tailored its action plan to low-literacy level [24]. Three interventions were delivered by research assistants [20,23,24] and one by a respiratory physician [23]. Four studies specified the language used to deliver the intervention; English or Spanish [20,24], 'native language' [23], Punjabi or Mandarin [22]. Only two studies specified the duration of the intervention: 20-minute video [21] or four 30-minute problem-solving sessions [20]. Length of follow-up ranged from five

weeks to a year [20-23]. One study assessed the immediate understanding of a pictorial asthma action plan [24] rather than longer-term outcomes

Quality assessment of the included studies

Only one study was at an overall low risk of bias [24] (Figure 3). The high risk of bias in the other four studies was typically due to no description of random sequence generation or blinding of outcome assessment. Other biases included no specified sample size [20,23] and use of non-validated tools to measure outcomes [22].

Quality of description and replication

All the studies described the rationale for the essential elements included in the intervention, but none were explicitly guided by a theoretical framework. Three studies lacked descriptions of how the intervention was provided:[21,22,24] for example; one report was unclear whether the video-based intervention was provided individually or in a group [21]. Brief descriptions of the interventions are in Table 3; see Supplementary file 5 for detailed information.

Effectiveness of interventions on primary and secondary outcomes

The study at low risk of bias did not report any of our primary outcomes. [24]. Two studies (at high risk of bias) reported health outcomes [20,23], one of which reported a positive outcome for unscheduled care [23]. None of the five studies reported on implementation outcomes (such as uptake/completion of the intervention). Findings are detailed in Table 3 and the key points described below.

1) Primary (Health outcomes): Asthma control and unscheduled care

- *Impact on asthma control.*

Two studies at high risk of bias measured asthma control using validated questionnaires (see table 3) [20,23]. Neither of the interventions had an effect on asthma control.

- *Impact on unscheduled care*

Three studies at high risk of bias measured the impact of the intervention on unscheduled care [20,21,23]. One study reduced emergency visits in the intervention group compared to control [23]. One study only reported within-group changes, stating that there was no between-group difference though no statistical comparison was provided [20].

2) Secondary outcomes:

- *Impact on knowledge*

The low risk of bias study reported a positive outcome on knowledge [24] while the other studies reported no effect [21] (see Table 3).

- *Impact on correct inhaler use*

A high risk of bias study did not provide sufficient details to gauge the impact of the intervention on correct inhaler use [22].

- *Impact on other practical self-management measures*

Other measures included in this review are perceived ease of action plan use, understanding of low-literacy AAP (low risk of bias) [24], perceived sense of asthma control [21], understanding of physician instruction [22] and adherence [20] (high risk of bias). All studies either reported no effect [20,24] or reported insufficient details to gauge effectiveness [21,22]. (see Table 3)

Identification of intervention components in relation to the behaviour change

Limited reporting and the lack of effectiveness in the included studies meant that it was not possible to map the components of BCW to effectiveness. The core components of behaviour and the intervention functions used in the included studies based on reported information are provided in Figure 4. Reports were sometimes

limited: for example, one intervention described providing 'patient skills' in its education video [21], with no further description of what was taught.

In terms of the use of the BCW core components of behaviour (COM-B), three studies only addressed 'capability' in their interventions [21,23,24]. Two studies, at high risk of bias, addressed a combination of capability and motivation [20,22].

In Figure 5, we used the published matrix [18] to plot the included studies according to the core components of behaviour change and intervention function. The low risk of bias study used only one intervention function (enablement) [24]. For the high risk of bias studies; two used three intervention functions [22,23], and two studies used two intervention functions [20,21].

Michie et al. (2011) suggest that the core components of behaviour can be linked to the interventions in more than one way. As an example, the use of a pictorial action plan by Ozyigit et al. [23] is 'education' as it increases the capability to understand asthma self-management. A pictorial action plan is also a form of 'enablement' as it reduces barriers (e.g. lack of knowledge/cognitive skills) to self-management of asthma in the event of deterioration. Most of the interventions concentrated on capability components of the behaviour model, and these interventions used functions such as education, training, persuasion and enabling interventions (action plans) to produce behaviour change.

DISCUSSION

Summary of findings

This review reports the synthesised findings from five randomised control trials. Four studies, at high risk of bias, concluded that their interventions were ineffective; the only study at low risk of bias did not report on health outcomes. The paucity of studies, limitations in study design and diversity of the interventions meant we are

unable to draw conclusions about overall effectiveness on any of our outcomes of interest.

Most studies [20,21,23,24] included in this review did not describe any theoretical framework underpinning the intervention development, although one conducted prior exploratory work to understand the impact of health literacy in the targeted population [22]. Education, training and enablement are the intervention functions used in these interventions, and the content and the method of delivery varied, including video-based [21,22] and a pictorial action plan [24]. All the interventions used components of behaviour change primarily directed at individuals' (physical and psychosocial) capabilities; two addressed self-motivation; none targeted opportunity).

Interpretation of the findings and comparison with previous findings.

The use of theory in developing a complex intervention

Health literacy is a complex concept, and as the concept has evolved, a number of definitions have been suggested by researchers and organisations [26]. Tools to measure the health literacy status of populations arise from these definitions and are similarly diverse, making studies in this area heterogeneous and more difficult to interpret. The use of health literacy as a dichotomous variable in many of these tools remained an inherent flaw, especially when health literacy is a spectrum which interacts in complex ways with the environment and socio-cultural factors. In this review, we used a systematically-defined definition by Sørensen et al. (2012) [8] which enabled us to include studies that employed other aspects of health literacy in their intervention, e.g. functional health literacy skills [23].

Only one study [22] in our review developed its intervention based on a recognised definition of health literacy (by Nutbeam et al. (2000) [27]). Pourslami et al. (2011),

aligned their asthma educational material with the definition of 'critical health literacy' which requires sufficient cognitive skills in order to understand, analyse and independently act on adversities in life to care for asthma [25]. In their prior qualitative work, language was found to be a barrier in understanding health information [28]. Thus, in the trial, the education material was delivered using the spoken languages of the participants and was designed to help participants learn and understand beliefs about asthma from the ethno-cultural point of view [22].

Four other studies [20,21,23,24] did not use specific health literacy definitions, although they used interventional designs which explicitly aimed to improve health literacy (e.g. pictograms) as defined by our operational definitions (see table 1). None of the studies described any theoretical framework that informed the development of their intervention, implying that the authors had not systematically considered the inter-related barriers among people who struggled with limited health literacy and identified factors which could overcome these barriers.

The Medical Research Council's framework for developing and evaluating complex interventions clearly outlines the importance of defining a theoretical concept as well as undertaking qualitative exploration [29,30]. A theoretical framework provides a roadmap for the programme of work. In its absence, it is challenging to visualise how the intervention operates to bring about change [31,32]. Interpreting effectiveness is difficult if it is not clear what works and why [29,30].

'Behaviour Change Wheel': using a theoretical approach to understand the process of change and to evaluate interventions.

The BCW provides an understanding of what needs to change and how to change it. Targeted behaviour is more likely to change if the specific intervention function is employed. As an example, education using video presentations improved

inhaler techniques across the three experimental groups in one study (though the lack of comparison with the control group means it is not possible to gauge effectiveness) [22].

A multi-component approach to change behaviour

Previous studies have concluded that the use of more than one strategy in an intervention increased the likelihood of it being effective [10,11]. A review reported that interventions which employed three to four self-management skills were more effective than those using fewer [10]. The five self-management skills considered in that review were problem-solving, taking action, decision making, partnership and resource utilisation [10]. Another review concluded that mixed-strategy interventions focusing on self-management reduced emergency visits, hospitalisations and disease severity in people with long term conditions [11]. Three of the quasi-experimental studies in this review included people with asthma [6,7,33], one of which reduced emergency department visits [6]. Multiple-components in a complex intervention incurs costs in terms of development and manpower [34,35]. However, designing a complex intervention without understanding the behaviour which it aims to change can lead to failure, which is also wasteful. A much criticised example of this is the ineffective UK public health campaign which focused on motivating responsible drinking but failed to reduce opportunity by addressing price and availability [36]. The other point to bring into this section is that the empty green cells of the matrix (see figure 5), are gaps that a future multi-component intervention could usefully address.

Strengths and limitations of this study

We followed Cochrane methodology to search systematically for trials of interventions addressing health literacy in the specific context of asthma self-management. All the stages in the review were duplicated, including the selection of

papers, risk of bias assessment and data extraction. Our decision not to search some LMIC-focused databases may mean we missed some relevant studies, though our initial scoping exercise in discussion with a medical librarian suggested this was unlikely. All the included studies were RCTs though we would have accepted other designs of controlled trials. We defined our outcomes with care, ensuring we looked for standardised measures of asthma symptom control and risk of attacks [37] and we included trials based on an evidence-based definition of limited health literacy [8,38]. We used the BCW, a validated framework to describe each of the intervention functions, and interpretation of the findings was conducted by a multidisciplinary team to ensure accuracy. The primary studies have small sample size and diverse in populations which makes it challenging to draw a conclusion from the reported results. Four studies did not use health literacy definitions or framework to map its interventional design. Unfortunately, less than half of interventions in this review reported on asthma control [20,23] or unscheduled care, [20,21,23] limiting the conclusions we could draw. For example, there were insufficient data to present our findings graphically (e.g. in a Harvest plot [12]) or to use the GRADE [14] approach to assess the quality of evidence. There was limited description of some of the interventions. We could not, for example be certain whether the 'patient skills' described as being included in educational videos in one trial, [21] covered behaviour change techniques such as demonstration of behaviour and/or instruction how to perform the task.

CONCLUSION

Despite the global importance of the problem, effective interventions addressing health literacy to improve asthma self-management have yet to be developed and evaluated. The studies that we found in this review were diverse,

generally at high risk of bias, poorly reported, lacked theoretical underpinning and were ineffective. In designing future interventions, researchers need to be able to identify and understand the factors, including social determinants of health that mediate behaviour change in different contexts (LMICs as well as high-income countries) [34,35]. Tailored asthma self-management interventions for people with limited health literacy should consider a multifaceted approach, including strategies that can be adapted to local needs [35,39], building on theoretical underpinning and careful planning especially in the development stage to optimise effectiveness and sustainability of the intervention.

Acknowledgements

We acknowledged the RESPIRE collaboration, which includes Aziz Sheikh, Harry Campbell, Steve Cunningham, Neneh Rowa-Dewar, Harish Nair, Aisha Holloway, Alimuddin Zumla, Kit Chan, Moira Whyte, Roberto Rabinovich, Rita Isaac, Ee Ming Khoo, Nik Sherina Hanafi, David Dockrell, Su May Liew, Norita Hussein, Tabish Hazir Ahmad Ihsan Bin Abu Bakar, Colin Simpson and Saturnino Luz.

We are grateful to Marshall Dozier, Academic Librarian at the University of Edinburgh, for her help in developing the search strategy.

Funding

HS and SNR are supported by PhD studentships from the NIHR Global Health Research Unit on Respiratory Health (RESPIRE). RESPIRE is funded by the National Institute of Health Research using Official Development Assistance (ODA) funding.

Disclaimer: The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. Neither the funder nor the sponsor (University of Edinburgh) contributed to protocol development.

Authorship contributions: HS, HP, IY, SSG, PYL contributed the concept and designing of this review. HS and SNR/KM conducted the literature search, screening, data extracting and quality assessment. HS and KM further analysed the data and completing the mapping of the BCW. HS and HP worked with the data synthesis with assistance from IY, SSG and PYL. All authors were involved in writing the manuscript and approved the final version.

Competing interest: The authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure_.pdf (available on request from the corresponding author) and declare no conflict of interest

Additional material: Supplementary documents are available online

REFERENCES

1. Pinnock H, Parke HL, Panagioti M, Daines L, Pearce G, Epiphaniou E, et al. Systematic meta-review of supported self-management for asthma: a healthcare perspective. *BMC Med.* 2017;15:64. Medline 28302126 doi: 10.1186/s12916-017-0823-7.
2. Pinnock H, Epiphaniou E, Pearce G, Parke H, Greenhalgh T, Sheikh A, et al. Implementing supported self-management for asthma: a systematic review and suggested hierarchy of evidence of implementation studies. *BMC Med.* 2015;13:127. Medline 26032941 doi: 10.1186/s12916-015-0361-0.
3. Pearce G, Parke HL, Pinnock H, Epiphaniou E, Bourne CL, Sheikh A, et al. The PRISMS taxonomy of self-management support: derivation of a novel taxonomy and initial testing of its utility. *J Health Serv Res Policy.* 2016;21:73-82. Medline 26377727 doi: 10.1177/1355819615602725.
4. Corrigan, J.M., Greiner, A.C., Adams, K. *The 1st Annual Crossing the Quality Chasm Summit: A Focus on Communities: Report of a Summit.* Washington: National Academies Press; 2004
5. Okan O, Bauer U, Levin - Zamir D, Pinheiro P, Sørensen K. *International Handbook of Health Literacy.* Bristol:Policy Press; 2019.
6. Robinson LD, Calmes DP, Bazargan Mn. The Impact of Literacy Enhancement on Asthma-Related Outcomes among Underserved Children. *J Natl Med Assoc.* 2008;100:892-6. Medline 18717138 doi:10.1016/s0027-9684(15)31401-2
7. Paasche-Orlow MK, Riekert KA, Bilderback A, Chanmugam A, Hill P, Rand CS, et al. Tailored education may reduce health literacy disparities in asthma self-management. *Am J Respir Crit Care Med.* 2005;172:980-6. Medline 16081544 doi: 10.1164/rccm.200409-1291OC
8. Sørensen K, Van den Broucke S, Fullam J, Doyle G, Pelikan J, Slonska Z, et al. Health literacy and public health: a systematic review and integration of definitions and models. *BMC Public Heal.* 2012;12:80. Medline 22276600 doi: 10.1186/1471-2458-12-80.
9. Paasche-Orlow MK, Wolf MS. The causal pathways linking health literacy to health outcomes. *Am J Health Behav.* 2007;31:S19-S26. Medline 17931132 doi: 10.5555/ajhb.2007.31.suppl.S19
10. Schaffler J, Leung K, Tremblay S, Merdsoy L, Belzile E, Lambrou A, et al. The Effectiveness of Self-Management Interventions for Individuals with Low Health Literacy and/or Low Income: A Descriptive Systematic Review. *J Gen Intern Med.* 2018;33:510-23. Medline 29427178 doi: 10.1007/s11606-017-4265-x
11. Sheridan SL, Halpern DJ, Viera AJ, Berkman ND, Donahue KE, Crotty K. Interventions for individuals with low health literacy: a systematic review. *J Health Commun.* 2011;16 Suppl 3:30-54. Medline 21951242 doi: 10.1080/10810730.2011.604391.
12. Salim H, Young I, Shariff Ghazali S, Lee PY, Ramdzan SN, Pinnock H. Protocol for a systematic review of interventions addressing health literacy to improve

- asthma self-management. *npj Prim Care Respir Med*. 2019;29:18. Medline 31068584 doi: 10.1038/s41533-019-0125-y.
13. Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0* The Cochrane Collaboration; 2011. Available: www.handbook.cochrane.org. Accessed: 25 September 2017.
 14. Schünemann HJ, Oxman AD, Brozek J, Glasziou P, Jaeschke R, Vist GE, et al. Grading quality of evidence and strength of recommendations for diagnostic tests and strategies. *BMJ*. 2008;336:1106-10. Medline 18483053 doi: 10.1136/bmj.39500.677199.AE.
 15. Cochrane Effective Practice and Organisation of Care (EPOC). EPOC Resources for review authors. Available: <http://epoc.cochrane.org/epoc-specific-resources-review-authors>. Accessed: 10 October 2018
 16. Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ*. 2014;348. Medline 24609605 doi: 10.1136/bmj.g1687.
 17. Review Manager (RevMan) [Computer program]. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014. Available: <https://community.cochrane.org/help/tools-and-software/revman-5>. Accessed: 11 August 2018.
 18. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci*. 2011;6:42. Medline 21513547 doi: 10.1186/1748-5908-6-42.
 19. Michie S. *The behaviour change wheel : a guide to designing interventions*. London: Silverback Publishing; 2014.
 20. Apter AJ, Wang X, Bogen DK, Rand CS, McElligott S, Polsky D, et al. Problem solving to improve adherence and asthma outcomes in urban adults with moderate or severe asthma: A randomized controlled trial. *J Allergy Clin Immunol*. 2011;128:516-23.e5. Medline 21704360 doi: 10.1016/j.jaci.2011.05.010.
 21. Macy ML, Davis MM, Clark SJ, Stanley RM. Parental health literacy and asthma education delivery during a visit to a community-based pediatric emergency department: a pilot study. *Pediatr Emerg Care*. 2011;27:469-74. Medline 21629152 doi: 10.1097/PEC.0b013e31821c98a8.
 22. Poureslami I, Nimmon L, Doyle-Waters M, Rootman I, Schulzer M, Kuramoto L, et al. Effectiveness of educational interventions on asthma self-management in Punjabi and Chinese asthma patients: a randomized controlled trial. *J Asthma*. 2012;49:542-51. Medline 22715910 doi: 10.3109/02770903.2012.682125.
 23. Ozyigit LP, Ozcelik B, Ciloglu SO, Erkan F. The effectiveness of a pictorial asthma action plan for improving asthma control and the quality of life in illiterate women. *J Asthma*. 2014;51:423-8. Medline 24200510 doi: 10.3109/02770903.2013.863331

24. Yin HS, Gupta RS, Mendelsohn AL, Dreyer B, van Schaick L, Brown CR, et al. Use of a low-literacy written action plan to improve parent understanding of pediatric asthma management: A randomized controlled study. *J Asthma*. 2017;54:919-29. Medline 28045551 doi: 10.1080/02770903.2016.1277542.
25. Poureslami I, Nimmon L, Doyle-Waters M, Fitzgerald J. Using community-based participatory research (CBPR) with ethno-cultural groups as a tool to develop culturally and linguistically appropriate asthma educational material. *Divers Equal Health Care*. 2011; 8: 203–215.
26. Sørensen K, Van den Broucke S, Pelikan JM, Fullam J, Doyle G, Slonska Z, et al. Measuring health literacy in populations: illuminating the design and development process of the European Health Literacy Survey Questionnaire (HLS-EU-Q). *BMC Public Health*. 2013;13:948. Medline 24112855 doi: 10.1186/1471-2458-13-948
27. Nutbeam D. Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health Promot Int*. 2000;15:259-67 doi: 10.1093/heapro/15.3.259
28. Poureslami I, Rootman I, Doyle-Waters MM, Nimmon L, FitzGerald JM. Health Literacy, Language, and Ethnicity-Related Factors in Newcomer Asthma Patients to Canada: A Qualitative Study. *J Immigr Minor Health*. 2011;13:315-22. Medline 20938742 doi:10.1007/s10903-010-9405-x.
29. Campbell M, Fitzpatrick R, Haines A, Kinmonth AL, Sandercock P, Spiegelhalter D, et al. Framework for design and evaluation of complex interventions to improve health. *BMJ*. 2000;321:694-6. Medline 10987780 doi: 10.1136/bmj.321.7262.694
30. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ*. 2008;337. Medline 18824488 doi: 10.1136/bmj.a1655.
31. Steed L, Sohanpal R, James W-Y, Rivas C, Jumbe S, Chater A, et al. Equipping community pharmacy workers as agents for health behaviour change: developing and testing a theory-based smoking cessation intervention. *BMJ*. 2017;7:e015637. Medline 28801403 doi: 10.1136/bmjopen-2016-015637.
32. Band R, Bradbury K, Morton K, May C, Michie S, Mair FS, et al. Intervention planning for a digital intervention for self-management of hypertension: a theory-, evidence- and person-based approach. *Implement Sci*. 2017;12:25. Medline 28231840 doi: 10.1186/s13012-017-0553-4.
33. Sobel RM, Paasche-Orlow MK, Waite KR, Rittner SS, Wilson EA, Wolf MS. Asthma 1-2-3: a low literacy multimedia tool to educate African American adults about asthma. *J Community Health*. 2009;34:321-7. Medline 19353250 doi: 10.1007/s10900-009-9153-9.
34. DiLiberto DD, Staedke SG, Nankya F, Maiteki-Sebuguzi C, Taaka L, Nayiga S, et al. Behind the scenes of the PRIME intervention: designing a complex intervention to improve malaria care at public health centres in Uganda.

- Glob Health Action. 2015;8:29067. Medline 26498744 doi: 10.3402/gha.v8.29067
35. Hoddinott P, Britten J, Pill R. Why do interventions work in some places and not others: A breastfeeding support group trial. *Soc Sci Med*. 2010;70:769-78. Medline 20005617 doi:10.1016/j.socscimed.2009.10.067
 36. Room R. Disabling the public interest: alcohol strategies and policies for England. *Addiction*. 2004;99:1083-9. Medline 15317624 doi: 10.1111/j.1360-0443.2004.00803.x
 37. Reddel HK, Taylor DR, Bateman ED, Boulet L-P, Boushey HA, Busse WW, et al. An official American Thoracic Society/European Respiratory Society statement: asthma control and exacerbations: standardizing endpoints for clinical asthma trials and clinical practice. *Am J Respir Crit Care Med*. 2009;180:59-99. Medline 19535666 doi:10.1164/rccm.200801-060ST.
 38. United Nations E, Social C. Health Literacy and the Millennium Development Goals: United Nations Economic and Social Council (ECOSOC) Regional Meeting Background Paper (Abstracted). *J Health Commun*. 2010;15:211-23. Medline 20845205 doi: 10.1080/10810730.2010.499996.
 39. Okwaro FM, Chandler CIR, Hutchinson E, Nabirye C, Taaka L, Kayendeke M, et al. Challenging logics of complex intervention trials: Community perspectives of a health care improvement intervention in rural Uganda. *Soc Sci Med*. 2015;131:10-7. Medline 25748110 doi: 10.1016/j.socscimed.2015.02.032.
 40. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *Eur Respir J*. 1999;14:902-7. Medline 10573240 doi: 10.1034/j.1399-3003.1999.14d29.x
 41. Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, Murray JJ, Pendergraft TB. Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol*. 2004;113:59-65. Medline 14713908 doi: 10.1016/j.jaci.2003.09.008

List of figures and tables:

Figure 1	PRISMA flowchart
Figure 2	Risk of bias summary
Figure 3	The Behaviour Change Wheel (BCW) model
Figure 4	The use of COM-B components in the included studies.
Figure 5	Mapping of core components of behaviour and intervention functions used in the included studies
Table 1	Definition of terms
Table 2	PICOS table
Table 3	Summary of impact of clinical and process outcomes categorised by health literacy status of the population
Supplementary file 1	GRADE table
Supplementary file 2	BCW mapping exercise
Supplementary file 3	Overview of population characteristics of included studies
Supplementary file 4	Overview of intervention characteristics
Supplementary file 5	TIDieR Checklist