



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Generalized hypergeometric functions and intersection theory for Feynman integrals

Citation for published version:

Abreu, S, Britto, R, Duhr, C, Gardi, E & Matthew, J 2020, 'Generalized hypergeometric functions and intersection theory for Feynman integrals', Paper presented at 14th International Symposium on Radiative Corrections, Avignon, France, 9/09/19 - 13/09/19. <https://doi.org/10.22323/1.375.0067>

Digital Object Identifier (DOI):

[10.22323/1.375.0067](https://doi.org/10.22323/1.375.0067)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Generalized hypergeometric functions and intersection theory for Feynman integrals

Samuel Abreu

Center for Cosmology, Particle Physics and Phenomenology (CP3), Université Catholique de Louvain, 1348 Louvain-La-Neuve, Belgium

E-mail: samuel.abreu@uclouvain.be

Ruth Britto*

School of Mathematics, Trinity College, Dublin 2, Ireland ;

Hamilton Mathematics Institute, Trinity College, Dublin 2, Ireland ;

Institut de Physique Théorique, Université Paris Saclay, CEA, CNRS, F-91191 Gif-sur-Yvette cedex, France

E-mail: britto@maths.tcd.ie

Claude Duhr

Theoretical Physics Department, CERN, Geneva, Switzerland ;

E-mail: claude.duhr@cern.ch

Einan Gardi

Higgs Centre for Theoretical Physics, School of Physics and Astronomy, The University of Edinburgh, Edinburgh EH9 3FD, Scotland, UK

E-mail: einan.gardi@ed.ac.uk

James Matthew

Higgs Centre for Theoretical Physics, School of Physics and Astronomy, The University of Edinburgh, Edinburgh EH9 3FD, Scotland, UK

E-mail: james.matthew@ed.ac.uk

Feynman integrals that have been evaluated in dimensional regularization can be written in terms of generalized hypergeometric functions. It is well known that properties of these functions are revealed in the framework of intersection theory. We propose a new application of intersection theory to construct a coaction on generalized hypergeometric functions. When applied to dimensionally regularized Feynman integrals, this coaction reproduces the coaction on multiple polylogarithms order by order in the parameter of dimensional regularization.

14th International Symposium on Radiative Corrections (RADCOR2019)

9-13 September 2019

Palais des Papes, Avignon, France

*Speaker.

1. Introduction

The algebraic structure known as *coaction* has recently played an interesting role in computing Feynman diagrams and understanding their analytic structure. In dimensional regularization with $D = n - 2\epsilon$ dimensions where n is even, large classes of loop integrals including all one-loop integrals are expressible in terms of multiple polylogarithms (MPLs) when expanded in ϵ . MPLs are equipped with a coaction [1, 2, 3, 4], a mathematical operation that exposes properties of MPLs through a decomposition into simpler functions. In particular, since the coaction is naturally compatible with the actions of differential operators and taking discontinuities across branch cuts, it can be considered as a computational tool.


There is a natural mathematical coaction on Feynman integrals [5, 6, 7]. The coaction captures information about discontinuities, which for Feynman integrals can be expressed in terms of unitarity cuts and their generalizations. Motivated by these observations, it was conjectured that there exists a diagrammatic coaction corresponding precisely to the well known coaction on MPLs, for the class of Feynman graphs admitting an expansion in MPLs. This conjecture was expressed in [8, 9], which further proposed a general form for a coaction on integrals:

$$\Delta \left(\int_{\gamma} \omega \right) = \sum_{ij} c_{ij} \int_{\gamma} \omega_i \otimes \int_{\gamma_j} \omega. \quad (1.1)$$

Here the ω_i are generators of a suitably defined cohomology group of which ω is also an element, and the γ_j are likewise generators of a suitable homology group of which γ is also an element. It is straightforward to check that eq. (1.1) satisfies the algebraic properties of a coaction, such as coassociativity.

In the case of Feynman integrals, the cohomology group contains the usual integrands constructed from Feynman rules. In turn, the homology group contains the original contour in addition to contours encircling poles giving rise to the residues associated to generalized unitarity cuts. The coefficients c_{ij} remain to be determined. In the case of one-loop Feynman integrals, a basis has been found in which the c_{ij} are known precisely [8, 9].¹

Consider for example the following simple scalar 1-loop integral, evaluated in $4 - 2\epsilon$ dimensions, where the thick lines represent a propagator of mass m and a non-null external leg of momentum p , while the thin lines represent massless propagators and external legs carrying null momenta:



$$= \frac{1}{p^2} \left[\frac{\log \left(\frac{m^2}{m^2 - p^2} \right)}{\epsilon} + \text{Li}_2 \left(\frac{p^2}{m^2} \right) + \log^2 \left(1 - \frac{p^2}{m^2} \right) + \log(m^2) \log \left(1 - \frac{p^2}{m^2} \right) \right] + \mathcal{O}(\epsilon). \quad (1.2)$$

The coaction of the Laurent expansion on right-hand side can be evaluated directly, to arbitrary orders in ϵ . The terms written explicitly can be treated with well-known formulas for the coaction of polylogarithms, such as

$$\begin{aligned} \Delta(\log z) &= 1 \otimes \log z + \log z \otimes 1, \\ \Delta(\log^2 z) &= 1 \otimes \log^2 z + 2 \log z \otimes \log z + \log^2 z \otimes 1, \\ \Delta(\text{Li}_2(z)) &= 1 \otimes \text{Li}_2(z) + \text{Li}_2(z) \otimes 1 + \text{Li}_1(z) \otimes \log z. \end{aligned}$$

¹For extensions to two loops, see the contribution of James Matthew to these proceedings.

We compare the result of that coaction with the diagrammatic coaction on the left-hand side, which was claimed in [8, 9] to be

$$\Delta \left[\text{triangle diagram} \right] = \text{circle diagram} \otimes \left(\text{triangle diagram} + \frac{1}{2} \text{triangle diagram with dashed lines} \right) + \text{bubble diagram} \otimes \text{triangle diagram} . \tag{1.3}$$

The claim is that when each of these graphs is evaluated according to Feynman rules for scalar fields, with specific prescriptions for cut lines and dimensionality, then, order by order in ϵ , the result will agree with the coaction on MPLs.

We now imagine summing the full Laurent series. The example shown above has a closed-form expression in terms of Gauss’s hypergeometric function,

$$\text{triangle diagram} = \frac{e^{\gamma\epsilon} \Gamma(1 + \epsilon)}{\epsilon(1 - \epsilon)} (m^2)^{-1-\epsilon} {}_2F_1 \left(1, 1 + \epsilon; 2 - \epsilon; \frac{p^2}{m^2} \right), \tag{1.4}$$

and each of the integrals on the right-hand side of eq. (1.3) is likewise a ${}_2F_1$ function, or a simpler function arising from a special configuration of the first three arguments. Many other Feynman integrals in dimensional regularization are known to evaluate to hypergeometric-type functions whose Laurent expansions in ϵ are given by MPLs. We now conjecture a coaction on generalized hypergeometric functions of this type that is compatible with the coaction on Feynman integrals, as well as with the coaction on MPLs order by order in the ϵ -expansion.

There are various ways to define generalized hypergeometric functions. Here we define them as integrals and use the framework of twisted (co)homology [10]. We propose a coaction of the form (1.1) in which the coefficients c_{ij} can be derived from so-called intersection numbers. In this article, we state our conjecture more explicitly and illustrate it with a couple of examples.

This contribution to the proceedings of RADCOR2019 is based on work that has since appeared in [11], which contains many more details and examples, and a fuller discussion.

2. Generalized hypergeometric functions and intersection numbers

We consider integrals of the form $\int_\gamma \omega$, where ω is a cohomology class represented by the differential n -form

$$\omega = d\mathbf{u} \prod_I P_I(\mathbf{u})^{\alpha_I}, \tag{2.1}$$

where $d\mathbf{u} = du_1 \wedge \dots \wedge du_n$, the P_I are irreducible polynomials in the variables u_i , and $\alpha_I \in \mathbb{C}$. We further assume that the exponents take the form $\alpha_I = n_I + a_I \epsilon$, where $n_I \in \mathbb{Z}$, $a_I \epsilon \in \mathbb{C}^*$, $\sum_I a_I \neq 0$, and where ϵ can be taken to be infinitesimally small. We define the decomposition $\omega = \Phi \varphi$ where

$$\Phi = \prod_I P_I(\mathbf{u})^{a_I \epsilon} \quad \text{and} \quad \varphi = d\mathbf{u} \prod_I P_I(\mathbf{u})^{n_I}. \tag{2.2}$$

Here Φ is a multivalued function, while φ is a single-valued differential form. The integration contour γ is chosen to have its boundary contained within the algebraic variety $\prod_I P_I(\mathbf{u}) = 0$.

The natural mathematical framework to discuss such integrals, which we identify as generalized hypergeometric functions, is that of twisted homology and cohomology [10].² In our notation introduced above, ω is an element of a twisted cohomology group, while γ is an element of a twisted homology group. The *twist* is the single-valued function $d \log \Phi$, and it is the multi-valuedness of Φ that gives rise to the twisted structure. The mathematical literature contains a great deal of information about these types of twisted homology and cohomology groups. We would like to identify bases $\{\gamma_j\}$ and $\{\omega_i\}$ of these groups in order to construct a coaction of the form (1.1). In general, this is a difficult problem. We restrict our attention to the case of so-called positive geometries [16], which are sufficient for the known examples of Feynman diagrams with MPL expansions. For these functions, it is possible to identify a basis of homology classes, represented by integration contours, for which one can construct corresponding *canonical* differential forms having logarithmic singularities and unit residues precisely at the boundary of the contour. In our examples where the $P_k(\mathbf{u})$ are mostly linear, there is a natural choice of the basis of integration contours $\{\gamma_j\}$, and it is straightforward to construct their canonical forms. In cases where the canonical forms can be written in the form $\bigwedge_i d \log y_i(\mathbf{u})$, we refer to them as dlog forms.

Stokes' theorem implies that $\int_\gamma \Phi \varphi = \int_\gamma \Phi(\varphi + \nabla_\Phi \xi)$ for an arbitrary smooth $(n-1)$ -form ξ , so φ is a twisted cohomology class. The *intersection number* pairing of twisted cohomology classes (forms) is defined by

$$\langle \varphi_i, \psi_j \rangle_\Phi = \frac{1}{(2\pi i)^2} \int \iota_\Phi(\varphi_i) \wedge \psi_j, \quad (2.3)$$

where ι_Φ denotes selection of a representative of the cohomology class with compact support. To compute intersection numbers in practice, it is preferable to use equivalent formulas based on residues [12, 13]. For the case of dlog forms in one variable, we have

$$\langle \varphi_i, \psi_j \rangle_\Phi = \sum_{u_p} \frac{\text{Res}_{u=u_p} \varphi_i \text{Res}_{u=u_p} \psi_j}{\text{Res}_{u=u_p} d \log \Phi}, \quad (2.4)$$

where the u_p are the poles of $d \log \Phi$.

To construct a coaction formula for a certain hypergeometric function with an integral representation $\int_\gamma \omega$, we use the following procedure.

- Check that ω can be written in the form of eq. (2.1) subject to the conditions expressed below the formula, and that the boundary of γ is contained within the algebraic variety $\prod_I P_I(\mathbf{u}) = 0$.
- Compute the dimensions of the nontrivial cohomology and homology groups, which are equal to each other. Let us call this number r . One method to determine it is to compute the number of critical points of the function Φ , i.e. the number of solutions to the equation $d \log \Phi = 0$. This number is an upper bound for r , which happens to be saturated for large classes of hypergeometric functions, including all of the examples we have studied.
- Identify a basis of cohomology $\{\varphi_i\}$ by selecting a set of r n -forms and computing the matrix of their intersection numbers. If the matrix has full rank, then the differential forms are linearly independent and can be taken as a basis.

²See also refs. [12, 13, 14, 15], which are featured in the contributions of Hjalte Frellesvig and Manoj Mandal to these proceedings.

- Identify a basis of homology $\{\gamma_j\}$ by selecting a set of r integration contours bounded by some of the algebraic varieties $P_l(\mathbf{u}) = 0$ and possibly also extending to infinity.³ We assume that we can choose these contours to be positive geometries. Therefore each γ_j has an associated canonical form $\Omega(\gamma_j)$. Linear independence of the homology classes can be checked by computing the matrix of the intersection numbers $\langle \Omega(\gamma_i), \Omega(\gamma_j) \rangle_{\Phi}$ and verifying that it has rank r .
- The intersection matrices in the previous steps were computed only to check linear independence of the proposed bases. We now use both bases to compute the intersection matrix needed to construct the coaction:

$$\langle \varphi_i, \Omega(\gamma_j) \rangle_{\Phi} \quad (2.5)$$

- The elements of the *inverse* of the intersection matrix (2.5) are the coefficients c_{ij} in the coaction formula (1.1) with the integrands $\omega_i = \Phi \varphi_i$ and the contours γ_j chosen above.

In this procedure, we have constructed two sets of differential forms: one consisting of the canonical forms of the basis of integration contours, and a second set chosen as a basis of integrands in twisted cohomology. The canonical forms of contours could also be taken directly as the basis of integrands. However, the examples that follow will show that making independent choices of the two bases can lead to intersection matrices that are more nearly diagonal, and thus to a coaction formula with fewer terms, if desired.

3. Coaction of ${}_2F_1$

The simplest nontrivial hypergeometric function is Gauss's ${}_2F_1$ function. This function admits an Euler-type integral representation of the form

$${}_2F_1(\alpha, \beta; \gamma; x) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int_0^1 du u^{\alpha-1} (1-u)^{\gamma-\alpha-1} (1-ux)^{-\beta}, \quad (3.1)$$

provided that the integral converges. As discussed above, we restrict our attention to the case where α, β, γ have the form $n_i + a_i \varepsilon$, with $n_i \in \mathbb{Z}$. Under these conditions, the Laurent expansion is given in terms of MPLs.

To construct the coaction, we first disregard the gamma-function prefactors and study the family of integrals of the form

$$\begin{aligned} \int_{\gamma} \omega &\equiv \int_0^1 du u^{n_0+a_0\varepsilon} (1-u)^{n_1+a_1\varepsilon} (1-ux)^{n_x+a_x\varepsilon}, \\ \omega &= \Phi \varphi, \\ \Phi &= u^{a_0\varepsilon} (1-u)^{a_1\varepsilon} (1-ux)^{a_x\varepsilon}, \\ \varphi &= du u^{n_0} (1-u)^{n_1} (1-ux)^{n_x}. \end{aligned}$$

³Hyperplanes at infinity can be included by going to projective space and then imposing the condition that the sum of all α_l be 0.

Here the algebraic variety $\prod_l P_l(\mathbf{u}) = 0$ is the set of finite branch points, 0, 1, and $1/x$. We see that φ is a single-valued differential form, and Φ is a multi-valued function, from which we construct the twist 1-form $d \log \Phi$:

$$d \log \Phi = a_0 \frac{du}{u} - a_1 \frac{du}{1-u} - xa_x \frac{du}{1-ux}.$$

The equation $d \log \Phi = 0$ has two solutions, so the upper bound on the dimensionality of the non-trivial (co)homology groups is 2. In this case it is easy to see that the bound is saturated. The fact that the space of differential forms related through integer shifts of the exponents is 2-dimensional can be understood as expressing Gauss's contiguity relations. One can also visibly identify sensible integration contours connecting pairs of the finite branch points, and see that only two of them can be linearly independent. Consider two such independent contours with boundaries at the branch points, for example $\gamma_1 = [0, 1]$, $\gamma_2 = [0, 1/x]$. From these contours, construct their associated canonical forms [17, 16]

$$\Omega(\gamma_1) = d \log \frac{u-1}{u}, \quad \Omega(\gamma_2) = d \log \frac{u-1/x}{u}.$$

Since it is possible to use these same forms as the basis of twisted cohomology, let us first simply set $\tilde{\varphi}_1 = \Omega(\gamma_1)$, $\tilde{\varphi}_2 = \Omega(\gamma_2)$. Then the entries of the intersection matrix are

$$\begin{aligned} \langle \tilde{\varphi}_1, \Omega(\gamma_1) \rangle_{\Phi} &= \frac{1}{a_0 \varepsilon} + \frac{1}{a_1 \varepsilon}, & \langle \tilde{\varphi}_1, \Omega(\gamma_2) \rangle_{\Phi} &= \frac{1}{a_0 \varepsilon}, \\ \langle \tilde{\varphi}_2, \Omega(\gamma_1) \rangle_{\Phi} &= \frac{1}{a_0 \varepsilon}, & \langle \tilde{\varphi}_2, \Omega(\gamma_2) \rangle_{\Phi} &= \frac{1}{a_0 \varepsilon} + \frac{1}{a_x \varepsilon}. \end{aligned}$$

The matrix can be inverted to produce a coaction formula.

However, a different choice of basis of differential forms leads to a cleaner coaction formula. Suppose that instead we choose φ_1 and φ_2 such that their dlog singularities overlap only with the upper boundaries of γ_1 and γ_2 respectively, leading to a diagonal intersection matrix. Moreover, let us normalize the differential forms so that the intersection matrix is in fact the identity. Concretely, we take

$$\varphi_1 = -a_1 \varepsilon d \log(1-u) = a_1 \varepsilon \frac{du}{1-u}, \quad \varphi_2 = -a_x \varepsilon d \log(1-xu) = a_x \varepsilon x \frac{du}{1-xu}. \quad (3.2)$$

Then the coaction can be written simply as

$$\Delta_{\varepsilon} \int_{\gamma_k} \Phi \varphi_l = \int_{\gamma_k} \Phi \varphi_1 \otimes \int_{\gamma_1} \Phi \varphi_l + \int_{\gamma_k} \Phi \varphi_2 \otimes \int_{\gamma_2} \Phi \varphi_l. \quad (3.3)$$

We have checked this coaction formula through order ε^4 , by verifying that order by order in ε we reproduce the coaction on MPLs.

If desired, the gamma-function prefactors in eq. (3.1) can be restored using the identity

$$\Delta(e^{\gamma \varepsilon} \Gamma(1+\varepsilon)) = e^{\gamma \varepsilon} \Gamma(1+\varepsilon) \otimes e^{\gamma \varepsilon} \Gamma(1+\varepsilon),$$

from which it follows that

$$\Delta(e^{\alpha \gamma \varepsilon} \Gamma(m+\alpha \varepsilon)) = e^{\alpha \gamma \varepsilon} \Gamma(1+\alpha \varepsilon) \otimes e^{\alpha \gamma \varepsilon} \Gamma(m+\alpha \varepsilon),$$

for integer values of m . The exponential factors will cancel between the numerator and the denominator of eq. (3.1). We thus obtain a coaction on ${}_2F_1$, in which each entry in turn can be expressed in terms of ${}_2F_1$ functions. The coaction is given explicitly by

$$\begin{aligned} \Delta\left({}_2F_1(\alpha, \beta; \gamma; x)\right) &= {}_2F_1(1+a\varepsilon, b\varepsilon; 1+c\varepsilon; x) \otimes {}_2F_1(\alpha, \beta; \gamma; x) \\ &\quad - \frac{b\varepsilon}{1+c\varepsilon} {}_2F_1(1+a\varepsilon, 1+b\varepsilon; 2+c\varepsilon; x) \\ &\quad \otimes \frac{\Gamma(1-\beta)\Gamma(\gamma)}{\Gamma(1-\beta+\alpha)\Gamma(\gamma-\alpha)} x^{1-\alpha} {}_2F_1\left(\alpha, 1+\alpha-\gamma; 1-\beta+\alpha; \frac{1}{x}\right), \end{aligned} \quad (3.4)$$

where $\alpha = n_\alpha + a\varepsilon$, $\beta = n_\beta + b\varepsilon$ and $\gamma = n_\gamma + c\varepsilon$.

4. Coaction of Appell F_3

We now consider the Appell F_3 function, which has the following two-dimensional integral representation:

$$\begin{aligned} F_3(\alpha, \alpha', \beta, \beta', \gamma; x, y) &= \frac{\Gamma(\gamma)}{\Gamma(\beta)\Gamma(\beta')\Gamma(\gamma-\beta-\beta')} \times \\ &\quad \int_0^1 dv \int_0^{1-v} du u^{\beta-1} v^{\beta'-1} (1-u-v)^{\gamma-\beta-\beta'-1} (1-ux)^{-\alpha} (1-vy)^{-\alpha'}. \end{aligned} \quad (4.1)$$

Thus we take

$$\begin{aligned} \omega &= \Phi \cdot u^{n_a} v^{n_b} (1-ux)^{n_c} (1-vy)^{n_d} (1-u-v)^{n_s} du \wedge dv. \\ \Phi &= u^{a\varepsilon} v^{b\varepsilon} (1-ux)^{c\varepsilon} (1-vy)^{d\varepsilon} (1-u-v)^{g\varepsilon}. \end{aligned}$$

The five factors $P_l(u, v)$ in ω are all linear, so the geometry underlying the Appell F_3 function is then determined by an arrangement of hyperplanes corresponding to $P_l(u, v) = 0$:

$$\begin{aligned} H_a &= \{u = 0\}, & H_b &= \{v = 0\}, & H_c &= \{1-xu = 0\}, \\ H_d &= \{1-yv = 0\}, & H_g &= \{1-u-v = 0\}, \end{aligned} \quad (4.2)$$

which we represent in Fig. 1 for $x > y > 1$. The dimension of the (co)homology groups can be determined by counting the critical points of Φ . Since the geometry at hand is an arrangement of hyperplanes that intersect only pairwise, there is also a natural basis of homology which is the set of *bounded chambers* (the finite connected regions in the complement of the hyperplanes) in Fig. 1. Either way, we see that the dimension is 4.

In order to get a convenient intersection matrix, we choose a basis of integration contours that are not exactly the bounded chambers mentioned above, but rather four independent regions that are all triangles. Denoting them by subscripts indicating the sides of the triangles, we select γ_{abg} which is the original integration contour γ used in the definition (4.1), γ_{bcg} , γ_{cdg} , and γ_{adg} . In two dimensions, a canonical form for a triangle bounded by the three lines $P_1 = 0, P_2 = 0, P_3 = 0$ can be constructed with the following formula [17]:

$$d \log \frac{P_1}{P_2} \wedge d \log \frac{P_2}{P_3}.$$




Figure 1: Appell F_3 is related to an arrangement of hyperplanes.

For a basis of integrands, let us take

$$\begin{aligned}\varphi_{ab} &= d \log u \wedge d \log v, \\ \varphi_{bc} &= d \log(1 - ux) \wedge d \log v, \\ \varphi_{cd} &= d \log(1 - ux) \wedge d \log(1 - vy), \\ \varphi_{da} &= d \log u \wedge d \log(1 - vy).\end{aligned}$$

Again, the notation indicates the location of the dlog singularities. This pair of bases is convenient because it leads to a diagonal intersection matrix. Since we are concerned with a nondegenerate arrangement of hyperplanes, i.e. the exponents and x and y are generic, and there are only normal crossings of divisors, one can read off the intersection numbers from the oriented intersections of the hyperplanes (and verify that this result agrees with the residue formula). Moreover, the intersection numbers also agree with the $\mathcal{O}(\varepsilon^{-2})$ terms of the integrals $\int_{\gamma} \Phi \varphi$: all nonvanishing terms arise from overlapping endpoint singularities. Explicitly, the intersection matrix is given by

$\langle \Omega(\gamma_i), \varphi_j \rangle$	φ_{ab}	φ_{bc}	φ_{cd}	φ_{ad}
γ_{abg}	$\frac{1}{abe^2}$	0	0	0
γ_{bcg}	0	$\frac{1}{bce^2}$	0	0
γ_{cdg}	0	0	$\frac{1}{cde^2}$	0
γ_{adg}	0	0	0	$\frac{1}{ade^2}$

We thus arrive at the following coaction formula:

$$\begin{aligned}\Delta \left(\int_{\gamma} \Phi \varphi \right) &= ab\varepsilon^2 \int_{\gamma} \Phi \varphi_{ab} \otimes \int_{\gamma_{abg}} \Phi \varphi + bc\varepsilon^2 \int_{\gamma} \Phi \varphi_{bc} \otimes \int_{\gamma_{bcg}} \Phi \varphi \\ &\quad + cd\varepsilon^2 \int_{\gamma} \Phi \varphi_{cd} \otimes \int_{\gamma_{cdg}} \Phi \varphi + ad\varepsilon^2 \int_{\gamma} \Phi \varphi_{ad} \otimes \int_{\gamma_{adg}} \Phi \varphi.\end{aligned}$$

We have explicitly checked that this coaction is compatible with the coaction on MPLs in the Laurent expansion for all entries of the period matrix $\int_{\gamma_i} \Phi \varphi_j$, through weight 4.

5. Summary and discussion

Based on the description of generalized hypergeometric functions in terms of their integral representations and considerations of twisted (co)homology, we have proposed an algebraic coaction on certain classes of these functions, including ${}_{p+1}F_p$, Appell F_1, F_2, F_3, F_4 , the Lauricella series F_A, F_B, F_D , and any hypergeometric functions with a basis of integration contours consisting of positive geometries. When the exponents in the integral representation are expanded around integer values, we claim that this coaction is compatible with the coaction on the MPLs in the Laurent expansion. We have checked this claim to several orders in ε for ${}_2F_1, {}_3F_2$, and the Appell functions F_1, F_2, F_3, F_4 . These examples, and the underlying theory, are discussed in more detail in ref. [11].

This coaction supports our examples of Feynman-diagrammatic coaction to all orders in the dimensional regularization parameter ε . We note that the hypergeometric functions appearing in known Feynman integrals actually violate a key assumption in the results related to twisted (co)homology and intersection numbers, namely that the exponents in the integral representations are nonzero and independent. However, we find that we are able to derive valid coaction formulas in degenerate limits, and we believe that these limits can be justified by a detailed analysis of twisted cycles.

A mathematical treatment of these ideas, applied to motivic versions of the Lauricella functions $F_D^{(n)}$, which includes ${}_2F_1$ and Appell F_1 , has recently been initiated in [18]. The motivic version does not make use of intersection numbers, since the first and second entries in the coaction are normalized independently. The coaction argument and the first entry are motivic, while the second entry is single-valued. It would be interesting to study the connection between that treatment and ours in detail.

Although it has not been emphasized in this brief article, the second entries in our coactions are also modified versions of the hypergeometric functions. The reason is that we require consistency with the Laurent expansion in terms of MPLs, but the second entries in the coaction of MPLs are equivalence classes modulo $2\pi i$. Therefore the closed-form hypergeometric functions in the second entry must carry a similar loss of information. A further, minor, point about the second entries of the coaction (1.1) is that while they fit our definition of generalized hypergeometric functions, it is not always obvious how to recognize them in terms of known functions when represented by integrals over different regions. However, it is true that the second entries in the coaction belong to the same class of function, as stated in the example of eq. (1.4). In the examples we have considered here, namely ${}_2F_1$ and Appell F_3 , it is straightforward to introduce a change of variables for each of our chosen elements of the basis of integration contours, such that the integral can be recognized as another ${}_2F_1$ or F_3 function, respectively.

In these proceedings we have looked only at cases with linear polynomials $P_l(\mathbf{u})$. Nonlinear polynomials appear in examples such as ${}_{p+1}F_p$ for $p > 1$, and Appell F_4 . These functions have been considered in [11]. Since they are associated to positive geometries, we have been able to construct coactions for them as well.

Acknowledgments

This work is supported by the ‘‘Fonds National de la Recherche Scientifique’’ (FNRS), Belgium

(SA), by the European Research Council under the European Union’s Horizon 2020 research and innovation programme through grants 647356 (RB) and 637019 (CD), and the STFC Consolidated Grant “Particle Physics at the Higgs Centre” (EG, JM).

References

- [1] A. B. Goncharov, *Multiple polylogarithms and mixed Tate motives*, *arXiv Mathematics e-prints* (Mar., 2001) [[math/0103059](#)].
- [2] A. B. Goncharov, *Galois symmetries of fundamental groupoids and noncommutative geometry*, *arXiv Mathematics e-prints* (Aug., 2002) [[math/0208144](#)].
- [3] F. C. S. Brown, *Mixed Tate Motives over $\text{Spec}(\mathbb{Z})$* , *Annals of Math.* **175** (2012), no. 1.
- [4] C. Duhr, *Hopf algebras, coproducts and symbols: an application to Higgs boson amplitudes*, *JHEP* **08** (2012) 043, [[1203.0454](#)].
- [5] F. Brown, *Notes on Motivic Periods*, *Commun. Num. Theor. Phys.* **11** (2017) 557–655, [[1512.06410](#)].
- [6] E. Panzer and O. Schnetz, *The Galois coaction on ϕ^4 periods*, *Commun. Num. Theor. Phys.* **11** (2017) 657–705, [[1603.04289](#)].
- [7] F. Brown, *Feynman amplitudes, coaction principle, and cosmic Galois group*, *Commun. Num. Theor. Phys.* **11** (2017) 453–556, [[1512.06409](#)].
- [8] S. Abreu, R. Britto, C. Duhr, and E. Gardi, *Algebraic Structure of Cut Feynman Integrals and the Diagrammatic Coaction*, *Phys. Rev. Lett.* **119** (2017), no. 5 051601, [[1703.05064](#)].
- [9] S. Abreu, R. Britto, C. Duhr, and E. Gardi, *Diagrammatic Hopf algebra of cut Feynman integrals: the one-loop case*, *JHEP* **12** (2017) 090, [[1704.07931](#)].
- [10] K. Aomoto and M. Kita, *Theory of Hypergeometric Functions*. Springer Monographs in Mathematics. Springer Japan, 2011.
- [11] S. Abreu, R. Britto, C. Duhr, E. Gardi, and J. Matthew, *From positive geometries to a coaction on hypergeometric functions*, [[1910.08358](#)].
- [12] S. Mizera, *Scattering Amplitudes from Intersection Theory*, *Phys. Rev. Lett.* **120** (2018), no. 14 141602, [[1711.00469](#)].
- [13] P. Mastrolia and S. Mizera, *Feynman Integrals and Intersection Theory*, *JHEP* **02** (2019) 139, [[1810.03818](#)].
- [14] H. Frellesvig, F. Gasparotto, M. K. Mandal, P. Mastrolia, L. Mattiazzi, and S. Mizera, *Vector Space of Feynman Integrals and Multivariate Intersection Numbers*, [[1907.02000](#)].
- [15] H. Frellesvig, F. Gasparotto, S. Laporta, M. K. Mandal, P. Mastrolia, L. Mattiazzi, and S. Mizera, *Decomposition of Feynman Integrals on the Maximal Cut by Intersection Numbers*, *JHEP* **05** (2019) 153, [[1901.11510](#)].
- [16] N. Arkani-Hamed, Y. Bai, and T. Lam, *Positive Geometries and Canonical Forms*, *JHEP* **11** (2017) 039, [[1703.04541](#)].
- [17] K. Matsumoto, *Intersection numbers for logarithmic k -forms*, *Osaka J. Math.* **35** (1998), no. 4 873–893.
- [18] F. Brown and C. Dupont, *Lauricella hypergeometric functions, unipotent fundamental groups of the punctured Riemann sphere, and their motivic coactions*, [[1907.06603](#)].