



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Earth's ice imbalance

Citation for published version:

Slater, T, Lawrence, IR, Ootosaka, IN, Shepherd, A, Gourmelen, N, Jakob, L, Tepes, P & Gilbert, L 2020 'Earth's ice imbalance' The Cryosphere, Copernicus Publications. <https://doi.org/10.5194/tc-2020-232>

Digital Object Identifier (DOI):

[10.5194/tc-2020-232](https://doi.org/10.5194/tc-2020-232)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Review Article: Earth's ice imbalance

Thomas Slater¹, Isobel R. Lawrence¹, Inès N. Otosaka¹, Andrew Shepherd¹, Noel Gourmelen², Livia Jakob³, Paul Tepes², Lin Gilbert⁴

¹Centre for Polar Observation and Modelling, School of Earth and Environment, University of Leeds, LS2 9JT, UK

5 ²School of GeoSciences, University of Edinburgh, Edinburgh, EH8 9XP, UK

³Earthwave Ltd, Edinburgh, EH9 3HJ, UK

⁴Mullard Space Science Laboratory, Department of Space & Climate Physics, University College London, WC1E 6BT, UK

Correspondence to: Thomas Slater (t.slater1@leeds.ac.uk)

Abstract. We combine satellite observations and numerical models to show that Earth lost 28 trillion tonnes of ice between
10 1994 and 2017. Arctic sea ice (7.6 trillion tonnes), Antarctic ice shelves (6.5 trillion tonnes), mountain glaciers (6.2 trillion
tonnes), the Greenland ice sheet (3.8 trillion tonnes), the Antarctic ice sheet (2.5 trillion tonnes), and Southern Ocean sea ice
(0.9 trillion tonnes) have all decreased in mass. Just over half (60 %) of the ice loss was from the northern hemisphere, and
the remainder (40 %) was from the southern hemisphere. The rate of ice loss has risen by 57 % since the 1990s – from 0.8 to
1.2 trillion tonnes per year – owing to increased losses from mountain glaciers, Antarctica, Greenland, and from Antarctic
15 ice shelves. During the same period, the loss of grounded ice from the Antarctic and Greenland ice sheets and mountain
glaciers raised the global sea level by 35.0 ± 3.2 mm. The majority of all ice losses from were driven by atmospheric melting
(68 % from Arctic sea ice, mountain glaciers ice shelf calving and ice sheet surface mass balance), with the remaining losses
(32 % from ice sheet discharge and ice shelf thinning) being driven by oceanic melting. Altogether, the cryosphere has taken
up 3.2 % of the global energy imbalance.

20 1 Introduction

Fluctuations in Earth's ice cover have been driven by changes in the planetary radiative forcing (Vaughan et al., 2013),
affecting global sea-level (The IMBIE Team, 2018, 2020; Zemp et al., 2019), oceanic conditions (Rahmstorf et al., 2015),
atmospheric circulation (Francis and Vavrus, 2012; Vellinga and Wood, 2002) and freshwater resources (Huss and Hock,
2018; Immerzeel et al., 2020). Earth's cryosphere is created as meteoric ice in Antarctica, Greenland, and in mountain
25 glaciers, and as frozen sea water in the Arctic and Southern oceans (Fig. 1). The polar ice-sheets store more than 99 % (30
million km³) of Earth's freshwater ice on land (Fretwell et al., 2013; Morlighem et al., 2017), and even modest losses raise
the global sea level (The IMBIE Team, 2018, 2020), increase coastal flooding (Vitousek et al., 2017) and disturb oceanic
currents (Golledge et al., 2019). Ice sheet mass balance is the net balance between mass losses associated with ice flow,
melting at the ice-ocean interface and the surface mass balance (the net difference between precipitation, sublimation and
30 meltwater runoff). Ice shelves are a major source of ocean fresh water (Jacobs et al., 1992), provide mechanical support for



grounded ice upstream (Weertman, 1974), and have been a persistent element of the climate system throughout the Holocene period (Domack et al., 2005). There are over 343 documented ice shelves around Antarctica (SCAR, 2020) containing an estimated 380 thousand km³ of ice (Fretwell et al., 2013), and fluctuations in their volume occur as a result of changes in their extent (Cook and Vaughan, 2010) and thickness (Adusumilli et al., 2020). Mountain glacier ice moderates global sea-level and regional hydrology (Huss and Hock, 2018), impacting on local communities who rely on it as a source of freshwater (Immerzeel et al., 2020). There are over 215 thousand glaciers worldwide (RGI Consortium, 2017) containing 160 thousand km³ of ice (Farinotti et al., 2019), and their retreat has accounted for 21 % of global sea-level rise between 1993 and 2017 (WCRP Global Sea Level Budget Group, 2018). Typically 15 to 25 million km² of the global ocean surface is covered in sea ice at any one time of year, though its thickness and extent vary seasonally and due to long-term changes in Earth's climate (Maksym, 2019). Sea ice plays a key role in the freshwater and energy budgets of the polar regions and impacts on their marine ecosystem (Stroeve and Notz, 2018), and its loss could influence oceanic and atmospheric circulation and affect weather patterns in the mid-latitudes (Maksym, 2019; Vihma, 2014).

Although in situ records of glacier mass balance date back to the 1890's (Zemp et al., 2015), substantial records of change for other components of the cryosphere did not begin until the advent of satellite observations in the 1970's. Ice shelf extent has been recorded episodically in satellite imagery since the 1940's (Cook and Vaughan, 2010), sea ice extent has been monitored by satellites since the late 1970's (Cavalieri et al., 1999), and ice sheet, ice shelf, sea ice, and glacier thickness changes have been recorded systematically in satellite altimetry since the 1990s (Gardner et al., 2013; Laxon et al., 2013; Shepherd et al., 2010; The IMBIE Team, 2018, 2020). Here, we combine satellite observations of changing ice sheet, ice shelf, glacier, and Arctic sea ice mass, with in situ and model-based estimates of glacier and Southern Ocean sea ice mass, to quantify trends in Earth's ice. We do not consider trends in other elements of the cryosphere which are small in comparison, such as permafrost, snow on land, and river and lake ice.

2 Mountain Glaciers

We combined 6 estimates of mass change from an extrapolation of local glaciological and geodetic measurements (Zemp et al., 2019, 2020), satellite gravimetry (Wouters et al., 2019), satellite swath altimetry (Foresta et al., 2016; Jakob et al., 2020; Tepes et al., 2020) and satellite synthetic differential aperture radar interferometry (DInSAR) (Braun et al., 2019), to produce a reconciled estimate of global glacier mass changes between 1962 and 2019 and over 19 glacier regions defined in the Randolph Glacier Inventory (RGI Consortium, 2017) (Fig. 2). Satellite gravimetry measures glacier mass change from fluctuations in Earth's gravitational field within local mass concentration units (Wouters et al., 2019). Satellite swath altimetry and satellite SAR interferometry both measure surface elevation change, which is converted to mass by assuming a fixed density of ice; the former uses the swath interferometric mode of CryoSat-2 to obtain a dense grid of repeated elevation measurements (Foresta et al., 2016), while the latter applies differential SAR interferometry to convert each radar scene into



a digital elevation model (DEM) and differences it to a reference DEM (Braun et al., 2019). For each region we aggregated annual mass change rates determined from the techniques available: each region includes between 2 and 4 estimates except for glaciers peripheral to Antarctica and Greenland, where only one is available. For studies in which time-varying mass change rates are not available, we assume the mass change rate to be linear over the period considered and scale the uncertainty by the square root of the number of years. We computed the cumulative mass change as the integral of the aggregated mass change rates and the associated uncertainty as the root sum square of the annual errors. We summed the regional estimates to derive the global glacier mass change and the overall uncertainty as the root-mean square of the regional errors.

We assessed the consistency of the in situ and satellite gravimetry and altimetry estimates between 2010 and 2015 in 7 regions (Arctic Canada North and South, Russian Arctic, Iceland, Svalbard and Jan Mayen, High Mountain Asia and Southern Andes) where measurements from all techniques overlap (Fig. 2). In general, there is close agreement between estimates determined using each technique and they agree to within 1 standard deviation of their mean between 2010 and 2015 with the exception of Iceland – where the altimeter estimates only provide a constant rate of mass change during the overlap period and reduce the variability. We find the greatest difference between estimates in High Mountain Asia, with a standard deviation of 214 Gt between the three techniques and those determined from altimetry being the most negative. Based on our reconciled estimate, glaciers have collectively lost $-10,130 \pm 1713$ Gt of ice between 1962 and 2019, raising the global mean sea-level by 28.1 ± 4.8 mm during this period. Glaciers peripheral to Greenland and in Alaska and the Southern Andes have experienced the largest losses (Fig. 2) – $5,697 \pm 648$ Gt between 1962 and 2019 – and account for more than half (56 %) of the global glacier mass loss over this period. Globally, the rate of glacier mass loss has increased from -120 ± 71 Gt yr⁻¹ in the 1970s to -346 ± 67 Gt yr⁻¹ between 2010 and 2019, peaking at -536 ± 198 Gt yr⁻¹ in 2018. Glacier mass loss is linked to increasing air temperatures; approximately 70 % of the global glacier mass loss has been attributed to anthropogenic forcing, and the remainder is due to natural climate variability (Marzeion et al., 2014).

3 Ice sheets

Ice sheets lose mass when ice discharge and melting at the surface and ice-ocean interface combined exceed snowfall. We use estimates of ice sheet mass balance derived from an ensemble of satellite altimetry, satellite gravimetry and input-output datasets which span the period 1992-2018. For the Antarctic (24 datasets) (The IMBIE Team, 2018) and Greenland (26 datasets) (The IMBIE Team, 2020) ice sheets, independently derived estimates of mass change from the three satellite geodetic techniques were combined into a single estimate of ice sheet mass balance. Estimates of ice sheet mass balance derived from these methods at the continental scale are similar and can be collated to reduce uncertainty (The IMBIE Team, 2018, 2020): satellite altimetry directly measures changes in ice sheet height (Otosaka et al., 2019; Sandberg Sørensen et al., 2018) converted into mass by assigning a specific density to the volume change (Shepherd et al., 2019) or by explicitly



accounting for snowfall fluctuations through firn modelling (Sørensen et al., 2011). Satellite gravimetry measures temporal variations in Earth's gravity field using spherical harmonic solutions (Velicogna et al., 2020) or through local mass concentration analysis (Luthcke et al., 2006). The input-output method removes ice discharge into the oceans (output), estimated from satellite observations of ice velocity and measurements of ice thickness, from the net snow accumulation (input) (Mouginot et al., 2019; Rignot et al., 2019) determined from regional climate modelling (Noël et al., 2018; van Wessem et al., 2018).

These satellite surveys (e.g. Fig. 1) show the Antarctic ice sheet lost $2,603 \pm 563$ Gt of ice between 1992 and 2017, and the Greenland ice sheet lost $3,902 \pm 342$ Gt of ice between 1992 and 2018. Since 2012, the rate of ice loss from Antarctica has tripled when compared to the previous two decades, owing to widespread glacier speedup (Mouginot et al., 2014) and thinning (Shepherd et al., 2019) in the Amundsen and Bellingshausen Sea sectors in response to the circulation of warm water under the region's ice shelves (Jacobs et al., 2011). Ice shelf collapse (Cook and Vaughan, 2010) (Fig. 3) and thinning at the Antarctic Peninsula has triggered speedup of glaciers upstream (Hogg et al., 2017) as a consequence of reduced ice shelf buttressing. Unlike in Antarctica, where almost all of the ice loss is associated with ice dynamical imbalance, just over half of Greenland's mass loss during this period arose due to increases in meltwater runoff (Enderlin et al., 2014) enhanced by atmospheric circulation during several warm summers (Bevis et al., 2019). The remaining ice loss was due to increased glacier discharge, primarily at Jakobshavn Isbræ (Holland et al., 2008) and at outlet glaciers in the southeast (Howat et al., 2008) and northwest (Moon et al., 2012). Both ice dynamic and surface processes in Greenland have led to widespread thinning at the ice sheet margins and within individual glacier catchments (McMillan et al., 2016) (Fig. 1). Altogether, ice losses from Antarctica and Greenland have caused global sea levels to rise by 17.8 ± 1.8 mm between 1992 and 2017 (The IMBIE Team, 2018, 2020).

4 Antarctic ice shelves

To compute trends in the volume of Antarctic ice shelves associated with changes in their extent, we combined satellite-based records of their thickness (Fretwell et al., 2013) and area change (Cook and Vaughan, 2010) over time, adjusted for changes in thickness where they have been recorded (Adusumilli et al., 2020). We restrict this calculation to ice shelves where a record of progressive retreat has been well-established (Fig. 3). Although area changes have been mapped since the late 1940's, comprehensive estimates of their thickness only began in the early 1990's. To estimate the thickness of icebergs calved prior to this period, we combined in situ, airborne, and satellite-derived measurements of ice thickness recorded prior to when the ice shelf calving took place (Fig. 3). Uncertainties in volume change associated with ice shelf retreat were computed as the product of errors in ice thickness, determined from the variance of the thickness data, and extent, determined from the precision of the satellite imagery (Cook and Vaughan, 2010). We then used multi-mission satellite radar altimetry to compute trends in the volume of Antarctic ice shelves due to changes in their thickness, and their associated



uncertainty, between 1994 and 2020 (Adusumilli et al., 2020). The altimeter range measurements were first adjusted for
125 changes in ocean surface height, including contributions due to the geoid, mean dynamic topography, ocean tide, ocean load
tide, atmospheric pressure, and sea-level rise. We constructed separate time-series of elevation change for each mission
using a space-time polynomial fit to data falling within regularly spaced 10 km grid cells. The individual elevation change
time-series were then merged by ensuring that rates of height-change during overlapping time periods were equal to the
average of rates from each mission. We calculated changes in ice shelf thickness, and the associated changes in volume, by
130 adjusting the height changes for changes in firm air content and using a hydrostatic buoyancy relationship, assuming values
of 917 and 1,028 kg/m³ for the densities of ice and ocean water, respectively.

Antarctic ice shelves have lost $8,667 \pm 1240$ Gt of their mass since 1994, 54 % of which has been due to reductions in their
extent and the remainder due to changes in their thickness. Although episodic iceberg calving is part of the natural cycle of
ice mass transport through the continent, there has been a 39,717 km² loss of ice shelf area at the Antarctic Peninsula (e.g.
135 (Cook and Vaughan, 2010)) where air temperatures have risen several times faster than the global trend (Vaughan et al.,
2003). Warmer air leads to increased surface melting, which can promote iceberg calving through hydraulic fracture of
crevasses (Scambos et al., 2013). At the same time, ocean-driven melting has caused some ice shelves to thin at their base,
particularly in the Amundsen and Bellingshausen Seas (Paolo et al., 2015; Shepherd et al., 2010) where warm circumpolar
deep water is present (Jacobs et al., 1996), but also at the Antarctic Peninsula (Shepherd et al., 2003). Ice shelf thinning can
140 promote instability by weakening their lateral margins (Vieli et al., 2007). Both processes – calving front retreat and basal
melting – have triggered speedup of inland ice (Rignot et al., 2004; Scambos et al., 2004; Shepherd et al., 2004) due to the
associated reduction in buttressing (Joughin et al., 2012), leading to global sea-level rise (The IMBIE Team, 2018) even
though ice shelves themselves are not a direct source of ocean mass. The ice shelf losses combined amount to 3 % of their
present volume, while those in the Amundsen and Bellingshausen Seas are now 10 to 18 % thinner (Paolo et al., 2015) and
145 those at the Antarctic Peninsula are 18 % smaller in extent (Cook and Vaughan, 2010).

5 Sea ice

We estimated trends in the mass of Arctic sea ice using a combination of sea-ice ocean modelling and satellite measurements
of thickness change: between 1980 and 2011 we used the Pan-Arctic Ice-Ocean Modelling and Assimilation System
(PIOMAS), a coupled sea ice-ocean model forced with atmospheric reanalyses (Zhang and Rothrock, 2003); from 2011, we
150 used CryoSat-2 satellite radar altimetry measurements of sea ice volume (Tilling et al., 2018). We converted PIOMAS
volume estimates to mass assuming a fixed density of 917 kg/m³ (Schweiger, personal comm). We divided CryoSat-2
monthly volume estimates into regions of multi-year and first-year ice and multiplied by densities of 882 kg/m³ and 916.7
kg/m³, respectively, to convert to mass (Tilling et al., 2018). The presence of melt ponds on the Arctic sea ice surface from
May to September make it difficult to discriminate between radar returns from leads and sea ice floes, preventing the



155 retrieval of summer sea ice thickness and volume from radar altimetry (Tilling et al., 2018). As a result, we computed the
winter-mean (October to April) mass trend across the Arctic for both CryoSat-2 and PIOMAS estimates to maintain
consistency: the difference between winter (October-April) and annual (January-December) PIOMAS mass trends during
1980-2011 is 19 Gt yr^{-1} , small (6 %) when compared to the magnitude of the overall 12-month trend (-324 Gt yr^{-1}). Since the
annual trend is slightly larger, we consider our winter-average mass trend to be a conservative estimate of the actual Arctic
160 sea ice mass loss. In the absence of an available satellite-derived Antarctic sea ice volume product, we used the Global Ice-
Ocean Modelling and Assimilation System (GIOMAS) (Zhang and Rothrock, 2003), the global equivalent to PIOMAS, to
estimate the trend. We gridded GIOMAS sea ice thickness data onto 0.2×0.5 degree grids, multiplied by cell area to retrieve
total volume and used a density of 917 kg/m^3 to convert to mass (Zhang, personal comm). Antarctic sea ice trends were
computed as annual averages between January and December. We assumed an error on monthly PIOMAS/GIOMAS volume
165 of 1800 km^3 (Schweiger et al., 2011; Schweiger & Zhang, personal comm), and converted to a mass error using a density of
 917 kg/m^3 . We estimated the uncertainty on monthly Arctic sea ice volume and mass from CryoSat-2 as a percentage error,
which varies from 14.5 % volume in October to 13 % volume in April (Tilling et al., 2018). The uncertainty on the winter-
average (Arctic) and annual-average (Antarctic) mass was propagated from the monthly errors. Finally, we estimated the
uncertainty associated with a rate of mass change over a given time period by dividing the total error by the number of years.

170 Between the winters of 1980 (October 1979 to April 1980) and 2019 (October 2018 to April 2019) Arctic sea ice mass
reduced by $230 \pm 27 \text{ Gt yr}^{-1}$, predominantly due to a decline in the lateral extent of the ice cover (Fig. 1) which accounts for
93 % of the variance in volume over the entire PIOMAS record. Arctic sea ice loss has been attributed to atmospheric
warming driven by anthropogenic CO_2 emissions (Stroeve and Notz, 2018), which has been enhanced in the Arctic when
compared to the mid-latitudes likely due to sea ice loss itself (Dai et al., 2019; Screen and Simmonds, 2010). Between 1980
175 and 2019, GIOMAS volume estimates, which incorporate observations of sea ice extent, show an increase in Antarctic sea
ice of $+43 \pm 17 \text{ Gt yr}^{-1}$. No consensus has been reached on whether trends in Antarctic sea ice cover are anthropogenically
driven, e.g. via depletion of the Ozone layer (Ferreira et al., 2015), or the result of natural climate variability (Meehl et al.,
2016; Zhang et al., 2019). Given the vastness of the continent it surrounds, regional analyses of Southern Ocean sea ice are
essential to understand the processes driving it. The overall trend is a combination of sea ice gains in the western Ross Sea
180 and losses in the Amundsen and Bellingshausen seas, with small increases elsewhere (Parkinson, 2019). In general, global
climate models predict a shrinking southern ice cap in response to climate change; projections from the latest coupled
climate models suggest that Antarctic sea ice will decline during the 21st century (Roach et al., 2020).

6 Earth's ice imbalance

To determine the global ice imbalance, we summed the mass change of each ice component computed at annual intervals
185 and estimated the combined uncertainty as the root sum square of the individual errors. Between 1994 and 2017, the Earth



lost 27.7 ± 2.0 Tt of ice (Fig. 4) – at an average rate of 1.2 ± 0.3 Tt per year (Table 1). Ice losses have been larger in the northern hemisphere, primarily owing to declining Arctic sea ice (-7559 ± 1021 Gt) followed by glacier retreat ($-5,247 \pm 617$ Gt) and Greenland ice sheet melt ($-3,821 \pm 323$ Gt). Ice in the southern hemisphere from the ice shelves ($-6,532 \pm 1221$ Gt), the Antarctic ice sheet ($-2,545 \pm 554$ Gt), glaciers (-972 ± 735 Gt), and sea ice in the Southern Ocean (-924 ± 674 Gt) has
190 been lost at a total rate of -478 ± 146 Gt yr⁻¹ – 34 % slower than in the northern hemisphere (-723 ± 207 Gt yr⁻¹). Earth’s ice can be categorised into its floating and on-land components; grounded ice loss from ice sheets and glaciers raises the global sea-level (The IMBIE Team, 2018, 2020; Zemp et al., 2019), influences oceanic circulation through freshwater input (Rahmstorf et al., 2015) and glacier retreat impacts local communities who rely on glaciers as a freshwater resource (Immerzeel et al., 2020). Grounded ice losses have raised the global mean sea-level by 25.2 ± 1.9 mm and 9.8 ± 2.6 mm in
195 the northern and southern hemispheres respectively, totalling 35.0 ± 3.2 mm over the 24-year period. Although the loss of floating sea ice and ice shelves does not contribute to global sea-level rise, sea ice decline increases habitat loss (Rode et al., 2014), coastal erosion (Overeem et al., 2011) and ocean circulation (Armitage et al., 2020), and may affect mid-latitude weather and climate (Blackport et al., 2019; Overland et al., 2016).

There is now widespread evidence that climate change has caused reductions in Earth’s ice. On average, the planetary
200 surface temperature has risen by 0.85 °C since 1880, and this signal has been amplified in the polar regions (Hartmann et al., 2013). Although this warming has led to higher snowfall in winter, it has also driven larger increases in summertime surface melting (Huss and Hock, 2018). The global oceans have warmed too (Hartmann et al., 2013), with significant impacts on tidewater glaciers (Hogg et al., 2017; Holland et al., 2008), on floating ice shelves (Shepherd et al., 2010), and on the ice streams which have relied on their buttressing (Rignot et al., 2004). Atmospheric warming – anthropogenic or otherwise – is
205 responsible for the recent and long-term reductions in mountain glacier ice (Marzeion et al., 2014), and ocean-driven melting of outlet glaciers has caused the vast majority of the observed ice losses from Antarctica (The IMBIE Team, 2018). Elsewhere, the picture is more complicated. In Greenland, for example, roughly half of all ice losses are associated with trends in surface mass balance, and the remainder is due to accelerated ice flow triggered by ocean melting at glacier termini (The IMBIE Team, 2020). Although the retreat and collapse of ice shelves at the Antarctic Peninsula has occurred in tandem
210 with a rapid regional atmospheric warming (Vaughan et al., 2003), warm circumpolar deep water has melted the base of ice shelves in the Amundsen and Bellingshausen Seas (Jacobs et al., 2011) and this now amounts to over half of their net loss. While the progressive retreat of Arctic sea ice has been driven by radiative forcing, this has been mediated in part by the increasing presence of open water (Perovich and Richter-Menge, 2009), and broader changes in oceanic conditions are expected to play an increasingly important role (Carmack et al., 2016). Finally, although the extent of Southern Ocean sea
215 ice has shown little overall change, there have been considerable regional variations owing to changes in both atmospheric and oceanic forcing (Hobbs et al., 2016). Attributing Arctic sea ice decline and ice shelf calving to increased radiative forcing, approximately 68 % of the recent global ice imbalance is due to atmospheric warming, and the remainder is due to ocean-driven melting. We determine the energy required to melt the total ice loss as:



$$E = M(L + c_p \Delta T), \quad (1)$$

220 where M is the mass of ice, ΔT is the rise in temperature required (we assume an initial ice temperature of $-20\text{ }^\circ\text{C}$), L is the latent heat of fusion for water (333 J/g), and c_p is specific heat capacity of water ($2108\text{ J/Kg }^\circ\text{C}$). Altogether, the ice loss amounts to an 8.9×10^{21} J sink of energy, or 3.2% of the global imbalance over the same period (Schuckmann et al., 2020).

7 Conclusions

Even though Earth's cryosphere has absorbed only a small fraction of the global energy imbalance, it has lost a staggering 28 trillion tonnes of ice between 1994 and 2017. The loss of grounded ice during this period has caused sea-levels to rise by 35.0 ± 3.2 mm, and the loss of floating ice has caused reductions in the planetary albedo (Thackeray and Hall, 2019), reductions in the buttressing of grounded ice (Rignot et al., 2004), ocean freshening (Jacobs et al., 1996), and ocean cooling (Bintanja et al., 2013). Our assessment is based primarily on observations; we use satellite measurements to determine Antarctic and Greenland ice sheet mass balance and to determine changes in the mass of Antarctic ice shelves associated with retreat and thinning, we use a combination of satellite observations and *in situ* measurements to determine changes in the mass of mountain glaciers, and we use a combination of numerical models and satellite observations to determine changes in the mass of sea ice. There is generally good agreement in mass trends derived from observations and models, where both are available. Only our estimate of Southern Ocean sea ice mass imbalance depends on modelling alone (Zhang and Rothrock, 2003), though satellite observations of changes in its extent (Parkinson, 2019) and *in situ* observations of changes in its thickness (Worby et al., 2008) suggest that little change has occurred in Antarctic sea ice cover. The overall rate of ice loss has increased by 49 % over the past 24 years compared to the 1990s, and *in situ* measurements of changes in glacier mass (Zemp et al., 2019) and satellite records of ice shelf extent (Cook and Vaughan, 2010) which pre-date the complete survey confirm this trend. Although a small fraction of mountain glacier losses are associated with retreat since the little ice age (Marzeion et al., 2014), there can be little doubt that the vast majority of Earth's ice loss is a direct consequence of climate warming.

Data availability

Mountain glacier mass change data from glaciological and geodetic observations are freely available at (<https://doi.org/10.5281/zenodo.1492141>) and elevation change fields from DInSAR are available via the World Data Center (<https://doi.org/10.1594/PANGAEA.893612>). Mass change data for the Antarctic and Greenland ice sheets are provided by the ice sheet mass balance intercomparison exercise (IMBIE) and are available at (<http://imbie.org/data-downloads/>). Changes in ice shelf extent can be downloaded from the Scientific Committee on Antarctic Research digital database



(<https://www.add.scar.org/>). PIOMAS/GIOMAS data are freely available from the University of Washington Polar Science Data Center (<http://psc.apl.uw.edu/data/>).

Author contributions

250 TS, IRL, INO and AS designed the study, performed the data analysis and wrote the manuscript. NG, LJ and PT prepared mountain glacier mass change estimates from CryoSat-2 satellite radar altimetry. LG prepared the ice sheet thickness change datasets from multi-mission satellite radar altimetry used in Fig. 1.

Competing interests

The authors declare that they have no conflict of interest.

255 Acknowledgements

We thank Axel Schweiger and Jinlun Zhang for their help with PIOMAS/GIOMAS data and Susheel Adusumilli for providing ice shelf thickness change data. This work was supported by the Natural Environment Research Council Centre for Polar Observation and Modelling (grant number cpom300001).

References

- 260 Adusumilli, S., Fricker, H. A., Medley, B., Padman, L. and Siegfried, M. B.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves, *Nature Geoscience*, doi:10.1038/s41561-020-0616-z, 2020.
- Armitage, T. W. K., Manucharyan, G. E., Petty, A. A., Kwok, R. and Thompson, A. F.: Enhanced eddy activity in the Beaufort Gyre in response to sea ice loss, *Nature Communications*, 11(1), 761, doi:10.1038/s41467-020-14449-z, 2020.
- 265 Bevis, M., Harig, C., Khan, S. A., Brown, A., Simons, F. J., Willis, M., Fettweis, X., Broeke, M. R. van den, Madsen, F. B., Kendrick, E., Caccamise, D. J., Dam, T. van, Knudsen, P. and Nylén, T.: Accelerating changes in ice mass within Greenland, and the ice sheet's sensitivity to atmospheric forcing, *PNAS*, 116(6), 1934–1939, doi:10.1073/pnas.1806562116, 2019.
- Bintanja, R., van Oldenborgh, G. J., Drijfhout, S. S., Wouters, B. and Katsman, C. A.: Important role for ocean warming and increased ice-shelf melt in Antarctic sea-ice expansion, *Nature Geoscience*, 6(5), 376–379, doi:10.1038/ngeo1767, 2013.
- 270 Blackport, R., Screen, J. A., van der Wiel, K. and Bintanja, R.: Minimal influence of reduced Arctic sea ice on coincident cold winters in mid-latitudes, *Nature Climate Change*, 9(9), 697–704, doi:10.1038/s41558-019-0551-4, 2019.
- Braun, M. H., Malz, P., Sommer, C., Fariás-Barahona, D., Sauter, T., Casassa, G., Soruco, A., Skvarca, P. and Seehaus, T. C.: Constraining glacier elevation and mass changes in South America, *Nature Climate Change*, 9(2), 130–136, doi:10.1038/s41558-018-0375-7, 2019.



- 275 Carmack, E. C., Yamamoto-Kawai, M., Haine, T. W. N., Bacon, S., Bluhm, B. A., Lique, C., Melling, H., Polyakov, I. V., Straneo, F., Timmermans, M.-L. and Williams, W. J.: Freshwater and its role in the Arctic Marine System: Sources, disposition, storage, export, and physical and biogeochemical consequences in the Arctic and global oceans, *Journal of Geophysical Research: Biogeosciences*, 121(3), 675–717, doi:10.1002/2015JG003140, 2016.
- 280 Cavaliere, D. J., Parkinson, C. L., Gloersen, P., Comiso, J. C. and Zwally, H. J.: Deriving long-term time series of sea ice cover from satellite passive-microwave multisensor data sets, *Journal of Geophysical Research: Oceans*, 104(C7), 15803–15814, doi:10.1029/1999JC900081, 1999.
- Cook, A. J. and Vaughan, D. G.: Overview of areal changes of the ice shelves on the Antarctic Peninsula over the past 50 years, *The Cryosphere*, 4(1), 77–98, doi:https://doi.org/10.5194/tc-4-77-2010, 2010.
- Dai, A., Luo, D., Song, M. and Liu, J.: Arctic amplification is caused by sea-ice loss under increasing CO₂, *Nature Communications*, 10(1), 121, doi:10.1038/s41467-018-07954-9, 2019.
- 285 Domack, E., Duran, D., Leventer, A., Ishman, S., Doane, S., McCallum, S., Amblas, D., Ring, J., Gilbert, R. and Prentice, M.: Stability of the Larsen B ice shelf on the Antarctic Peninsula during the Holocene epoch, *Nature*, 436(7051), 681–685, doi:10.1038/nature03908, 2005.
- Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M.-J., Angelen, J. H. van and Broeke, M. R. van den: An improved mass budget for the Greenland ice sheet, *Geophysical Research Letters*, 41(3), 866–872, doi:10.1002/2013GL059010, 2014.
- 290 Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F. and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nature Geoscience*, 12(3), 168–173, doi:10.1038/s41561-019-0300-3, 2019.
- Ferreira, D., Marshall, J., Bitz, C. M., Solomon, S. and Plumb, A.: Antarctic Ocean and Sea Ice Response to Ozone Depletion: A Two-Time-Scale Problem, *J. Climate*, 28(3), 1206–1226, doi:10.1175/JCLI-D-14-00313.1, 2015.
- 295 Foresta, L., Gourmelen, N., Pálsson, F., Nienow, P., Björnsson, H. and Shepherd, A.: Surface elevation change and mass balance of Icelandic ice caps derived from swath mode CryoSat-2 altimetry, *Geophysical Research Letters*, 43(23), 12,138–12,145, doi:10.1002/2016GL071485, 2016.
- Francis, J. A. and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes, *Geophysical Research Letters*, 39(6), doi:10.1029/2012GL051000, 2012.
- 300 Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippon, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C. and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7(1), 375–393, doi:https://doi.org/10.5194/tc-7-375-2013, 2013.
- 310 Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J., Berthier, E., Hock, R., Pfeffer, W. T., Kaser, G., Ligtenberg, S. R. M., Bolch, T., Sharp, M. J., Hagen, J. O., Broeke, M. R. van den and Paul, F.: A Reconciled Estimate of Glacier Contributions to Sea Level Rise: 2003 to 2009, *Science*, 340(6134), 852–857, doi:10.1126/science.1234532, 2013.



- 315 Golledge, N. R., Keller, E. D., Gomez, N., Naughten, K. A., Bernales, J., Trusel, L. D. and Edwards, T. L.: Global environmental consequences of twenty-first-century ice-sheet melt, *Nature*, 566(7742), 65–72, doi:10.1038/s41586-019-0889-9, 2019.
- 320 Hartmann, D. L., Klein Tank, A. M. G., Rusticucci, M., Alexander, L. V., Brönnimann, S., Charabi, Y., Dentener, F. J., Dlugokencky, E. J., Easterling, D. R., Kaplan, A., Soden, B. J., Thorne, P. W., Wild, M. and Zhai, P. M.: Observations: Atmosphere and Surface, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, pp. 159–254, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- Hobbs, W. R., Massom, R., Stammerjohn, S., Reid, P., Williams, G. and Meier, W.: A review of recent changes in Southern Ocean sea ice, their drivers and forcings, *Global and Planetary Change*, 143, 228–250, doi:10.1016/j.gloplacha.2016.06.008, 2016.
- 325 Hogg, A. E., Shepherd, A., Cornford, S. L., Briggs, K. H., Gourmelen, N., Graham, J. A., Joughin, I., Mouginot, J., Nagler, T., Payne, A. J., Rignot, E. and Wuite, J.: Increased ice flow in Western Palmer Land linked to ocean melting, *Geophysical Research Letters*, 44(9), 4159–4167, doi:10.1002/2016GL072110, 2017.
- Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H. and Lyberth, B.: Acceleration of Jakobshavn Isbræ triggered by warm subsurface ocean waters, *Nature Geoscience*, 1(10), 659–664, doi:10.1038/ngeo316, 2008.
- 330 Howat, I. M., Joughin, I., Fahnestock, M., Smith, B. E. and Scambos, T. A.: Synchronous retreat and acceleration of southeast Greenland outlet glaciers 2000–06: ice dynamics and coupling to climate, *Journal of Glaciology*, 54(187), 646–660, doi:10.3189/002214308786570908, 2008.
- Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nature Climate Change*, 8(2), 135–140, doi:10.1038/s41558-017-0049-x, 2018.
- 335 Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B. J., Elmore, A. C., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P. D. A., Kulkarni, A. V., Mayewski, P. A., Nepal, S., Pacheco, P., Painter, T. H., Pellicciotti, F., Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A. B., Viviroli, D., Wada, Y., Xiao, C., Yao, T. and Baillie, J. E. M.: Importance and vulnerability of the world’s water towers, *Nature*, 577(7790), 364–369, doi:10.1038/s41586-019-1822-y, 2020.
- 340 Jacobs, S. S., Helmer, H. H., Doake, C. S. M., Jenkins, A. and Frolich, R. M.: Melting of ice shelves and the mass balance of Antarctica, *Journal of Glaciology*, 38(130), 375–387, doi:10.3189/S0022143000002252, 1992.
- Jacobs, S. S., Hellmer, H. H. and Jenkins, A.: Antarctic Ice Sheet melting in the southeast Pacific, *Geophysical Research Letters*, 23(9), 957–960, doi:10.1029/96GL00723, 1996.
- 345 Jacobs, S. S., Jenkins, A., Giulivi, C. F. and Dutrieux, P.: Stronger ocean circulation and increased melting under Pine Island Glacier ice shelf, *Nature Geosci*, 4(8), 519–523, doi:10.1038/ngeo1188, 2011.
- Jakob, L., Gourmelen, N., Ewart, M. and Plummer, S.: Ice loss in High Mountain Asia and the Gulf of Alaska observed by CryoSat-2 swath altimetry between 2010 and 2019, *The Cryosphere Discussions*, 1–29, doi:https://doi.org/10.5194/tc-2020-176, 2020.
- 350 Joughin, I., Alley, R. B. and Holland, D. M.: Ice-Sheet Response to Oceanic Forcing, *Science*, 338(6111), 1172–1176, doi:10.1126/science.1226481, 2012.



- Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S. and Davidson, M.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophysical Research Letters*, 40(4), 732–737, doi:10.1002/grl.50193, 2013.
- 355 Luthcke, S. B., Zwally, H. J., Abdalati, W., Rowlands, D. D., Ray, R. D., Nerem, R. S., Lemoine, F. G., McCarthy, J. J. and Chinn, D. S.: Recent Greenland Ice Mass Loss by Drainage System from Satellite Gravity Observations, *Science*, 314(5803), 1286–1289, doi:10.1126/science.1130776, 2006.
- Maksym, T.: Arctic and Antarctic Sea Ice Change: Contrasts, Commonalities, and Causes, *Annual Review of Marine Science*, 11(1), 187–213, doi:10.1146/annurev-marine-010816-060610, 2019.
- 360 Marzeion, B., Cogley, J. G., Richter, K. and Parkes, D.: Attribution of global glacier mass loss to anthropogenic and natural causes, *Science*, 345(6199), 919–921, doi:10.1126/science.1254702, 2014.
- McMillan, M., Leeson, A., Shepherd, A., Briggs, K., Armitage, T. W. K., Hogg, A., Munneke, P. K., Broeke, M. van den, Noël, B., Berg, W. J. van de, Ligtenberg, S., Horwath, M., Groh, A., Muir, A. and Gilbert, L.: A high-resolution record of Greenland mass balance, *Geophysical Research Letters*, 43(13), 7002–7010, doi:10.1002/2016GL069666, 2016.
- 365 Meehl, G. A., Arblaster, J. M., Bitz, C. M., Chung, C. T. Y. and Teng, H.: Antarctic sea-ice expansion between 2000 and 2014 driven by tropical Pacific decadal climate variability, *Nature Geoscience*, 9(8), 590–595, doi:10.1038/ngeo2751, 2016.
- Moon, T., Joughin, I., Smith, B. and Howat, I.: 21st-Century Evolution of Greenland Outlet Glacier Velocities, *Science*, 336(6081), 576–578, doi:10.1126/science.1219985, 2012.
- 370 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M. and Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, 44(21), 11,051–11,061, doi:10.1002/2017GL074954, 2017.
- 375 Mouginot, J., Rignot, E. and Scheuchl, B.: Sustained increase in ice discharge from the Amundsen Sea Embayment, West Antarctica, from 1973 to 2013, *Geophysical Research Letters*, 41(5), 1576–1584, doi:10.1002/2013GL059069, 2014.
- Mouginot, J., Rignot, E., Bjørk, A. A., Broeke, M. van den, Millan, R., Morlighem, M., Noël, B., Scheuchl, B. and Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *PNAS*, 116(19), 9239–9244, doi:10.1073/pnas.1904242116, 2019.
- 380 Noël, B., Berg, W. J. van de, Wessem, J. M. van, Meijgaard, E. van, As, D. van, Lenaerts, J. T. M., Lhermitte, S., Kuipers Munneke, P., Smeets, C. J. P. P., Ulf, L. H. van, Wal, R. S. W. van de and Broeke, M. R. van den: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 – Part 1: Greenland (1958–2016), *The Cryosphere*, 12(3), 811–831, doi:https://doi.org/10.5194/tc-12-811-2018, 2018.
- Otosaka, I., Shepherd, A. and McMillan, M.: Ice Sheet Elevation Change in West Antarctica From Ka-Band Satellite Radar Altimetry, *Geophysical Research Letters*, 46(22), 13135–13143, doi:10.1029/2019GL084271, 2019.
- 385 Overeem, I., Anderson, R. S., Wobus, C. W., Clow, G. D., Urban, F. E. and Matell, N.: Sea ice loss enhances wave action at the Arctic coast, *Geophysical Research Letters*, 38(17), doi:10.1029/2011GL048681, 2011.



- Overland, J. E., Dethloff, K., Francis, J. A., Hall, R. J., Hanna, E., Kim, S.-J., Screen, J. A., Shepherd, T. G. and Vihma, T.: Nonlinear response of mid-latitude weather to the changing Arctic, *Nature Climate Change*, 6(11), 992–999, doi:10.1038/nclimate3121, 2016.
- 390 Paolo, F. S., Fricker, H. A. and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348(6232), 327–331, doi:10.1126/science.aaa0940, 2015.
- Parkinson, C. L.: A 40-y record reveals gradual Antarctic sea ice increases followed by decreases at rates far exceeding the rates seen in the Arctic, *Proc Natl Acad Sci USA*, 116(29), 14414, doi:10.1073/pnas.1906556116, 2019.
- 395 Perovich, D. K. and Richter-Menge, J. A.: Loss of Sea Ice in the Arctic, *Annual Review of Marine Science*, 1(1), 417–441, doi:10.1146/annurev.marine.010908.163805, 2009.
- Rahmstorf, S., Box, J. E., Feulner, G., Mann, M. E., Robinson, A., Rutherford, S. and Schaffernicht, E. J.: Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation, *Nature Clim Change*, 5(5), 475–480, doi:10.1038/nclimate2554, 2015.
- 400 RGI Consortium: Randolph Glacier Inventory – A dataset of global glacier outlines: Version 6.0: Technical Report, Global Land Ice Measurements from Space, Colorado, USA, doi:doi:10.7265/N5-RGI-60, 2017.
- Rignot, E., Casassa, G., Gogineni, P., Krabill, W., Rivera, A. and Thomas, R.: Accelerated ice discharge from the Antarctic Peninsula following the collapse of Larsen B ice shelf, *Geophysical Research Letters*, 31(18), doi:10.1029/2004GL020697, 2004.
- 405 Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J. and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proc Natl Acad Sci USA*, 116(4), 1095, doi:10.1073/pnas.1812883116, 2019.
- Roach, L. A., Dörr, J., Holmes, C. R., Massonnet, F., Blockley, E. W., Notz, D., Rackow, T., Raphael, M. N., O’Farrell, S. P., Bailey, D. A. and Bitz, C. M.: Antarctic Sea Ice Area in CMIP6, *Geophysical Research Letters*, 47(9), e2019GL086729, doi:10.1029/2019GL086729, 2020.
- 410 Sandberg Sørensen, L., Simonsen, S. B., Forsberg, R., Khvorostovsky, K., Meister, R. and Engdahl, M. E.: 25 years of elevation changes of the Greenland Ice Sheet from ERS, Envisat, and CryoSat-2 radar altimetry, *Earth and Planetary Science Letters*, 495, 234–241, doi:10.1016/j.epsl.2018.05.015, 2018.
- 415 Scambos, T., Hulbe, C. and Fahnestock, M.: Climate-Induced Ice Shelf Disintegration in the Antarctic Peninsula, in *Antarctic Peninsula Climate Variability: Historical and Paleoenvironmental Perspectives*, pp. 79–92, American Geophysical Union (AGU), 2013.
- Scambos, T. A., Bohlander, J. A., Shuman, C. A. and Skvarca, P.: Glacier acceleration and thinning after ice shelf collapse in the Larsen B embayment, Antarctica, *Geophysical Research Letters*, 31(18), doi:10.1029/2004GL020670, 2004.
- SCAR: Antarctic Digital Database, Version 7.2, 2020.
- 420 Schuckmann, K. von, Cheng, L., Palmer, M. D., Tassone, C., Aich, V., Adusumilli, S., Beltrami, H., Boyer, T., Cuesta-Valero, F. J., Desbruyères, D., Domingues, C., García-García, A., Gentine, P., Gilson, J., Gorfer, M., Haimberger, L., Ishii, M., Johnson, G. C., Killik, R., King, B. A., Kirchengast, G., Kolodziejczyk, N., Lyman, J., Marzeion, B., Mayer, M., Monier, M., Monselesan, D. P., Purkey, S., Roemmich, D., Schweiger, A., Seneviratne, S. I., Shepherd, A., Slater, D. A., Steiner, A. K., Straneo, F., Timmermans, M.-L. and Wijffels, S. E.: Heat stored in the Earth system: Where does the energy



- 425 go? The GCOS Earth heat inventory team, *Earth System Science Data Discussions*, 1–45, doi:https://doi.org/10.5194/essd-2019-255, 2020.
- Schweiger, A., Lindsay, R., Zhang, J., Steele, M., Stern, H. and Kwok, R.: Uncertainty in modeled Arctic sea ice volume, *Journal of Geophysical Research: Oceans*, 116(C8), doi:10.1029/2011JC007084, 2011.
- Screen, J. A. and Simmonds, I.: The central role of diminishing sea ice in recent Arctic temperature amplification, *Nature*, 464(7293), 1334–1337, doi:10.1038/nature09051, 2010.
- 430 Shepherd, A., Wingham, D., Payne, T. and Skvarca, P.: Larsen Ice Shelf Has Progressively Thinned, *Science*, 302(5646), 856–859, doi:10.1126/science.1089768, 2003.
- Shepherd, A., Wingham, D. and Rignot, E.: Warm ocean is eroding West Antarctic Ice Sheet, *Geophysical Research Letters*, 31(23), doi:10.1029/2004GL021106, 2004.
- 435 Shepherd, A., Wingham, D., Wallis, D., Giles, K., Laxon, S. and Sundal, A. V.: Recent loss of floating ice and the consequent sea level contribution, *Geophysical Research Letters*, 37(13), doi:10.1029/2010GL042496, 2010.
- Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E. and Engdahl, M. E.: Trends in Antarctic Ice Sheet Elevation and Mass, *Geophysical Research Letters*, 46(14), 8174–8183, doi:10.1029/2019GL082182, 2019.
- 440 Sørensen, L. S., Simonsen, S. B., Nielsen, K., Lucas-Picher, P., Spada, G., Adalgeirsdottir, G., Forsberg, R. and Hvidberg, C. S.: Mass balance of the Greenland ice sheet (2003–2008) from ICESat data – the impact of interpolation, sampling and firn density, *The Cryosphere*, 5(1), 173–186, doi:https://doi.org/10.5194/tc-5-173-2011, 2011.
- Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, *Environ. Res. Lett.*, 13(10), 103001, doi:10.1088/1748-9326/aade56, 2018.
- 445 Tepes, P., Gourmelen, N., Nienow, P., Tsamados, M., Shepherd, A. and Weissgerber, F.: Dynamic and surface mass imbalance of Arctic glaciers and ice caps, submitted to *Remote Sensing of Environment*, 2020.
- Thackeray, C. W. and Hall, A.: An emergent constraint on future Arctic sea-ice albedo feedback, *Nature Climate Change*, 9(12), 972–978, doi:10.1038/s41558-019-0619-1, 2019.
- The IMBIE Team: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558(7709), 219–222, doi:10.1038/s41586-018-0179-y, 2018.
- 450 The IMBIE Team: Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, doi:10.1038/s41586-019-1855-2, 2020.
- Tilling, R. L., Ridout, A. and Shepherd, A.: Estimating Arctic sea ice thickness and volume using CryoSat-2 radar altimeter data, *Advances in Space Research*, 62(6), 1203–1225, doi:10.1016/j.asr.2017.10.051, 2018.
- 455 Vaughan, D. G., Marshall, G. J., Connolley, W. M., Parkinson, C., Mulvaney, R., Hodgson, D. A., King, J. C., Pudsey, C. J. and Turner, J.: Recent Rapid Regional Climate Warming on the Antarctic Peninsula, *Climatic Change*, 60(3), 243–274, doi:10.1023/A:1026021217991, 2003.
- Vaughan, D. G., Comiso, J. C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray, T., Paul, F., Ren, J., Rignot, E., Solomina, O., Steffen, K. and Zhang, T.: Observations: Cryosphere, in *Climate Change 2013: The Physical Science*



- 460 Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, pp. 317–382, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- Velicogna, I., Mohajerani, Y., A. G., Landerer, F., Mouginot, J., Noel, B., Rignot, E., Sutterley, T., Broeke, M. van den, Wessem, M. van and Wiese, D.: Continuity of Ice Sheet Mass Loss in Greenland and Antarctica From the GRACE and GRACE Follow-On Missions, *Geophysical Research Letters*, 47(8), e2020GL087291, doi:10.1029/2020GL087291, 2020.
- 465 Vellinga, M. and Wood, R. A.: Global Climatic Impacts of a Collapse of the Atlantic Thermohaline Circulation, *Climatic Change*, 54(3), 251–267, doi:10.1023/A:1016168827653, 2002.
- Vieli, A., Payne, A. J., Shepherd, A. and Du, Z.: Causes of pre-collapse changes of the Larsen B ice shelf: Numerical modelling and assimilation of satellite observations, *Earth and Planetary Science Letters*, 259(3), 297–306, doi:10.1016/j.epsl.2007.04.050, 2007.
- 470 Vihma, T.: Effects of Arctic Sea Ice Decline on Weather and Climate: A Review, *Surv Geophys*, 35(5), 1175–1214, doi:10.1007/s10712-014-9284-0, 2014.
- Vitousek, S., Barnard, P. L., Fletcher, C. H., Frazer, N., Erikson, L. and Storlazzi, C. D.: Doubling of coastal flooding frequency within decades due to sea-level rise, *Sci Rep*, 7(1), 1–9, doi:10.1038/s41598-017-01362-7, 2017.
- WCRP Global Sea Level Budget Group: Global sea-level budget 1993–present, *Earth System Science Data*, 10(3), 1551–1590, doi:https://doi.org/10.5194/essd-10-1551-2018, 2018.
- 475 Weertman, J.: Stability of the Junction of an Ice Sheet and an Ice Shelf, *Journal of Glaciology*, 13(67), 3–11, doi:10.3189/S0022143000023327, 1974.
- van Wessem, J. M., Berg, W. J. van de, Noël, B. P. Y., Meijgaard, E. van, Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J. T. M., Lhermitte, S., Ligtenberg, S. R. M., Medley, B., Reijmer, C. H., Tricht, K. van, Trusel, L. D., Ulf, L. H. van, Wouters, B., Wuite, J. and Broeke, M. R. van den: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 – Part 2: Antarctica (1979–2016), *The Cryosphere*, 12(4), 1479–1498, doi:https://doi.org/10.5194/tc-12-1479-2018, 2018.
- 480 Worby, A. P., Geiger, C. A., Paget, M. J., Woert, M. L. V., Ackley, S. F. and DeLiberty, T. L.: Thickness distribution of Antarctic sea ice, *Journal of Geophysical Research: Oceans*, 113(C5), doi:10.1029/2007JC004254, 2008.
- 485 Wouters, B., Gardner, A. S. and Moholdt, G.: Global Glacier Mass Loss During the GRACE Satellite Mission (2002-2016), *Front. Earth Sci.*, 7, doi:10.3389/feart.2019.00096, 2019.
- Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlström, A. P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L. N., Cáceres, B. E., Casassa, G., Cobos, G., Dávila, L. R., Granados, H. D., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Hagen, J. O., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V., Severskiy, I., Sigurdsson, O., Soruco, A., Usabaliyev, R. and Vincent, C.: Historically unprecedented global glacier decline in the early 21st century, *Journal of Glaciology*, 61(228), 745–762, doi:10.3189/2015JoG15J017, 2015.
- 490 Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S. and Cogley, J. G.: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568(7752), 382–386, doi:10.1038/s41586-019-1071-0, 2019.



Zemp, M., Huss, M., Eckert, N., Thibert, E., Paul, F., Nussbaumer, S. U. and Gärtner-Roer, I.: Brief communication: Ad hoc estimation of glacier contributions to sea-level rise from the latest glaciological observations, *The Cryosphere*, 14(3), 1043–1050, doi:<https://doi.org/10.5194/tc-14-1043-2020>, 2020.

500 Zhang, J. and Rothrock, D. A.: Modeling Global Sea Ice with a Thickness and Enthalpy Distribution Model in Generalized Curvilinear Coordinates, *Mon. Wea. Rev.*, 131(5), 845–861, doi:10.1175/1520-0493(2003)131<0845:MGSIIWA>2.0.CO;2, 2003.

Zhang, L., Delworth, T. L., Cooke, W. and Yang, X.: Natural variability of Southern Ocean convection as a driver of observed climate trends, *Nature Climate Change*, 9(1), 59–65, doi:10.1038/s41558-018-0350-3, 2019.

505




	1980s	1990s*	2000s	2010s**	1994-2017
Arctic Sea Ice	-156 ± 88	-298 ± 88	-360 ± 88	-94 ± 119	-329 ± 44
Antarctic Sea Ice	+196 ± 67	-27 ± 67	+71 ± 67	-83 ± 75	-40 ± 29
Ice Shelves Calving	-140 ± 15	-125 ± 25	-176 ± 57	-250 ± 68	-155 ± 36
Ice Shelves Thinning	-	-19 ± 52	-233 ± 57	-53 ± 71	-129 ± 39
<i>Total Floating Ice</i>	-	-469 ± 125	-698 ± 137	-480 ± 172	-653 ± 75
Antarctic	-	-55 ± 38	-78 ± 37	-206 ± 47	-111 ± 24
Greenland	-	-34 ± 24	-166 ± 21	-247 ± 23	-166 ± 14
Glaciers	-62 ± 68	-206 ± 64	-250 ± 64	-346 ± 67	-270 ± 42
<i>Total Grounded Ice</i>	-	-296 ± 79	-494 ± 77	-798 ± 85	-547 ± 50
Total	-	-764 ± 147	-1191 ± 157	-1278 ± 192	-1200 ± 90

*1990s: the decade is not entirely surveyed but starts from 1994 for Ice Shelves Thinning, and from 1993 for Antarctica and from 1992 for Greenland

**2010s: the decade is not entirely surveyed but covers up to 2016 for the Antarctic ice sheet, up to 2017 for Greenland, and up to 2019 for sea ice, glaciers and ice shelf calving.

Table 1 Average mass change rates (Gt yr^{-1}) of the different global ice components, total floating ice, total grounded ice and global total per decade and over the common period 1994-2017.




510


515

520


Figure 1 Average rate of ice thickness change in the (left) southern and (right) northern hemispheres. Changes in Antarctic (1992-2017) and Greenland ice sheet (1992-2018) thickness were estimated using repeat satellite altimetry following the methods of (Shepherd et al., 2019). Sea ice thickness trends between 1990 and 2019 are determined from numerical sea ice and ocean modelling (Zhang and Rothrock, 2003), as well as the average minimum of sea ice extent in February (Antarctic) and September (Arctic) (purple lines) for each decade during the same period. Glacier thickness change between 1992 and 2018 for glacier regions defined in the Randolph Glacier Inventory (RGI Consortium, 2017) (black boundaries) are from mass change estimates (Braun et al., 2019; Foresta et al., 2016; Jakob et al., 2020; Tepes et al., 2020; Wouters et al., 2019; Zemp et al., 2019) which have been converted to a thickness change assuming an ice density of 850 kg/m³. The black circle at the south pole indicates the southern limit of the orbit of ERS and ENVISAT satellite altimeters, which were in operation between 1992 and 2010. The area between 81.5° and 88° S has been covered by CryoSat-2, which launched in 2010.



525 Figure 2 (a) Cumulative mass change for glacier regions (Gt) between 1962 and 2019. Outlines of the glacier regions (RGI 6.0) are
 marked by black lines and glacierized areas are indicated in orange: ACN, Arctic Canada North (area 105,110 km²); ACS, Arctic
 530 Canada South (40,888 km²); ALA, Alaska (86,725 km²); ANT, Antarctic and Subantarctic (132,867 km²); CAU, Caucasus and
 Middle East (1,307 km²); CEU, Central Europe (2,092 km²); GRL, Greenland (89,717 km²); HMA, High Mountain Asia (97,606
 km²); ISL, Iceland (11,059 km²); NZL, New Zealand (1,161 km²); RUA, Russian Arctic (51,591 km²); SAN, Southern Andes (29429
 km²); SCA, Scandinavia (2,949 km²); SJM, Svalbard and Jan Mayen (33,958 km²); TRP, Low Latitudes (2,341 km²); WNA,
 535 Western Canada and USA (14,524 km²). (b) Glacier rate of mass change (Gt yr⁻¹) in regions where estimates from different
 techniques are available, including satellite altimetry (Foresta et al., 2016; Jakob et al., 2020; Tepes et al., 2020), extrapolation of
 in-situ glaciological and geodetic data (Zemp et al., 2019, 2020), satellite gravimetry (Wouters et al., 2019), and satellite InSAR
 (Braun et al., 2019) over the period 2010-2015. The reconciled estimate (calculated as the average of the estimates available in a
 given region and year) is shown in grey.



540 **Figure 3** Change in Antarctic ice shelf barrier position (left) and thickness (right) over time. Barrier positions are derived from episodic satellite imagery (Cook and Vaughan, 2010), and barrier thicknesses are derived from airborne ice penetrating radar (light grey lines) and satellite radar altimetry (Fretwell et al., 2013).



545 **Figure 4** Global ice mass change between 1994 and 2017 partitioned into the different floating (blues) and grounded (purples) components. The sea-level contribution due to the loss of grounded ice from Antarctica, Greenland and mountain glaciers is shown in the y-axis on the right hand side.