



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

An ecological momentary assessment study of the role of emotional dysregulation in co-occurring ADHD and internalising symptoms in adulthood

Citation for published version:

Murray, AL, Wong, S-C, Obsuth, I, Rhodes, S, Eisner, M & Ribeaud, D 2020, 'An ecological momentary assessment study of the role of emotional dysregulation in co-occurring ADHD and internalising symptoms in adulthood', *Journal of Affective Disorders (JAD)*. <https://doi.org/10.1016/j.jad.2020.11.086>

Digital Object Identifier (DOI):

[10.1016/j.jad.2020.11.086](https://doi.org/10.1016/j.jad.2020.11.086)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Affective Disorders (JAD)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



An ecological momentary assessment study of the role of emotional dysregulation in co-occurring ADHD and internalising symptoms in adulthood

Aja Louise Murray¹, Siu-Ching Wong², Ingrid Obsuth², Sinead Rhodes³, Manuel Eisner^{4,5},
Denis Ribeaud⁵

¹Department of Psychology, University of Edinburgh

²Clinical Psychology Department, University of Edinburgh

³Centre for Clinical Brain Sciences, University of Edinburgh

⁴Institute of Criminology, University of Cambridge

⁵Jacobs Center for Productive Youth Development, University of Zurich

Corresponding author at Aja.Murray@ed.ac.uk; 7 George Square, University of Edinburgh
Department of Psychology, Edinburgh, UK, EH8 9JZ

Abstract

Previous ecological momentary assessment (EMA) research in children has suggested that emotional dysregulation problems are commonly associated with ADHD symptoms and may help explain their relatively strong co-occurrence with anxiety and depression (collectively referred to as ‘internalising problems’); however, this has yet to be replicated for adults. In this study, we used data from a n=260 longitudinal cohort and EMA study, to evaluate the hypothesis that emotional dysregulation mediates the association between ADHD symptoms and internalising problems in emerging adulthood. Emotional dysregulation was based on affect data collected in near real time and in ecological context over a 14-day period, providing a measure of emotional lability in the context of participants’ daily lives. Cross-sectional mediation was tested using structural equation modelling. Emotional lability significantly mediated the association between ADHD symptoms and internalising problems. Results suggest that interventions that address the emotional dysregulation aspects of ADHD are likely to be beneficial for preventing and managing secondary internalising symptoms.

Keywords: attention-deficit/hyperactivity disorder; emotional dysregulation; depression; anxiety; internalising problems; ecological momentary assessment

Emotional lability, defined as sudden and marked shifts in emotion that are inappropriate to setting and/or developmental stage, is recognised as a common associated feature of ADHD symptoms (Posner et al., 2014; Shaw et al., 2015). It is assumed to reflect emotional dysregulation problems and has been proposed to play a significant role in many of the difficulties associated with ADHD in childhood, including anxiety and depression ; collectively referred to as internalising symptoms (Anastopoulos et al., 2011; Rosen & Factor, 2015). However, despite the fact that ADHD symptoms and their sequelae remain common in adulthood (Kessler et al., 2006; Simon et al., 2009), less is known about the role of emotional lability in co-occurring internalising symptoms during this developmental phase, with studies investigating its role using emotion data collected in the context of daily life particularly lacking. In this study, we use data from the community-ascertained n=260 D2M study (an ecological momentary assessment sub-study of the Zurich Project on Social Development from Childhood to Adulthood; z-proso) to evaluate whether emotional dysregulation, as indicated by higher levels of emotional lability, mediates the association between ADHD symptoms and a common sequela, namely internalising problems in emerging adulthood.

Emotional dysregulation is commonly associated with ADHD symptoms in childhood. Estimates vary but suggest that approximately 25% children with ADHD show severe emotional dysregulation, and another 50% show moderate emotional dysregulation problems (Shaw et al., 2015; Sobanski et al., 2010). Such is its prevalence in individuals with ADHD that there have been ongoing debates concerning whether emotional dysregulation should be considered a core rather than associated feature of the disorder (Faraone et al., 2019; Caroline Skirrow & Asherson, 2013). While it previously served as a marker for ADHD in older editions of the Diagnostic and Statistical Manual of Mental Disorders (DSM),

in the most recent editions, only behavioral dysregulation is recognised as a core feature of ADHD (American Psychiatric Association, 2013).

The significance of emotional dysregulation in ADHD is also in its proposed role in many of the behavioural, emotional and psychosocial sequelae of ADHD, including internalising problems (Anastopoulos et al., 2011; Rosen & Factor, 2015). Both anxiety and depression are frequently associated with ADHD symptoms (Angold et al., 1999; Biederman et al., 2008; Jarrett & Ollendick, 2008; Schatz & Rostain, 2006). A commonly advanced explanation is that the psychosocial problems associated with ADHD, such as peer victimisation/rejection and academic/occupational failures undermine self-esteem and increase the risk of developing anxiety and/or depression (Roy et al., 2015; Schatz & Rostain, 2006; Yip et al., 2013). However, there is preliminary evidence that issues with emotional regulation also play an important role in co-occurring internalising problems (Anastopoulos et al., 2011; Leaberry et al., 2017; Rosen & Factor, 2015).

Numerous measures have been developed to assess emotional dysregulation (e.g., see Faraone et al., 2019 for a review); however, those that capture emotional dysregulation in the context of daily life (e.g., Helfer et al., 2019) are especially valuable in illuminating its role in internalising problems. In particular, ecological momentary assessment (EMA) studies can provide measures of emotional functioning in the context of real life and ‘in the moment’, that are more ecologically valid and that are considerably less vulnerable to issues such as recall bias that commonly affect questionnaire measures (e.g., Bolger & Laurenceau, 2013). Ecological momentary assessment studies involve administering small sets of items frequently (e.g., 3-10 times a day over several days or weeks), often via participants’ own smartphones, in order to capture momentary affective states, cognitions, behaviours, and experiences as they occur in a real-life context. There is a growing literature on using EMA methodologies to measure manifestations of emotional dysregulation; however, their

application in ADHD research has been relatively limited (Miguel-Fernandez et al., 2018). Only a handful of studies have employed this methodology to evaluate its role in internalising symptoms associated with ADHD symptoms (Rosen et al., 2013, 2015) and all studies to date have explored these links in child samples.

Leaberry et al. (2017) used EMA to evaluate the relations between emotional lability and comorbid internalising problems in $n=58$ 8-12 year olds. They found that participants with ADHD and a comorbid internalising disorder showed more EMA-derived emotional lability than those with ADHD alone. However, Rosen et al., (2015), in a sample of $n=102$ 8-12 year olds ($n=56$ with ADHD) found that emotional lability was associated with internalising problems but not directly with ADHD diagnostic status. In another study, in a sample of $n=27$ 8-12 year-olds with ADHD, Rosen and Factor (2015) found that EMA-derived emotional impulsivity was correlated with internalising problems at $r=.54$ ($p<.001$) based on parent reports. Thus, further work is required to clarify the role of emotional lability in the relation between ADHD and internalising symptoms. Furthermore, given the risk of Berkson's bias (i.e., the over-estimation of symptom co-occurrence) in clinical samples (e.g., Pearce & Richiardi, 2014), studies utilising community-ascertained samples will be an important complement to the clinically diagnosed studies used in previous studies. Finally, although there have been informative EMA studies of emotion dysregulation in adults (Skirrow et al., 2014), none have yet addressed its role in co-occurring internalising problems. Internalising problems remain a common issue associated with ADHD symptoms in adulthood (e.g., Krone & Newcorn, 2015), that may be partially explained by the emotional dysregulation features of ADHD. In this study we thus evaluated the hypothesis that emotional dysregulation mediates the relation between ADHD symptoms and internalising problems in adults using a measure of emotional dysregulation based on an EMA design that captures emotional dysregulation as it occurs in the context of daily life.

Given that ADHD symptoms and internalising symptoms show meaningful variation both above and below clinical cut-offs, we used a general population sample.

Methods

Ethics

Ethical approval was obtained from the Ethics Committee from the Faculty of Arts and Social Sciences of the University of Zurich (ECFASSUZH). Participants provided informed consent before participating. The study information instructed participants not to answer prompts during times where it would not be safe to do so (e.g., while driving).

Procedure

Data come from the ‘Decade-to-minutes’ (D2M) study which is a combined longitudinal and EMA study. Specifically, the D2M sample is a subsample of the z-proso longitudinal study (e.g., M. Eisner & Ribeaud, 2007; N. L. Eisner et al., 2018). Data were collected at the age 20 measurement wave of z-proso. The main age 20 z-proso data collection consisted of a questionnaire that included measures of ADHD symptoms and internalising problems. These were administered in the context of a range of other constructs related to psychosocial development, including substance use, behavioural problems and criminality, sexual experiences, and victimisation. A separate EMA survey was conducted to measure emotional symptoms in the context of daily life.

The EMA survey was conducted over a 14-day period with measurements occurring four times a day at quasi-random intervals between 10am and 10pm. Measures were administered via an application provided by *LifeDataCorp LLC*, which participants downloaded on their smartphone device. At each measurement instance, participants received a notification that directed them to a brief set of items. Incentives were scaled to level of compliance, with participants receiving a maximum of 50 CHF for a response rate of at least

70% over both week 1 and week 2. Fieldwork was carried out by Decision Science Laboratory (DeSciL) at the ETH Zurich: <https://www.descil.ethz.ch/>. As well as the measures utilised in the current study, the survey included measures of momentary substance use, aggression, provocations, stress, and context (i.e., who a person was with and what they were doing at the time of responding to a prompt).

Participants

D2M participants were a $n=260$ ($n=100$ male) subsample of the z-proso study: <https://www.jacobscenter.uzh.ch/en/research/zproso/aboutus.html> with a median age of 20. The z-proso sample is a school-year cohort sample, first recruited when entering first grade (median age 7) and followed up at ages 8,9,10,11,12,13,15,17 and 20. All participants entering first grade in the year 2004 from across 56 schools in the Zurich, Switzerland area at age 7 were invited to participate. At the latest wave, $n=1620$ had provided data for at least one measurement wave (from a target sample of $n=1675$). Previous studies have suggested that the sample is mostly representative of the underlying same-aged population (Eisner et al., 2018). The main exception was an initial slight under-representation of participants whose parents did not speak German as their first language; however, this effect diminished after consent procedures switched from being parent-based to participant-based.

An inclusion criterion for D2M was that z-proso participants were required to have access to a smartphone running iOS or android; however, as this accounts for 99.6% of all smartphones (Thai & Page-Gould, 2017), this was not thought to be a significant biasing factor. Most importantly in the context of the current study, levels of ADHD symptoms ($p=.17$) and internalising symptoms ($p=.20$) did not differ between D2M and non-D2M participants i.e., those participating in the broader z-proso study but not the D2M sub-study. We did not impose compliance thresholds as doing so is likely to introduce missingness

related to the constructs of interest, such as internalising problems (Silvia et al., 2013) and thus bias parameter estimates by creating non-random missingness (Rubin, 1976).

Measures

ADHD symptoms

ADHD symptoms were measured using an adapted self-reported version of the ADHD subscale from the Social Behavior Questionnaire (SBQ; Tremblay et al., 1991), modified to be more developmentally appropriate for emerging adults. This was collected as part of the main age 20 z-proso survey. The scale included nine items measuring both inattention and hyperactivity/impulsivity symptoms, including restlessness, being easily distracted, having difficulties concentrating, acting without thinking, forgetfulness, difficulties paying attention, being hectic and fidgety, being unable to settle to anything for long, and internal feelings of restlessness. Responses were recorded on a five-point Likert-type scale from *never* to *very often*. The psychometric properties of versions of this questionnaire have been examined in previous studies, supporting its reliability, structural validity, and criterion validity of its scores (Murray, Eisner, Obsuth, et al., 2017; Murray, Eisner, & Ribeaud, 2017; Murray, Obsuth, et al., 2017; Tremblay et al., 1991). In the current sample, omega reliability (McDonald, 1999) was good at .87.

Internalising problems

Internalising problems were also measured using the SBQ (Tremblay et al., 1991) administered as part of the main age 20 wave of data collection. The internalising subscale of the SBQ includes 14 items measuring both anxiety and depression. Items refer to symptoms including boredom, anhedonia, crying frequently, lethargy, feeling fearful, feeling unhappy, feeling as if one is doing everything wrong, loneliness, insomnia, feeling useless, feeling sad without reason, worrying, hating oneself, and self-harm. Responses were recorded on a five-

point Likert-type scale from *never* to *very often*. Previous research using similar versions of the SBQ has also supported the reliability and validity of the internalising scores (Murray, Eisner, Obsuth, et al., 2017; Murray, Eisner, & Ribeaud, 2017; Murray, Obsuth, et al., 2017; Tremblay et al., 1991). In the current sample, omega reliability was good at .92.

Emotional lability

Emotional lability scores were derived from the ecological momentary assessment component of D2M using an abbreviated version of the negative affect schedule of the *Positive Affect Negative Affect Schedule Expanded* (PANAS-X; Watson & Clark, 1999). Specifically, data were collected on the following affective states: afraid, scared, nervous, hostile, guilty, ashamed, and distressed. Participants were asked to record on a five-point scale from *Very slightly or not at all* to *Extremely*. The prompt-wise response rates for these variables was 67%. Mean squared successive differences (MSSDs; Von Neumann et al., 1941) were calculated for each of the affect items separately based on the data available for each individual, to obtain a measure of the variability in each affective state for each individual over the ecological momentary assessment period. These were then used as indicators in a confirmatory factor analysis (described in the *Statistical Procedure* section) to derive a measure of overall emotional lability.

Statistical Procedure

A structural equation model was estimated to test the mediating role of emotional lability in the relations between ADHD symptoms and internalising problems. Emotional lability, ADHD, and internalising problems were all specified as latent variables. The indicators for the emotional lability latent variable were the above-described MSSDs for each negative affect item. The indicators for ADHD, internalising problems were the above-

described SBQ items. Scaling and identification were achieved by fixing the loading for one reference indicator per latent factor to one.

A mediation model was specified with the ADHD latent variable as the predictor, the emotional lability latent variable as the mediator, and the internalising latent variable as the outcome. Direct effects of ADHD on the outcome were also modelled. The model is summarised in Figure 1. The statistical significance of the indirect effects of ADHD internalising problems was evaluated using the bootstrapped confidence intervals for 1) the product of the regression of internalising problems on emotional lability and the regression of emotional lability on ADHD. Missingness was dealt with using full information maximum likelihood estimation. All models were fit using lavaan (Rosseel, 2012) in R Statistical software.

Results

The initial model fit did not fit well according to incremental fit indices, though it fit reasonably well according to SRMR and RMSEA (CFI=.83, TLI=.84 , RMSEA=.08 , SRMR=.07), Modification indices and expected parameter changes suggested that residual covariances between some of the emotional lability items and a cross-loading of the internalising factor on an ADHD item would improve fit. The cross-loading was included to reduce the risk that the latent internalising-ADHD association would be inflated. The residual covariances were also included for the emotional lability items as they pertained to lability in emotional states that could feasibly show excess correlations, over and above a generalised emotional lability tendency.

The modified structural equation model fit reasonably well (CFI=.91, TLI=.90, RMSEA=.06, SRMR=.06). Fully standardised parameter estimates are provided in Figure 1 (residual variances and covariances and factor variances are not shown for visual clarity). All

parameters were significant at $p < .05$. Results show that there was a significant direct effect of ADHD symptoms on internalising problems ($r = .54$), and a significant effect of emotional lability on internalising problems ($r = .27$) and a significant effect of ADHD symptoms on emotional lability ($r = .37$). Based on bootstrapped confidence intervals, emotional lability significantly mediated the association between ADHD symptoms and internalising problems ($b = .07$, 95% CI = .03-.14; standardised $b = .10$). The mediation was partial and accounted for 16% of the total effect of ADHD symptoms on internalising problems.

Discussion

Emotional dysregulation issues have previously been shown to be associated with ADHD symptoms and have been proposed to contribute to the tendency for individuals with ADHD symptoms to experience co-occurring anxiety and depression (Rosen & Factor, 2015). In this study we evaluated this hypothesis in an $n = 260$ sample of community-ascertained adults, using EMA to obtain measures of emotional dysregulation as it occurs in near real time and in the context of daily life. Our results suggest that emotional dysregulation partially mediates the association between ADHD symptoms and internalising problems.

Our findings are consistent with previous EMA studies in children which have shown that emotional dysregulation problems are associated with higher levels of internalising problems (Leaberry et al., 2017; Rosen & Factor, 2015). However, despite the continuing prevalence of ADHD symptoms and co-occurring internalising problems in adults, equivalent studies in adults have not been conducted prior to the current study. Our results thus provide important evidence that emotional dysregulation remains an important factor in the co-occurrence of anxiety/depression with ADHD symptoms in adulthood.

Our results suggest that interventions targeting emotional dysregulation problems could help reduce the overall symptom burden associated with ADHD by preventing the development of secondary internalising problems. There is already some evidence that emotional dysregulation symptoms respond positively to standard ADHD treatments; however, the effects are not as large as for symptoms that are currently regarded as ‘core’ ADHD symptoms (Faraone et al., 2019; Posner et al., 2014). In a systematic review and meta-analysis of the effects of pharmacological treatments for ADHD in adults, the standardised effect sizes for methylphenidate, atomoxetine, and lisdexamfetamine were in only the small-to-moderate range (Lenzi et al., 2018). Thus, approaches more specifically tailored to the management of emotional dysregulation symptoms could have even stronger effects. There is some preliminary evidence showing positive effects of mindfulness-based interventions for emotional dysregulation in children with ADHD (Huguet et al., 2019). However, a range of therapeutic approaches could be relevant, including dialectal behaviour therapy, which includes emotion regulation skill components (Neacsiu et al., 2014).

As regards pharmacological interventions, it is not clear whether medications for internalising symptoms (antidepressants and mood stabilisers) are more promising for reducing emotional dysregulation and/or associated internalising problems in unmedicated adults with ADHD symptoms than ADHD medications themselves, especially considering the potential side effects of medications typically prescribed for anxiety and depression (Lenzi et al., 2018). Further research will be required to directly compare the impacts of these different types of medication on emotional dysregulation and internalising symptoms in the context of ADHD, as well as identify factors that may moderate treatment impact, such as ADHD symptom profiles (inattentive, hyperactive/impulsive or combined) and severity.

Interventions targeting emotional dysregulation in ADHD will also benefit from an improved understanding of which of the mechanisms underlying emotional control are

particularly impaired in ADHD. As several authors have previously noted, a multitude of processes could be involved, including both those involved in the generation of affective states and those involved in the management of existing affective states (Faraone et al., 2019; Posner et al., 2014).

Our findings also indicate that both ADHD symptoms and internalising problems are frequently associated with emotional dysregulation problems and thus support previous suggestions that adults presenting with emotional dysregulation issues be screened for ADHD (Skirrow & Asherson, 2013). They similarly suggest that patients presenting with either ADHD symptoms or internalising problems could benefit from being screened for emotional dysregulation issues: doing so could identify a potentially important source of functional impairment and help make better sense of symptoms. Given that smartphone use is embedded in the daily lives of many adults and that user-friendly EMA applications are becoming increasingly available (e.g., Hofmann & Patel, 2015; Thai & Page-Gould, 2017), it would be feasible to conduct these screens via similar methods as employed in the current study i.e., over a period of a couple of weeks or more, using participants' own smartphones. This would help to better capture emotional functioning as it occurs in the context of a person's daily life, as compared to using questionnaire measures which have more limited ecological validity and rely on retrospective recall.

However, the fact that the mediation of the ADHD-internalising problems relation by emotional dysregulation was only partial suggests that there are additional mechanisms linking ADHD symptoms to internalising problems. Several such mechanisms have previously been proposed. For example, it has been suggested that the negative psychosocial sequelae of ADHD can foster a negative self-concept, placing an individual at greater risk of developing anxiety or depression (Schatz & Rostain, 2006). Consistent with this, several studies have found parent or peer problems to be mediators of the ADHD-internalising

associations (Humphreys et al., 2013; Ostrander & Herman, 2006; Roy et al., 2015; Yip et al., 2013). For example, Yip et al. (2013) found that peer problems mediated the association between attention problems and anxiety, while Humphreys et al. (2013) found that the ADHD-depression relation was mediated by both peer and parent-child problems. It is also likely that the ADHD-internalising problems association at least partly reflects genetic, neurobiological, and cognitive factors that are common to both ADHD symptoms and internalising problems (e.g., Levy, 2004). For example, previous research has suggested that young people with depression and young people with ADHD have similar memory and executive functioning deficits (Matthews et al., 2008; Rhodes et al., 2012). Whether these factors also help account for the shared emotion dysregulation issues between internalising problems and ADHD is not yet clear.

Future Directions

While emotional dysregulation may appear to have particular relevance for internalising problems, where emotional issues are prominent, it is likely that the impairing effects of emotional dysregulation are much broader in terms of ADHD sequelae. Future research could assess the mediating role of EMA-derived measures of emotional dysregulation in the relation between ADHD symptoms and other adulthood outcomes robustly associated with ADHD symptoms, e.g., substance use, intimate partner problems, and behaviour problems (Buitelaar et al., 2015; Graziano et al., 2015; Lee et al., 2011). Further illumination of the relations between emotional dysregulation and cognitive factors with established relations to ADHD, such as mind-wandering and neuropsychological impairments will also be valuable. It is not clear, for example, whether cognitive factors in ADHD underpin emotional dysregulation, or are independent or interactive factors which – together with emotional dysregulation – contribute to the risk of outcomes such as internalising problems and other sequelae. The development and evaluation of interventions

for emotion dysregulation in the context of ADHD symptoms also represents an important future direction.

Limitations

It is important to consider the limitations of the current study. First, while part of a longitudinal study, the measures available in the current study allowed only cross-sectional mediation tests. Longitudinal EMA studies remain rare but as more longitudinal EMA data becomes available, future replications using repeated measures of ADHD symptoms, emotional dysregulation, and internalising problems will be valuable to provide a longitudinal test of our hypotheses. Second, while our sample did not differ from its (same-aged population-representative parent sample) on ADHD or internalising problems, it was not a random draw from this sample. Future replications in population-representative samples will be helpful to ensure the generalisability of results.

Conclusions

Emotional dysregulation is associated with both ADHD symptoms and internalising problems and partially mediates their association. Interventions targeting emotional dysregulation could, therefore, be beneficial for reducing the development of anxiety and depression secondary to ADHD symptoms.

References

- Anastopoulos, A. D., Smith, T. F., Garrett, M. E., Morrissey-Kane, E., Schatz, N. K., Sommer, J. L., Kollins, S. H., & Ashley-Koch, A. (2011). Self-regulation of emotion, functional impairment, and comorbidity among children with AD/HD. *Journal of Attention Disorders, 15*(7), 583–592.
- Angold, A., Costello, E. J., & Erkanli, A. (1999). Comorbidity. *The Journal of Child Psychology and Psychiatry and Allied Disciplines, 40*(1), 57–87.
- Association, A. P. (2013). *Diagnostic and statistical manual of mental disorders (DSM-5®)*. American Psychiatric Pub.
- Biederman, J., Ball, S. W., Monuteaux, M. C., Mick, E., Spencer, T. J., McCreary, M., Cote, M., & Faraone, S. V. (2008). New insights into the comorbidity between ADHD and major depression in adolescent and young adult females. *Journal of the American Academy of Child, 47*(4), 426–434. <https://doi.org/10.1097/CHI.0b013e31816429d3>
- Bolger, N., & Laurenceau, J.-P. (2013). *Intensive longitudinal methods: An introduction to diary and experience sampling research*. Guilford Press.
- Buitelaar, N. J., Posthumus, J. A., & Buitelaar, J. K. (2015). ADHD in childhood and/or adulthood as a risk factor for domestic violence or intimate partner violence: A systematic review. *Journal of Attention Disorders, 1087054715587099*.
- Eisner, M., & Ribeaud, D. (2007). Conducting a criminological survey in a culturally diverse context: Lessons from the Zurich Project on the Social Development of Children. *European Journal of Criminology, 4*(3), 271–298.
- Eisner, N. L., Murray, A. L., Eisner, M., & Ribeaud, D. (2018). A practical guide to the analysis of non-response and attrition in longitudinal research using a real data example. *International Journal of Behavioral Development, 0165025418797004*.

- Faraone, S. V., Rostain, A. L., Blader, J., Busch, B., Childress, A. C., Connor, D. F., & Newcorn, J. H. (2019). Practitioner Review: Emotional dysregulation in attention-deficit/hyperactivity disorder—implications for clinical recognition and intervention. *Journal of Child Psychology and Psychiatry*, *60*(2), 133–150.
- Graziano, P. A., Reid, A., Slavec, J., Paneto, A., McNamara, J. P., & Geffken, G. R. (2015). ADHD symptomatology and risky health, driving, and financial behaviors in college: The mediating role of sensation seeking and effortful control. *Journal of Attention Disorders*, *19*(3), 179–190.
- Helfer, B., Cooper, R. E., Bozhilova, N., Maltezos, S., Kuntsi, J., & Asherson, P. (2019). The effects of emotional lability, mind wandering and sleep quality on ADHD symptom severity in adults with ADHD. *European Psychiatry*, *55*, 45–51.
- Hofmann, W., & Patel, P. V. (2015). SurveySignal: A convenient solution for experience sampling research using participants' own smartphones. *Social Science Computer Review*, *33*(2), 235–253.
- Huguet, A., Eguren, J. I., Miguel-Ruiz, D., Vallés, X. V., & Alda, J. A. (2019). Deficient Emotional Self-Regulation in Children with Attention Deficit Hyperactivity Disorder: Mindfulness as a Useful Treatment Modality. *Journal of Developmental & Behavioral Pediatrics*, *40*(6), 425–431.
- Humphreys, K. L., Katz, S. J., Lee, S. S., Hammen, C., Brennan, P. A., & Najman, J. M. (2013). The association of ADHD and depression: Mediation by peer problems and parent–child difficulties in two complementary samples. *Journal of Abnormal Psychology*, *122*(3), 854.
- Jarrett, M. A., & Ollendick, T. H. (2008). A conceptual review of the comorbidity of attention-deficit/hyperactivity disorder and anxiety: Implications for future research and practice. *Clinical Psychology Review*, *28*(7), 1266–1280.

- Kessler, R. C., Adler, L., Barkley, R., Biederman, J., Conners, C. K., Demler, O., Faraone, S. V., Greenhill, L. L., Howes, M. J., & Secnik, K. (2006). The prevalence and correlates of adult ADHD in the United States: Results from the National Comorbidity Survey Replication. *American Journal of Psychiatry*, *163*(4), 716–723.
- Krone, B., & Newcorn, J. H. (2015). Comorbidity of ADHD and anxiety disorders. *Attention-Deficit Hyperactivity Disorder in Adults and Children*, *98*.
- Leaberry, K. D., Rosen, P. J., Fogleman, N. D., Walerius, D. M., & Slaughter, K. E. (2017). Comorbid internalizing and externalizing disorders predict lability of negative emotions among children With ADHD. *Journal of Attention Disorders*, 1087054717734647.
- Lee, S. S., Humphreys, K. L., Flory, K., Liu, R., & Glass, K. (2011). Prospective association of childhood attention-deficit/hyperactivity disorder (ADHD) and substance use and abuse/dependence: A meta-analytic review. *Clinical Psychology Review*, *31*(3), 328–341.
- Lenzi, F., Cortese, S., Harris, J., & Masi, G. (2018). Pharmacotherapy of emotional dysregulation in adults with ADHD: A systematic review and meta-analysis. *Neuroscience and Biobehavioral Reviews*, 359–367.
<https://doi.org/10.1016/j.neubiorev.2017.08.010>
- Levy, F. (2004). Synaptic gating and ADHD: A biological theory of comorbidity of ADHD and anxiety. *Neuropsychopharmacology*, *29*(9), 1589–1596.
- Matthews, K., Coghill, D., & Rhodes, S. (2008). Neuropsychological functioning in depressed adolescent girls. *Journal of Affective Disorders*, *111*(1), 113–118.
- McDonald, R. P. (1999). *Test theory: A unified treatment*.
- Migueluez-Fernandez, C., de Leon, S. J., Baltasar-Tello, I., Peñuelas-Calvo, I., Barrigon, M. L., Capdevila, A. S., Delgado-Gómez, D., Baca-García, E., & Carballo, J. J. (2018).

- Evaluating attention-deficit/hyperactivity disorder using ecological momentary assessment: A systematic review. *ADHD Attention Deficit and Hyperactivity Disorders*, 10(4), 247–265.
- Murray, A. L., Eisner, M., Obsuth, I., & Ribeaud, D. (2017). Situating violent ideations within the landscape of mental health: Associations between violent ideations and dimensions of mental health. *Psychiatry Research*, 249, 70–77.
- Murray, A. L., Eisner, M., & Ribeaud, D. (2017). Can the Social Behavior Questionnaire help meet the need for dimensional, transdiagnostic measures of childhood and adolescent psychopathology? *European Journal of Psychological Assessment*.
- Murray, A. L., Obsuth, I., Eisner, M., & Ribeaud, D. (2017). Evaluating longitudinal invariance in dimensions of mental health across adolescence: An analysis of the Social Behavior Questionnaire. *Assessment*, 1073191117721741.
- Neacsiu, A. D., Bohus, M., & Linehan, M. M. (2014). *Dialectical behavior therapy: An intervention for emotion dysregulation*.
- Ostrander, R., & Herman, K. C. (2006). Potential cognitive, parenting, and developmental mediators of the relationship between ADHD and depression. *Journal of Consulting and Clinical Psychology*, 74(1), 89–98. <https://doi.org/10.1037/0022-006X.74.1.89>
- Pearce, N., & Richiardi, L. (2014). Commentary: Three worlds collide: Berkson's bias, selection bias and collider bias. *International Journal of Epidemiology*, 43(2), 521–524.
- Posner, J., Kass, E., & Hulvershorn, L. (2014). Using stimulants to treat ADHD-related emotional lability. *Current Psychiatry Reports*, 16(10), 478.
- Rhodes, S. M., Park, J., Seth, S., & Coghill, D. R. (2012). A comprehensive investigation of memory impairment in attention deficit hyperactivity disorder and oppositional defiant disorder. *Journal of Child Psychology and Psychiatry*, 53(2), 128–137.


- Rosen, P. J., Epstein, J. N., & Van Orden, G. (2013). I know it when I quantify it: Ecological momentary assessment and recurrence quantification analysis of emotion dysregulation in children with ADHD. *ADHD Attention Deficit and Hyperactivity Disorders, 5*(3), 283–294.
- Rosen, P. J., & Factor, P. I. (2015). Emotional impulsivity and emotional and behavioral difficulties among children with ADHD: An ecological momentary assessment study. *Journal of Attention Disorders, 19*(9), 779–793.
- Rosen, P. J., Walerius, D. M., Fogleman, N. D., & Factor, P. I. (2015). The association of emotional lability and emotional and behavioral difficulties among children with and without ADHD. *ADHD Attention Deficit and Hyperactivity Disorders, 7*(4), 281–294.
- Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling and more. Version 0.5–12 (BETA). *Journal of Statistical Software, 48*(2), 1–36.
- Roy, A., Hartman, C. A., Veenstra, R., & Oldehinkel, A. J. (2015). Peer dislike and victimisation in pathways from ADHD symptoms to depression. *European Child & Adolescent Psychiatry, 24*(8), 887–895.
- Rubin, D. B. (1976). Inference and missing data. *Biometrika, 63*(3), 581–592.
- Schatz, D. B., & Rostain, A. L. (2006). ADHD with comorbid anxiety: A review of the current literature. *Journal of Attention Disorders, 10*(2), 141–149.
- Shaw, P., Stringaris, A., Nigg, J., & Leibenluft, E. (2015). Emotion dysregulation in attention deficit hyperactivity disorder. *Focus, 14*(1), 127–144.
- Silvia, P. J., Kwapil, T. R., Eddington, K. M., & Brown, L. H. (2013). Missed beeps and missing data: Dispositional and situational predictors of nonresponse in experience sampling research. *Social Science Computer Review, 31*(4), 471–481.

- Simon, V., Czobor, P., Bálint, S., Mészáros, A., & Bitter, I. (2009). Prevalence and correlates of adult attention-deficit hyperactivity disorder: Meta-analysis. *The British Journal of Psychiatry, 194*(3), 204–211.
- Skirrow, C., Ebner-Priemer, U., Reinhard, I., Malliaris, Y., Kuntsi, J., & Asherson, P. (2014). Everyday emotional experience of adults with attention deficit hyperactivity disorder: Evidence for reactive and endogenous emotional lability. *Psychological Medicine, 44*(16), 3571–3583.
- Skirrow, Caroline, & Asherson, P. (2013). Emotional lability, comorbidity and impairment in adults with attention-deficit hyperactivity disorder. *Journal of Affective Disorders, 147*(1–3), 80–86.
- Sobanski, E., Banaschewski, T., Asherson, P., Buitelaar, J., Chen, W., Franke, B., Holtmann, M., Krumm, B., Sergeant, J., & Sonuga-Barke, E. (2010). Emotional lability in children and adolescents with attention deficit/hyperactivity disorder (ADHD): Clinical correlates and familial prevalence. *Journal of Child Psychology and Psychiatry, 51*(8), 915–923.
- Thai, S., & Page-Gould, E. (2017). *ExperienceSampler: An open-source scaffold for building smartphone apps for experience sampling*.
- Tremblay, R. E., Loeber, R., Gagnon, C., Charlebois, P., Larivee, S., & LeBlanc, M. (1991). Disruptive boys with stable and unstable high fighting behavior patterns during junior elementary school. *Journal of Abnormal Child Psychology, 19*(3), 285–300.
- Von Neumann, J., Kent, R. H., Bellinson, H. R., & Hart, B. t. (1941). The mean square successive difference. *The Annals of Mathematical Statistics, 12*(2), 153–162.
- Watson, D., & Clark, L. A. (1999). *The PANAS-X: Manual for the positive and negative affect schedule-expanded form*.

Yip, V. T., Ang, R. P., Ooi, Y. P., Fung, D. S., Mehrotra, K., Sung, M., & Lim, C. G. (2013). The association between attention problems and internalizing and externalizing problems: The mediating role of peer problems. *Child & Youth Care Forum, 42*, 571–584.

Figures

Figure 1: Structural equation model of ADHD, emotional lability and internalising problem relations



Note. Int= internalising problems; EL= emotional lability. All structural paths were significant at $p < .05$. Residual factors not shown for visual clarity.