



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Ocean Acidification and the Permo-Triassic Mass Extinction

Citation for published version:

Clarkson, MO, Kasemann, S, Wood, R, Lenton, TM, Daines, SJ, Richoz, S, Ohnemüller, F, Poulton, SW & Tipper, ET 2015, 'Ocean Acidification and the Permo-Triassic Mass Extinction', *Science*, vol. 348, no. 6321, pp. 229-232 . <https://doi.org/10.1126/science.aaa0193>

Digital Object Identifier (DOI):

[10.1126/science.aaa0193](https://doi.org/10.1126/science.aaa0193)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Science

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Title: Ocean Acidification and the Permo-Triassic Mass Extinction**

2 **Authors:** Clarkson, M.O.^{1*†}; Kasemann, S.A.²; Wood, R.¹; Lenton, T.M.³; Daines, S.J.³;
3 Richoz, S.⁴; Ohnemüller, F.²; Meixner, A.²; Poulton, S.W.⁵ and Tipper, E.T.⁶

4

5 **Affiliations:**

6 ¹ School of Geosciences, University of Edinburgh, West Mains Road, Edinburgh, EH9 3JW,
7 UK

8 ² Faculty of Geosciences and MARUM-Center for Marine Environmental Sciences,
9 University of Bremen, 28334 Bremen, Germany

10 ³ College of Life and Environmental Sciences, University of Exeter, Laver Building, North
11 Parks Road, Exeter, EX4 4QE, UK.

12 ⁴ Institute of Earth Sciences, NAWI Graz, University of Graz, Heinrichstraße 26, 8010 Graz,
13 Austria

14 ⁵ School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK

15 ⁶ Dept. of Earth Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EQ,
16 UK

17 *Correspondence to: matthew.clarkson@otago.ac.nz

18 †Current address: Department of Chemistry, University of Otago, Union Street, Dunedin,
19 9016, PO Box 56, New Zealand.

20

21 **Abstract:** Ocean acidification triggered by Siberian Trap volcanism was a possible kill
22 mechanism for the Permian Triassic Boundary (PTB) mass extinction, but direct evidence for
23 an acidification event is lacking. We present a high resolution seawater pH record across this
24 interval, utilizing boron isotope data combined with a quantitative modeling approach. In the
25 latest Permian, increased ocean alkalinity, primed the Earth system with a low level of

26 atmospheric CO₂ and a high ocean buffering capacity. The first phase of extinction was
27 coincident with a slow injection of carbon into the atmosphere and ocean pH remained stable.
28 During the second extinction pulse, however, a rapid and large injection of carbon caused an
29 abrupt acidification event that drove the preferential loss of heavily calcified marine biota.

30

31 **One Sentence Summary:** Ocean acidification caused the second phase of mass extinction in
32 the Permo-Triassic, due to a rapid and large injection of carbon which overwhelmed the
33 buffering capacity of the ocean.

34

35 **Main Text:** The Permian Triassic Boundary (PTB) mass extinction, at ~ 252 Ma, represents
36 the most catastrophic loss of biodiversity in geological history, and played a major role in
37 dictating the subsequent evolution of modern ecosystems (1). The PTB extinction event
38 spanned ~60 thousand years (2) and can be resolved into two distinct marine extinction
39 pulses (3). The first occurred in the latest Permian (Extinction Pulse 1; EP1) and was
40 followed by an interval of temporary recovery before the second pulse (EP2) which occurred
41 in the earliest Triassic. The direct cause of the mass extinction is widely debated with a
42 diverse range of overlapping mechanisms proposed, including widespread water column
43 anoxia (4), euxinia (5), global warming (6) and ocean acidification (7).

44 Models of PTB ocean acidification suggest that a massive, and rapid, release of CO₂ from
45 Siberian Trap volcanism, acidified the ocean (7). Indirect evidence for acidification comes
46 from the interpretation of faunal turnover records (3, 8), potential dissolution surfaces (9) and
47 Ca isotope data (7). A rapid input of carbon is also potentially recorded in the negative
48 carbon isotope excursion (CIE) that characterizes the PTB interval (10, 11). The
49 interpretation of these records is, however, debated (12-16), and is of great importance to
50 understanding the current threat of anthropogenically-driven ocean acidification (11).

51 To test the ocean acidification hypothesis we have constructed a proxy record of
52 ocean pH across the PTB, using the boron isotope composition of marine carbonates
53 ($\delta^{11}\text{B}_{\text{carb}}$) (17). We then employ a carbon cycle model (*supplementary online text*) to explore
54 ocean carbonate chemistry and pH scenarios that are consistent with our $\delta^{11}\text{B}$ data and
55 published records of carbon cycle disturbance and environmental conditions. Through this
56 combined geochemical, geological and modelling approach we are able to produce an
57 envelope that encompasses the most realistic range in pH, which then allows us to resolve
58 three distinct chronological phases of carbon cycle perturbation, each with very different
59 environmental consequences for the Late Permian-Early Triassic Earth system.

60 We analyzed boron and carbon isotope data from two complementary transects in a
61 shallow marine, open water carbonate succession from the United Arab Emirates (U.A.E.),
62 where depositional facies and $\delta^{13}\text{C}_{\text{carb}}$ are well constrained (18). During the PTB interval the
63 U.A.E. formed an expansive carbonate platform that remained connected to the central Neo-
64 Tethyan Ocean (15) (Fig 1A). Conodont stratigraphy and the distinct $\delta^{13}\text{C}_{\text{carb}}$ curve are used
65 to constrain the age model (17).

66 The run-up to PTB in the Tethys is characterized by two negative $\delta^{13}\text{C}$ excursions
67 interrupted by a short-term positive event (10). There is no consensus as to the cause of this
68 rebound event and so we instead focus on the broader $\delta^{13}\text{C}$ trend. Our $\delta^{13}\text{C}$ transect (Fig. 1B)
69 starts in the Changhsingian (Late Permian) with a gradual decreasing trend, interrupted by the
70 first negative shift in $\delta^{13}\text{C}$ at EP1 (at 53 m, ~251.96 Ma, Fig. 1B and 2). This is followed by
71 the minor positive 'rebound' event (at 54 m, ~251.95 Ma, Fig. 1B and 2) prior to the minima
72 of the second phase of the negative CIE (58-60 m, ~251.92 Ma, Figs. 1B and 2) that marks
73 the PTB itself. After the CIE minimum, $\delta^{13}\text{C}$ gradually increases to ~1.8‰ and remains
74 relatively stable during the earliest Triassic and across EP2.

75 Our boron isotope record shows a different pattern to the carbon isotope excursion. $\delta^{11}\text{B}$
76 is persistently low (Fig. 1C) at the start of our record during the late-Changhsingian, with an
77 average of $10.9 \pm 0.9\text{‰}$ (1σ). This is in agreement with $\delta^{11}\text{B}$ values (average of $10.6 \pm 0.6\text{‰}$,
78 1σ) reported for early-Permian brachiopods (19). Further up section (at ~40 m, ~252.04 Ma,
79 Fig. 1C), there is a stepped increase in $\delta^{11}\text{B}$ to $15.3 \pm 0.8\text{‰}$ (average $\pm 2\sigma$), and by
80 implication an increase in ocean pH of ~0.4-0.5 units (Fig. 2). $\delta^{11}\text{B}$ values then remain
81 relatively stable, scattering around $14.8 \pm 1.0\text{‰}$ (1σ) and implying variations within 0.1-0.2
82 pH units, into the Early Griesbachian (Early Triassic) and hence across EP1 and the period of
83 carbon cycle disturbance (Figs. 1 and 2).

84 After the $\delta^{13}\text{C}$ increase and stabilization (at ~85 m, ~251.88 Ma, Fig. 1), $\delta^{11}\text{B}$ begins
85 to decrease rapidly to $8.2 \pm 1.2\text{‰}$ (2σ), implying a sharp drop in pH of ~0.6-0.7 units. The
86 $\delta^{11}\text{B}$ minimum is coincident with the interval identified as EP2. This ocean acidification
87 event is short-lived (~10 thousand years) and $\delta^{11}\text{B}$ values quickly recover toward the more
88 alkaline values evident during EP1 (average of ~14‰).

89 The initial rise in ocean pH of ~0.4-0.5 units during the Late Permian (Fig. 2)
90 suggests a large increase in carbonate alkalinity (20). We are able to simulate the observed
91 rise in $\delta^{11}\text{B}$ and pH through different model combinations of increasing silicate weathering,
92 increased pyrite deposition (21), an increase in carbonate weathering, and a decrease in
93 shallow marine carbonate depositional area (supplementary online text). Both silicate
94 weathering and pyrite deposition result in a large drop in $p\text{CO}_2$ (and temperature) for a given
95 increase in pH and saturation state (Ω). There is no evidence for a large drop in $p\text{CO}_2$, and
96 independent proxy data indicate only a minor temperature decrease of a few degrees C during
97 the Changhsingian (22), suggesting that these mechanisms alone cannot explain the pH
98 increase (Fig. S5). Conversely, an increase in carbonate input or a reduction in rates of

99 carbonate deposition both result in increases in Ω , with a greater impact on pH per unit
100 decrease in $p\text{CO}_2$ and temperature (Fig. S6).

101 A decrease in carbonate sedimentation is consistent with the decrease in depositional
102 shelf area that occurred due to the 2nd order regression of the Late Permian (23). With the
103 added expansion of anoxia into shelf environments (24), this would effectively create both
104 bottom-up and top-down pressures to reduce the area of potential carbonate sedimentation.
105 Sea level fall also exposed carbonates to weathering (23), which would have further
106 augmented the alkalinity influx. The pH increase event supports the CO_2Lo initialization
107 scenario (*supplementary online text*; $\text{CO}_2 \sim 3$ PAL (Present Atmospheric Levels), pH ~ 8 ,
108 $\delta^{11}\text{B}_{\text{Sw}} \sim 34\text{‰}$) as the simulated CO_2 and temperature decrease is much reduced, and
109 therefore more consistent with independent proxy data (22), compared to CO_2Hi ($\text{CO}_2 \sim 10$
110 PAL, pH ~ 7.5 $\delta^{11}\text{B}_{\text{Sw}} \sim 36.8\text{‰}$) (Fig. 2D).

111 Prior to EP1, $\delta^{13}\text{C}_{\text{carb}}$ values began to decrease before reaching the minimum of the
112 globally recognized negative CIE at the PTB (Fig. 1). At this time both $\delta^{11}\text{B}$ and ocean pH
113 remained stable. Hypotheses to explain the negative CIE require the input of isotopically light
114 carbon, such as from volcanism (14, 25) with the assimilation of very light organic carbon
115 from the surrounding host rock (26), methane destabilization (27), collapse of the biological
116 pump (15), and/or a decrease in the burial of terrestrial carbon (16). We can simulate the
117 observed drop in $\delta^{13}\text{C}_{\text{carb}}$, whilst remaining within the uncertainty of the $\delta^{11}\text{B}$ data (Fig. 2), by
118 combining a cessation of terrestrial carbon burial with a relatively slow (50 thousand years)
119 carbon injection from any of the above sources (Fig. S8). A small source of methane (3.2
120 $\times 10^{17}$ mol C with $\delta^{13}\text{C} = -50\text{‰}$) gives the least change in $\delta^{11}\text{B}$ and pH, whilst either a larger
121 source of organic carbon ($\sim 6.5 \times 10^{17}$ mol C with $\delta^{13}\text{C} = -25\text{‰}$) or a mixture of mantle and
122 lighter carbon sources ($\sim 1.3 \times 10^{18}$ mol C with $\delta^{13}\text{C} = -12.5\text{‰}$) are still within the measured
123 uncertainty in $\delta^{11}\text{B}$.

124 This relatively slow addition of carbon minimizes the tendency for a transient decline
125 in surface ocean pH in an ocean that was already primed with a high Ω and hence high
126 buffering capacity from the Late Permian. The global presence of microbial and abiotic
127 carbonate fabrics after EP1 (28) are indicative that this high Ω was maintained across the
128 CIE. The carbon injection triggers an increase in $p\text{CO}_2$, temperature and silicate weathering,
129 thereby creating an additional counterbalancing alkalinity flux, which is consistent with
130 independent proxy data (6). The alkalinity source may have been further increased through
131 soil loss (29), the emplacement of easily-weathered Siberian Trap basalt, or the impact of
132 acid rain (30) that would have increased weathering efficiency.

133 The negative $\delta^{11}\text{B}_{\text{carb}}$ excursion at 251.88 Ma represents a calculated pH decrease of
134 up to 0.7 pH units. This pH decrease coincides with the second pulse of the extinction (Fig.
135 1), which preferentially affected the heavily calcifying, physiologically un-buffered and
136 sessile organisms (3). This was also accompanied by the temporary loss of abiotic and
137 microbial carbonates throughout the Tethys (31, 32), thereby suggesting a coeval decrease in
138 Ω . To overwhelm the buffering capacity of the ocean and decrease pH in this way requires a
139 second, more abrupt injection of carbon into the atmosphere, yet remarkably, the acidification
140 event occurs after the decline in $\delta^{13}\text{C}$, when $\delta^{13}\text{C}$ has rebounded somewhat and is essentially
141 stable (Fig. 1).

142 Unlike the first carbon injection, the lack of change in $\delta^{13}\text{C}$ at this time rules out very
143 ^{13}C -depleted carbon sources, because no counterbalancing strongly ^{13}C -enriched source
144 exists. Instead, it requires a carbon source near $\sim 0\%$. A plausible scenario for this is the
145 decarbonation of overlying carbonate host rock, into which the Siberian Traps intruded (26)
146 or the direct assimilation of carbonates and evaporites into the melt (33). Host carbonates
147 would have had $\delta^{13}\text{C} \sim +2\text{-}4\%$, which when mixed with mantle carbon ($\sim -5\%$), potentially
148 produces a source near 0% . We can simulate the sharp drop in pH and stable $\delta^{13}\text{C}$ values

149 (Fig. 2) through a large and rapid carbon release of 2×10^{18} mol C over 10 thousand years (Fig.
150 S8). This is undoubtedly a massive injection of 24,000 PgC at a rapid rate of 2.4 PgC/yr, but
151 it is physically plausible given existing estimates of the volume of carbonate host sediments
152 subject to contact metamorphism and postulated mechanisms of carbon release
153 (*supplementary online text*). This second rapid carbon release produces a sharp rise in $p\text{CO}_2$
154 to ~20 PAL and warming of ~15°C, consistent with the observation of peak temperatures
155 after EP1 (22). Initialization of the carbon cycle model under CO_2Hi cannot generate the
156 magnitude of $\delta^{11}\text{B}$ drop (Fig. 2A) because the non-linear relation between pH and $\delta^{11}\text{B}$
157 fractionation sets a lower limit of $\delta^{11}\text{B}$ at ~10‰ in this case (Fig. S3). Thus low initial CO_2 of
158 ~3 PAL in the late Permian (CO_2Lo) is more consistent with our data.

159 An acidification event of ~10 thousand years is consistent with the modelled
160 timescale required to replenish the ocean with alkalinity, as carbonate deposition is reduced
161 and weathering is increased under higher $p\text{CO}_2$ and global temperatures. Increased silicate
162 weathering rates drive further CO_2 drawdown resulting in stabilization (Fig. S7). High global
163 temperature (6) and increased silicate weathering are consistent with a sudden increase in
164 both $^{87}\text{Sr}/^{86}\text{Sr}$ (34) and sedimentation rates (29) in the Griesbachian.

165 The PTB was a time of extreme environmental change, and our combined data and
166 modeling approach falsifies several of the mechanisms currently proposed. Whilst the
167 coincident stresses of anoxia, increasing temperature, and ecosystem restructuring were
168 important during this interval, the $\delta^{11}\text{B}$ record strongly suggests that widespread ocean
169 acidification was not a factor in the first phase of the mass extinction, but did drive the
170 second pulse. The carbon release required to drive the observed acidification event must have
171 occurred at a rate comparable to the current anthropogenic perturbation, but exceeds it in
172 expected magnitude. Specifically, the required model perturbation of 24,000 PgC exceeds the
173 ~5000 PgC of conventional fossil fuels and is at the upper end of the range of estimates of

174 unconventional fossil fuels (e.g. methane hydrates). We show that such a rapid and large
175 release of carbon is critical to causing the combined synchronous decrease in both pH and
176 saturation state that defines an ocean acidification event (*II*).

177

178

179 **References and Notes**

- 180 1. D. H. Erwin, The Permo-Triassic Extinction. *Nature* **367**, 231-236 (1994).
- 181 2. S. D. Burgess, S. A. Bowring, S. Z. Shen, High-precision timeline for Earth's most severe
182 extinction. *Proceedings of the National Academy of Sciences of the United States of*
183 *America* **111**, 3203-3204 (2014).
- 184 3. H. J. Song, P. B. Wignall, J. A. Tong, Y. Hongfu, Two pulses of extinction during the
185 Permian-Triassic crisis. *Nat. Geosci.* **6**, 52-56 (2012).
- 186 4. P. B. Wignall, R. J. Twitchett, Oceanic anoxia and the end Permian mass extinction.
187 *Science* **272**, 1155-1158 (1996).
- 188 5. K. Grice, C. Q. Cao, G. D. Love, M. E. Bottcher, R. J. Twitchett, E. Grosjean, R. E. Summons, S.
189 C. Turgeon, W. Dunning, Y. G. Jin, Photic zone euxinia during the Permian-Triassic
190 superanoxic event. *Science* **307**, 706-709 (2005); (10.1126/science.1104323).
- 191 6. Y. D. Sun, M. M. Joachimski, P. B. Wignall, C. B. Yan, Y. L. Chen, H. S. Jiang, L. N. Wang, X. L.
192 Lai, Lethally Hot Temperatures During the Early Triassic Greenhouse. *Science* **338**, 366-
193 370 (2012); (10.1126/science.1224126).
- 194 7. J. L. Payne, A. V. Turchyn, A. Paytan, D. J. DePaolo, D. J. Lehrmann, M. Y. Yu, J. Y. Wei,
195 Calcium isotope constraints on the end-Permian mass extinction. *Proceedings of the*
196 *National Academy of Sciences of the United States of America* **107**, 8543-8548 (2010);
197 (10.1073/pnas.0914065107).
- 198 8. A. H. Knoll, R. K. Barnbach, J. L. Payne, S. Pruss, W. W. Fischer, Paleophysiology and end-
199 Permian mass extinction. *Earth and Planetary Science Letters* **256**, 295-313 (2007);
200 (10.1016/j.epsl.2007.02.018).
- 201 9. J. L. Payne, D. J. Lehrmann, D. Follett, M. Seibel, L. R. Kump, A. Riccardi, D. Altiner, H.
202 Sano, J. Wei, Erosional truncation of uppermost Permian shallow-marine carbonates and
203 implications for Permian-Triassic boundary events. *Geol. Soc. Am. Bull.* **119**, 771-784
204 (2007); (10.1130/b26091.1).
- 205 10. C. Korte, H. W. Kozur, Carbon-isotope stratigraphy across the Permian-Triassic
206 boundary: A review. *Journal of Asian Earth Sciences* **39**, 215-235 (2010);
207 (10.1016/j.jseaes.2010.01.005).
- 208 11. B. Hönisch, A. Ridgwell, D. N. Schmidt, E. Thomas, S. J. Gibbs, A. Sluijs, R. Zeebe, L. Kump,
209 R. C. Martindale, S. E. Greene, W. Kiessling, J. Ries, J. C. Zachos, D. L. Royer, S. Barker, T. M.
210 Marchitto, Jr., R. Moyer, C. Pelejero, P. Ziveri, G. L. Foster, B. Williams, The Geological
211 Record of Ocean Acidification. *Science* **335**, 1058-1063 (2012);
212 (10.1126/science.1208277).
- 213 12. C. L. Blätter, H. C. Jenkyns, L. M. Reynard, G. M. Henderson, Significant increases in global
214 weathering during Oceanic Anoxic Events 1a and 2 indicated by calcium isotopes. *Earth*
215 *and Planetary Science Letters* **309**, 77-88 (2011).
- 216 13. P. B. Wignall, S. Kershaw, P. Y. Collin, S. Crasquin-Soleau, Erosional truncation of
217 uppermost Permian shallow-marine carbonates and implications for Permian-Triassic

- 218 boundary events: Comment. *Geol. Soc. Am. Bull.* **121**, 954-956 (2009);
219 (10.1130/b26424.1).
- 220 14. R. A. Berner, Examination of hypotheses for the Permo-Triassic boundary extinction by
221 carbon cycle modeling. *Proceedings of the National Academy of Sciences of the United*
222 *States of America* **99**, 4172-4177 (2002); (10.1073/pnas.032095199).
- 223 15. M. R. Rampino, K. Caldeira, Major perturbation of ocean chemistry and a 'Strangelove
224 Ocean' after the end-Permian mass extinction. *Terra Nova* **17**, 554-559 (2005);
225 (10.1111/j.1365-3121.2005.00648.x).
- 226 16. W. S. Broecker, S. Peacock, An ecologic explanation for the Permo-Triassic carbon and
227 sulfur isotope shifts. *Global Biogeochemical Cycles* **13**, 1167-1172 (1999).
- 228 17. Materials and methods are available on Science online
- 229 18. M. O. Clarkson, S. Richoz, R. A. Wood, F. Maurer, L. Krystyn, D. J. McGurty, D. Astratti, A
230 new high-resolution delta C-13 record for the Early Triassic: Insights from the Arabian
231 Platform. *Gondwana Research* **24**, 233-242 (2013); (10.1016/j.gr.2012.10.002).
- 232 19. M. M. Joachimski, L. Simon, R. van Geldern, C. Lecuyer, Boron isotope geochemistry of
233 Paleozoic brachiopod calcite: Implications for a secular change in the boron isotope
234 geochemistry of seawater over the Phanerozoic. *Geochimica et Cosmochimica Acta* **69**,
235 4035-4044 (2005); (10.1016/j.gca.2004.11.017).
- 236 20. The alternative way to drive an increase in pH would be through a removal of carbon,
237 however this would be evident in the $\delta^{13}\text{C}$ record so we can rule it out.
- 238 21. Bacterial Sulfate Reduction (BSR) is a net source of alkalinity if the generated H_2S is
239 buried as pyrite. Pyrite deposition is seen widely in certain settings during the Late
240 Permian to PTB. See SM for further information.
- 241 22. M. M. Joachimski, X. L. Lai, S. Z. Shen, H. S. Jiang, G. M. Luo, B. Chen, J. Chen, Y. D. Sun,
242 Climate warming in the latest Permian and the Permian-Triassic mass extinction.
243 *Geology* **40**, 195-198 (2012); (Doi 10.1130/G32707.1).
- 244 23. H. Yin, H. Jiang, W. Xia, Q. Feng, N. Zhang, J. Shen, The end-Permian regression in South
245 China and its implication on mass extinction. *Earth-Sci. Rev.* **137**, 19-33 (2014);
246 (<http://dx.doi.org/10.1016/j.earscirev.2013.06.003>).
- 247 24. P. B. Wignall, R. J. Twitchett, Extent, duration, and nature of the Permian-Triassic
248 superanoxic event. *Catastrophic Events and Mass Extinctions: Impacts and Beyond*, 395-
249 413 (2002).
- 250 25. C. Korte, P. Pande, P. Kalia, H. W. Kozur, M. M. Joachimski, H. Oberhaensli, Massive
251 volcanism at the Permian-Triassic boundary and its impact on the isotopic composition
252 of the ocean and atmosphere. *Journal of Asian Earth Sciences* **37**, 293-311 (2010);
253 (10.1016/j.jseaes.2009.08.012).
- 254 26. H. Svensen, S. Planke, A. G. Polozov, N. Schmidbauer, F. Corfu, Y. Y. Podladchikov, B.
255 Jamtveit, Siberian gas venting and the end-Permian environmental crisis. *Earth and*
256 *Planetary Science Letters* **277**, 490-500 (2009); (10.1016/j.epsl.2008.11.015).

- 257 27. E. S. Krull, G. J. Retallack, $\delta^{13}\text{C}$ depth profiles from paleosols across the Permian-Triassic
258 boundary: Evidence for methane release. *Geol. Soc. Am. Bull.* **112**, 1459-1472 (2000);
259 (10.1130/0016-7606(2000)112<1459:CDPFPA>2.0.CO;2).
- 260 28. A. D. Woods, Assessing Early Triassic Paleooceanographic conditions via unusual
261 sedimentary fabrics and features. *Earth-Sci. Rev.*, (2013);
262 (<http://dx.doi.org/10.1016/j.earscirev.2013.08.015>).
- 263 29. T. J. Algeo, Z. Q. Chen, M. L. Fraiser, R. J. Twitchett, Terrestrial-marine teleconnections in
264 the collapse and rebuilding of Early Triassic marine ecosystems. *Palaeogeography*
265 *Palaeoclimatology Palaeoecology* **308**, 1-11 (2011); (10.1016/j.palaeo.2011.01.011).
- 266 30. B. A. Black, J. F. Lamarque, C. A. Shields, L. T. Elkins-Tanton, J. T. Kiehl, Acid rain and
267 ozone depletion from pulsed Siberian Traps magmatism. *Geology* **42**, 67-70 (2014); (Doi
268 10.1130/G34875.1).
- 269 31. A. Baud, S. Richoz, S. Pruss, The lower Triassic anachronistic carbonate facies in space
270 and time. *Global and Planetary Change* **55**, 81-89 (2007);
271 (10.1016/j.gloplacha.2006.06.008).
- 272 32. S. Richoz, L. Krystyn, A. Baud, R. Brandner, M. Horacek, P. Mohtat-Aghai, Permian-
273 Triassic boundary interval in the Middle East (Iran and N. Oman): Progressive
274 environmental change from detailed carbonate carbon isotope marine curve and
275 sedimentary evolution. *Journal of Asian Earth Sciences* **39**, 236-253 (2010);
276 (10.1016/j.jseaes.2009.12.014).
- 277 33. B. A. Black, L. T. Elkins-Tanton, M. C. Rowe, I. U. Peate, Magnitude and consequences of
278 volatile release from the Siberian Traps. *Earth and Planetary Science Letters* **317**, 363-
279 373 (2012); (Doi 10.1016/J.Epsl.2011.12.001).
- 280 34. C. Korte, H. W. Kozur, M. M. Joachimski, H. Strauss, J. Veizer, L. Schwark, Carbon, sulfur,
281 oxygen and strontium isotope records, organic geochemistry and biostratigraphy across
282 the Permian/Triassic boundary in Abadeh, Iran. *International Journal of Earth Sciences*
283 **93**, 565-581 (2004); (Doi 10.1007/S00531-004-0406-7).
- 284 35. G. M. Stampfli, G. D. Borel, A plate tectonic model for the Paleozoic and Mesozoic
285 constrained by dynamic plate boundaries and restored synthetic oceanic isochrons.
286 *Earth and Planetary Science Letters* **196**, 17-33 (2002); (10.1016/s0012-
287 821x(01)00588-x).
- 288 36. S. Z. Shen, C. Q. Cao, H. Zhang, S. A. Bowring, C. M. Henderson, J. L. Payne, V. I. Davydov, B.
289 Chen, D. X. Yuan, Y. C. Zhang, W. Wang, Q. F. Zheng, High-resolution delta C-13(carb)
290 chemostratigraphy from latest Guadalupian through earliest Triassic in South China and
291 Iran. *Earth and Planetary Science Letters* **375**, 156-165 (2013); (Doi
292 10.1016/J.Epsl.2013.05.020).
- 293 37. B. Koehler, M. Zeller, T. Aigner, M. Poepfelreiter, P. Milroy, H. Forke, S. Al-Kindi, Facies
294 and stratigraphic framework of a Khuff outcrop equivalent: Saiq and Mahil formations,
295 Al Jabal al-Akhdar, Sultanate of Oman. *GeoArabia* **15**, 91-156 (2010).
- 296 38. F. Maurer, R. Martini, R. Rettori, H. Hillgartner, S. Cirilli, The geology of Khuff outcrop
297 analogues in the Musandam Peninsula, United Arab Emirates and Oman. *GeoArabia* **14**,
298 125-158 (2009).

- 299 39. S. Kasemann, A. Meixner, A. Rocholl, T. Vennemann, M. Rosner, A. K. Schmitt, M.
300 Wiedenbeck, Boron and oxygen isotope composition of certified reference materials
301 NIST SRM 610/612 and reference materials JB-2 and JR-2. *Geostandards Newsletter-the*
302 *Journal of Geostandards and Geoanalysis* **25**, 405-416 (2001); (Doi 10.1111/J.1751-
303 908x.2001.Tb00615.X).
- 304 40. S. A. Kasemann, D. N. Schmidt, J. Bijma, G. L. Foster, In situ boron isotope analysis in
305 marine carbonates and its application for foraminifera and palaeo-pH. *Chem. Geol.* **260**,
306 138-147 (2009); (Doi 10.1016/J.Chemgeo.2008.12.015).
- 307 41. S. A. Kasemann, C. J. Hawkesworth, A. R. Prave, A. E. Fallick, P. N. Pearson, Boron and
308 calcium isotope composition in Neoproterozoic carbonate rocks from Namibia: evidence
309 for extreme environmental change. *Earth and Planetary Science Letters* **231**, 73-86
310 (2005); (Doi 10.1016/J.Epsl.2004.12.006).
- 311 42. S. A. Kasemann, A. R. Prave, A. E. Fallick, C. J. Hawkesworth, K. H. Hoffmann,
312 Neoproterozoic ice ages, boron isotopes, and ocean acidification: Implications for a
313 snowball Earth. *Geology* **38**, 775-778 (2010); (10.1130/g30851.1).
- 314 43. P. B. Wignall, A. Hallam, Facies change and the end-Permian mass extinction in SE
315 Sichuan, China. *Palaios* **11**, 587-596 (1996).
- 316 44. L. Breesch, R. Swennen, B. Dewever, F. Roure, B. Vincent, Diagenesis and fluid system
317 evolution in the northern Oman Mountains, United Arab Emirates: Implications for
318 petroleum exploration. *GeoArabia* **16**, 111-148 (2011).
- 319 45. G. Paris, A. Bartolini, Y. Donnadieu, V. Beaumont, J. Gaillardet, Investigating boron
320 isotopes in a middle Jurassic micritic sequence: Primary vs. diagenetic signal. *Chem. Geol.*
321 **275**, 117-126 (2010); (10.1016/j.chemgeo.2010.03.013).
- 322 46. J. Veizer, D. Ala, K. Azmy, P. Bruckschen, D. Buhl, F. Bruhn, G. A. F. Carden, A. Diener, S.
323 Ebner, Y. Godderis, T. Jasper, C. Korte, F. Pawellek, O. G. Podlaha, H. Strauss, Sr-87/Sr-
324 86, delta C-13 and delta O-18 evolution of Phanerozoic seawater. *Chem. Geol.* **161**, 59-88
325 (1999); (Doi 10.1016/S0009-2541(99)00081-9).
- 326 47. S. B. Jacobsen, A. J. Kaufman, The Sr, C and O isotopic evolution of Neoproterozoic
327 seawater. *Chem. Geol.* **161**, 37-57 (1999); (Doi 10.1016/S0009-2541(99)00080-7).
- 328 48. P. K. Swart, M. J. Kennedy, Does the global stratigraphic reproducibility of delta C-13 in
329 Neoproterozoic carbonates require a marine origin? A Pliocene-Pleistocene comparison.
330 *Geology* **40**, 87-90 (2012); (Doi 10.1130/G32538.1).
- 331 49. A. M. Oehlert, P. K. Swart, Interpreting carbonate and organic carbon isotope covariance
332 in the sedimentary record. *Nature Communications* **5**, 7 (2014);
333 (10.1038/ncomms5672).
- 334 50. F. Ohnemüller, A. R. Prave, A. E. Fallick, S. A. Kasemann, Ocean acidification in the
335 aftermath of the Marinoan glaciation. *Geology* **42**, 1103-1106 (2014);
336 (10.1130/g35937.1).
- 337 51. D. Lemarchand, J. Gaillardet, E. Lewin, C. J. Allegre, The influence of rivers on marine
338 boron isotopes and implications for reconstructing past ocean pH. *Nature* **408**, 951-954
339 (2000).

- 340 52. A. J. Spivack, J. M. Edmond, Boron Isotope Exchange between Seawater and the Oceanic-
341 Crust. *Geochimica et Cosmochimica Acta* **51**, 1033-1043 (1987); (Doi 10.1016/0016-
342 7037(87)90198-0).
- 343 53. A. G. Dickson, Thermodynamics of the dissociation of boric-acid in synthetic seawater
344 from 273.15-k to 318.15-k. *Deep-Sea Research Part a-Oceanographic Research Papers* **37**,
345 755-766 (1990).
- 346 54. M. Schobben, M. M. Joachimski, D. Korn, L. Leda, C. Korte, Palaeotethys seawater
347 temperature rise and an intensified hydrological cycle following the end-Permian mass
348 extinction. *Gondwana Research*, (2013); (<http://dx.doi.org/10.1016/j.gr.2013.07.019>).
- 349 55. J. T. Kiehl, C. A. Shields, Climate simulation of the latest Permian: Implications for mass
350 extinction. *Geology* **33**, 757-760 (2005); (Doi 10.1130/G21654.1).
- 351 56. R. A. Locarnini, A. V. Mishonov, J. I. Antonov, T. P. Boyer, H. E. Garcia, in *NOAA Atlas*
352 *NESDIS 61*, S. Levitus, Ed. (U.S. Government Printing Office, Washington, D.C, 2006), vol.
353 1: Temperature.
- 354 57. G. L. Foster, Seawater pH, $p\text{CO}_2$ and $[\text{CO}_3^{2-}]$ variations in the Caribbean Sea over the last
355 130 kyr: A boron isotope and B/Ca study of planktic foraminifera. *Earth and Planetary*
356 *Science Letters* **271**, 254-266 (2008); (10.1016/j.epsl.2008.04.015).
- 357 58. N. G. Hemming, B. Honisch, A critical review and recent advances in the boron isotope
358 paleo-pH proxy. *Geochimica et Cosmochimica Acta* **69**, A129-A129 (2005).
- 359 59. B. Hönisch, N. G. Hemming, Surface ocean pH response to variations in $p\text{CO}_2$ through
360 two full glacial cycles. *Earth and Planetary Science Letters* **236**, 305-314 (2005); (Doi
361 10.1016/j.epsl.2005.04.027).
- 362 60. N. G. Hemming, G. N. Hanson, Boron isotopic composition and concentration in modern
363 marine carbonates. *Geochimica et Cosmochimica Acta* **56**, 537-543 (1992).
- 364 61. K. Klochko, A. J. Kaufman, W. S. Yao, R. H. Byrne, J. A. Tossell, Experimental measurement
365 of boron isotope fractionation in seawater. *Earth and Planetary Science Letters* **248**, 276-
366 285 (2006); (10.1016/j.epsl.2006.05.034).
- 367 62. E. Lewis, D. W. R. Wallace. (Oak Ridge, 1998).
- 368 63. J. W. B. Rae, G. L. Foster, D. N. Schmidt, T. Elliott, Boron isotopes and B/Ca in benthic
369 foraminifera: Proxies for the deep ocean carbonate system. *Earth and Planetary Science*
370 *Letters* **302**, 403-413 (2011).
- 371 64. G. L. Foster, P. A. E. Pogge von Strandmann, J. W. B. Rae, Boron and magnesium isotopic
372 composition of seawater. *Geochemistry Geophysics Geosystems* **11**, 10 (2010);
373 (10.1029/2010gc003201).
- 374 65. A. Ridgwell, A Mid Mesozoic Revolution in the regulation of ocean chemistry. *Marine*
375 *Geology* **217**, 339-357 (2005); (Doi 10.1016/J.Margeo.2004.10.036).
- 376 66. A. Sanyal, M. Nugent, R. J. Reeder, J. Buma, Seawater pH control on the boron isotopic
377 composition of calcite: Evidence from inorganic calcite precipitation experiments.
378 *Geochimica et Cosmochimica Acta* **64**, 1551-1555 (2000).

- 379 67. M. Pagani, D. Lemarchand, A. Spivack, J. Gaillardet, A critical evaluation of the boron
380 isotope-pH proxy: The accuracy of ancient ocean pH estimates. *Geochimica et*
381 *Cosmochimica Acta* **69**, 953-961 (2005); (Doi 10.1016/J.Gca.2004.07.029).
- 382 68. K. Klochko, G. D. Cody, J. A. Tossell, P. Dera, A. J. Kaufman, Re-evaluating boron speciation
383 in biogenic calcite and aragonite using B-11 MAS NMR. *Geochimica et Cosmochimica Acta*
384 **73**, 1890-1900 (2009); (10.1016/j.gca.2009.01.002).
- 385 69. R. H. Byrne, W. S. Yao, K. Klochko, J. A. Tossell, A. J. Kaufman, Experimental evaluation of
386 the isotopic exchange equilibrium $B-10(OH)(3)+B-11(OH)(4)(-) = B-11(OH)(3)+B-$
387 $10(OH)(-)(4)$ in aqueous solution. *Deep-Sea Research Part I-Oceanographic Research*
388 *Papers* **53**, 684-688 (2006); (10.1016/j.dsr.2006.01.005).
- 389 70. J. Payne, L. Kump, Evidence for recurrent Early Triassic massive volcanism from
390 quantitative interpretation of carbon isotope fluctuations. *Earth and Planetary Science*
391 *Letters* **256**, 264-277 (2007); (10.1016/j.epsl.2007.01.034).
- 392 71. A. Ridgwell, R. E. Zeebe, The role of the global carbonate cycle in the regulation and
393 evolution of the Earth system. *Earth and Planetary Science Letters* **234**, 299-315 (2005);
394 (Doi 10.1016/J.Epsl.2005.03.006).
- 395 72. R. E. Locklair, A. Lerman, A model of Phanerozoic cycles of carbon and calcium in the
396 global ocean: Evaluation and constraints on ocean chemistry and input fluxes. *Chem.*
397 *Geol.* **217**, 113-126 (2005); (10.1016/j.chemgeo.2004.12.010).
- 398 73. Y. Cui, L. R. Kump, Global warming and the end-Permian extinction event: Proxy and
399 modeling perspectives. *Earth-Science Reviews*, (2014);
400 (<http://dx.doi.org/10.1016/j.earscirev.2014.04.007>).
- 401 74. K. Caldeira, J. F. Kasting, The life span of the biosphere revisited. *Nature* **360**, 721-723
402 (1992).
- 403 75. J. M. Bergman, T. M. Lenton, A. J. Watson, COPSE: A new model of biogeochemical cycling
404 over Phanerozoic time. *American Journal of Science* **304**, (2004).
- 405 76. A. F. Hofmann, K. Soetaert, J. J. Middelburg, F. J. R. Meysman, AquaEnv : An Aquatic Acid-
406 Base Modelling Environment in R. *Aquatic Geochemistry* **16**, 507-546 (2010);
407 (10.1007/s10498-009-9084-1).
- 408 77. R. E. Zeebe, P. Westbrooke, A simple model for the CaCO₃ saturation state of the ocean:
409 The "Strangelove," the "Neritan," and the "Cretan" Ocean. *Geochemistry Geophysics*
410 *Geosystems* **4**, (2003).
- 411 78. J. Zhang, P. D. Quay, D. O. Wilbur, Carbon isotope fractionation during gas-water
412 exchange and dissolution of CO₂. *Geochimica et Cosmochimica Acta* **59**, 107-114 (1995).
- 413 79. H. D. Holland, in *Treatise on Geochemistry: Vol 6 The oceans and marine geochemistry*, H.
414 D. Holland, K. K. Turekian, Eds. (Elsevier Academic Press, 2003), pp. 583-625.
- 415 80. H. Song, J. Tong, T. J. Algeo, H. Song, H. Qiu, Y. Zhu, L. Tian, S. Bates, T. W. Lyons, G. Luo, L.
416 R. Kump, Early Triassic seawater sulfate drawdown. *Geochimica et Cosmochimica Acta*
417 **128**, 95-113 (2014); (10.1016/j.gca.2013.12.009).
- 418 81. G. Luo, L. R. Kump, Y. Wang, J. Tong, M. a. Arthur, H. Yang, J. Huang, H. Yin, S. Xie, Isotopic
419 evidence for an anomalously low oceanic sulfate concentration following end-Permian

- 420 mass extinction. *Earth and Planetary Science Letters* **300**, 101-111 (2010);
421 (10.1016/j.epsl.2010.09.041).
- 422 82. R. E. Zeebe, LOSCAR: Long-term Ocean-atmosphere-Sediment Carbon cycle Reservoir
423 Model v2.0.4. *Geoscientific Model Development* **5**, 149-166 (2012); (10.5194/gmd-5-149-
424 2012).
- 425 83. S. Ben-Yaakov, M. B. Goldhaber, The influence of sea water composition on the apparent
426 constants of the carbonate system. *Deep Sea Research and Oceanographic Abstracts* **20**,
427 87-99 (1973); (10.1016/0011-7471(73)90044-2).
- 428 84. Y. Cui, L. R. Kump, Global warming and the end-Permian extinction event: Proxy and
429 modeling perspectives. (2014); (doi:10.1016/j.earscirev.2014.04.007).
- 430 85. Y. Cui, L. R. Kump, A. Ridgwell, Initial assessment of the carbon emission rate and
431 climatic consequences during the end-Permian mass extinction. *Palaeogeography*
432 *Palaeoclimatology Palaeoecology* **389**, 128-136 (2013).
- 433 86. T. J. Algeo, K. Kuwahara, H. Sano, S. Bates, T. Lyons, E. Elswick, L. Hinnov, B. Ellwood, J.
434 Moser, J. B. Maynard, Spatial variation in sediment fluxes, redox conditions, and
435 productivity in the Permian–Triassic Panthalassic Ocean. *Palaeogeography,*
436 *Palaeoclimatology, Palaeoecology* **308**, 65-83 (2011); (10.1016/j.palaeo.2010.07.007).
- 437 87. T. J. Algeo, R. J. Twitchett, Anomalous Early Triassic sediment fluxes due to elevated
438 weathering rates and their biological consequences. *Geology* **38**, 1023-1026 (2010);
439 (10.1130/g31203.1).
- 440 88. G. A. Brennecka, A. D. Herrmann, T. J. Algeo, A. D. Anbar, Rapid expansion of oceanic
441 anoxia immediately before the end-Permian mass extinction. *Proceedings of the National*
442 *Academy of Sciences of the United States of America* **108**, 17631-17634 (2011);
443 (10.1073/pnas.1106039108).
- 444 89. S. E. Grasby, B. Beauchamp, Latest Permian to Early Triassic basin-to-shelf anoxia in the
445 Sverdrup Basin, Arctic Canada. *Chem. Geol.* **264**, 232-246 (2009); (Doi
446 10.1016/J.Chemgeo.2009.03.009).
- 447 90. C. Winguth, A. M. E. Winguth, Simulating Permian-Triassic oceanic anoxia distribution:
448 Implications for species extinction and recovery. *Geology* **40**, 127-130 (2012);
449 (10.1130/g32453.1).
- 450 91. T. J. Algeo, C. M. Henderson, J. N. Tong, Q. L. Feng, H. F. Yin, R. V. Tyson, Plankton and
451 productivity during the Permian-Triassic boundary crisis: An analysis of organic carbon
452 fluxes. *Global and Planetary Change* **105**, 52-67 (2013); (Doi
453 10.1016/J.Gloplacha.2012.02.008).
- 454 92. E. Tziperman, I. Halevy, D. T. Johnston, A. H. Knoll, D. P. Schrag, Biologically induced
455 initiation of Neoproterozoic snowball-Earth events. *Proceedings of the National Academy*
456 *of Sciences of the United States of America*, 1-6 (2011); (10.1073/pnas.1016361108).
- 457 93. H. Song, J. Tong, T. J. Algeo, H. Song, H. Qiu, Y. Zhu, L. Tian, S. Bates, T. W. Lyons, G. Luo, L.
458 R. Kump, Early Triassic seawater sulfate drawdown. *Geochimica et Cosmochimica Acta*,
459 (2013); (doi: <http://dx.doi.org/10.1016/j.gca.2013.12.009>).
- 460 94. G. Luo, Y. Wang, K. Grice, S. Kershaw, T. J. Algeo, X. Ruan, H. Yang, C. Jia, S. Xie, Microbial-
461 algal community changes during the latest Permian ecological crisis: Evidence from lipid

- 462 biomarkers at Cili, South China. *Global and Planetary Change* **105**, 36-51 (2013);
463 (10.1016/j.gloplacha.2012.11.015).
- 464 95. K. M. Meyer, M. Yu, A. B. Jost, B. M. Kelley, J. L. Payne, $\delta^{13}\text{C}$ evidence that high primary
465 productivity delayed recovery from end-Permian mass extinction. *Earth and Planetary*
466 *Science Letters* **302**, 378-384 (2011); (10.1016/j.epsl.2010.12.033).
- 467 96. H. J. Song, J. N. Tong, Y. L. Xiong, D. Y. Sun, L. Tian, H. Y. Song, The large increase of
468 $\delta^{13}\text{C}(\text{carb})$ -depth gradient and the end-Permian mass extinction. *Sci. China-Earth Sci.*
469 **55**, 1101-1109 (2012); (10.1007/s11430-012-4416-1).
- 470 97. D. E. Ogden, N. H. Sleep, Explosive eruption of coal and basalt and the end-Permian mass
471 extinction. *Proceedings of the National Academy of Sciences of the United States of*
472 *America* **109**, 59-62 (2012); (10.1073/pnas.1118675109).
- 473 98. M. K. Reichow, A. D. Saunders, R. V. White, M. S. Pringle, A. I. Al'Mukhamedov, A. I.
474 Medvedev, N. P. Kirida, Ar-40/Ar-39 dates from the West Siberian Basin: Siberian flood
475 basalt province doubled. *Science* **296**, 1846-1849 (2002).
- 476 99. M. B. Harfoot, J. a. Pyle, D. J. Beerling, End-Permian ozone shield unaffected by oceanic
477 hydrogen sulphide and methane releases. *Nat. Geosci.* **1**, 247-252 (2008);
478 (10.1038/ngeo154).
- 479 100. C. Ganino, N. T. Arndt, Climate changes caused by degassing of sediments during the
480 emplacement of large igneous provinces. *Geology* **37**, 323-326 (2009);
481 (10.1130/G25325A.1).

482

483 **Acknowledgements:** MOC acknowledges funding from the Edinburgh University Principal's
484 Career Development Scholarship, the International Centre for Carbonate Reservoirs and The
485 Marsden Fund (U001314). RW, TL and SWP acknowledge support from NERC through the
486 'Co-evolution of Life and the Planet' scheme (NE/I005978). TL and SD were supported by
487 the Leverhulme Trust (RPG-2013-106). SK and AM acknowledge support from the German
488 Research Foundation (DFG Major Research Instrumentation Program). This is a contribution
489 to IGCP 572 with SR sponsored for fieldwork by the Austrian National Committee (Austrian
490 Academy of Sciences) for IGCP. We are grateful to R. Newton and A. Thomas for helpful
491 discussions, L. Krystyn for field assistance, F. Maurer for providing photomicrographs, and
492 B. Mills for assisting with model studies. Data are available online in the Supplementary
493 Materials and at www.pangaea.de

494

495 **Fig. 1.** Site locality and high resolution carbon and boron isotope data. **(A)** Paleogeographic
496 reconstruction for the Late Permian showing the studied section Wadi Bih, in the Musandam
497 Mountains of U.A.E. that formed an extensive carbonate platform in the Neo-Tethyan Ocean.
498 Modified from (35). **(B)** Shallow water $\delta^{13}\text{C}$ record (18). **(C)** Boron isotope ($\delta^{11}\text{B}$) record
499 (propagated uncertainty given as $2\sigma_f$) and average Early Permian brachiopod value (n=5)
500 (19). See **(A)** for lithology, biota and transect key. Only *Hindeodus parvus* has been found so
501 far in this section and the conodont zones with dashed line are identified from the $\delta^{13}\text{C}$ record
502 (36-38).

503

504 **Fig. 2.** Model results of carbon cycle parameters for the two end-member CO_2 scenarios;
505 CO_2Hi and CO_2Lo (17). **(A)** Model reproduced $\delta^{11}\text{B}$ vs data. **(B)** Modelled $\delta^{13}\text{C}$ vs data. **(C)**
506 Modelled pH envelope incorporating uncertainty of seawater B isotope composition ($\delta^{11}\text{B}_{\text{sw}}$)
507 and dynamic temperatures. **(D)** Calculated atmospheric CO_2 .

508 **Supplementary Materials:**

509 Materials and Methods

510 Supplementary Text

511 Figures S1-S9

512 Tables S1-S10

513 References (39-100)