



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Synthesis and summary of patient-reported outcomes (PRO) to inform the development of a core outcome set in colorectal cancer surgery

Citation for published version:

McNair, AGK, Whistance, RN, Forsythe, R, Rees, J, Jones, JE, Pullyblank, AM, Avery, KNL, Brookes, ST, Thomas, M, Sylvester, P, Russell, A, Oliver, A, Morton, D, Kennedy, R, Jayne, DG, Huxtable, R, Hackett, R, Dutton, SJ, Coleman, MG, Card, M, Brown, J & Blazeby, JM 2015, 'Synthesis and summary of patient-reported outcomes (PRO) to inform the development of a core outcome set in colorectal cancer surgery', *Colorectal Disease*. <https://doi.org/10.1111/codi.13021>

Digital Object Identifier (DOI):

[10.1111/codi.13021](https://doi.org/10.1111/codi.13021)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Colorectal Disease

Publisher Rights Statement:

Available under Open Access

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Received Date : 16-Mar-2015

Revised Date : 24-Apr-2015

Accepted Date : 05-May-2015

Article type : Systematic Review

Synthesis and summary of patient-reported outcome measures (PROMs) to inform the development of a core outcome set in colorectal cancer surgery

McNair AGK^{1,5}, Whistance RN^{1,2}, Forsythe RO^{1,2}, Rees J¹, Jones JE³, Pullyblank AM⁴, Avery KNL¹, Brookes ST¹, Thomas MG⁶, Sylvester PA⁶, Russell A⁷, Oliver A⁷, Morton D⁸, Kennedy R⁹, Jayne DG¹⁰, Huxtable R¹¹, Hackett R¹², Dutton SJ¹³, Coleman MG¹⁴, Card M⁶, Brown J¹⁵, Blazeby JM^{1,2}

On behalf of the CONSENSUS-CRC (Core Outcomes and iNformation SEts iN SURgical Studies – ColoRectal Cancer) working group.

¹Centre for Surgical Research, School of Social and Community Medicine, University of Bristol, Bristol, UK

²Division of Surgery Head and Neck, University Hospitals Bristol NHS Foundation Trust, Bristol, UK

³Colorectal Cancer Patient Representative, North Bristol NHS Trust, Bristol, UK

⁴Department of General Surgery, North Bristol NHS Trust, Bristol, UK

⁵Severn School of Surgery, Bristol, UK

⁶Colorectal Surgery Unit, University Hospitals Bristol NHS Foundation Trust, Bristol, UK

⁷Colorectal Consumer Liaison Group, National Cancer Research Institute, London, UK

⁸Academic Department of Surgery, University of Birmingham, Birmingham, UK

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1111/codi.13021

This article is protected by copyright. All rights reserved.

⁹Department of Surgery, St Mark's Hospital and Academic Institute, Harrow, UK

¹⁰Academic Surgical Unit, St James' University Hospital NHS Trust, Leeds, UK

¹¹Centre for Ethics in Medicine, University of Bristol, Bristol, UK

¹²Colorectal Network Site Specific Group, Avon, Somerset & Wiltshire Cancer Services, UK

¹³Centre for Statistics in Medicine and Oxford Clinical Trials Research Unit, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK

¹⁴Department of Colorectal Surgery, Plymouth Hospitals NHS Trust, Plymouth, UK

¹⁵Clinical Trials Research Unit, University of Leeds, Leeds, UK

Corresponding author:

Professor Jane M Blazeby MD FRCS

Centre for Surgical Research

School of Social and Community Medicine

University of Bristol

Canynges Hall

39 Whatley Road

Bristol

BS8 2PS

United Kingdom

Email: j.m.blazeby@bristol.ac.uk

Tel: +44 177 928 3495

Keywords: Colorectal cancer; surgery; patient-reported outcomes; core outcome set; systematic review

Abstract

Aim: Patient reported outcome measures (PROMs) are standard in the assessment of colorectal cancer (CRC) treatment, but the range and complexity of available PROMs may

This article is protected by copyright. All rights reserved.

Accepted Article

be hindering evidence synthesis. This systematic review aimed to 1) summarise PROMs in studies of CRC surgery and 2) categorise PRO content to inform the future development of an agreed minimum 'core' outcome set to be measured in all trials.

Method: All PROMs were identified from a systematic review of prospective CRC surgical studies. The type and frequency of PROMs in each study were summarized, and numbers of items documented. All items were extracted and independently categorized by content by two researchers into 'health domains' and discrepancies discussed with a patient and expert. Domain popularity and distribution of items were summarized.

Results: 58 different PROMs were identified from the 104 included studies. There were 23 generic, 4 cancer specific, 11 disease and 16 symptom specific questionnaires, and 3 *ad hoc* measures. The most frequently used PROM was the EORTC QLQ-C30 (50 studies), and most PROMs (40,69%) were used in only one study. Detailed examination of the 50 available measures identified 917 items, which were categorized into 51 domains. The domains comprising the most items were 'anxiety' (n=85,9.2%), 'fatigue' (n=67,7.3%), and 'physical function' (n=63,6.9%). No domains were included in all PROMs

Conclusion: There is major heterogeneity of PRO measurement and wide variation in content assessed by PROMs available for CRC. A core outcome set will improve PRO outcome measurement and reporting in CRC trials.

What does this paper adds to the literature?

The review summarises colorectal cancer surgical PROMs and demonstrates major heterogeneity in PRO measurement in trials that hinders evidence synthesis and meta-analysis. PROMs were categorized to inform the development of a core outcome set to resolve this problem.

This article is protected by copyright. All rights reserved.

Introduction

The measurement of patient reported outcomes (PROs) has become standard in the assessment of colorectal cancer treatments, and their use is recommended by funding and regulatory agencies¹. Many patient-reported outcome measures (PROMs) have therefore been developed for a variety of purposes². Some are generic, and allow comparisons between patients with other conditions (e.g. SF-36, EQ-5D). Others are designed for patients with cancer (e.g. EORTC QLQ-C30, FACT-G), and some are specific for colorectal cancer (e.g. EORTC-CR29, FACT-C). To add further complexity, each of these PROMs typically consist of a number of questions (items), which are often grouped together to represent similar concepts (scales). For example, two questions regarding activities of daily living and leisure activities in the EORTC QLQ-C30 measure are grouped into a single “role function” scale. There are therefore a multitude of ways to measure PROs to evaluate treatment for colorectal cancer, and this creates problems that may influence the conduct and clinical impact of trials.

Trials may use different PROMs^{3,4} making it impossible to synthesize data across trials or undertake meta-analyses. The multiplicity of results available from trials means that it is difficult to interpret findings in the context of clinical practice because of lack of familiarity with the number of measures, scales and items². For example, the scale “physical function” exists in several different PROMs, but individual items in these scales vary considerably between questionnaires. This is confusing for clinicians, who may not be aware of the differences between PROMs and it is likely to limit the meaningful use of the data in practice. Finally, the opportunities of multiple outcome measurement may lead to selective reporting of significant findings. This can generate bias and influence clinical interpretation of trials⁵.

A proposed solution to these issues are “core outcome sets”. Core outcomes are the minimum set of outcomes that patients and professionals agree to be measured in all trials of a certain condition⁶. They aim to facilitate comparisons between trials and aid meta-analysis by standardising outcome measurement, including PROs. The use of core sets may also facilitate clinical communication of data. Many core outcome sets have now been developed in different clinical areas including rheumatology⁷, paediatrics⁸ and obstetrics⁹, but not in colorectal cancer surgery. This systematic review aims to examine the measurement of PROs in colorectal cancer (CRC) surgical studies, and use the data to inform the development of the core outcome set.

Method

A systematic review of prospective colorectal cancer surgical studies measuring PROs was undertaken to 1) summarise PRO measurement in CRC surgical studies, and 2) examine each PROM in detail, and categorize analogous concepts into domains to inform the future development of a core outcome set.

Systematic search and data extraction

This systematic review adhered to a pre-defined protocol (available on request from authors). Validated terms relating to ‘surgery’, ‘colorectal cancer’ and ‘prospective studies’ (Table 4) were used to search the OVID SP versions of MEDLINE, EMBASE and the Cochrane Central Register of Controlled Trials. A validated filter for ‘prospective studies’ was used because PRO data are typically collected prospectively. The search was limited to studies conducted in humans aged 18 years and over, reported in the English language between January 2009 and December 2010. Previous reviews have considered PROs of colorectal cancer surgery in terms of elderly patients¹⁰, methodological challenges in

This article is protected by copyright. All rights reserved.

measuring PROs in colorectal cancer¹¹, laparoscopic surgery¹², long-term survivors¹³, rectal cancer³, and colorectal cancer before 2009¹⁴. The studies identified in these reviews were included. All citations were collated with Reference Manager 12 (Thomson Reuters, New York, USA) and the duplicates removed.

Titles and abstracts of identified publications were screened by one researcher. If there was uncertainty about the eligibility of a publication, the full paper was also accessed. Articles were included if they were original research papers reporting PROs of CRC surgery (curative or palliative), with or without neoadjuvant or adjuvant therapies, or systematic reviews of such publications. PROs were defined as endpoints provided by patients themselves and not interpreted by observers. Excluded were studies of non-biomedical interventions (e.g. alternative medicine), palliative treatments that did not include a surgical component (e.g. palliative chemotherapy), screening studies, treatment of colorectal metastases and molecular and genetic prognostic studies. Studies of more than one cancer site or of mixed benign and malignant disease were included provided the data for CRC patients was presented separately to that of the other diseases.

Data extraction included participant demographics (number, age and gender); treatment received (surgery, neoadjuvant radiotherapy/chemoradiation and adjuvant chemotherapy); treatment intent (curative or palliative); the study design (randomized trial, case-control study, cohort study, cross-sectional study, prospective case series or other design); the PRO questionnaire used; and the individual items included in each questionnaire. When the individual questionnaires were not available in publications, internet searches and direct contact with authors were used to obtain the information. All data were entered into a Microsoft Access (Microsoft, Washington, USA) database to facilitate data management and

This article is protected by copyright. All rights reserved.

analyses. The data extraction was checked by a second reviewer (ROF) for a sample of included articles (n=25) and any disagreements were discussed and resolved with the senior author (JMB).

1) Summary of PRO measurement in CRC surgery

The number of publications reporting each PROM was tabulated and descriptive statistics used to summarise PRO measurement. Popularity of PROMs was assessed by comparing the frequency of use in studies. A summary of each PROM is provided in terms of the numbers of items, scales and whether a total score is used. The distribution of items among PROMs was examined by calculating the median number and range of items per PROM. Questionnaires were categorized as 1) generic (for use in all patients); 2) cancer-specific (for use in all cancer patients); 3) CRC-specific (for use in colorectal cancer patients); 4) symptom-specific (to assess a single symptom e.g. pain) or 5) ad-hoc.

2) Examination of PROs and domain categorization

Individual items from all questionnaires were extracted and formed into a long-list before categorization into health domains by two researchers (RNW and JR). Both were kept masked as to which PRO questionnaire the items were derived from. Two patient representatives (JEJ and GS) and one consultant colorectal surgeon (AMP) subsequently checked this process. Discrepancies were discussed and resolved with the senior author (JMB).

Categorization was summarised using descriptive statistics to explore the distribution of items and PROMs between domains. Numbers of items included in each domain were

This article is protected by copyright. All rights reserved.

counted, as were the number of PROMs from which they were sourced. The contribution of each source PROM was demonstrated by calculating the median number and range of items included from the measures.

Results

A total of 5644 titles and abstracts was identified, of which 2127 were duplicates. The remaining 3517 were screened and 29 original research articles included. In addition to this, six systematic reviews of PROs in CRC surgery identified a further 72 original research articles (Figure 1). In total, 102 original publications including 25 RCTs (25%) and 77 non-randomised studies (75%) reporting the outcome for 66,386 patients with CRC¹⁵⁻¹¹⁷ were included. Studies are summarised in Table 1.

1) Summary of PRO measurement in CRC surgery

Fifty eight different PRO questionnaires were identified and these were reported 184 times in the included publications (Table 2). There were 23 (39.7%) generic questionnaires, 4 (6.9%) cancer-specific questionnaires, 11 (19.0%) CRC-specific questionnaires and 17 (29.3%) symptom-specific questionnaires. Three ad-hoc questionnaires (those devised specifically for the study) were not categorized.

Most questionnaires were reported only once (n=40, 69.0%). The most frequently reported were the European Organisation for the Research and Treatment of Cancer (EORTC) QLQ-C30 (50 studies, 48%), the EORTC QLQ-CR38 (33 studies, 32%) and the Medical Outcome Study Short Form-36 (n=21, 21%). The median number of items per PROM was 14, and ranged from one (five PROMs: Visual Analogue Scale (overall, pain and wound satisfaction),

This article is protected by copyright. All rights reserved.

Satisfaction with Sexual Function, and Present Pain Intensity Index) to 65 (the Profile of Mood States). Some 159 scales were evident, and most PROMs (48 [83%]) included a total score.

2) Examination of PROs and domain categorization

Fifty (86.2%) full questionnaires were available. Reasons for unavailability were because of inability to obtain the questionnaires from authors or web searches (n=6) or lack of an English language translation (n=2). The 50 questionnaires comprised some 917 individual items, and were categorised into 51 domains as described above (Table 3). The full categorization is presented in Table 5 (online supplement). The domains comprising the most items were 'anxiety' (n=85, 9.2%), 'fatigue' (n=67, 7.2%), and 'physical function' (n=63, 6.8%). The disease specific domains comprising most items were 'faecal incontinence' (n=53, 5.7%) and 'stoma problems' (n=52, 5.6%). Most domains (27, 53%) contained 10 or more items.

There was little evidence of consistency between PROMs. No domains were measured in all the PROMs. The two domains that were best represented were 'anxiety' and 'social function', each measured by 22 (44%) PROMs. Otherwise, most domains (39, 76%) were measured by less than a quarter of PROMs, highlighting further heterogeneity. There were two domains with a high median number of items included per PROM: 'Stoma problems', which contained 52 items from only five PROMs (median 7 items per PROM) and 'Satisfaction with care', which featured six items from just one PROM. This may reflect specialization of PROMs, with some measures focusing on very specific concepts.

Discussion

This systematic review aimed to summarise PRO measurement in contemporary CRC surgical studies and categorise PRO items into analogous concepts to inform the development of a core outcome set. There was evidence of significant heterogeneity of PRO measurement in the included studies. Fifty eight different PROMs were used to assess patient experience of colorectal surgery. Most (40, [69.0%]) were only ever used once, and even the most common (EORTC QLQ-C30) was measured in less than half of studies. PROMs also varied greatly in terms of their content, with some as simple as a single item, and others including up to 65. Most (52%) of PROMs were not designed to be specific to CRC surgery or symptoms thereof and, although this may bring benefits in terms of comparison between diseases, they may not be sensitive enough to issues that are of specific importance to patients with CRC. Over 900 individual questionnaire items were evident from 50 PROMs and, through a rigorous process, these were categorized into 51 'health domains'. This demonstrated a further lack of consistency, with no domains being measured in all the PROMs, and most health domains only being measured by less than a quarter of PROMs. All of this highlights potentially major questions for evidence synthesis and clinical interpretation of results in studies of CRC surgery, and demonstrates the need for a standardized core outcome set.

Other studies have highlighted the problem with outcome heterogeneity for clinical and PRO data. A recently published systematic review identified 194 studies of colorectal cancer surgery that measured 766 different clinical outcomes, with no single outcome reported in all¹¹⁸. Even considering a seemingly simple outcome such as mortality, there were over 84 different ways that this was defined and measured. The same problem has been highlighted in studies of oesophageal cancer surgery¹¹⁹, where a review of 122 articles reported 210 unique complications and 10 different measures of operative mortality, and breast

This article is protected by copyright. All rights reserved.

Accepted Article

reconstruction following mastectomy for cancer¹²⁰, which identified 134 studies reporting 950 unique complications. The problem with multiplicity of PRO measures has also been described previously in oesophageal surgery¹²¹, but there is no evidence of this issue in trials of CRC surgery.

This study is the largest systematic review of PROs in colorectal cancer and was conducted with rigorous methodology, but there are some limitations. The review covers published colorectal cancer studies in English up until 2010. A more exhaustive search over a more recent period of time, or inclusion of unpublished data or non-English publications may have yielded further PROs, but all the most commonly used PROs were captured by these inclusion criteria and extending the review is likely to have only identified additional rare PROs. The categorization process could be criticized as arbitrary, but efforts were made to objectify the process. First, two researchers categorized the questionnaire items independently and blinded to the other. Second, categorization was checked for face validity by a patient representative. Finally, there has been full disclosure of the categorization in this article to allow scientific scrutiny of the process.

Having identified all the potential patient reported health domains measured in CRC surgical studies, the next phase of this research is to gain a consensus on which outcomes are essential to measure in all trials. Recommended methods to achieve this have been defined by the international Core Outcome Measurement in Effectiveness Trials (COMET) group⁶. Domains will be combined with clinical outcomes generated from a previous systematic review¹¹⁸ to create an exhaustive long list of all potential outcomes. Key stakeholders, including patients and professionals, will then consider the importance of these outcomes and undertake a prioritization exercise called a Delphi process. This will allow the outcomes

This article is protected by copyright. All rights reserved.

of lower importance to be discarded from the core set. Finally, when the number of outcomes has been reduced, face-to-face meeting will be conducted to allow for debate about their relative merits, before the final core set is agreed.

In conclusion, this systematic review of CRC surgery demonstrated significant PRO measurement heterogeneity that may hinder comparisons between studies, limit meta-analysis and allow outcome reporting bias. A long-list of patient reported 'health domains' was generated using robust methodology to inform the development of a core outcome set.

Table 1

Summary of included articles.

	All studies n=102	Randomised trials n=25	Non-randomised studies n=77
Number of participants	66,386	7,172	59,214
Age range of participants (years)	18 – 99	29 – 89	18 – 99
Number of participating centres (%):			
Single	58 (57)	13 (52)	45 (58)
Multiple	44 (43)	12 (48)	32 (42)
IRB or ethical approval reported^a (%)	52 (51)	14 (56)	38 (49)
Tumour site (%):			
Colon	10 (9)	2 (8)	7 (9)
Rectum	54 (53)	11 (44)	44 (55)
Mixed colon and rectum	38 (38)	12 (48)	26 (34)
Surgical approach (%):			
Laparoscopy	1 (1)	0	1 (1)
Hand-assisted laparoscopy	0	0	0
Open	5 (5)	1 (4)	4 (5)
Mixed	13 (12)	9 (36)	5 (6)
Not reported or incomplete information reported	83 (81)	15 (60)	67 (87)

Neoadjuvant treatment^b (%)			
Radiotherapy alone	20 (20)	9 (36)	11 (14)
Chemotherapy alone	0	0	0 (0.0)
Chemoradiotherapy	22 (22)	3 (12)	18 (24)
None	7 (7)	1 (4)	6 (8)
Not reported or incomplete information reported	53 (51)	12 (48)	42 (54)
Adjuvant treatment^b (%)			
Chemotherapy or chemoradiotherapy	56 (55)	16 (64)	40 (52)
None	0	0	0
Not reported or incomplete information reported	46 (45)	9 (36)	37 (48)
Number of PROMs reported			
1	43	10	33
2	47	12	35
3	6	2	4
4	4	1	3
5	2	0	2

^a IRB= Institutional Review Board.

^b Some studies included patients with or without neoadjuvant therapy, some patients undergoing different neoadjuvant or adjuvant treatment within the same study.

Table 2 Summary of identified patient-reported outcome measures (questionnaires) N=58

Name of generic questionnaire (n=23)	Number of items	Number of scales	Overall score	Frequency (n=184)
Short Form-36	36	8	No	21
EuroQol-5D	6	6	Yes	3
Rotterdam Symptom Checklist	35	4	No	3
Gastrointestinal Quality of Life Index	36	0	Yes	2
Functional Difficulty Index	15	0	Yes	2
Illness Impact Scale	9	0	Yes	2
Visual Analogue Scale (overall health)	1	0	Yes	2
Self-rated health*	-	-	-	1
Freiburger Illness Coping Strategies questionnaire*	-	-	-	1
Brief Symptom Inventory-18	18	3	Yes	1

This article is protected by copyright. All rights reserved.

Constructed Meaning Scale	8	0	Yes	1
Surgical Recovery Score	31	0	Yes	1
Nottingham Health Profile	45	6	Yes	1
Duke Generic Instrument	17	11	Yes	1
Instrumental Activities of Daily Living	7	0	Yes	1
Profile of Moods States	65	6	Yes	1
Health and Activities Limitation Index	8	2	Yes	1
Health Utility Index	7	7	Yes	1
Spitzer Quality of Life Index	5	5	Yes	1
Global Quality of Life*	-	-	-	1
Multidimensional Functional Assessment Questionnaire*	-	-	-	1
Symptom Experience Scale	24	6	Yes	1
Ad-hoc satisfaction questionnaire#	6	6	Yes	1
Name of cancer-specific questionnaire (n=4)				
EORTC QLQ-C30	30	15	No	50
Cancer-related Health Worries Scale	4	0	Yes	2
Quality of Life – Cancer Survivors	41	4	No	1
Cancer Problems in Living Scale	31	0	Yes	1
Name of disease-specific questionnaire (n=11)				
EORTC QLQ-CR38	38	9	No	33
Functional Assessment of Cancer Therapy-Colorectal	37	5	Yes	5
Modified City of Hope Quality of life-Ostomy	41	6	Yes	2
EORTC QLQ-CR29	34	4	No	1
University of Padova Bowel Function questionnaire	8	0	Yes	1
Bowel Function questionnaire	8	0	Yes	1
Bowel Problems Scale	7	7	No	1
Late Effects Normal Tissue – Subjective, objective, management, analytic scale*	-	-	-	1
Quality of Life Index for Colostomy Patients	23	3	No	1
Colorectal Cancer Quality of Life	62	4	Yes	1
COloREctal Functional Outcome Questionnaire	26	5	Yes	1
Name of symptom-specific questionnaire (n=17)				
International Index of Erectile Function	15	5	Yes	4

This article is protected by copyright. All rights reserved.

Faecal Incontinence Quality of Life questionnaire	29	4	No	3
Wexner Incontinence Scale	5	0	Yes	3
Visual Analogue Scale (pain)	1	0	Yes	3
Center for Epidemiologic Studies – Depression	20	6	Yes	3
Hospital Anxiety and Depression Scale	14	0	Yes	2
Holschneider questionnaire	8	0	Yes	1
Internation Index of Erectile Function-5	5	5	Yes	1
Body Image Questionnaire	10	2	No	1
Body Image Scale	10	0	Yes	1
Faecal Incontinence Scoring System	5	0	Yes	1
Patient Assessment of Constipation Symptom Scale	12	3	Yes	1
Present Pain Intensity Index	1	0	Yes	1
Satisfaction with Sexual Function	1	0	Yes	1
Visual Analogue Scale (wound satisfaction)	1	0	Yes	1
Symptom Distress Scale	15	0	Yes	1
Multidimensional Fatigue Inventory-20	20	5	No	1
Name of questionnaire (n=3)				
Ad hoc QOL questionnaire A*#	-	-	-	1
Ad-hoc QOL questionnaire B*#	-	-	-	1
Ad-hoc QOL questionnaire C*#	-	-	-	1

Abbreviations: EORTC: European Organisation for Research and Treatment of Cancer *Questionnaire not available # PROM not validated

Table 3 Summary of domain categorization including number of items per domain, numbers of PROMs, and median items per PROM

PRO Domain (n=51)	Number of items n=917 (%)	Number of PROMs n=50 (%)	Median items per source PROM (range)
Psychological domains			
Anxiety	85 (9.2)	22 (44)	2.5 (1-12)
Fatigue	67 (7.2)	21 (42)	1.0 (1-23)
Depression	47 (5.1)	16 (32)	1.5 (1-12)
Body image	37 (4.0)	13 (26)	1.0 (1-10)
Frustration/irritability	15 (1.6)	7 (14)	1.0 (1-9)
Outlook on life	13 (1.4)	5 (10)	2.0 (1-6)
Self-esteem	11 (1.2)	6 (12)	2.0 (1-3)
Coping	10 (1.1)	6 (12)	1.0 (1-3)
Spiritual	7 (0.7)	2 (4)	3.5 (3-4)
Regret	5 (0.5)	2 (4)	2.5 (1-4)
Control	3 (0.3)	3 (6)	1.0 (1)
Functional domains			
Physical function	63 (6.8)	19 (38)	1.0 (1-9)
Role function	51 (5.5)	20 (40)	2.0 (1-7)
Social function	50 (5.4)	22 (44)	2.0 (1-8)
Sexual function	44 (4.7)	13 (26)	1.0 (1-15)
Cognitive function	30 (3.2)	14 (28)	1.0 (1-7)
Symptom domains			
Faecal incontinence	53 (5.7)	12 (24)	2.0 (1-27)
Stoma problems	52 (5.6)	5 (10)	7.0 (7-21)
Pain	50 (5.4)	18 (36)	1.5 (1-8)
Insomnia	18 (1.9)	13 (26)	1.0 (1-4)
Appetite/eating problems	17 (1.8)	10 (20)	1.5 (1-3)

Faecal frequency	14 (1.5)	8 (16)	2.0 (1-3)
Nausea/vomiting	12 (1.3)	8 (16)	1.0 (1-3)
Faecal Urgency	11 (1.2)	8 (16)	1.0 (1-2)
Flatulence or Gas	11 (1.2)	7 (14)	1.0 (1-3)
Treatment problems	11 (1.2)	7 (14)	1.0 (1-3)
Rectal blood or mucus	10 (1.1)	8 (16)	1.0 (1-2)
Bloating	7 (0.7)	6 (12)	1.0 (1-2)
Diarrhoea	7 (0.7)	7 (14)	1.0 (1)
Tenesmus	7 (0.7)	4 (8)	2.0 (1-2)
Constipation	6 (0.6)	5 (10)	1.0 (1-2)
Shortness of breath	5 (0.5)	5 (10)	1.0 (1)
Urinary Frequency	5 (0.5)	3 (6)	2.0 (1-2)
Faint or Dizzy	4 (0.4)	4 (8)	1.0 (1)
Hair Problems	4 (0.4)	4 (8)	1.0 (1)
Discrimination	3 (0.3)	3 (6)	1.0 (1)
Dry Mouth	3 (0.3)	3 (6)	1.0 (1)
Menstruation	3 (0.3)	3 (6)	1.0 (1)
Taste	3 (0.3)	3 (6)	1.0 (1)
Duration of bowel movement	2 (0.2)	2 (4)	1.0 (1)
Dyspepsia	2 (0.2)	2 (4)	1.0 (1)
Dysphagia	2 (0.2)	1 (2)	2.0 (2)
Dysuria	1 (0.1)	1 (2)	1.0 (1)
Urinary Incontinence	1 (0.1)	1 (2)	1.0 (1)
Global domains			
Global quality of life	12 (1.3)	9 (18)	1.0 (1-2)
Self-care	10 (1.1)	10 (20)	1.0 (1)
Financial	8 (0.9)	5 (10)	1.0 (1-4)
Satisfaction with care	6 (0.6)	1 (2)	6.0 (6)
Information needs	1 (0.1)	1 (2)	1.0 (1)

Table 4: OvidSP version of Medline search strategy

Search criteria	Search terms
Colorectal cancer	<ol style="list-style-type: none">1. exp Colonic Neoplasms/2. exp Rectal Neoplasms/3. ((colorect\$ or colon or colonic or rect\$) adj3 (cancer\$ or tumor\$ or neoplasm\$ or carcinoma\$ or adenocarcinoma\$ or malignan\$)).tw.4. or/1-3
Surgery	<ol style="list-style-type: none">1. exp Specialties, Surgical/2. surg\$.tw.3. operat\$.tw.4. intervention\$.tw.5. procedur\$.tw.6. resect\$.tw.7. or/1-6
Randomised controlled trials/prospective studies	<ol style="list-style-type: none">1. randomized controlled trial.pt.2. controlled clinical trial.pt.3. randomized controlled trials.sh.4. random allocation.sh.5. double blind method.sh.6. single-blind method.sh.7. or/1-68. exp animals/ not human/9. 7 not 810. clinical trial.pt.11. exp clinical trials/12. (clin\$ adj25 trial\$.ti,ab.13. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj25 (blind\$ or mask\$)).ti,ab.14. placebos.sh.15. placebo\$.ti,ab.16. random\$.ti,ab.17. research design.sh.18. or/10-1719. 18 not 820. 19 not 921. comparative study.sh.22. exp evaluation studies/23. follow up studies.sh.24. prospective studies.sh.25. (control\$ or 18prospective\$ or volunteer\$).ti,ab.26. or/21-2527. 26 not 828. 27 not (9 or 20)29. 9 or 20 or 28

Acronyms

COS – Core outcome set

CRC – Colorectal cancer

PRO – Patient-reported outcome

References

1. Speight J, Barendse SM. FDA guidance on patient reported outcomes. *Bmj* 2010; **340**: c2921.
2. McNair AG, Blazeby JM. Health-related quality-of-life assessment in GI cancer randomized trials: improving the impact on clinical practice. *Expert review of pharmacoeconomics & outcomes research* 2009; **9**(6): 559-67.
3. Pachler J, Wille-Jorgensen P. Quality of life after rectal resection for cancer, with or without permanent colostomy. *Cochrane Db Syst Rev* 2012; (12).
4. Parameswaran R, McNair A, Avery KN, et al. The role of health-related quality of life outcomes in clinical decision making in surgery for esophageal cancer: a systematic review. *Annals of surgical oncology* 2008; **15**(9): 2372-9.
5. Kirkham JJ, Dwan KM, Altman DG, et al. The impact of outcome reporting bias in randomised controlled trials on a cohort of systematic reviews. *Bmj* 2010; **340**: c365.
6. Williamson PR, Altman DG, Blazeby JM, et al. Developing core outcome sets for clinical trials: issues to consider. *Trials* 2012; **13**: 132.
7. Boers M, Brooks P, Strand CV, Tugwell P. The OMERACT filter for Outcome Measures in Rheumatology. *The Journal of rheumatology* 1998; **25**(2): 198-9.
8. McGrath PJ, Walco GA, Turk DC, et al. Core outcome domains and measures for pediatric acute and chronic/recurrent pain clinical trials: PedIMMPACT recommendations. *The journal of pain : official journal of the American Pain Society* 2008; **9**(9): 771-83.
9. Devane D, Begley CM, Clarke M, Horey D, C OB. Evaluating maternity care: a core set of outcome measures. *Birth* 2007; **34**(2): 164-72.
10. Sanoff HK, Goldberg RM, Pignone MP. A systematic review of the use of quality of life measures in colorectal cancer research with attention to outcomes in elderly patients. *Clinical colorectal cancer* 2007; **6**(10): 700-9.
11. Efficace F, Bottomley A, Vanvoorden V, Blazeby JM. Methodological issues in assessing health-related quality of life of colorectal cancer patients in randomised controlled trials. *European Journal of Cancer* 2004; **40**: 187-97.

This article is protected by copyright. All rights reserved.

12. Bartels SA, Vlug MS, Ubbink DT, Bemelman WA. Quality of life after laparoscopic and open colorectal surgery: a systematic review. *World journal of gastroenterology : WJG* 2010; **16**(40): 5035-41.
13. Jansen L, Koch L, Brenner H, Arndt V. Quality of life among long-term (≥ 5 years) colorectal cancer survivors - Systematic review. *European Journal of Cancer* 2010; **46**(16): 2879-88.
14. Gujral S, Avery KN, Blazeby JM. Quality of life after surgery for colorectal cancer: clinical implications of results from randomised trials. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer* 2008; **16**(2): 127-32.
15. Allal AS, Bieri S, Pelloni A, et al. Sphincter-sparing surgery after preoperative radiotherapy for low rectal cancers: feasibility, oncologic results and quality of life outcomes. *British journal of cancer* 2000; **82**(6): 1131-7.
16. Allal AS, Gervaz P, Gertsch P, et al. Assessment of quality of life in patients with rectal cancer treated by preoperative radiotherapy: a longitudinal prospective study. *International journal of radiation oncology, biology, physics* 2005; **61**(4): 1129-35.
17. Anthony T, Jones C, Antoine J, Sivess-Franks S, Turnage R. The effect of treatment for colorectal cancer on long-term health-related quality of life. *Annals of surgical oncology* 2001; **8**(1): 44-9.
18. Anthony T, Long J, Hynan LS, et al. Surgical complications exert a lasting effect on disease-specific health-related quality of life for patients with colorectal cancer. *Surgery* 2003; **134**(2): 119-25.
19. Arndt V, Merx H, Stegmaier C, Ziegler H, Brenner H. Quality of life in patients with colorectal cancer 1 year after diagnosis compared with the general population: a population-based study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2004; **22**(23): 4829-36.
20. Arndt V, Merx H, Stegmaier C, Ziegler H, Brenner H. Restrictions in quality of life in colorectal cancer patients over three years after diagnosis: a population based study. *Eur J Cancer* 2006; **42**(12): 1848-57.
21. Bailey C, Corner J, Addington-Hall J, Kumar D, Haviland J. Older patients' experiences of treatment for colorectal cancer: an analysis of functional status and service use. *European journal of cancer care* 2004; **13**(5): 483-93.
22. Baker F, Denniston M, Smith T, West MM. Adult cancer survivors: How are they faring? *Cancer* 2005; **104**(11): 2565-76.
23. Bloemen JG, Visschers RG, Truin W, Beets GL, Konsten JL. Long-term quality of life in patients with rectal cancer: association with severe postoperative complications and presence of a stoma. *Diseases of the colon and rectum* 2009; **52**(7): 1251-8.

24. Borrás JM, Sanchez-Hernandez A, Navarro M, et al. Compliance, satisfaction, and quality of life of patients with colorectal cancer receiving home chemotherapy or outpatient treatment: a randomised controlled trial. *Brit Med J* 2001; **322**(7290): 826-8A.
25. Braga M, Frasson M, Vignali A, Zuliani W, Civelli V, Di Carlo V. Laparoscopic vs. open colectomy in cancer patients: long-term complications, quality of life, and survival. *Diseases of the colon and rectum* 2005; **48**(12): 2217-23.
26. Bruheim K, Guren MG, Skovlund E, et al. Late side effects and quality of life after radiotherapy for rectal cancer. *International journal of radiation oncology, biology, physics* 2010; **76**(4): 1005-11.
27. Bruheim K, Tveit KM, Skovlund E, et al. Sexual function in females after radiotherapy for rectal cancer. *Acta oncologica* 2010; **49**(6): 826-32.
28. Caffo O, Amichetti M, Romano M, Maluta S, Tomio L, Galligioni E. Evaluation of toxicity and quality of life using a diary card during postoperative radiotherapy for rectal cancer. *Diseases of the colon and rectum* 2002; **45**(4): 459-65; discussion 65-7.
29. Camilleri-Brennan J, Steele RJ. Objective assessment of morbidity and quality of life after surgery for low rectal cancer. *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland* 2002; **4**(1): 61-6.
30. Canda AE, Terzi C, Gorken IB, Oztop I, Sokmen S, Fuzun M. Effects of preoperative chemoradiotherapy on anal sphincter functions and quality of life in rectal cancer patients. *International journal of colorectal disease* 2010; **25**(2): 197-204.
31. Celentano V, Fabbrocile G, Luglio G, Antonelli G, Tarquini R, Bucci L. Prospective study of sexual dysfunction in men with rectal cancer: feasibility and results of nerve sparing surgery. *International journal of colorectal disease* 2010; **25**(12): 1441-5.
32. Chambers SK, Lynch BM, Aitken J, Baade P. Relationship Over Time Between Psychological Distress and Physical Activity in Colorectal Cancer Survivors. *Journal of Clinical Oncology* 2009; **27**(10): 1600-6.
33. Chau I, Norman AR, Cunningham D, et al. Longitudinal quality of life and quality adjusted survival in a randomised controlled trial comparing six months of bolus fluorouracil/leucovorin vs. twelve weeks of protracted venous infusion fluorouracil as adjuvant chemotherapy for colorectal cancer. *Eur J Cancer* 2005; **41**(11): 1551-9.
34. Chen RC, Mamon HJ, Chen YH, et al. Patient-reported acute gastrointestinal symptoms during concurrent chemoradiation treatment for rectal cancer. *Cancer* 2010; **116**(8): 1879-86.
35. Courneya KS, Friedenreich CM, Quinney HA, Fields AL, Jones LW, Fairey AS. A randomized trial of exercise and quality of life in colorectal cancer survivors. *European journal of cancer care* 2003; **12**(4): 347-57.
36. Deimling GT, Bowman KF, Sterns S, Wagner LJ, Kahana B. Cancer-related health worries and psychological distress among older adult, long-term cancer survivors. *Psycho-Oncol* 2006; **15**(4): 306-20.

This article is protected by copyright. All rights reserved.

37. Deimling GT, Schaefer ML, Kahana B, Bowman KF, Reardon J. Racial differences in the health of older-adult long-term cancer survivors. *J Psychosoc Oncol* 2002; **20**(4): 71-94.
38. Deimling GT, Sterns S, Bowman KF, Kahana B. The health of older-adult, long-term cancer survivors. *Cancer nursing* 2005; **28**(6): 415-24.
39. Dobrowolski S, Hac S, Kobiela J, Sledzinski Z. Should we preserve the inferior mesenteric artery during sigmoid colectomy? *Neurogastroent Motil* 2009; **21**(12).
40. Engel J, Kerr J, Schlesinger-Raab A, Eckel R, Sauer H, Holzel D. Quality of life in rectal cancer patients: a four-year prospective study. *Annals of surgery* 2003; **238**(2): 203-13.
41. Fields AL, Keller A, Schwartzberg L, et al. Adjuvant therapy with the monoclonal antibody Edrecolomab plus fluorouracil-based therapy does not improve overall survival of patients with stage III colon cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2009; **27**(12): 1941-7.
42. Fucini C, Gattai R, Urena C, Bandettini L, Elbetti C. Quality of life among five-year survivors after treatment for very low rectal cancer with or without a permanent abdominal stoma. *Annals of surgical oncology* 2008; **15**(4): 1099-106.
43. Fujii S, Ota M, Ichikawa Y, et al. Comparison of short, long-term surgical outcomes and mid-term health-related quality of life after laparoscopic and open resection for colorectal cancer: a case-matched control study. *International journal of colorectal disease* 2010; **25**(11): 1311-23.
44. Furst A, Burghofer K, Hutzel L, Jauch KW. Neorectal reservoir is not the functional principle of the colonic J-pouch: the volume of a short colonic J-pouch does not differ from a straight coloanal anastomosis. *Diseases of the colon and rectum* 2002; **45**(5): 660-7.
45. Given B, Given C, Azzouz F, Stommel M. Physical functioning of elderly cancer patients prior to diagnosis and following initial treatment. *Nursing research* 2001; **50**(4): 222-32.
46. Gosselink MP, Busschbach JJ, Dijkhuis CM, Stassen LP, Hop WC, Schouten WR. Quality of life after total mesorectal excision for rectal cancer. *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland* 2006; **8**(1): 15-22.
47. Grumann MM, Noack EM, Hoffmann IA, Schlag PM. Comparison of quality of life in patients undergoing abdominoperineal extirpation or anterior resection for rectal cancer. *Annals of surgery* 2001; **233**(2): 149-56.
48. Guillou PJ, Quirke P, Thorpe H, et al. Short-term endpoints of conventional versus laparoscopic-assisted surgery in patients with colorectal cancer (MRC CLASICC trial): multicentre, randomised controlled trial. *Lancet* 2005; **365**(9472): 1718-26.
49. Guren MG, Dueland S, Skovlund E, Fossa SD, Poulsen JP, Tveit KM. Quality of life during radiotherapy for rectal cancer. *Eur J Cancer* 2003; **39**(5): 587-94.
50. Guren MG, Eriksen MT, Wiig JN, et al. Quality of life and functional outcome following anterior or abdominoperineal resection for rectal cancer. *European journal of surgical oncology :*

This article is protected by copyright. All rights reserved.

the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology 2005; **31**(7): 735-42.

51. Hamashima C. Long-term quality of life of postoperative rectal cancer patients. *Journal of gastroenterology and hepatology* 2002; **17**(5): 571-6.
52. Harisi R, Bodoky G, Borsodi M, Flautner L, Weltner J. Rectal cancer therapy: decision making on basis of quality of life? *Zentralblatt fur Chirurgie* 2004; **129**(2): 139-48.
53. Hirche C, Mrak K, Kneif S, et al. Perineal colostomy with spiral smooth muscle graft for neosphincter reconstruction following abdominoperineal resection of very low rectal cancer: long-term outcome. *Diseases of the colon and rectum* 2010; **53**(9): 1272-9.
54. Bramhall SR, Hallissey MT, Whiting J, et al. Marimastat as maintenance therapy for patients with advanced gastric cancer: a randomised trial. *British journal of cancer* 2002; **86**(12): 1864-70.
55. Hoerske C, Weber K, Goehl J, Hohenberger W, Merkel S. Long-term outcomes and quality of life after rectal carcinoma surgery. *The British journal of surgery* 2010; **97**(8): 1295-303.
56. Holzner B, Kemmler G, Greil R, et al. The impact of hemoglobin levels on fatigue and quality of life in cancer patients. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2002; **13**(6): 965-73.
57. Janson M, Lindholm E, Anderberg B, Haglund E. Randomized trial of health-related quality of life after open and laparoscopic surgery for colon cancer. *Surgical endoscopy* 2007; **21**(5): 747-53.
58. Jayne DG, Guillou PJ, Thorpe H, et al. Randomized trial of laparoscopic-assisted resection of colorectal carcinoma: 3-year results of the UK MRC CLASICC Trial Group. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2007; **25**(21): 3061-8.
59. Jess P, Christiansen J, Bech P. Quality of life after anterior resection versus abdominoperineal extirpation for rectal cancer. *Scandinavian journal of gastroenterology* 2002; **37**(10): 1201-4.
60. Kang SB, Park JW, Jeong SY, et al. Open versus laparoscopic surgery for mid or low rectal cancer after neoadjuvant chemoradiotherapy (COREAN trial): short-term outcomes of an open-label randomised controlled trial. *The lancet oncology* 2010; **11**(7): 637-45.
61. Kerr J, Engel J, Schlesinger-Raab A, Sauer H, Holzel D. Doctor-patient communication: results of a four-year prospective study in rectal cancer patients. *Diseases of the colon and rectum* 2003; **46**(8): 1038-46.
62. Kim DJ, Kim TI, Suh JH, et al. Oral tegafur-uracil plus folinic acid versus intravenous 5-fluorouracil plus folinic acid as adjuvant chemotherapy of colon cancer. *Yonsei medical journal* 2003; **44**(4): 665-75.

63. Kim NK, Min JS, Park JK, et al. Intravenous 5-fluorouracil versus oral doxifluridine as preoperative concurrent chemoradiation for locally advanced rectal cancer: prospective randomized trials. *Japanese journal of clinical oncology* 2001; **31**(1): 25-9.
64. King PM, Blazeby JM, Ewings P, et al. Randomized clinical trial comparing laparoscopic and open surgery for colorectal cancer within an enhanced recovery programme. *The British journal of surgery* 2006; **93**(3): 300-8.
65. King PM, Blazeby JM, Ewings P, Kennedy RH. Detailed evaluation of functional recovery following laparoscopic or open surgery for colorectal cancer within an enhanced recovery programme. *International journal of colorectal disease* 2008; **23**(8): 795-800.
66. Ko CY, Maggard M, Livingston EH. Evaluating health utility in patients with melanoma, breast cancer, colon cancer, and lung cancer: A nationwide, population-based assessment. *J Surg Res* 2003; **114**(1): 1-5.
67. Koda K, Yasuda H, Hirano A, et al. Evaluation of Postoperative Damage to Anal Sphincter/Levator Ani Muscles with Three-Dimensional Vector Manometry after Sphincter-Preserving Operation for Rectal Cancer. *J Am Coll Surgeons* 2009; **208**(3): 362-7.
68. Kopp I, Bauhofer A, Koller M. Understanding quality of life in patients with colorectal cancer: comparison of data from a randomised controlled trial, a population based cohort study and the norm reference population. *Inflammation research : official journal of the European Histamine Research Society [et al]* 2004; **53 Suppl 2**: S130-5.
69. Krouse RS, Herrinton LJ, Grant M, et al. Health-related quality of life among long-term rectal cancer survivors with an ostomy: manifestations by sex. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2009; **27**(28): 4664-70.
70. Kurtz ME, Kurtz JC, Stommel M, Given CW, Given B. Predictors of depressive symptomatology of geriatric patients with colorectal cancer: a longitudinal view. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer* 2002; **10**(6): 494-501.
71. Kuzu MA, Topcu O, Ucar K, et al. Effect of sphincter-sacrificing surgery for rectal carcinoma on quality of life in Muslim patients. *Diseases of the colon and rectum* 2002; **45**(10): 1359-66.
72. Lundy JJ, Coons SJ, Wendel C, et al. Exploring household income as a predictor of psychological well-being among long-term colorectal cancer survivors. *Qual Life Res* 2009; **18**(2): 157-61.
73. Marijnen CA, van de Velde CJ, Putter H, et al. Impact of short-term preoperative radiotherapy on health-related quality of life and sexual functioning in primary rectal cancer: report of a multicenter randomized trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2005; **23**(9): 1847-58.
74. Mik M, Narbutt P, Tchorzewski M, Dziki A. Functional outcomes after anterior resection for rectal cancer. *Clinical and Experimental Medical Letters* 2009; **50**(3): 193-6.

This article is protected by copyright. All rights reserved.

75. Mosconi P, Apolone G, Barni S, Secondino S, Sbanotto A, Filiberti A. Quality of life in breast and colon cancer long-term survivors: an assessment with the EORTC QLQ-C30 and SF-36 questionnaires. *Tumori* 2002; **88**(2): 110-6.
76. Mosher CE, Sloane R, Morey MC, et al. Associations Between Lifestyle Factors and Quality of Life Among Older Long-Term Breast, Prostate, and Colorectal Cancer Survivors. *Cancer* 2009; **115**(17): 4001-9.
77. Ong J, Ho KS, Chew MH, Eu KW. Prospective randomised study to evaluate the use of DERMABOND ProPen (2-octylcyanoacrylate) in the closure of abdominal wounds versus closure with skin staples in patients undergoing elective colectomy. *International journal of colorectal disease* 2010; **25**(7): 899-905.
78. Park JG, Lee MR, Lim SB, et al. Colonic J-pouch anal anastomosis after ultralow anterior resection with upper sphincter excision for low-lying rectal cancer. *World journal of gastroenterology : WJG* 2005; **11**(17): 2570-3.
79. Perez Lara FJ, Navarro Pinero A, de la Fuente Perucho A. Study of factors related to quality of life in patients with locally advanced rectal cancer. *Revista espanola de enfermedades digestivas : organo oficial de la Sociedad Espanola de Patologia Digestiva* 2004; **96**(11): 746-57.
80. Persson CR, Johansson BB, Sjoden PO, Glimelius BL. A randomized study of nutritional support in patients with colorectal and gastric cancer. *Nutrition and cancer* 2002; **42**(1): 48-58.
81. Phipps E, Braitman LE, Stites S, Leighton JC. Quality of life and symptom attribution in long-term colon cancer survivors. *J Eval Clin Pract* 2008; **14**(2): 254-8.
82. Pietrangeli A, Pugliese P, Perrone M, Sperduti I, Cosimelli M, Jandolo B. Sexual dysfunction following surgery for rectal cancer - a clinical and neurophysiological study. *Journal of experimental & clinical cancer research : CR* 2009; **28**: 128.
83. Pollack J, Holm T, Cedermark B, et al. Late adverse effects of short-course preoperative radiotherapy in rectal cancer. *The British journal of surgery* 2006; **93**(12): 1519-25.
84. Pucciarelli S, Del Bianco P, Toppan P, et al. Health-related quality of life outcomes in disease-free survivors of mid-low rectal cancer after curative surgery. *Annals of surgical oncology* 2008; **15**(7): 1846-54.
85. Ramsey SD, Berry K, Moinpour C, Giedzinska A, Andersen MR. Quality of life in long term survivors of colorectal cancer. *The American journal of gastroenterology* 2002; **97**(5): 1228-34.
86. Rauch P, Miny J, Conroy T, Neyton L, Guillemin F. Quality of life among disease-free survivors of rectal cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2004; **22**(2): 354-60.
87. Ravasco P, Monteiro-Grillo I, Vidal PM, Camilo ME. Cancer: disease and nutrition are key determinants of patients' quality of life. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer* 2004; **12**(4): 246-52.

88. Ravasco P, Monteiro-Grillo I, Vidal PM, Camilo ME. Dietary counseling improves patient outcomes: a prospective, randomized, controlled trial in colorectal cancer patients undergoing radiotherapy. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2005; **23**(7): 1431-8.
89. Ross L, Abild-Nielsen AG, Thomsen BL, Karlsen RV, Boesen EH, Johansen C. Quality of life of Danish colorectal cancer patients with and without a stoma. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer* 2007; **15**(5): 505-13.
90. Sailer M, Fuchs KH, Fein M, Thiede A. Randomized clinical trial comparing quality of life after straight and pouch coloanal reconstruction. *The British journal of surgery* 2002; **89**(9): 1108-17.
91. Sammour T, Kahokehr A, Hayes J, Hulme-Moir M, Hill AG. Warming and humidification of insufflation carbon dioxide in laparoscopic colonic surgery: a double-blinded randomized controlled trial. *Annals of surgery* 2010; **251**(6): 1024-33.
92. Sapp AL, Trentham-Dietz A, Newcomb PA, Hampton JM, Moinpour CM, Remington PL. Social networks and quality of life among female long-term colorectal cancer survivors. *Cancer* 2003; **98**(8): 1749-58.
93. Scarpa M, Erroi F, Ruffolo C, et al. Minimally invasive surgery for colorectal cancer: quality of life, body image, cosmesis, and functional results. *Surgical Endoscopy and Other Interventional Techniques* 2009; **23**(3): 577-82.
94. Schmidt C, Daun A, Malchow B, Kuchler T. Sexual impairment and its effects on quality of life in patients with rectal cancer. *Deutsches Arzteblatt international* 2010; **107**(8): 123-30.
95. Schmidt CE, Bestmann B, Kuchler T, Longo WE, Kremer B. Ten-year historic cohort of quality of life and sexuality in patients with rectal cancer. *Diseases of the colon and rectum* 2005; **48**(3): 483-92.
96. Schmidt CE, Bestmann B, Kuchler T, Longo WE, Kremer B. Impact of age on quality of life in patients with rectal cancer. *World journal of surgery* 2005; **29**(2): 190-7.
97. Schwenk W, Studiengruppe LC. [The LAPDIV-CAMIC Study. Multicenter prospective randomized study of short-term and intermediate-term outcome of laparoscopic and conventional sigmoid resection in diverticular disease]. *Der Chirurg; Zeitschrift fur alle Gebiete der operativen Medizen* 2004; **75**(7): 706-7.
98. Siassi M, Weiss M, Hohenberger W, Losel F, Matzel K. Personality Rather Than Clinical Variables Determines Quality of Life After Major Colorectal Surgery. *Diseases of the Colon & Rectum* 2009; **52**(4): 662-8.
99. Sideris L, Zenasni F, Vernerey D, et al. Quality of life of patients operated on for low rectal cancer: impact of the type of surgery and patients' characteristics. *Diseases of the colon and rectum* 2005; **48**(12): 2180-91.

100. Song PH, Yun SM, Kim JH, Moon KH. Comparison of the erectile function in male patients with rectal cancer treated by preoperative radiotherapy followed by surgery and surgery alone. *International journal of colorectal disease* 2010; **25**(5): 619-24.
101. Stamopoulos P, Theodoropoulos GE, Papailiou J, et al. Prospective evaluation of sexual function after open and laparoscopic surgery for rectal cancer. *Surgical Endoscopy and Other Interventional Techniques* 2009; **23**(12): 2665-74.
102. Stephens RJ, Thompson LC, Quirke P, et al. Impact of short-course preoperative radiotherapy for rectal cancer on patients' quality of life: data from the Medical Research Council CR07/National Cancer Institute of Canada Clinical Trials Group C016 randomized clinical trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2010; **28**(27): 4233-9.
103. Sultan S, Fisher DA, Voils CI, Kinney AY, Sandler RS, Provenzale D. Impact of functional support on health-related quality of life in patients with colorectal cancer. *Cancer* 2004; **101**(12): 2737-43.
104. Tiv M, Puyraveau M, Mineur L, et al. Long-term quality of life in patients with rectal cancer treated with preoperative (chemo)-radiotherapy within a randomized trial. *Cancer radiotherapie : journal de la Societe francaise de radiotherapie oncologique* 2010; **14**(6-7): 530-4.
105. Tjandra JJ, Reading DM, McLachlan SA, et al. Phase II clinical trial of preoperative combined chemoradiation for T3 and T4 resectable rectal cancer: preliminary results. *Diseases of the colon and rectum* 2001; **44**(8): 1113-22.
106. Trentham-Dietz A, Remington PL, Moinpour CM, Hampton JM, Sapp AL, Newcomb PA. Health-related quality of life in female long-term colorectal cancer survivors. *The oncologist* 2003; **8**(4): 342-9.
107. Trninic Z, Vidacak A, Vrhovac J, Petrov B, Setka V. Quality of life after colorectal cancer surgery in patients from University Clinical Hospital Mostar, Bosnia and Herzegovina. *Collegium antropologicum* 2009; **33 Suppl 2**: 1-5.
108. Tsunoda A, Nakao K, Tsunoda Y, Watanabe M, Matsui N. Health-related quality of life of colorectal cancer patients receiving oral UFT plus leucovorin compared with those with surgery alone. *International journal of clinical oncology* 2010; **15**(2): 153-60.
109. van den Brink M, van den Hout WB, Kievit J, et al. The impact of diagnosis and treatment of rectal cancer on paid and unpaid labor. *Diseases of the colon and rectum* 2005; **48**(10): 1875-82.
110. Velenik V, Oblak I, Anderluh F. Long-term results from a randomized phase II trial of neoadjuvant combined-modality therapy for locally advanced rectal cancer. *Radiation oncology* 2010; **5**: 88.
111. Vironen JH, Kairaluoma M, Aalto AM, Kellokumpu IH. Impact of functional results on quality of life after rectal cancer surgery. *Diseases of the colon and rectum* 2006; **49**(5): 568-78.

112. Weeks JC, Nelson H, Gelber S, Sargent D, Schroeder G, Clinical Outcomes of Surgical Therapy Study G. Short-term quality-of-life outcomes following laparoscopic-assisted colectomy vs open colectomy for colon cancer: a randomized trial. *JAMA : the journal of the American Medical Association* 2002; **287**(3): 321-8.
113. Whistance RN, Conroy T, Chie W, et al. Clinical and psychometric validation of the EORTC QLQ-CR29 questionnaire module to assess health-related quality of life in patients with colorectal cancer. *European Journal of Cancer* 2009; **45**(17): 3017-26.
114. Whistance RN, Gilbert R, Fayers P, et al. Assessment of body image in patients undergoing surgery for colorectal cancer. *International journal of colorectal disease* 2010; **25**(3): 369-74.
115. Yau T, Watkins D, Cunningham D, Barbachano Y, Chau I, Chong G. Longitudinal Assessment of Quality of Life in Rectal Cancer Patients With or Without Stomas Following Primary Resection. *Diseases of the Colon & Rectum* 2009; **52**(4): 669-77.
116. Yau T, Watkins D, Cunningham D, Barbachano Y, Chau I, Chong G. Longitudinal assessment of quality of life in rectal cancer patients with or without stomas following primary resection. *Diseases of the colon and rectum* 2009; **52**(4): 669-77.
117. Yoo HJ, Kim JC, Eremenco S, Han OS. Quality of life in colorectal cancer patients with colectomy and the validation of the Functional Assessment of Cancer Therapy-Colorectal (FACT-C), Version 4. *Journal of pain and symptom management* 2005; **30**(1): 24-32.
118. Whistance RN, Forsythe RO, McNair AG, et al. A systematic review of outcome reporting in colorectal cancer surgery. *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland* 2013; **15**(10): e548-60.
119. Blencowe NS, Strong S, McNair AG, et al. Reporting of short-term clinical outcomes after esophagectomy: a systematic review. *Annals of surgery* 2012; **255**(4): 658-66.
120. Potter S, Brigid A, Whiting PF, et al. Reporting clinical outcomes of breast reconstruction: a systematic review. *Journal of the National Cancer Institute* 2011; **103**(1): 31-46.
121. Macefield RC, Jacobs M, Korffage IJ, et al. Developing core outcomes sets: methods for identifying and including patient-reported outcomes (PROs). *Trials* 2014; **15**: 49.

Figure 1

PRISMA diagram of studies considered for the systematic review.

