



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Frailty screening in critical care at scale

Citation for published version:

Pugh, R & Lone, NI 2021, 'Frailty screening in critical care at scale', *Chest Journal*, vol. 160, no. 4, pp. 1165-1166. <https://doi.org/10.1016/j.chest.2021.06.031>

Digital Object Identifier (DOI):

[10.1016/j.chest.2021.06.031](https://doi.org/10.1016/j.chest.2021.06.031)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Chest Journal

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





If there is Online Only content that cannot be converted to a Word processing format, you may have to click the Supplemental Files icon on the menu bar in your Reviewer Center to access.

Frailty screening in critical care at scale

(Editorial for: Routine frailty screening in critical illness- a population-based cohort study in Australia and New Zealand)

Journal:	<i>CHEST</i>
Manuscript ID	CHEST-21-2317
Article Type:	Editorial
Date Submitted by the Author:	17-Jun-2021
Complete List of Authors:	Pugh, Richard; Betsi Cadwaladr University Health Board, Department of Anaesthetics, Glan Clwyd Hospital Lone , Nazir; University of Edinburgh, Usher Institute ; NHS Lothian,
Keywords:	CRITICAL CARE, AGING, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abbreviation List

APACHE III – Acute Physiology and Chronic Health Evaluation III

CFS – Clinical Frailty Scale

ICU – Intensive Care Unit

CONFIDENTIAL

1
2
3 *Word Count: 986*
4
5

6 ***To accompany manuscript:*** Darvall JN, Bellomo R, Paul E, et al. Routine
7 *frailty screening in critical illness- a population-based cohort study in*
8 *Australia and New Zealand. Chest. 2021*
9
10
11
12
13

14 **Editorial**
15
16
17

18 **Frailty screening in critical care at scale**
19
20
21

22 **Dr Richard J. Pugh**
23 **Dr Nazir I. Lone**
24
25
26

27 richard.pugh@wales.nhs.uk
28 nazir.lone@ed.ac.uk
29
30
31
32
33
34

35 **Dr Richard Pugh**
36 **Consultant in Intensive Care Medicine**
37 **Department of Anaesthetics**
38 **Glan Clwyd Hospital**
39 **Bodelwyddan**
40 **Denbighshire**
41 **LL18 5UJ**
42
43
44
45
46
47
48
49
50

51 **Correspondence to:**
52 **Dr Richard Pugh**
53 **richard.pugh@wales.nhs.uk**
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conflict of interest statements for all authors -

The authors declare no conflict of interest.

CONFIDENTIAL

Frailty screening in critical care at scale

The concept of frailty as a determinant of outcome from critical illness has resonated hugely with the critical care community since its description a decade ago.¹ Having long appreciated the importance of “physiological” as opposed to “chronological” age, frailty is now widely understood to represent a state of diminished physiological reserve more prevalent with age, which is associated with increased vulnerability towards adverse outcomes, but which is distinct from co-morbidity.² The Clinical Frailty Scale (CFS) lends itself to screening for frailty in the critical care setting; it has “face validity,” combines clinical judgement with objective measurement, can be readily applied without adaptation, and has demonstrable reliability³ and predictive validity.⁴ Given global demographic trends and potential critical care resource requirements of an ageing population, the added value of frailty assessment to clinical discussions and decision-making, and to risk-adjusted outcome reporting have been highlighted previously.⁵ However, until now there has been no published research regarding the feasibility and prognostic value of population-scale screening for frailty in the critically ill.

Darvall and colleagues’ study published in this edition of Chest is therefore an important addition to the literature.⁶ In a well-designed study, they report successful implementation of routine frailty screening using the CFS across diverse ICUs in Australia and New Zealand. Frailty was common in this cohort, approaching 1 in 5 patients when assessed at the time of admission. Consistent with findings in other studies, increasing frailty was associated with higher hospital mortality. Furthermore, addition of CFS score to the APACHE III-j risk prediction model improved model performance as measured by discriminant function (area under the receiver operating characteristics curve).

Importantly, the authors looked beyond mortality and investigated whether frailty was associated with a number of additional person-centered outcomes. The presence of frailty increased length of ICU and hospital stay and increased the likelihood of discharge to a nursing home or chronic care facility. Furthermore, patients with frailty were more likely to experience delirium and pressure injuries within ICU. These findings provide a more complete picture of the likely consequences of ICU admission in the context of frailty to inform clinicians and patients.

A significant strength of the study was the extremely large sample size (n= 234,568). The authors were able to conduct analyses using the Australian and New Zealand Intensive Care Society Adult Patient Database, a bi-national audit database. This allowed associations to be reported with precision for each category of the CFS, rather than the dichotomization that other studies have needed to undertake. Furthermore, this larger sample size allowed subgroup analyses to be undertaken in a younger cohort of patients, confirming that the presence of frailty was associated with worse outcomes.

An important limitation was the proportion of missing data relating to CFS recording. With 1 in 3 observations missing a value for the primary exposure, the potential for bias could have been substantial. Ideally, evaluating the impact of missing data requires an understanding of the mechanism by which missing data has arisen, along with sensitivity

1
2
3 analyses to impute missing values if the mechanism of missingness is random.⁷ However,
4 the authors undertook a number of additional analyses which were reassuring, indicating
5 that the likelihood of significant bias due to missing data was low. This included a
6 comparison of baseline characteristics in those with and without missing data, and
7 subgroup analyses on ICUs with low levels of missing data.
8
9

10 Training of assessors was not mandated and the accuracy of CFS recording was not
11 evaluated. Although the authors have suggested that lack of individualized training was not
12 a significant barrier to implementation, this may have contributed to measurement error.
13 As long as such error is non-differential (not systematically biased towards over- or under-
14 estimating CFS), this would tend to reduce the strength of association between exposure
15 and outcome.⁸
16
17

18 Lastly, the CFS was originally developed for a cohort of patients aged 65 years or more.⁹
19 The inter-relationship between age, frailty and comorbidity is complex, and identification
20 of “frailty” may have different biological and clinical implications across the age spectrum.
21^{10,11} As such, this study provides much needed insight into the implications of frailty for
22 younger people in critical care. 23% patients were aged less than 50 years, among whom
23 frailty was identified in 6% (versus 23% in those 50 years and over). With the exceptions
24 of chronic cardiovascular disease and metastatic cancer, the distribution of chronic illness
25 followed a similar pattern according to frailty among younger and older patients. Variations
26 in processes of care were evident, with an apparent decreasing tendency to undergo
27 invasive ventilation with increasing frailty not observed in the younger cohort. However,
28 the relationship between frailty and main outcome measures (hospital mortality, length of
29 stay and discharge disposition) appeared consistent between age cohorts. Interestingly, risk
30 of readmission among frail patients was recently reported to decrease with age when frailty
31 was identified using administrative data;¹² longer-term outcome data which might
32 evidence rate and extent of recovery from critical illness according to CFS, age cohort and
33 comorbidity is therefore still greatly needed.
34
35
36
37

38 As with all studies using data collected in critical care audit databases, generalizing
39 findings to other settings requires an understanding of critical care service organization,
40 admission practices and prevailing cultural attitudes in wider society to critical care
41 admission. This means that the prevalence of frailty and its association with outcomes in
42 Australia and New Zealand may not necessarily translate to other settings. Similarly, whilst
43 a dose-response relationship was demonstrated between CFS and a range of outcomes,
44 these should be interpreted as being associations rather than ascribing causal inference.
45
46

47 To conclude, Darvall and colleagues have answered important questions as to the
48 feasibility and prognostic value of large-scale frailty screening in the critically ill.
49 Crucially, they have been able to evaluate the implications of frailty assessment among
50 younger patients and provide assurance of the validity of routine frailty assessment using
51 CFS among unselected patients admitted to adult critical care units.
52
53
54
55
56
57
58
59
60

References

1. McDermid RC, Stelfox HT, Bagshaw SM. Frailty in the critically ill: a novel concept. *Crit Care*. 2011;15(1):301.
2. Fried LP, Ferrucci L, Darer J, Williamson JD, Anderson G. Untangling the concepts of disability, frailty, and comorbidity: implications for improved targeting and care. *J Gerontol A Biol Sci Med Sci*. 2004;59(3):255-263.
3. Pugh RJ, Battle CE, Thorpe C, et al. Reliability of frailty assessment in the critically ill: a multicentre prospective observational study. *Anaesthesia*. 2019;74(6):758-764.
4. Muscedere J, Waters B, Varambally A, et al. The impact of frailty on intensive care unit outcomes: a systematic review and meta-analysis. *Intensive Care Medicine*. 2017;43(8):1105-1122.
5. De Biasio JC, Mittel AM, Mueller AL, Ferrante LE, Kim DH, Shaefi S. Frailty in Critical Care Medicine: A Review. *Anesth Analg*. 2020;130(6):1462-1473.
6. Darvall JN, Bellomo R, Paul E, et al. Routine frailty screening in critical illness- a population-based cohort study in Australia and New Zealand. *Chest*. 2021.
7. Sterne JA, White IR, Carlin JB, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ*. 2009;338:b2393.
8. Hutcheon JA, Chiolero A, Hanley JA. Random measurement error and regression dilution bias. *BMJ*. 2010;340:c2289.
9. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005;173(5):489-495.
10. Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS. Frailty and pre-frailty in middle-aged and older adults and its association with multimorbidity and mortality: a prospective analysis of 493 737 UK Biobank participants. *Lancet Public Health*. 2018;3(7):e323-e332.
11. Spiers GF, Kunonga TP, Hall A, et al. Measuring frailty in younger populations: a rapid review of evidence. *BMJ Open*. 2021;11(3):e047051.
12. Hill AD, Fowler RA, Wunsch H, Pinto R, Scales DC. Frailty and long-term outcomes following critical illness: A population-level cohort study. *J Crit Care*. 2021;62:94-100.