



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

On the fixation or nonfixation of inversions under epistatic selection

Citation for published version:

Charlesworth, B & Flatt, T 2021, 'On the fixation or nonfixation of inversions under epistatic selection', *Molecular Ecology*. <https://doi.org/10.1111/mec.16026>

Digital Object Identifier (DOI):

[10.1111/mec.16026](https://doi.org/10.1111/mec.16026)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Molecular Ecology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



On the fixation or nonfixation of inversions under epistatic selection

Brian Charlesworth¹  | Thomas Flatt² 

¹School of Biological Sciences, Institute of Evolutionary of Evolutionary Biology, University of Edinburgh, Edinburgh, UK

²Department of Biology, University of Fribourg, Fribourg, Switzerland

Correspondence

Brian Charlesworth, Institute of Evolutionary of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, UK.
Email: Brian.Charlesworth@ed.ac.uk

Thomas Flatt, Department of Biology, University of Fribourg, Fribourg, Switzerland
Email: thomas.flatt@unifr.ch

Abstract

Several recent publications have stated that epistatic fitness interactions cause the fixation of inversions that suppress recombination among the loci involved. Under this type of selection, however, the suppression of recombination in an inversion heterozygote can create a form of heterozygote advantage, which prevents the inversion from becoming fixed by selection. This process has been explicitly modelled by previous workers.

KEYWORDS

balanced polymorphism, epistasis, fitness, inversions, linkage disequilibrium recombination suppression

There is a growing interest in the evolutionary role of inversions and other types of chromosome rearrangements, with several recent review papers (e.g., Faria et al., 2019; Huang & Rieseberg, 2020; Kapun & Flatt, 2019; Wellenreuther & Bernatchez, 2018) and a special issue of *Molecular Ecology* devoted to this topic (Wellenreuther et al., 2019, and references therein). As these papers have pointed out, there are multiple ways in which natural selection can act on inversions. In order to discriminate between different hypotheses, it is necessary to have a clear understanding of their observable consequences. Unfortunately, there appears to be a serious misconception about one process that has been proposed as providing a selective advantage to an inversion. This involves Dobzhansky's (e.g., Dobzhansky, 1949, 1950, 1951) concept of "coadaptation" among polymorphic loci that interact in their effects on fitness, an idea that traces back to Fisher (1930, pp. 102–104).

In its simplest form, this model invokes two loci, A and B, each segregating for a pair of alleles (A_1 vs. A_2 , B_1 vs. B_2) in a diploid, randomly mating population. Assume that the fitness effects of the two loci are epistatic, in the sense that the fitnesses of the nine possible diploid genotypes at the two loci deviate from those predicted by additive combinations of the effects of the alleles at each locus

(Fisher, 1918). It is then possible for linkage disequilibrium (LD) to be maintained at a stable equilibrium in the face of recombination, where both loci are polymorphic, and fitter combinations of alleles are in excess of the frequencies expected by randomly combining alleles according to their frequencies (Fisher, 1930; Karlin, 1975; Kimura, 1956; Lewontin & Kojima, 1960). There is then selection for modifiers that reduce the rate of recombination between the two loci (Feldman, 1972; Fisher, 1930; Kimura, 1956). The same principle applies to more general multilocus systems in randomly mating populations (Charlesworth, 1976; Zhivotovsky et al., 1994), but not necessarily to partially inbreeding populations (Charlesworth et al., 1979). In particular, an inversion that arises on a haplotype that is present in excess of random expectation (and is thus fitter than average) experiences a selective advantage if recombination is suppressed in heterozygotes for the inversion, simply because the inversion maintains its association with a high-fitness genotype (Kimura, 1956; Charlesworth & Charlesworth, 1973).

The misconception is that this process causes the inversion to spread to fixation, implying that it cannot explain the balanced inversion polymorphisms that have been the subject of so much recent attention. This claim appears to have originated in table 1 of

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Molecular Ecology* published by John Wiley & Sons Ltd.

Kirkpatrick and Barton (2006), and has been repeated in review papers by Hoffmann and Rieseberg (2008), Kapun and Flatt (2019) and Huang and Rieseberg (2020). However, it overlooks the fact that crossing over is suppressed only in inversion heterozygotes, so that the selective benefits of recombination suppression confer a heterozygote advantage to the inversion, and hence prevent its fixation. This scenario was first modelled by Kimura (1956) and examined in more detail by Charlesworth (1974), who showed that the exchange of alleles among inverted and standard arrangements by gene conversion or double crossing over does not prevent the establishment and maintenance of an inversion polymorphism when there is epistatic selection. An example of this mechanism for maintaining inversion polymorphisms is provided by the epistatic interactions among the different components of segregation distorter systems, and probably explains the frequent association of such systems with inversions (Fuller et al., 2020). Direct effects of inversions in causing segregation distortion should, therefore, not be assumed without further evidence. Of course, there are processes other than epistatic selection that can select for inversions, as described in the reviews cited above, and we do not mean to imply that epistatic selection is necessarily involved in the maintenance of inversion polymorphisms.

AUTHOR CONTRIBUTIONS

Both authors wrote the paper.

DATA AVAILABILITY STATEMENT

No new data were generated in this study.

ORCID

Brian Charlesworth  <https://orcid.org/0000-0002-2706-355X>

Thomas Flatt  <https://orcid.org/0000-0002-5990-1503>

REFERENCES

- Charlesworth, B., & Charlesworth, D. (1973). Selection of new inversions in multi-locus genetic systems. *Genetical Research*, 21, 167–183.
- Charlesworth, B. (1974). Inversion polymorphism in a two-locus genetic system. *Genetical Research*, 23, 259–280. <https://doi.org/10.1017/S0016672300014919>
- Charlesworth, B. (1976). Recombination modification in a fluctuating environment. *Genetics*, 83, 181–195. <https://doi.org/10.1093/genetics/83.1.181>
- Charlesworth, D., Charlesworth, B., & Strobeck, C. (1979). Selection for recombination in partially self-fertilizing species. *Genetics*, 93, 237–244. <https://doi.org/10.1093/genetics/93.1.237>
- Dobzhansky, T. (1949). Observations and experiments on natural selection in *Drosophila*. *Heredity*, 35(S1), 210–224. <https://doi.org/10.1111/j.1601-5223.1949.tb03334.x>
- Dobzhansky, T. (1950). Genetics of natural populations. XIX. Origin of heterosis through natural selection in populations of *Drosophila pseudoobscura*. *Genetics*, 35, 288–302. <https://doi.org/10.1093/genetics/35.3.288>
- Dobzhansky, T. (1951). *Genetics and the Origin of Species* (3rd edn). Columbia University Press.

- Faria, R., Johannesson, K., Butlin, R. K., & Westram, A. M. (2019). Evolving inversions. *Trends in Ecology and Evolution*, 34, 239–248. <https://doi.org/10.1016/j.tree.2018.12.005>
- Feldman, M. W. (1972). Selection for linkage modification: I. Random mating populations. *Theoretical Population Biology*, 3, 324–346. [https://doi.org/10.1016/0040-5809\(72\)90007-X](https://doi.org/10.1016/0040-5809(72)90007-X)
- Fisher, R. A. (1918). The correlation between relatives on the supposition of Mendelian inheritance. *Transactions of the Royal Society of Edinburgh*, 52, 399–433. <https://doi.org/10.1017/S0080456800012163>
- Fisher, R. A. (1930). *The genetical theory of natural selection*. Clarendon Press.
- Fuller, Z. L., Koury, S. A., Leonard, C. J., Young, R. E., Ikegami, K., Westlake, J., Richards, S., Schaeffer, S. W., & Phadnis, N. (2020). Extensive recombination suppression and epistatic selection causes chromosome-wide differentiation of a selfish sex chromosome in *Drosophila pseudoobscura*. *Genetics*, 216, 205–226. <https://doi.org/10.1534/genetics.120.303460>
- Hoffmann, A. A., & Rieseberg, L. H. (2008). Revisiting the impact of inversions in evolution: from population genetic markers to drivers of adaptive shifts and speciation? *Annual Review of Ecology, Evolution, and Systematics*, 39, 21–42. <https://doi.org/10.1146/annurev.ecolsys.39.110707.173532>
- Huang, K., & Rieseberg, L. H. (2020). Frequency, origins, and evolutionary role of chromosomal inversions in plants. *Frontiers in Plant Science*, 1, 296. <https://doi.org/10.3389/fpls.2020.00296>
- Kapun, M., & Flatt, T. (2019). The adaptive significance of chromosomal inversion polymorphisms in *Drosophila melanogaster*. *Molecular Ecology*, 28, 1263–1282. <https://doi.org/10.1111/mec.14871>
- Karlin, S. (1975). General two-locus selection models: Some objectives, results and interpretations. *Theoretical Population Biology*, 7, 364–398. [https://doi.org/10.1016/0040-5809\(75\)90025-8](https://doi.org/10.1016/0040-5809(75)90025-8)
- Kimura, M. (1956). A model of a genetic system which leads to closer linkage by natural selection. *Evolution*, 10, 278–287. <https://doi.org/10.1111/j.1558-5646.1956.tb02852.x>
- Kirkpatrick, M., & Barton, N. (2006). Chromosome inversions, local adaptation and speciation. *Genetics*, 173, 419–434. <https://doi.org/10.1534/genetics.105.047985>
- Lewontin, R. C., & Kojima, K.-I. (1960). The evolutionary dynamics of complex polymorphisms. *Evolution*, 14, 458–472. <https://doi.org/10.1111/j.1558-5646.1960.tb03113.x>
- Wellenreuther, M., & Bernatchez, L. (2018). Eco-evolutionary genomics of chromosomal inversions. *Trends in Ecology and Evolution*, 33, 427–440. <https://doi.org/10.1016/j.tree.2018.04.002>
- Wellenreuther, M., Mérot, C., Berdan, E., & Bernatchez, L. (2019). Going beyond SNPs: The role of structural genomic variants in adaptive evolution and species diversification. *Molecular Ecology*, 28, 1203–1209. <https://doi.org/10.1111/mec.15066>
- Zhivotovsky, L. A., Feldman, M. W., & Christiansen, F. B. (1994). Evolution of recombination among multiple selected loci: a generalized reduction principle. *Proceedings of the National Academy of Sciences USA*, 91, 1079–1083. <https://doi.org/10.1073/pnas.91.3.1079>

How to cite this article: Charlesworth B, Flatt T. On the fixation or nonfixation of inversions under epistatic selection. *Mol Ecol*. 2021;00:1–2. <https://doi.org/10.1111/mec.16026>