



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

ALTERED PROTEIN COMPOSITION OF PORCINE FOLLICULAR FLUID DUE TO A HIGH-FIBRE DIET AND THE POTENTIAL FOR OPTIMISATION OF IN VITRO CULTURE MEDIA

Citation for published version:

Jarrett, S, Gill, AC, Thekkedath Kurian, D, Ferguson, EM & Ashworth, CJ 2015, 'ALTERED PROTEIN COMPOSITION OF PORCINE FOLLICULAR FLUID DUE TO A HIGH-FIBRE DIET AND THE POTENTIAL FOR OPTIMISATION OF IN VITRO CULTURE MEDIA', *Reproduction, Fertility and Development*, vol. 28, no. 2, pp. 170. <https://doi.org/10.1071/RDv28n2Ab81>

Digital Object Identifier (DOI):

[10.1071/RDv28n2Ab81](https://doi.org/10.1071/RDv28n2Ab81)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Reproduction, Fertility and Development

Publisher Rights Statement:

This is the author's peer-reviewed manuscript as accepted for publication

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Altered protein composition of porcine follicular fluid due to a high fibre diet and the potential for optimisation of *in vitro* culture media.

S. Jarrett¹, A. C. Gill¹, D. Kurian¹, E. M. Ferguson² and C. J. Ashworth¹

¹The Roslin Institute and R(D)SVS, The University of Edinburgh, Easter Bush, Midlothian, EH25 9RG, Scotland, UK;

²Aberdeen Maternity Hospital, Foresterhill, Aberdeen, AB25 2ZL, Scotland, UK

This study reports a proteomic analyses on porcine follicular fluid (FF) obtained from a previous nutritional trial, where oocytes from gilts fed a high fibre (HF) diet for the first 19 days of their third oestrous cycle produced blastocysts with more cells following *in vitro* maturation (IVM) and IVF compared with oocytes from control-fed (CON) pigs. Oocytes were matured in TCM-199 supplemented with LH and FSH at 0.5 mg mL⁻¹ and 10% of the animals' own pooled FF. Following IVF, resultant embryos were cultured in NCSU-23 medium for 6 to 7 days. We hypothesize that FF protein composition is altered by the HF diet and that this confers the reproductive benefits previously observed. The FF had previously been stored at 80°C after the IVF trials and was thawed for the current study, which compared the protein composition of pooled Day 19 FF from 12 CON pigs and 12 HF pigs. These gilts were a subset of the pigs described above with the largest FF volumes. The protein composition of pooled FF from 6 CON pigs whose oocytes produced blastocysts was compared with FF from 6 CON pigs whose oocytes did not produce blastocysts. The same analysis was carried out with the 6 HF pigs that produced blastocysts and the 6 HF pigs that did not produce blastocysts. Equal numbers of samples from animals were selected for experimental balance. The proteomic study was carried out in duplicate. Abundant proteins were depleted from FF by Proteominer enrichment. Samples were labelled by isotopic di-methylation, where in each analysis, one sample was labelled with a heavy methyl group, the other with a light methyl group. Proteins were detected by liquid chromatography tandem mass spectrometry. Protein identifications were filtered using a 1% false discovery threshold and a requirement for two or more peptides detected for each protein. Differentially expressed proteins (DEPs) were identified as having heavy/light ratios greater than 1.2 or less than 0.8, which are recognised cut-off points for differential expression in proteomics. Over 140 DEPs were detected between CON and HF samples, indicating a nutritional influence on FF protein composition. Over one-third (37%) of these DEPs were also differentially expressed in the blastocyst versus no blastocyst analyses, suggesting that the altered FF protein composition may affect IVF outcome. DEPs were submitted into Ingenuity Pathway Analysis to highlight associated canonical pathways and upstream regulators. Top ranking canonical pathways detected included coagulation system, acute phase response, and LXR/RXR activation pathways. Potential upstream regulators detected by IPA included transforming growth factor beta, tumour protein P53, and beta-oestradiol. These pathways and upstream regulators could serve as potential avenues for elucidating the mechanism(s) by which the HF diet results in the reproductive benefits and could lead to the refinement of IVM and IVF culture conditions.

This study was funded by AHDB Pork and BBSRC.