



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Sense and sensibility**

Of synthetic biology and the redesign of bioreporter circuits

**Citation for published version:**

Belkin, S & Wang, B 2021, 'Sense and sensibility: Of synthetic biology and the redesign of bioreporter circuits', *Microbial biotechnology*. <https://doi.org/10.1111/1751-7915.13955>

**Digital Object Identifier (DOI):**

[10.1111/1751-7915.13955](https://doi.org/10.1111/1751-7915.13955)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Microbial biotechnology

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Editorial

## Sense and sensibility: of synthetic biology and the redesign of bioreporter circuits

It is tempting to speculate that sixty years ago, when Jacob and Monod presented their model of the *lac* operon (Jacob and Monod, 1961), they already had a glimpse of the future of the *lacZ* gene, not only as encoding a cleaver of disaccharides, nor as a component in a beautiful and groundbreaking model of gene regulation, but also as a universal reporter of gene activation. Indeed, reporter gene technology rapidly became a basic tool in studying the regulation of gene expression; several decades had to pass, however, before the same approach has led to the first report of a microorganism genetically engineered to perform an accurate, specific and sensitive analysis of an environmental pollutant (King *et al.*, 1990). The term ‘whole cell biosensor’ soon entered into use, accompanied by some semantic controversy: purists view the term ‘biosensor’ as a hardware device, in which the biological entity (e.g. enzyme, antibody, oligonucleotide or a live cell) serves as its sensing component (IUPAC, 2017); according to this view, a microbial strain, notwithstanding the complexity of its re-engineering, may be called a ‘sensor strain’ or a ‘bioreporter’, but never a ‘biosensor’. Long before this linguistic polemic became an issue, however, a pioneering article from the Sayler group (King *et al.*, 1990) described a bioluminescent *Pseudomonas*-based sensor of naphthalene. This publication was trailed by the first *E. coli*-based mercury sensor (Selifonova *et al.*, 1993), soon to be followed by numerous others, all sharing the same basic structure: a gene promoter induced by the target compound (directly, or via the removal of a repressor), fused downstream of a reporter gene. The latter could code for a traceable protein (e.g. GFP) or – more often – for an enzyme, the activity of which could be monitored quantitatively in real time (van der Meer and Belkin, 2010). When necessary, regulatory elements had to be cloned as well, especially when the gene promoter acting as the sensing element was not native to the host organism. In view of the practically infinite number of gene promoters and regulatory proteins available as candidate sensor elements, the scope of possible sensing targets of such sensors is exceptionally broad. In parallel to the development of microbial sensors of specific compounds, bioreporter strains have also been

described for the detection of global sample characteristics such as toxicity or genotoxicity/mutagenicity, parameters of importance for environmental health as well as for chemicals’ safety. The commercial SOS Chromotest (Quillardet *et al.*, 1982), the forerunner of this group of assays, was followed by the *umu*-test (Oda *et al.*, 1985). In both cases, the activation of gene promoters from the *E. coli* SOS repair regulon by DNA damaging agents was chromogenically monitored with *lacZ* as a reporter gene.

Looking back over the last 15 years, possibly the most powerful innovator of microbial biosensor design was the coming of age of synthetic biology. While the term has been introduced to the scientific literature over a century ago (Leduc, 1910), its meaning has slowly changed over the years. Following the introduction of the Jacob and Monod model, microbial biotechnology horizons opened up with the advent of increasingly more sophisticated molecular tools, including numerous enzymes derived from diverse microorganisms and viruses, harnessed and retrained to perform cutting, pasting and editing tricks. The same horizons practically exploded when thermophilic variants of these enzymes were ingeniously employed in the invention of PCR technology, and turned essentially limitless when genome sequencing was made trivial and bioinformatic data (and tools for its analysis) became freely available to all. These advances have prepared the ground for the invasion of practitioners of additional disciplines into the realm of whole cell sensor design; when engineers, physicists and computer scientists started to practice biology in earnest, things have started to become truly interesting. In 2004, van der Meer *et al.* have claimed that one of the reasons current bioreporters’ performance cannot comply with environmental detection standards is the ‘lack of engineering principles’. More or less at the same time, the ‘Bibricks’ concept has been presented (Knight, 2003), aiming to provide ‘a set of standard and reliable engineering mechanisms to remove much of the tedium and surprise during assembly of genetic components into larger systems’. The trend embodied by these two examples paved the ground for engineering school graduates to advise ‘classical’ molecular biologists involved in

microbial bioreporter design that the time of simplistic promoter-reporter fusions is over; more complex (and hopefully, more efficient and diverse) molecular sensor circuits can be designed, for both *in vivo* and *in vitro* expression, by employing an engineering-like point of view.

Indeed, synthetic biology adopts engineering principles (e.g. standardization, modularization and modelling) to facilitate complex genetic circuit construction, particularly using 'Lego-like' standardized building blocks (Endy, 2005). Though the blocks alone do not perform advanced actions, they can generate bespoke coordinated functions when connected. Hence, synthetic biology offers new tools to precisely manipulate cells for achieving customized tasks, using engineered gene circuits of varying scales and complexity. The developments in synthetic biology have permitted both fine-tuning the performance of existing microbial biosensors, and creating new ones with unique functionalities in a more predictable and rapid manner.

Synthetic microbial biosensors typically comprise three exchangeable modules: an input sensing block, an internal signal processing block, and an output reporting block (Wang and Barahona, 2013). In contrast to traditional microbial sensors consisting of a genetic reporter fused to an inducible promoter to control the expression of a detectable output, synthetic biology enables biosensor designs to incorporate additional complex signal processing circuits. Accordingly, the sensing unit triggers more sophisticated actions before activating reporter expression, in order to enhance a sensor's performance or perform additional functions. Such circuits include toggle switches (Gardner and Cantor, 2000), logic gates (Anderson *et al.*, 2006; Wang *et al.*, 2011), transcriptional amplifiers (Wang and Barahona, 2014) and memory circuits (Courbet *et al.*, 2015; Riglar *et al.*, 2017). Furthermore, microbial sensor cell arrays could be designed to display an easy-to-interpret output pattern corresponding to specific input analyte levels without the use of specialist lab equipment (Wan *et al.*, 2019).

As many early stage microbial biosensors are inadequate to meet practical requirements in detection limit, specificity and output amplitude, various gene circuit-based optimization strategies have recently been developed to improve their sensing performance. In contrast to traditional optimization methods such as random mutagenesis, these synthetic biology-enabled optimization tools are based on rational design, and are thus more predictable and faster to achieve the desired sensing specifications. For example, integrating multiple signal inputs using genetic AND gates have been shown to be effective in increasing microbial sensors' specificity (Wang *et al.*, 2013), and rationally tuning the intracellular levels of the receptor proteins can drastically improve

sensors' detection limits (Wang and Barahona, 2015). In addition, a toggle switch (Wu *et al.*, 2009) and a post-translational regulation device (Wan *et al.*, 2019) have been designed to lower microbial sensors' background expression and detection limits. Amplification of the transduced sensor signal is another powerful strategy to further boost the sensor's performance, using strategies such as positive feedback loops (Jia *et al.*, 2019) or transcription signal amplifiers (Wan *et al.*, 2019).

Albeit successful proof-of-concept laboratory demonstrations of a number of synthetic microbial sensors, very few have made it into the market. Several barriers remain to be overcome, including an insufficient number of sensory building blocks, poor sensing performance, long-term stability issues, risk of releasing genetically modified microorganisms (GMMs), and lack of practical experience in acceptance by professional stakeholders (Hicks and Bachmann, 2020). Nevertheless, synthetic biology has contributed novel strategies to address these limitations. For example, different approaches have been applied to keep biosensor cells alive and active for longer term including freeze-drying of cells, and encapsulating cells within polymers (Bjerketorp *et al.*, 2006; Liu *et al.*, 2018; Wan *et al.*, 2019; Shemer *et al.*, 2020). Recent advances demonstrated the potential of harnessing the amazing sensing capabilities of microbes for versatile applications, for example as wearable sensors for biomarker analysis in sweat to achieve non-invasive *in situ* real-time physiological state monitoring (Liu *et al.*, 2018; Nguyen *et al.*, 2021), or the standoff detection of buried landmines (Belkin *et al.*, 2017). However, biosafety concerns regarding the usage of GMMs remain an issue associated with field and *in vivo* applications, including potential horizontal gene transfer and disruption of natural ecosystems. Accordingly, different biocontainment strategies have been proposed to mitigate biosafety concerns such as replacing antibiotics resistance with toxin-antitoxin systems (Wright *et al.*, 2015), incorporating conditional kill switches (Chan *et al.*, 2016) and non-canonical amino acid substitution (Fredens *et al.*, 2019). Furthermore, chromosome-free bacterial chassis such as SimCells (Fan *et al.*, 2020) can be considered. Notably, cell-free expression systems have become increasingly popular as a new sensor platform, by avoiding biosafety concerns associated with using living cells. Cell-free biosensors lend faster responses, higher sensitivity and an enhanced compatibility to toxic samples (Lopreside *et al.*, 2019; Silverman and Karim, 2020). Moreover, cell-free extracts comprising genetic sensors could be embedded on paper, providing a portable platform for easy-to-use and cost-effective on-site screening (Pardee *et al.*, 2016) or in hydrogels acting as smart stimuli-responsive biomaterials (Whitfield *et al.*, 2020).

The latest developments in synthetic biology enable a fast design-build-test cycle for sensor construction and response optimization, to address the limitations of microbial biosensors. Yet, challenges remain to be addressed both within and beyond the scope of technical developments. For instance, environmental, food and health monitoring necessitate sensor cell exposure to complex samples, and therefore require complex signal processing circuits and likely multiple input modules. Notably, for biomedical applications involving complex media compositions such as tumours, non-specific localization of sensor cells prevents accurate diagnosis and biotherapy. Consequently, engineering microbes for sensing and reporting at designated spatial locations will be critical (Chien *et al.*, 2021). Considering that a single microbial cell has a limited capacity in resources, and that large complex circuits tend to burden host cells, cell consortia comprising multiple communicating sensor strains may be used to facilitate multiplex detection and reconfigurability of sensor function (Wang *et al.*, 2013; Khatun *et al.*, 2018). Innovative designs with fewer time-consuming signal propagation steps such as a transcription-only design, with RNA as the reporter entity, or engineered ligand-responsive fluorescent reporter proteins, could significantly shorten the response time of microbial biosensors, while alternative reporting formats such as direct bioelectronic signal output may lead to increasing seamless interfacing with conventional electronic devices. Further, integrating engineered microbial biosensors into various materials will lead to programmable living materials with bespoke functionalities such as self-healing.

In summary, we have witnessed a new wave of microbial sensors development in the rising era of synthetic biology, and expect this trend to continue and probably grow stronger in the coming decades. While living bioreporters presently face certain restrictions, synthetic biology offers new tools and strategies to accelerate the development, enhance the performance and address the current limitations of microbial biosensors; this will facilitate their future adoption and uptake as promising alternative analytical devices in diverse settings.

### Funding Information

B.W. acknowledges support by the UK Research and Innovation Future Leaders Fellowship [MR/S018875/1], Leverhulme Trust research project grant [RPG-2020-241] and US Office of Naval Research Global grant [N62909-20-1-2036]. S.B. was partially supported by the Minerva Center for Bio-Hybrid Complex Systems.

### Conflict of interest

None declared.

Shimshon Belkin<sup>1\*</sup>  and Baojun Wang<sup>2,3\*\*</sup> 

<sup>1</sup>*Institute of Life Sciences, the Hebrew University of Jerusalem, Jerusalem, 9190401, Israel;* <sup>2</sup>*School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3FF, UK and* <sup>3</sup>*Hangzhou Innovation Center and College of Chemical and Biological Engineering, Zhejiang University, Hangzhou, 311200, China*

\*E-mail: sb@mail.huji.ac.il

\*\*E-mail: baojun.wang@ed.ac.uk

### References

- Anderson, J.C., Clarke, E.J., Arkin, A.P., and Voigt, C.A. (2006) Environmentally controlled invasion of cancer cells by engineered bacteria. *J Mol Biol* **355**: 619–627.
- Belkin, S., Yagur-Kroll, S., Kabessa, Y., Korouma, V., Septon, T., Anati, Y., *et al.* (2017) Remote detection of buried landmines using a bacterial sensor. *Nat Biotechnol* **35**: 308–310.
- Bjerketorp, J., Håkansson, S., Belkin, S., and Jansson, J.K. (2006) Advances in preservation methods: keeping biosensor microorganisms alive and active. *Curr Opin Biotechnol* **17**: 1–7.
- Chan, C.T.Y., Lee, J.W., Cameron, D.E., Bashor, C.J., and Collins, J.J. (2016) 'Deadman' and 'Passcode' microbial kill switches for bacterial containment. *Nat Chem Biol* **12**: 82–86.
- Chien, T., Harimoto, T., Kepecs, B., Gray, K., Coker, C., Pu, K., *et al.* (2021) Multiplexed biosensors for precision bacteria tropism *in vivo*. *Nat Biomed Eng*. <https://doi.org/10.1038/s41551-021-00772-3>
- Courbet, A., Endy, D., Renard, E., Molina, F., and Bonnet, J. (2015) Detection of pathological biomarkers in human clinical samples via amplifying genetic switches and logic gates. *Sci Transl Med* **7**: 289ra283.
- Endy, D. (2005) Foundations for engineering biology. *Nature* **438**: 449–453.
- Fan, C., Davison, P.A., Habgood, R., Zeng, H., Decker, C.M., Gesell Salazar, M., *et al.* (2020) Chromosome-free bacterial cells are safe and programmable platforms for synthetic biology. *Proc Natl Acad Sci USA* **117**: 6752–6761.
- Fredens, J., Wang, K., de la Torre, D., Funke, L.F.H., Robertson, W.E., Christova, Y., *et al.* (2019) Total synthesis of *Escherichia coli* with a recoded genome. *Nature* **569**: 514–518.
- Gardner, T.S., Cantor, C.R., and Collins, J.J. (2000) Construction of a genetic toggle switch in *Escherichia coli*. *Nature* **403**: 339–342.
- Hicks, M., Bachmann, T.T., and Wang, B. (2020) Synthetic biology enables programmable cell-based biosensors. *ChemPhysChem* **21**: 132–144.
- IUPAC (2017) *Compendium of Chemical Terminology*, 2nd ed. (the "Gold Book"). Compiled by A. D. McNaught and A. Wilkinson. Blackwell Scientific Publications, Oxford (1997). Online version (2019-) created by S. J. Chalk. ISBN 0-9678550-9-8. <https://goldbook.iupac.org/terms/view/B00663>.
- Jacob, F., and Monod, J. (1961) On the regulation of gene activity. *Cold Spring Harb Symp Quant Biol* **26**: 193–211.

#### 4 Editorial

- Jia, X.Q., Bu, R.R., Zhao, T.T., and Wu, K. (2019) Sensitive and specific whole-cell biosensor for arsenic detection. *Appl Environ Microbiol* **85**: e00694–e619.
- Khatun, M.A., Hoque, M.A., Zhang, Y., Lu, T., Cui, L., Zhou, N.Y., and Feng, Y. (2018) Bacterial consortium-based sensing system for detecting organophosphorus pesticides. *Anal Chem* **90**: 10577–10584.
- King, J.M., Digrazia, P.M., Applegate, B., Burlage, R., Sanseverino, J., Dunbar, P., *et al.* (1990) Rapid, sensitive bioluminescent reporter technology for naphthalene exposure and biodegradation. *Science* **249**: 778–781.
- Knight, T. (2003) *Idempotent Vector Design for Standard Assembly of Biobricks*. MIT Artificial Intelligence Laboratory; MIT Synthetic Biology Working Group. <http://hdl.handle.net/1721.1/21168>
- Leduc, S. (1910) *Théorie physico-chimique de la vie et générations spontanées*. Paris: A. Poinat.
- Liu, X., Yuk, H., Lin, S., Parada, G.A., Tang, T.-C., Tham, E., *et al.* (2018) 3D printing of living responsive materials and devices. *Adv Mater* **30**: 1704821.
- Lopreside, A., Wan, X., Michelini, E., Roda, A., and Wang, B. (2019) Comprehensive profiling of diverse genetic reporters with application to whole-cell and cell-free biosensors. *Anal Chem* **91**: 15284–15292.
- van der Meer, J.R., and Belkin, S. (2010) Where microbiology meets microengineering: design and applications of reporter bacteria. *Nat Rev Microbiol* **8**: 511–522.
- Nguyen, P.Q., Soenksen, L.R., Donghia, N.M., Angenent-Mari, N.M., de Puig, H., Huang, A., *et al.* (2021) Wearable materials with embedded synthetic biology sensors for biomolecule detection. *Nat Biotech.* <https://doi.org/10.1038/s41587-021-00950-3>
- Oda, Y., Nakamura, S., Oki, I., Kato, T., and Shinagawa, H. (1985) Evaluation of the new system (*umu*-test) for the detection of environmental mutagens and carcinogens. *Mutat Res* **147**: 219–229.
- Pardee, K., Green, A.A., Takahashi, M.K., Braff, D., Lambert, G., Lee, J.W., *et al.* (2016) Rapid, low-cost detection of Zika virus using programmable biomolecular components. *Cell* **165**: 1255–1266.
- Quillardet, P., Huisman, O., D'Ari, R., and Hofnung, M. (1982) SOS chromotest, a direct assay of induction of an SOS function in *Escherichia coli* K-12 to measure genotoxicity. *Proc Natl Acad Sci USA* **79**: 5971–5975.
- Riglar, D.T., Giessen, T.W., Baym, M., Kerns, S.J., Niederhuber, M.J., Bronson, R.T., *et al.* (2017) Engineered bacteria can function in the mammalian gut long-term as live diagnostics of inflammation. *Nat Biotechnol* **35**: 653–658.
- Selifonova, O., Burlage, R., and Barkay, T. (1983) Bioluminescent sensors for detection of bioavailable Hg(II) in the environment. *Appl Environ Microbiol* **59**: 3083–3090.
- Shemer, B., Shpigel, E., Hazan, C., Kabessa, Y., Agranat, A.J., and Belkin, S. (2020) Detection of buried explosives with immobilized bacterial bioreporters. *Microb Biotechnol* **14**: 251–261.
- Silverman, A.D., Karim, A.S., and Jewett, M.C. (2020) Cell-free gene expression: an expanded repertoire of applications. *Nat Rev Genet* **21**: 151–170.
- Van Der Meer, J.R., Tropel, D., and Jaspers, M. (2004) Illuminating the detection chain of bacterial bioreporters. *Environ Microbiol* **6**: 1005–1020.
- Wan, X., Volpetti, F., Petrova, E., French, C., Maerkl, S.J. & Wang, B. (2019) Cascaded amplifying circuits enable ultrasensitive cellular sensors for toxic metals. *Nat Chem Biol*, **15**, 540–548.
- Wang, B., Barahona, M., and Buck, M. (2013) A modular cell-based biosensor using engineered genetic logic circuits to detect and integrate multiple environmental signals. *Biosens Bioelectron* **40**: 368–376.
- Wang, B., Barahona, M., and Buck, M. (2014) Engineering modular and tunable genetic amplifiers for scaling transcriptional signals in cascaded gene networks. *Nucleic Acids Res* **42**: 9484–9492.
- Wang, B., Barahona, M., and Buck, M. (2015) Amplification of small molecule-inducible gene expression via tuning of intracellular receptor densities. *Nucleic Acids Res* **43**: 1955–1964.
- Wang, B., Kitney, R.I., Joly, N., and Buck, M. (2011) Engineering modular and orthogonal genetic logic gates for robust digital-like synthetic biology. *Nat Commun* **2**: 508.
- Whitfield, C.J., Banks, A.M., Dura, G., Love, J., Fieldsend, J.E., Goodchild, S.A., *et al.* (2020) Cell-free protein synthesis in hydrogel materials. *ChemComm* **56**: 7108–7111.
- Wright, O., Delmans, M., Stan, G.B., and Elis, T. (2015) GeneGuard: a modular plasmid system designed for biosafety. *ACS Synth Biol* **4**: 307–316.
- Wu, C.H., Le, D., Mulchandani, A., and Chen, W. (2009) Optimization of a whole-cell cadmium sensor with a toggle gene circuit. *Biotechnol Prog* **25**: 898–903.