



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Capsule Commentary on Reese et al., Two Randomized Controlled Pilot Trials of Social Forces to Improve Statin Adherence Among Patients with Diabetes

Citation for published version:

Wolters, MK 2016, 'Capsule Commentary on Reese et al., Two Randomized Controlled Pilot Trials of Social Forces to Improve Statin Adherence Among Patients with Diabetes', *Journal of General Internal Medicine*, vol. 31, no. 4, pp. 416-416. <https://doi.org/10.1007/s11606-016-3599-0>

Digital Object Identifier (DOI):

[10.1007/s11606-016-3599-0](https://doi.org/10.1007/s11606-016-3599-0)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of General Internal Medicine

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Capsule Commentary on Reese et al., Two Randomized Controlled Pilot Trials of Social Forces to Improve Statin Adherence Among Patients with Diabetes

Maria K. Wolters, Dr. phil. Dipl-Inform

Informatics & Psychology, , University of Edinburgh, Edinburgh, UK.

J Gen Intern Med

DOI: 10.1007/s11606-016-3599-0

© The Author(s) 2016. This article is published with open access at Springerlink.com

At first glance, readers might disagree with the decision to publish a paper that reports not one, but two negative results. However, it is absolutely vital for scientific progress that such findings are shared with the research and the clinical community, because publication bias skews the quality of the evidence.¹

As this paper reports two carefully designed and executed randomised controlled trials, its lessons are particularly important. In both trials, all participants greatly increased their medication adherence, despite being in a high risk group, older, and of lower socioeconomic status.² All they did was take part in a trial where they received an electronic pill bottle. Additional social feedback, as provided in the intervention arms of the trials, was not needed.

While the authors list a number of possible reasons for this finding, it would have been much better to explore it through a programme of qualitative interviews and focus groups. The data from such qualitative work are key for translating findings from randomised controlled trials into practice.³

In this study, the Hawthorne effect—that observation changes behaviour, a phenomenon well-documented in clinical trials⁴—may have been augmented by the fact that all study arms received a special device, the electronic pill bottle. Providing such bottles in itself constitutes an intervention that improves medication adherence.⁵ A well-designed qualitative interview study with a small proportion of the participants in

each arm could have helped clarify to what extent those effects might have been present. It would also have provided valuable data on the participants' motivation for taking part in the study, and highlighted the extent to which the study itself might have been a valuable social force.

Corresponding Author: Maria K. Wolters, Dr. phil. Dipl-Inform; Informatics & Psychology, University of Edinburgh, Edinburgh, UK (e-mail: Maria.wolters@ed.ac.uk).

Compliance with Ethical Standards:

Conflict of Interest: The author has no conflict with any of the material in this manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

REFERENCES

1. Ioannidis JPA. Why most published research findings are false. *PLoS Med.* 2005;2:124. doi:10.1371/journal.pmed.0020124.
2. Reese PP, Kessler JB, Doshi JA, Friedman J, Mussell AS, Carney C, Zhu J, Wang W, Troxel A, Young P, Lawnicki V, Rajpathak S, Volpp K. Two randomized controlled pilot trials of social forces to improve statin adherence among patients with diabetes. *J Gen Intern Med.* 2015. doi:10.1007/s11606-015-3540-y.
3. Greenhalgh T, Howick J, Maskrey N. Evidence based medicine: a movement in crisis? *BMJ.* 2014;348:g3725.
4. De Amici D, Klersy C, Ramajoli F, Brustia L, Politi P. Impact of the hawthorne effect in a longitudinal clinical study. *Control Clin Trials.* 2000;21:103–14.
5. Haynes RB, Ackloo E, Sahota N, McDonald HP, Yao X. Interventions for enhancing medication adherence. *Cochrane Database Syst Rev.* 2008;16(2):CD000011.

This comment refers to the article available at: <http://dx.doi.org/10.1007/s11606-015-3540-y>.
