



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

11 β -Hydroxysteroid Dehydrogenase type 1 is expressed in neutrophils and restrains an inflammatory response in male mice

Citation for published version:

Coutinho, AE, Kipari, TMJ, Zhang, Z, Esteves, CL, Lucas, CD, Gilmour, JS, Webster, SP, Walker, BR, Hughes, J, Savill, JS, Seckl, JR, Rossi, AG & Chapman, KE 2016, '11 β -Hydroxysteroid Dehydrogenase type 1 is expressed in neutrophils and restrains an inflammatory response in male mice', *Endocrinology*, vol. 157, no. 7, pp. 2928-2936. <https://doi.org/10.1210/en.2016-1118>

Digital Object Identifier (DOI):

[10.1210/en.2016-1118](https://doi.org/10.1210/en.2016-1118)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Endocrinology

Publisher Rights Statement:

Author's final peer-reviewed manuscript as accepted for publication

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



SUPPLEMENTARY FIGURES

Supplementary Figure 1. Purity of mouse neutrophil preparations

Peritoneal cells were incubated with Ly6G-phycoerythrin (PE) antibody followed by incubation with anti-PE secondary antibody magnetic beads. (A) Flow cytometry with Ly6G staining was used to quantify the purity of isolated neutrophils, which was generally $\geq 96\%$. (B) Neutrophil purity was confirmed by May-Geimsa staining of cytocentrifuged cells; left panel, total peritoneal cells prior to neutrophil isolation; centre panel, flow-through cells (Ly6G⁻) and right panel, neutrophils (Ly6G⁺ cells), images captured at x40 magnification.

Supplementary Figure 2. 11 β -HSD1 staining in peritoneal cells

Flow cytometry was used to measure the number of 11 β -HSD1⁺ cells and to quantify levels of 11 β -HSD1 (by mean fluorescence intensity, MFI). Representative dot plots of peritoneal cells collected 4h after thioglycollate injection (from 7-8 mice; 5×10^5 cells/sample), showing 11 β -HSD1⁺ cells after incubation with 11 β -HSD1 antibody (left panel), control IgG (centre panel) and cells incubated with secondary antibody only (right panel).

Supplementary Figure 3. Monocyte numbers and 11 β -HSD1 expression in monocytes from BM and blood during peritonitis

Monocyte (Ly6G⁻, 7/4⁺, CD11b⁺) cell number was determined by flow cytometry, in freshly isolated (A) BM and (C) blood at various times following thioglycollate injection. 11 β -HSD1 expression (mean fluorescence intensity, MFI) was measured in BM monocytes (B) and the following subsets of blood monocytes: (D) 7/4^{hi}, (E) 7/4^{med} and (F) 7/4^{lo}. Data are means \pm SEM and were analysed by 1-way ANOVA followed by Dunnett's post-hoc test; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, $n = 6-8$.

Supplementary Figure 4. UE2316 inhibits 11 β -HSD1 in peritoneal cells in vitro


Peritoneal cells (2×10^5) isolated 4h following thioglycollate injection were incubated with ^3H -cortisone vehicle (solid line, circles) or 1 μM UE2316 (dashed line, squares). Data are means of 2 samples each assayed in duplicate and show conversion (cpm) of ^3H -cortisone to ^3H -cortisol over time.

Supplementary Figure 5. The effect of 11 β -HSD1 inhibition upon CD11b expression in bone marrow and blood neutrophils during 4h peritonitis


CD11b mean fluorescence intensity (MFI) was measured by flow cytometry performed on BM cells and blood collected from mice pretreated with vehicle (Veh, black bars) or UE2316 (white bars), 4h following injection of thioglycollate. (A) BM neutrophils, (B) blood neutrophils. Data are means \pm SEM and were analysed by student's t-test, n=9-10.

Supplementary Figure 1


A









B




Supplementary Figure 2



Supplementary Figure 3



Supplementary Figure 4



Supplementary Figure 5

