



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Genetic parameters of calcium, phosphorus, magnesium, and potassium serum concentrations during the first 8 days after calving in Holstein cows

Citation for published version:

Tsiamadis, V, Banos, G, Panousis, N, Kritsepi-Konstantinou, M, Arsenos, G & Valergakis, GE 2016, 'Genetic parameters of calcium, phosphorus, magnesium, and potassium serum concentrations during the first 8 days after calving in Holstein cows', *Journal of Dairy Science*, vol. 99, no. 7, pp. 5535-5544. <https://doi.org/10.3168/jds.2015-10787>

Digital Object Identifier (DOI):

[10.3168/jds.2015-10787](https://doi.org/10.3168/jds.2015-10787)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Published In:

Journal of Dairy Science

Publisher Rights Statement:

This is the author's final peer-reviewed manuscript as accepted for publication

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Genetic Parameters of Calcium, Phosphorus, Magnesium and Potassium Serum**
2 **Concentrations during the First Eight Days after Calving in Holstein Cows**
3 **V. Tsiamadis, * G. Banos*§, N. Panousis, † M. Kritsepi-Konstantinou, ‡ G. Arsenos, * G.**
4 **E. Valergakis***

5 *Department of Animal Production, Faculty of Veterinary Medicine, Box 393, Aristotle University of
6 Thessaloniki, GR-54124 Thessaloniki, Greece

7 §Scotland's Rural College/Roslin Institute, Edinburgh, Scotland, UK

8 †Clinic of Farm Animals, Department of Clinics, Faculty of Veterinary Medicine, Aristotle University of
9 Thessaloniki, GR-54124 Thessaloniki, Greece

10 ‡Diagnostic Laboratory, Department of Clinics, Faculty of Veterinary Medicine, Aristotle University of
11 Thessaloniki, GR-54124 Thessaloniki, Greece

12 ¹Corresponding author: Georgios E. Valergakis, phone: +30 2310 99 98 50, fax: 2310 99 98 92,

13 email: geval@vet.auth.gr

14 ***Interpretive Summary***

15 Macromineral-related disorders immediately after calving are of great importance for the health
16 and productivity of dairy cows. They predispose animals to other major diseases, increase culling
17 rate and impair production. Our objective was to estimate the genetic parameters of
18 macrominerals' concentrations during the first 8 days after calving in Holstein cows. Repeated
19 measurements of blood serum macrominerals concentrations from 986 cows, in 9 commercial
20 farms located in Northern Greece were analyzed with random regression models. Results
21 revealed the presence of significant genetic variation. Achieving and maintaining normal

22 macromineral concentrations through genetic selection could contribute towards reduction of the
23 related disorders.

24

25 **ABSTRACT**

26 Calcium (Ca), magnesium (Mg), phosphorus (P) and potassium (K) are of great importance for
27 the health and productivity of dairy cows after calving. So far genetic studies have focused on
28 clinical hypocalcemia, leaving the genetic parameters of these macroelements unstudied. Our
29 objective was to estimate the genetic parameters of Ca, Mg, P and K serum concentrations and
30 their changes during the first 8 days after calving. The study was conducted in 9 herds located in
31 Northern Greece, with 1,021 Holstein cows enrolled from November 2010 until November 2012.
32 No herd used any kind of preventive measures for hypocalcemia. Pedigree information for all
33 cows was available. A total of 35 cows were diagnosed and treated for periparturient paresis and,
34 therefore, excluded from the study. The remaining 986 cows were included in genetic analysis.
35 The distribution of cows across parities was 459 (parity 1), 234 (parity 2), 158 (parity 3) and 135
36 (parity 4 and above). A sample of blood was taken from each cow on day 1, 2, 4 and 8 after
37 calving and serum concentrations of Ca, P, Mg and K were measured in each sample. A final
38 data set of 15,390 biochemical records was created consisting of 3,903 Ca, 3,902 P, 3,903 Mg
39 and 3,682 K measurements. Moreover, changes of these concentrations between day 1 and 4 as
40 well as day 1 and 8 after calving were calculated and treated as different traits. Random
41 regression models were used to analyze the data. Results showed that daily heritabilities of Ca, P
42 and Mg concentrations traits were moderate to high (0.20 – 0.43; $P < 0.05$), while those of K were
43 low to moderate (0.12 – 0.23; $P < 0.05$). Regarding concentration changes, only Mg change
44 between day 1 and day 8 after calving had a significant heritability of 0.18. Genetic correlations

45 between Ca, P, Mg and K concentrations and their concentration changes from days 1-4 and 1-8
46 after calving were not significantly different from zero. Most phenotypic correlations among Ca,
47 P, Mg, and K concentrations were positive and low (0.09 – 0.16; P<0.05), while the correlation
48 between P and Mg was negative and low (-0.16; P<0.05). Phenotypic correlations among
49 macromineral concentrations on day 1 and their changes from day 1 to 4 and 1 to 8 after calving
50 varied for each macromineral. This study revealed that genetic selection for normal Ca, P, Mg
51 and K concentrations in the first week of lactation is possible and could facilitate the
52 management of their deficiencies during the early stages of lactation.

53 **Key words:** macrominerals, genetic parameters

54

55

INTRODUCTION

56 During the first critical days after calving, calcium (Ca), phosphorus (P), magnesium (Mg) and
57 potassium (K) blood serum concentrations are of great importance for the health and productivity
58 of the dairy cow. Possible deviations from normal levels of these macrominerals are interrelated
59 (Goff and Horst, 1997; Goff, 2000; Lean et al., 2013).

60

61 Calcium plays a key role at the onset of lactation (DeGaris and Lean, 2008). Hypocalcaemia
62 (serum Ca<8.3 mg/dL) is the most important macromineral disorder of the transition dairy cow
63 (Oetzel, 2011; Goff, 2014; Martinez et al., 2014) and is associated with health disorders
64 including retained fetal membranes, mastitis, uterine infection, displaced abomasum and ketosis
65 (Correa et al., 1990; Gröhn and Bruss, 1990; DeGaris and Lean, 2008), as well as reduced dry
66 matter intake and milk production (Rajala-Schultz et al., 1999).

67

68 Phosphorus and Mg play important roles in the etiology of hypocalcemia, as well.
69 Hypophosphatemia (serum P<4.0 mg/dL) is involved in the manifestation of the alert downer
70 cow syndrome, while elevated phosphorus concentrations increase the risk of milk fever (Lean et
71 al., 2013; Grünberg, 2014). Hypomagnesaemia (serum Mg<1.8 mg/dL) reduces parathormone
72 (PTH) secretion, tissue sensitivity to PTH and synthesis of 1,25-dihydroxycholecalciferol
73 (Littledike et al., 1983; Rude, 1998). Moreover, mild hypomagnesaemia (serum Mg between 1.3
74 and 1.8 mg/dL) is common in anorectic fresh cows and in most cases is accompanied by mild
75 hypophosphatemia (serum P between 2 and 4 mg/dL) and mild hypokalemia (serum K between
76 2.6 and 3.9 mmol/L) (Peek and Divers, 2008).

77

78 Potassium homeostasis in transition dairy cows is affected by numerous factors. Off-feed fresh
79 cows, increased milk production and concurrent diseases predispose to hypokalemia (serum K
80 <3.9 mmol/L) (Pradhan and Hemken, 1968; Sattler et al., 1998; Sattler and Fecteau, 2014).

81

82 Blood Ca concentration is considered to reach its minimum 12 to 24 hours after calving and then
83 it increases gradually (Goff, 2014). Relative estimates for the other three macrominerals are
84 lacking from the literature.

85

86 Serum Ca, P, Mg and K concentrations are influenced by environmental factors, mainly nutrition
87 (NRC, 2001; Kronqvist, 2011). Nutritional and management strategies for the prevention of
88 these macromineral deficiencies have been developed (Bethard et al., 1998; Tauriainen et al.,

89 2003; Rérat et al., 2009). However, there is also a genetic component to these traits, as reported
90 for serum Ca concentration by Tveit et al. (1991).

91

92 Genetic studies so far have focused on heritability estimates of clinical hypocalcemia (milk
93 fever) (Dyrendahl et al., 1972; Lin et al., 1989; Abdel-Azim et al., 2005) and genetic and
94 phenotypic correlations between milk fever and various disease (Lin et al., 1989) and production
95 traits (Lyons et al., 1991; Uribe et al., 1995; Heringstad et al., 2005). Tveit et al. (1991) reported
96 heritability estimates for post-partum serum Ca concentrations in first lactation Norwegian cows.
97 However, genetic studies of serum Ca, P, Mg and K concentrations in fresh Holstein dairy cows
98 are lacking.

99

100 Therefore, the objective of this study was to estimate the genetic parameters of Ca, Mg, P and K
101 serum concentrations and their changes in Holstein cows during the first 8 days after calving.

102

103

MATERIALS AND METHODS

104 The research was conducted in compliance with institutional guidelines and approved by the
105 Research Committee of the Aristotle University of Thessaloniki, Thessaloniki, Greece. All
106 farmers gave informed consent for the cows to be included in the study and the testing
107 procedures.

108

Animals and Management

110 A total of 1,021 Holstein cows from 9 commercial free-stall dairy herds in Northern Greece were
111 included in the study. The distribution across parities was 466, 242, 165 and 148 cows for

112 parities 1, 2, 3 and 4 and above, respectively. Farms were visited regularly between November
113 2010 and November 2012 for data collection. No herd used any kind of preventive measures for
114 hypocalcemia. Total mixed rations (TMR) were formulated to meet or exceed net energy and
115 metabolizable protein requirements according to National Research Council recommendations
116 (NRC, 2001).

117

118 ***Clinical Examination, Blood Sampling and Analyses***

119 Each animal was clinically examined and blood sampled on day 1, 2, 4 and 8 after calving, by
120 the first author. Blood samples, in all herds, were collected between 08:00 – 10:00 a.m., after the
121 morning milking. Moreover, to standardize sampling and handling procedures, all samplings
122 were performed in absence of unusual stressors and in proper containment systems that minimize
123 stress and pain of the animal.

124

125 Blood sampling was performed by coccygeal venipuncture into 10-ml vacuum glass tubes
126 without anticoagulant (BD Vacutainer[®], Plymouth, United Kingdom) for serum macromineral
127 measurements. Samples were placed in a cooler, transported to the Diagnostic Laboratory of the
128 Faculty of Veterinary Medicine and centrifuged immediately upon arrival (3,000 x g for 15 min).
129 Serum was transferred into polyethylene tubes and stored at -80°C until assay. All sera were
130 analyzed for total Ca and Mg concentrations using flame atomic absorption spectrophotometry
131 (Perkin ElmerAAAnalyst 100, Perkin Elmer Co, Norwalk, CT, USA), according to manufacturer's
132 instructions. Serum inorganic phosphorus concentrations were determined photometrically using
133 a Flexor E autoanalyzer (Vital Scientific, Netherlands), according to the procedure described by
134 Daly and Ertingshausen (1972), with the use of standard commercial reagents (Thermo Fisher
135 Scientific Inc. USA). Potassium serum concentrations were measured using an ion-selective

136 electrode according to manufacturer's instructions (Electrolyte Analyzer 9180, Roche Austria).

137 The intra- and inter-assay coefficients of variation for all the above analyses were less than 3%.

138

139 ***Data set***

140 Considering that pedigree information was available for all cows, the total population increased

141 to 4,262 animals, spanning the last 5 generations. Calving date, parity number, calving ease and

142 twinning was recorded.

143

144 A total of 35 cows were diagnosed with periparturient paresis, treated appropriately with

145 intravenous Ca and excluded from the study. Therefore, the remaining 986 cows were finally

146 included in the genetic analysis. The distribution across parities was 459, 234, 158 and 135 cows

147 for parities 1, 2, 3 and 4 and above, respectively.

148

149 Following all analyses, a data set of 15,390 biochemical records was created (Table 1),

150 consisting of 3,903 Ca, 3,902 P, 3,903 Mg and 3,682 K serum concentration measurements.

151 Moreover, changes of these concentrations between day 1 and day 4 as well as day 1 and day 8

152 were calculated and treated as different traits.

153

154 ***Statistical Analysis***

155 Repeated cow records of Ca, Mg, P and K serum concentrations were analyzed with a random

156 regression model which accounted for the covariance between successive records of the same

157 animal; each trait was analyzed separately:

158
$$Y_{ijkm} = HYS_i + L_j + M_k + a_1 \cdot age + \sum_{n=0}^2 b_n P_n D_m + \sum_{n=0}^2 A_{km} P_n D_m + e_{ijkm}$$

159 (1)

160 where:

161 Y_{ijkm} is the macromineral concentration of cow k on day from calving m ;

162 HYS_i is the fixed effect of herd-year-season of calving i (72 levels);

163 L_j the fixed effect of number of lactation (4 levels);

164 M_k the fixed effect of calendar month when the record was taken p (12 levels);

165 a_1 the linear regression coefficient on age at calving (age);

166 D_m the number of days from calving;

167 b_m the fixed regression coefficient on days from calving;

168 A_{km} the random regression coefficient on day from calving associated with the additive
 169 genetic effect of cow k including all pedigree data (4,262 animals);

170 P_m the m th orthogonal polynomial of day from calving (m the order of polynomial);

171 e_{ijkm} the random residual term.

172

173 The fixed effects in the model were fitted after preliminary analyses had confirmed their
 174 statistically significant effect ($P < 0.05$) on the traits. The final order of the random polynomial
 175 (third for either trait) was determined with the use of the log-likelihood test in sequential
 176 analyses of gradually increasing orders. The final order choice was also confirmed with the
 177 Akaike Information Criterion test. Four measurement error classes were defined using the time
 178 relative to calving as day 1, 2, 4 and 8. The definition of these classes, even at this small time

179 span, aimed to capture the day-to-day differences in health events at the beginning of lactation.

180 Covariances between the error classes were assumed to be zero.

181

182 Estimates of variance components from model 1 were used to calculate heritabilities for each
183 trait and day after calving.

184

185 Variance components and heritability estimated for Ca, K, P and Mg serum concentrations were
186 also calculated across all days from calving using the following model:

187

$$188 \quad Y_{ijkm} = HYS_i + L_j + a_1 \cdot age + D_m + A_k + e_{ijkm} \quad (2)$$

189

190 where: Y_{ijkm} is the macromineral concentration change of cow k; A_k is the additive genetic effect
191 of cow k and all effects are as in model 1.

192

193 Serum concentration changes between day 1 and day 4 (days 1-4), as well as day 1 and day 8
194 (days 1-8) after calving were analyzed with the following model:

195

$$196 \quad Y_{ijk} = HYS_i + L_j + age + A_k + e_{ijk} \quad (3)$$

197

198 where: Y_{ijk} is the macromineral concentration change of cow k; All other effects are as in
199 Model 2.

200

201 Genetic and phenotypic correlations among all traits analyzed with the above models were
202 estimated with a series of bivariate analyses.

203

204 All analyses were conducted using the statistical software package ASREML (Gilmour and
205 Gogel, 2006).

206

207

RESULTS

208 *Mean Macromineral Serum Concentrations and Prediction Lines for Concentrations*

209 Mean serum Ca concentration increased gradually from day 1 to day 8 after calving ($P < 0.001$).
210 In 1st and 2nd lactation cows, mean Ca concentration remained above the 8.3 mg/dL threshold
211 throughout the sampling period, whereas in older cows it was below the threshold on days 1 and
212 2 after calving. On the contrary, mean serum P, Mg and K concentrations decreased from day 1
213 to day 8 after calving ($P < 0.001$). Descriptive statistics and analysis of variance results by parity
214 are presented in Table 1. Fixed curves of serum macromineral concentrations, across all
215 lactations, during the first 8 days after calving from the random regression model analysis
216 (Model 1) are shown in Figure 1. These curves are adjusted for all other effects included in
217 Model 1.

218

219 *Serum Macromineral Concentrations Variances and Heritabilities Estimates*

220 Estimates of day-to-day phenotypic, genetic and residual variances, and heritabilities for serum
221 Ca, P, Mg and K concentrations are presented in Table 2. All estimates were statistically greater
222 than zero ($P < 0.001$). During the first 8 days after calving the estimated phenotypic (σ_p^2) and
223 residual variances (σ_r^2) for Ca and P serum concentrations were high, while those of Mg and K

224 were low. During the same period, the estimated genetic variance (σ_a^2) for Ca and P serum
225 concentration was moderate and high, respectively, while for Mg and K was low. Day-to-day
226 heritabilities of serum Ca, P and Mg concentrations were moderate ($h^2 = 0.20 - 0.43$), while
227 heritability estimates of K serum concentrations were low ($h^2 = 0.12 - 0.15$) except on day 8 after
228 calving ($h^2 = 0.23$) (Figure 2).

229

230 Heritability estimates of serum Ca, P, Mg, and K concentrations across all days using Model 2
231 are in Table 3. Although smaller, they were comparable with the ones derived with the random
232 regression model analysis. Regarding concentration changes, only Mg change between day 1 and
233 day 8 after calving had a significant ($P < 0.05$) heritability of 0.18.

234

235 *Serum Macromineral Concentrations Correlations*

236 Significant genetic correlations between serum Ca, P, Mg and K concentrations and their
237 concentration changes from days 1-4 and 1-8 after calving were not detected in the present study.

238

239 Statistically significant ($P < 0.010 - 0.001$) phenotypic correlations among Ca, P, Mg, and K
240 serum concentrations are shown in Table 3. Most correlations were positive and low ($r_p = 0.09 -$
241 0.16), while the P – Mg correlation was negative and low ($r_p = -0.16 \pm 0.03$).

242

243 Significant phenotypic correlations among serum macromineral concentrations on day 1 and
244 their changes from day 1 to 4 and 1 to 8 after calving are shown in Table 4. On day 1, there was
245 a low positive correlation between Ca and P, Ca and K, as well as P and K; there was also a low
246 negative correlation between P and Mg. Calcium and Mg serum concentrations on day 1 had

247 moderate negative correlations with both their changes from day 1 to 4 and 1 to 8. Phosphorus
248 serum concentration on day 1 had moderate negative correlation with its change from day 1 to 8,
249 while K serum concentration at day 1 had a moderate positive correlation with its change from
250 day 1 to 8. Phosphorus serum concentration on day 1 had a low positive correlation with both
251 Mg changes (days 1 – 4 and 1 – 8) and a low negative one with both K changes (days 1 – 4 and 1
252 – 8). Phosphorus change from day 1 to 4 had a low negative correlation with both Mg changes.
253 Both P changes (days 1 – 4 and 1 – 8) had a low positive correlation with both K changes (days 1
254 – 4 and 1 – 8). For each macromineral, its serum concentration changes between day 1 to 4 and 1
255 to 8 were positively and moderately correlated.

256

257

DISCUSSION

258 The present study was designed to estimate the genetic parameters of serum Ca, P, Mg and K
259 concentrations immediately after calving.

260

261 Normally, serum Ca concentration is maintained within a narrow range, between 8.3 and 10.4
262 mg/dL (Goff, 2014). During the first 12 to 24 hours after calving, Ca concentration reaches the
263 lower value and then gradually increases (Goff, 2014). In the present study, an increase across all
264 lactations in serum Ca concentrations from day 1 to day 8 after calving was observed. Mean Ca
265 serum concentrations from days 1 to 8 were different, depending on parity number and days after
266 calving. Response of cows to the decreased serum Ca concentration was not similar across
267 lactations. The homeorhetic mechanisms that determine the Ca balance (parathormone,
268 cholocalciferol and calcitonin) restored Ca serum concentration in most 1st and 2nd parity cows.
269 However, in older cows (3rd and 4th+ parities) the same homeorhetic mechanisms that affect the

270 Ca concentration did not react as efficiently, putting these animals in a profound hypocalcaemic
271 status just after calving (day 1).

272

273 The prediction curve generated with the random regression model denotes that there was a
274 significant rise in Ca concentration from day 1 to day 8 across all lactations. This is in agreement
275 with results from studies dealing with Ca physiology after calving (Littledike and Goff, 1987;
276 Goff, 2000; DeGaris and Lean, 2008). Furthermore, mean serum P, Mg and K concentrations
277 were within reference ranges (P: 4.2 – 7.7 mg/dL, Mg: 1.8 – 2.4 mg/dL, K: 3.9 – 5.8 mmol/L;
278 Peek and Divers, 2008; Goff, 2008) during the 1st day after calving and then gradually decreased,
279 but always remaining within those ranges. The prediction curves denote that there was a
280 significant decline in P, Mg and K concentrations from day 1 to day 8 across all lactations.
281 Serum Ca and P concentrations are regulated by the same hormones. The main regulatory
282 hormone is PTH, which increases Ca and decreases P concentration, within normal ranges. The
283 increase in PTH mobilization due to decreased Ca levels can explain the concurrent fall in P
284 concentration observed in the present study. Regarding Mg and K, since there is no major
285 hormonal control for these macrominerals (Kaneko et al., 2008), the observed decrease in their
286 concentrations is difficult to explain but may be attributed to the demands of the increasing milk
287 production.

288

289 Large scale field studies on Ca, P, Mg and K serum concentrations during the first week after
290 calving are lacking in literature. Recently, Reinhardt et al. (2011) conducted a field study for
291 hypocalcaemia in 1,462 cows, with only one Ca measurement within 48 h postpartum. To our
292 knowledge this is the first time that repeated measurements of Ca, P, Mg, and K concentrations

293 during the first 8 days after calving are reported. The observed variation allowed the
294 development of Ca, P, Mg and K serum concentration prediction lines with the use of random
295 regression model.

296

297 The estimated day-to-day heritabilities for serum Ca concentration were moderate (0.23 – 0.32).
298 So far, genetic studies have focused on the estimation of clinical hypocalcemia (milk fever)
299 heritability. Some studies reported moderate to high estimates (0.30 – 0.47) (Lin et al., 1989;
300 Lyons et al., 1991, Abdel-Azim et al., 2005), while others (Dyrendahl et al., 1972; Pryce et al.,
301 1997; Van Dorp et al., 1998; Heringstad et al., 2005) reported low ones (0.04 – 0.13), depending
302 on lactation number, method of statistical analysis and method of data collection, with higher
303 estimates being observed in later lactations. Heritability estimates for serum Ca concentration in
304 Holsteins after calving are lacking. Only one study investigated the genetic variation of Ca
305 concentration in Norwegian Reds cows and reported a low heritability (0.11±0.09) that was not
306 statistically different from zero (Tveit et al., 1991).

307

308 Similarly, the estimated day-to-day heritabilities for serum P and Mg concentrations in the
309 present study were moderate to high (0.30 – 0.43 and 0.20 – 0.39, respectively), while those for
310 K were low to moderate (0.12 – 0.23). To our knowledge this is the first time that such estimates
311 are reported. So far, only Kadarmideen et al. (2000) reported heritability estimates (0.004±0.004)
312 for clinical hypomagnesaemia in dairy cattle, which was not statistically different than zero.
313 Moreover, the information for hypomagnesaemia cases in that study was based on subjective
314 clinical observations made by farmers and was not confirmed by serum Mg concentration
315 measurements.

316

317 Genetic variance estimates of Ca and P were high (0.28 to 0.44 and 0.40 to 0.70, respectively),
318 indicating high influence of additive genetic effects on these traits. Their serum concentrations
319 are regulated mainly by PTH, 1,25-dihydroxyvitamin D and calcitonin (Kaneko et al., 2008). The
320 existence of the above major hormonal mechanism that regulates Ca and P concentrations can
321 help explain the moderate to high heritability estimates of these two elements. It was an early
322 belief that milk fever resulted from the failure of parathyroid glands to respond to the reduced Ca
323 concentration soon after calving. However, it has been shown that such cows have very high
324 blood PTH concentrations. Therefore, this finding implies that PTH's target tissues cannot
325 respond to its action (Goff, 2014). The main target of PTH is the skeleton. In humans the
326 RANK/RANKL/OPG system is well known for its osteoclastic function. This axis has a genetic
327 control and is hormonally stimulated by PTH and calcitonin, both of which control serum Ca and
328 P concentrations (Asagiri and Takayanagi, 2007; Cappariello et al., 2014). Further investigation
329 is needed in order to clarify whether this axis is also functional to dairy cows and whether is
330 involved in the etiology of hypocalcemia at the genetic level.

331

332 Genetic variance estimates for Mg and K were low (0.03 to 0.07 and 0.03 to 0.05, respectively).
333 In humans, PTH contributes towards a small increase of Mg concentration (Swaminathan, 2000).
334 Moreover, aldosterone is the only known hormone that partly regulates K concentration. The
335 absence of any major hormonal mechanism that regulates the serum concentration of Mg and K
336 may help explain the low genetic variances. The high precision of the diagnostic methods for Mg
337 and K measurements strongly contributed to our heritability estimates.

338

339 Our results indicate that genetic improvement is possible for these traits, probably to the same
340 degree with traits such as milk yield ($h^2= 0.20 - 0.50$; Castillo-Juarez et al., 2000; Windig et al.,
341 2006; Bastin et al., 2011) or BCS ($h^2= 0.34 - 0.79$; Berry et al., 2003; Banos et al., 2005;
342 Oikonomou et al., 2008), which are already included in breeding programs worldwide. Both the
343 amount of genetic variance and size of heritability for macromineral concentrations suggest that
344 selection could be effective during the first critical days after calving. Especially for Ca, whose
345 role in health status and disease development is of great importance (Goff and Horst, 1997), this
346 genetic improvement could favor animal welfare and productivity. In the meantime, appropriate
347 management and nutritional strategies during the close up part of the transition period are vital in
348 order to establish normal macromineral concentrations at parturition.

349

350 In the present study, no genetic correlations among serum Ca, P, Mg and K concentrations and
351 their changes from days 1-4 and 1-8 after calving were detected. If there are no genetic
352 correlations, this probably denotes that there are no competitive mechanisms at genetic level that
353 regulate the concentrations of macrominerals. Further research is needed in order to clarify this
354 issue.

355

356 Although small, significant positive phenotypic correlations were found between Ca and P and
357 Ca and K. These correlations are not easy to explain; e.g. one might expect that the action of
358 PTH would result in a negative correlation between Ca and P. However, at the onset of lactation
359 large amounts of macrominerals are excreted in the milk which are maintained almost constant,
360 regardless of serum concentrations in the dam, so that adequate mineral supply can be offered to
361 the newborn calf (Grünberg, 2014). This could explain the observed positive phenotypic

362 correlations. Moreover, the role of calcitonin in decreasing Ca and P blood concentration is well
363 established (Allen and Sansom, 1985; Goff, 2000). Calcitonin actually counteracts PTH and,
364 thus, it protects skeleton against major Ca losses during periods of intense Ca mobilization, such
365 as pregnancy and, especially, lactation. It is likely that this might also explain the observed
366 phenotypic correlation.

367

368 An interesting finding was the negative phenotypic correlations of P with Mg. In humans, the
369 presence of Mg ions in the binding regions of adenylate cyclase and phospholipase C –two
370 intracellular molecules that are activated after the binding of PTH to its cell receptors– is
371 essential for the full activation of these two secondary messengers and the manifestation of PTH
372 action on target tissues (Rude, 1998; Potts and Gardella, 2007). Therefore, hypomagnesaemia
373 reduces the secretion of PTH and decreases the sensitivity of tissues to PTH (Littledike et al.,
374 1983; Goff, 2014). Consequently, this PTH reduction could contribute towards increasing serum
375 P concentration. Moreover, in humans, PTH action in distal tubules reduces Mg renal excretion
376 and contributes towards increased serum Mg levels, while at the same time decreases P
377 concentration (Rude, 1998; Swaminathan, 2000). It remains uncertain whether these mechanisms
378 apply to dairy cows, as well.

379

380 Other interesting findings included the high negative correlations of Ca, P, Mg and K
381 concentrations on day 1 with the respective changes between day 1 and 4 and day 1 and 8. This
382 indicates that the higher the serum concentration on day 1 the smaller is the expected change
383 during the following days (always within normal range). This seems to be particularly interesting
384 especially for Ca. These observations imply that Ca homeostasis was effective, at a population

385 level and support the need for proper nutritional and management strategies during the transition
386 period. Correlations between Ca serum concentration on day 1 and P serum changes corroborate
387 the previous assumptions. Correlations between P serum concentration on day 1 and Ca serum
388 changes follows the same pattern: high concentrations of P in plasma, at levels greater than 6.0
389 mg/dL, inhibit the action of renal 1 α -hydroxylase 25-(OH)-D₃, decreasing Ca reabsorption and
390 thus limiting serum Ca concentration increase (Goff, 2014).

391

392 Phenotypic correlations between Mg serum concentration on day 1 and Ca changes from day 1 to
393 8 and P changes from day 1 to 4 and 1 to 8, as well as K serum concentrations at day 1 and P
394 changes from day 1 to 4 and 1 to 8 are difficult to interpret, as they usually remain within normal
395 ranges. Cluster analysis may be the appropriate statistical method to analyze these phenomena.

396

397

CONCLUSIONS

398 In the present study, significant genetic variation was found in serum macromineral
399 concentrations immediately after calving. During the first 8 days post-partum, day-to-day
400 heritabilities of serum Ca, P and Mg concentrations traits were moderate to high, while those of
401 K were low to moderate. Genetic evaluation of dairy cows for these traits seems possible and this
402 would contribute to the selection of animals that are less prone to macromineral-related
403 deficiencies during the early stages of lactation that can compromise health and productivity. As
404 these results are the first of their kind, independent validation on different cattle populations
405 would be desirable. Further studies should also focus on the identification of specific genomic
406 regions affecting these traits.

407

408

REFERENCES

- 409 Abdel-Azim, G.A., A.E. Freeman, M.E. Kehrli, S.C. Kelm, J.L. Burton, A.L. Kuck, and S.
410 Schnell. 2005. Genetic basis and risk factors for infectious and noninfectious diseases in US
411 Holsteins. I. Estimation of genetic parameters for single diseases and general health. *J.*
412 *Dairy Sci.* 88:1199–1207. doi:10.3168/jds.S0022-0302(05)72786-7.
- 413 Allen, W.M., and B.F. Sansom. 1985. Milk fever and calcium metabolism. *J. Vet. Pharmacol.*
414 *Ther.* 8:19–29.
- 415 Asagiri, M., and H. Takayanagi. 2007. The molecular understanding of osteoclast differentiation.
416 *Bone.* 40:251–64. doi:10.1016/j.bone.2006.09.023.
- 417 Banos, G., S. Brotherstone, and M.P. Coffey. 2005. Genetic profile of total body energy content
418 of Holstein cows in the first three lactations. *J. Dairy Sci.* 88:2616–2623.
419 doi:10.3168/jds.S0022-0302(05)72938-6.
- 420 Bastin, C., N. Gengler, and H. Soyeurt. 2011. Phenotypic and genetic variability of production
421 traits and milk fatty acid contents across days in milk for Walloon Holstein first-parity
422 cows. *J. Dairy Sci.* 94:4152–4163. doi:10.3168/jds.2010-4108.
- 423 Berry, D.P., F. Buckley, P. Dillon, R.D. Evans, M. Rath, and R.F. Veerkamp. 2003. Genetic
424 relationships among body condition score, body weight, milk yield, and fertility in dairy
425 cows. *J. Dairy Sci.* 86:2193–2204. doi:10.3168/jds.S0022-0302(03)73809-0.
- 426 Bethard, G., R. Verbeck, and J.F. Smith. 1998. Technical Report 31 - Controlling Milk Fever and
427 Hypocalcemia in Dairy Cattle : Use of Dietary Cation-Anion Difference (DCAD) in
428 Formulating Dry Cow Rations. 1–5.
- 429 Cappariello, A., A. Maurizi, V. Veeriah, and A. Teti. 2014. The Great Beauty of the osteoclast.

430 *Arch. Biochem. Biophys.* 558:70–78. doi:10.1016/j.abb.2014.06.017.

431 Castillo-Juarez, H., P.A. Oltenacu, R.W. Blake, C.E. McCulloch, and E.G. Cienfuegos-Rivas.
432 2000. Effect of herd environment on the genetic and phenotypic relationships among milk
433 yield, conception rate, and somatic cell score in Holstein cattle. *J. Dairy Sci.* 83:807–814.
434 doi:10.3168/jds.S0022-0302(00)74943-5.

435 Correa, M.T., C.R. Curtis, H.N. Erb, J.M. Scarlett, and R.D. Smith. 1990. An ecological analysis
436 of risk factors for postpartum disorders of Holstein-Friesian cows from thirty-two New
437 York farms. *J. Dairy Sci.* 73:1515–24. doi:10.3168/jds.S0022-0302(90)78819-4.

438 DeGaris, P.J., and I.J. Lean. 2008. Milk fever in dairy cows: A review of pathophysiology and
439 control principles. *Vet. J.* 176:58–69. doi:10.1016/j.tvjl.2007.12.029.

440 Van Dorp, T.E., J.C. Dekkers, S.W. Martin, and J.P. Noordhuizen. 1998. Genetic parameters of
441 health disorders, and relationships with 305-day milk yield and conformation traits of
442 registered Holstein cows. *J. Dairy Sci.* 81:2264–2270. doi:10.3168/jds.S0022-
443 0302(98)75806-0.

444 Dyrendahl, I., B. Henricson, and G. Jönsson. 1972. Clinical puerperal paresis and hypocalcaemia
445 in cattle. A statistical and genetic investigation. *Zentralbl. Veterinarmed. A.* 19:621–638.

446 Gilmour, A.R., and B.J. Gogel. 2006. ASReml User Guide. 2005-2007 pp.

447 Goff, J.P. 2000. Pathophysiology of calcium and phosphorus disorders. *Vet. Clin. North Am.*
448 *Food Anim. Pract.* 16:319–337, vii.

449 Goff, J.P. 2008. The monitoring, prevention, and treatment of milk fever and subclinical
450 hypocalcemia in dairy cows. *Vet. J.* 176:50–57. doi:10.1016/j.tvjl.2007.12.020.

451 Goff, J.P. 2014. Calcium and magnesium disorders. *Vet. Clin. North Am. Food Anim. Pract.*
452 30:359–81, vi. doi:10.1016/j.cvfa.2014.04.003.

453 Goff, J.P., and R.L. Horst. 1997. Physiological changes at parturition and their relationship to
454 metabolic disorders. *J. Dairy Sci.* 80:1260–1268. doi:10.3168/jds.S0022-0302(97)76055-7.

455 Gröhn, Y.T., and M.L. Bruss. 1990. Effect of diseases, production, and season on traumatic
456 reticuloperitonitis and ruminal acidosis in dairy cattle. *J. Dairy Sci.* 73:2355–2363.
457 doi:10.3168/jds.S0022-0302(90)78918-7.

458 Grünberg, W. 2014. Treatment of Phosphorus Balance Disorders. *Vet. Clin. North Am. - Food*
459 *Anim. Pract.* 30:383–408. doi:10.1016/j.cvfa.2014.03.002.

460 Heringstad, B., Y.M. Chang, D. Gianola, and G. Klemetsdal. 2005. Genetic analysis of clinical
461 mastitis, milk fever, ketosis, and retained placenta in three lactations of Norwegian red
462 cows. *J. Dairy Sci.* 88:3273–3281. doi:10.3168/jds.S0022-0302(05)73010-1.

463 Kadarmideen, H.N., R. Thompson, G. Simm, and M. Eh. 2000. Linear and threshold model
464 genetic parameters for disease , fertility and milk production in dairy cattle. 411–419.

465 Kaneko, J.J., J.W. Harvey, and M.L. Bruss. 2008. *Clinical Biochemistry of Domestic Animals.*
466 Elsevier. 623-634 pp.

467 Kronqvist, C. 2011. *Minerals to Dairy Cows with Focus on Calcium and Magnesium Balance.*

468 Lean, I.J., R. Van Saun, and P.J. Degaris. 2013. Energy and protein nutrition management of
469 transition dairy cows. *Vet. Clin. North Am. Food Anim. Pract.* 29:337–66.
470 doi:10.1016/j.cvfa.2013.03.005.

471 Lin, H.K., P.A. Oltenacu, L.D. Van Vleck, H.N. Erb, and R.D. Smith. 1989. Heritabilities of and

472 genetic correlations among six health problems in Holstein cows. *J. Dairy Sci.* 72:180–186.
473 doi:10.3168/jds.S0022-0302(89)79095-0.

474 Littledike, E.T., and J. Goff. 1987. Interactions of calcium, phosphorus, magnesium and vitamin
475 D that influence their status in domestic meat animals. *J. Anim. Sci.* 65:1727–1743.

476 Littledike, E.T., J.A. Stuedemann, S.R. Wilkinson, and R.L. Horst. 1983. Grass tetany syndrome.
477 *In Proceedings of John Lee Pratt International Symposium on the Role of Magnesium in*
478 *Animal Nutrition.* Virginia Polytechnic Institute and State University, Blacksburg, Virginia,
479 VA, USA. 173.

480 Lyons, D.T., A.E. Freeman, and A.L. Kuck. 1991. Genetics of health traits in Holstein cattle. *J.*
481 *Dairy Sci.* 74:1092–1100. doi:10.3168/jds.S0022-0302(91)78260-X.

482 Martinez, N., L.D.P. Sinedino, R.S. Bisinotto, E.S. Ribeiro, G.C. Gomes, F.S. Lima, L.F. Greco,
483 C.A. Risco, K.N. Galvão, D. Taylor-Rodriguez, J.P. Driver, W.W. Thatcher, and J.E.P.
484 Santos. 2014. Effect of induced subclinical hypocalcemia on physiological responses and
485 neutrophil function in dairy cows. *J. Dairy Sci.* 97:874–87. doi:10.3168/jds.2013-7408.

486 NRC. 2001. Nutrient Requirements of Dairy Cattle Seventh Revised Edition , 2001. 1-333 pp.

487 Oetzel, G.R. 2011. Diseases of Dairy Animals | Non-Infectious Diseases: Milk Fever. *In*
488 *Encyclopedia of Dairy Sciences (Second Edition).* J.W. Fuquay, editor. Academic Press,
489 San Diego. 239–245.

490 Oikonomou, G., G.E. Valergakis, G. Arsenos, N. Roubies, and G. Banos. 2008. Genetic profile
491 of body energy and blood metabolic traits across lactation in primiparous Holstein cows. *J.*
492 *Dairy Sci.* 91:2814–2822. doi:10.3168/jds.2007-0965.

493 Peek, S.F., and T.J. Divers. 2008. Chapter 14 - Metabolic Diseases. *In* Rebhun's Diseases of
494 Dairy Cattle (Second Edition). T.J.D.F. Peek, editor. W.B. Saunders, Saint Louis. 590–605.

495 Perkin Elmer. 1996. Perkin Elmer AAnalyst 100.

496 Potts, J.T., and T.J. Gardella. 2007. Progress, paradox, and potential: parathyroid hormone
497 research over five decades. *Ann. N. Y. Acad. Sci.* 1117:196–208.
498 doi:10.1196/annals.1402.088.

499 Pradhan, K., and R.W. Hemken. 1968. Potassium depletion in lactating dairy cows. *J. Dairy Sci.*
500 51:1377–81. doi:10.3168/jds.S0022-0302(68)87198-X.

501 Pryce, J.E., R.F. Veerkamp, R. Thompson, W.G. Hill, and G. Simm. 1997. Genetic aspects of
502 common health disorders and measures of fertility in Holstein Friesian dairy cattle. *Anim.*
503 *Sci.* 65:353–360. doi:10.1017/S1357729800008559.

504 Rajala-Schultz, P.J., Y.T. Gröhn, and C.E. McCulloch. 1999. Effects of milk fever, ketosis, and
505 lameness on milk yield in dairy cows. *J. Dairy Sci.* 82:288–94. doi:10.3168/jds.S0022-
506 0302(99)75235-5.

507 Reinhardt, T.A., J.D. Lippolis, B.J. McCluskey, J.P. Goff, and R.L. Horst. 2011. Prevalence of
508 subclinical hypocalcemia in dairy herds. *Vet. J.* 188:122–124.
509 doi:10.1016/j.tvjl.2010.03.025.



510 Rérat, M., A. Philipp, H.D. Hess, and A. Liesegang. 2009. Effect of different potassium levels in
511 hay on acid-base status and mineral balance in periparturient dairy cows. *J. Dairy Sci.*
512 92:6123–6133. doi:10.3168/jds.2009-2449.

513 Rude, R.K. 1998. Magnesium deficiency: a cause of heterogeneous disease in humans. *J. Bone*



- 514 *Miner. Res.* 13:749–58. doi:10.1359/jbmr.1998.13.4.749.
- 515 Sattler, N., and G. Fecteau. 2014. Hypokalemia Syndrome in Cattle. *Vet. Clin. North Am. - Food*
516 *Anim. Pract.* 30:351–357. doi:10.1016/j.cvfa.2014.04.004.
- 517 Sattler, N., G. Fecteau, C. Girard, and Y. Couture. 1998. Description of 14 cases of bovine
518 hypokalaemia syndrome. *Vet. Rec.* 143:503–507. doi:10.1136/vr.143.18.503.
- 519 Swaminathan, R. 2000. Disorders of magnesium metabolism. *CPD Bull. Clin. Biochem.* 2:3–12.
520 doi:10.1016/B978-0-323-04883-5.50036-2.
- 521 Tauriainen, S., S. Sankari, S. Pyörälä, and L. Syrjälä-Qvist. 2003. Effect of anionic salts on some
522 blood and urine minerals, acid-base balance and udder oedema of dry pregnant cows. *Agric.*
523 *Food Sci. Finl.* 12:83–93.
- 524 Tveit, B., M. Svendsen, and K. Hove. 1991. Heritability of hypocalcemia at first parturition in
525 Norwegian cattle: genetic correlations with yield and weight. *J. Dairy Sci.* 74:3561–3567.
526 doi:10.3168/jds.S0022-0302(91)78548-2.
- 527 Uribe, H.A., B.W. Kennedy, S.W. Martin, and D.F. Kelton. 1995. Genetic parameters for
528 common health disorders of Holstein cows. *J. Dairy Sci.* 78:421–430.
529 doi:10.3168/jds.S0022-0302(95)76651-6.
- 530 Windig, J.J., M.P.L. Calus, B. Beerda, and R.F. Veerkamp. 2006. Genetic correlations between
531 milk production and health and fertility depending on herd environment. *J. Dairy Sci.*
532 89:1765–1775. doi:10.3168/jds.S0022-0302(06)72245-7.

533

GENETIC PARAMETERS OF Ca, P, Mg AND K



534




535

536 **Tsiamadis Figure 1.**

537

GENETIC PARAMETERS OF Ca, P, Mg AND K



538

539 **Tsiamadis Figure 2.**

1 **Figure captures**

2

3 **Figure 1.** Fixed curves for serum Calcium (Ca), Phosphorus (P), Magnesium (Mg) and
4 Potassium (K) concentrations across all lactations during the first 8 days after calving from
5 random regression model analyses.

6

7 **Figure 2.** Heritability estimates of serum Calcium (Ca), Phosphorus (P), Magnesium (Mg) and
8 Potassium (K) concentrations during the first 8 days after calving.

Table 1. Least square means for Calcium (Ca), Phosphorus (P), Magnesium (Mg) and Potassium (K) serum concentrations and analysis of variance by parity number and days after calving

	1 st Day after calving					2 nd Day after calving					4 th Day after calving					8 th Day after calving					P	
	Parity number	Mean	SEM	St. dev	Min-Max	Mean	SEM	St. dev	Min-Max	Mean	SEM	St. dev	Min-Max	Mean	SEM	St. dev	Min-Max	Mean	SEM	St. dev		Min-Max
Ca (mg/dL)	1	9.14 ^a	0.05	1.02	3.90-12.90	8.84 ^b	0.04	0.95	4.45-11.65	9.06 ^c	0.05	1.06	4.10-13.90	9.45 ^c	0.05	1.09	5.75-12.85					**
	2	8.49 ^a	0.07	1.10	5.55-11.15	8.54 ^b	0.08	1.19	4.50-12.60	9.10 ^b	0.07	1.08	5.75-12.30	9.42 ^c	0.07	1.14	6.00-13.25					**
	3	8.30 ^a	0.10	1.29	5.25-11.85	8.20 ^a	0.10	1.31	4.05-11.85	8.98 ^b	0.09	1.13	5.15-11.30	9.21 ^b	0.10	1.18	4.45-11.50					***
	4	7.94 ^a	0.11	1.26	3.95-10.90	8.42 ^b	0.10	1.14	5.65-12.00	9.00 ^c	0.10	1.20	5.60-12.20	9.29 ^c	0.11	1.26	6.05-12.95					**
P (mg/dL)	1	5.54 ^a	0.06	1.18	2.20-9.30	5.26 ^b	0.06	1.19	2.10-10.20	5.14 ^b	0.05	1.10	2.70-10.00	4.98 ^c	0.05	1.05	2.10-8.70					*
	2	5.32 ^a	0.09	1.35	2.60-9.30	4.98 ^b	0.09	1.36	2.00-9.50	5.08 ^{ab}	0.08	1.25	2.00-9.40	5.13 ^{ab}	0.08	1.15	2.60-8.60					**
	3	5.18 ^a	0.11	1.44	2.30-9.80	5.14 ^b	0.11	1.35	2.20-10.30	5.18 ^{ab}	0.11	1.35	2.40-10.10	4.76 ^c	0.09	1.10	2.20-7.70					**
	4	5.29 ^a	0.13	1.53	1.40-9.60	5.46 ^{ab}	0.12	1.41	3.00-10.20	5.65 ^b	0.11	1.30	3.00-10.50	5.34 ^{ab}	0.12	1.34	2.50-9.30					*
Mg (mg/dL)	1	2.37 ^a	0.02	0.35	1.50-3.90	2.35 ^a	0.02	0.35	0.95-3.55	2.21 ^b	0.02	0.46	0.38-7.20	2.18 ^b	0.01	0.31	1.10-3.35					***
	2	2.31 ^a	0.03	0.46	1.30-7.10	2.22 ^b	0.03	0.40	1.20-3.45	2.09 ^c	0.02	0.37	1.20-3.35	2.19 ^b	0.03	0.39	0.36-3.65					*
	3	2.41 ^a	0.03	0.44	1.30-4.15	2.31 ^b	0.03	0.42	1.40-3.90	2.07 ^c	0.03	0.36	1.40-3.55	2.13 ^c	0.03	0.35	1.10-3.20					***
	4	2.38 ^a	0.03	0.41	1.55-4.00	2.27 ^a	0.03	0.38	1.30-3.60	2.10 ^b	0.03	0.35	1.40-3.35	2.14 ^b	0.03	0.38	0.40-3.20					*
K (mmol/L)	1	4.7 ^a	0.03	0.59	2.8-6.3	4.6 ^b	0.01	0.55	2.5-6.3	4.5 ^c	0.01	0.47	2.9-6.3	4.4 ^c	0.01	0.47	2.8-6.3					*
	2	4.8 ^a	0.01	0.57	3.6-6.3	4.6 ^b	0.01	0.53	3.2-6.2	4.5 ^{ab}	0.01	0.50	3.5-6.2	4.5 ^c	0.01	0.54	3.0-6.2					**
	3	4.7 ^a	0.01	0.53	3.4-6.3	4.7 ^a	0.01	0.55	3.4-6.2	4.6 ^a	0.01	0.55	3.3-6.3	4.4 ^b	0.01	0.53	2.9-6.3					**
	4	4.7 ^a	0.10	0.63	3.5-6.3	4.7 ^a	0.10	0.61	3.3-6.3	4.6 ^a	0.01	0.54	3.5-6.3	4.6 ^a	0.10	0.65	3.2-6.3					NS

^{a-c} Means in the same row having different superscripts differ significantly.

NS: Non-significant.

* P<0.05, ** P<0.001, *** P<0.0001.

Table 2. Variances and heritability estimates of Calcium (Ca), Phosphorus (P), Magnesium (Mg) and Potassium (K) serum concentrations by days after calving from random regression model analyses. All estimates were statistically greater than zero at P<0.001 level

Trait	Day after calving	σ_p^2	σ_a^2	σ_r^2	h^2
Ca	1 st	1.40 (0.06)	0.44 (0.05)	0.96 (0.06)	0.32 (0.03)
	2 nd	1.26 (0.05)	0.37 (0.04)	0.89 (0.05)	0.29 (0.03)
	4 th	1.22 (0.05)	0.28 (0.03)	0.94 (0.05)	0.23 (0.02)
	8 th	1.30 (0.06)	0.35 (0.08)	0.95 (0.08)	0.27 (0.06)
P	1 st	1.91 (0.08)	0.70 (0.07)	1.21 (0.07)	0.37 (0.03)
	2 nd	1.48 (0.06)	0.57 (0.05)	0.92 (0.05)	0.38 (0.03)
	4 th	1.31 (0.06)	0.40 (0.03)	0.91 (0.05)	0.30 (0.02)
	8 th	1.05 (0.05)	0.45 (0.08)	0.60 (0.07)	0.43 (0.07)
Mg	1 st	0.17 (0.01)	0.07 (0.01)	0.11 (0.01)	0.39 (0.03)
	2 nd	0.16 (0.01)	0.06 (0.01)	0.10 (0.01)	0.35 (0.03)
	4 th	0.19 (0.01)	0.04 (0.00)	0.15 (0.01)	0.20 (0.02)
	8 th	0.12 (0.01)	0.03 (0.01)	0.10 (0.01)	0.21 (0.06)
K	1 st	0.34 (0.02)	0.04 (0.01)	0.30 (0.02)	0.12 (0.03)
	2 nd	0.29 (0.01)	0.04 (0.01)	0.26 (0.01)	0.12 (0.02)
	4 th	0.21 (0.01)	0.03 (0.01)	0.18 (0.01)	0.15 (0.02)
	8 th	0.22 (0.01)	0.05 (0.01)	0.17 (0.01)	0.23 (0.06)

* Phenotypic (σ_p^2), genetic (σ_a^2), residual variances (σ_r^2) and heritability (h^2) estimates (standard errors in parentheses)

Table 3. Heritability estimates of Calcium (Ca), Phosphorus (P), Magnesium (Mg) and Potassium (K) serum concentrations across days (diagonals) and statistically greater than zero phenotypic correlations (above diagonal); standard error in parentheses

Trait	Ca	P	Mg	K
Ca	0.20 (0.02)**	0.16 (0.03)**	NS	0.09 (0.03)*
P		0.25 (0.02)**	-0.16 (0.03)^{a**}	0.20 (0.03)**
Mg			0.21 (0.02)**	NS
K				0.10 (0.02)**

^a Bold letters indicate undesirable correlations.

NS: Non-significant.

* P<0.01, ** P<0.0001.

Table 4. Phenotypic correlations of Calcium, Phosphorus, Magnesium and Potassium serum concentrations on day 1 and corresponding change in days 1-4 and 1-8 after calving; standard error in parentheses

	Ca Change_1-4	Ca Change_1-8	P_1	P Change_1-4	P Change_1-8	Mg_1	Mg Change_1-4	Mg Change_1-8	K_1	K Change_1-4	K Change_1-8
Ca_1	-0.56 (0.02)	-0.45 (0.03)	0.12 (0.03)	-0.12 (0.03)	NS	NS	NS	NS	0.08 (0.03)	NS	NS
Ca Change_1-4		0.53 (0.02)	-0.16 (0.03)	0.15 (0.03)	NS	NS	-0.06 (0.03)	NS	NS	0.09 (0.03)	NS
Ca Change_1-8			NS	NS	-0.11 (0.03)	-0.08 (0.03)	NS	NS	NS	0.07 (0.03)	NS
P_1				NS	-0.52 (0.02)	-0.14 (0.03)	0.18 (0.03)	0.11 (0.03)	0.19 (0.03)	-0.12 (0.03)	-0.08 (0.03)
P Change_1-4					0.66 (0.02)	0.09 (0.03)	-0.07 (0.03)	NS	-0.20 (0.03)	0.14 (0.03)	0.10 (0.03)
P Change_1-8						0.09 (0.03)	NS	NS	-0.16 (0.03)	0.09 (0.03)	0.09 (0.03)
Mg_1							-0.57 (0.02)	-0.55 (0.02)	NS	NS	NS
Mg Change_1-4								0.52 (0.02)	NS	NS	NS
K_1									NS	NS	-0.57 (0.02)
K Change_1-4											0.54 (0.02)

NS: Non-significant.

Ca/P/Mg/K_1 = serum Calcium/ Phosphorus/ Magnesium/ Potassium concentration on day 1 after calving.

Ca/P/Mg/K Change_1-4 = serum Calcium/ Phosphorus/ Magnesium/ Potassium concentration change from days 1 to 4 after calving.

Ca/P/Mg/K Change_1-8 = serum Calcium/ Phosphorus/ Magnesium/ Potassium concentration change from days 1 to 8 after calving.