



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Prevalence of Sarcopenia in A Longitudinal UK Cohort Study Using EWGSOP2 Criteria Varies Widely Depending on Which Measures of Muscle Strength and Performance are Used

Citation for published version:

Kilgour, A, Redmond, P, Taylor, A, Deary, IJ, Starr, J & Shenkin, SD 2020, 'Prevalence of Sarcopenia in A Longitudinal UK Cohort Study Using EWGSOP2 Criteria Varies Widely Depending on Which Measures of Muscle Strength and Performance are Used', British Geriatrics Society Autumn Meeting 2019, 6/11/19 - 8/11/19 pp. i22-i23. <https://doi.org/10.1093/ageing/afz188.01>

Digital Object Identifier (DOI):

[10.1093/ageing/afz188.01](https://doi.org/10.1093/ageing/afz188.01)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Scientific Research — Bone, Muscle and Rheumatology

70 PREVALENCE OF SARCOPENIA IN A LONGITUDINAL UK COHORT STUDY USING EWGSOP2 CRITERIA VARIES WIDELY DEPENDING ON WHICH MEASURES OF MUSCLE STRENGTH AND PERFORMANCE ARE USED

A.H.M. Kilgour^{1,2}, P.Redmond^{1,3}, A. Taylor³, I.J. Deary^{1,3}, J.M. Starr^{1,2}, S.D. Shenkin^{1,2}

¹Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, UK

²Geriatric Medicine Unit, University of Edinburgh, UK

³Department of Psychology, University of Edinburgh, UK

Background: The European Working Group on Sarcopenia in Older People 2 (EWGSOP2) guidelines have recently been published to aid diagnosis of sarcopenia in the clinical setting and to allow for better comparison between research studies. The guidelines recommend several different tests for diagnosing sarcopenia. We hypothesised that the prevalence of sarcopenia might vary depending on which tests are used.

Methods: We used data from Wave 3 of the Lothian Birth Cohort 1936 study, a longitudinal ageing study of healthy, community dwelling older adults (n= 697, 52%

men, mean age 76y), to assess the prevalence of sarcopenia using the suggested cut-offs in the EWGSOP2 guidelines. Probable sarcopenia was defined as low muscle strength (measured by handgrip strength and 5x chair stand test), confirmed sarcopenia was defined as low muscle strength + low lean mass (measured by bioimpedance analysis), and severe sarcopenia was defined as confirmed sarcopenia + low muscle performance (measured by gait speed and short physical performance battery score). SPSS version 24.0 was used for statistical analysis.

Results: The maximum prevalence of probable sarcopenia was 24.2% in men and 24.8% in women, of confirmed sarcopenia was 7.4% in men and 11.0% in women, and of severe sarcopenia was 4.6% in men and 5.9% in women, when either of the cut-offs for muscle strength +/- muscle performance were met. When using only one measure of muscle strength +/- performance, rates of probable sarcopenia ranged from 7.7% to 21.1% in men and 5.9% to 21.3% in women; rates of confirmed sarcopenia ranged from 3.9% to 5.3% in men and 5.1% to 9% in women; and rates of severe sarcopenia ranged from 1.4% to 3.9% in men and from 2.0% to 5.1% in women.

Conclusions: In a UK-based longitudinal ageing study we found that the prevalence of probable, confirmed and severe sarcopenia varied widely using the EWGSOP2 guidelines depending on which identifying tests were used. We found that the cut-off points suggested for some of the measures in the guidelines are not comparable and may lead to differing groups being identified as sarcopenic between different trials. We suggest modification of the cut-offs to adjust for this.