



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Canine and ovine tick-borne pathogens in camels, Nigeria

**Citation for published version:**

Lorusso, V, Wijnveld, M, Latrofa, MS, Fajinmi, A, Majekodunmi, AO, Dogo, AG, Igweh, AC, Otranto, D, Jongejan, F, Welburn, SC & Picozzi, K 2016, 'Canine and ovine tick-borne pathogens in camels, Nigeria', *Veterinary Parasitology*, vol. 228, pp. 90-92. <https://doi.org/10.1016/j.vetpar.2016.08.014>

**Digital Object Identifier (DOI):**

[10.1016/j.vetpar.2016.08.014](https://doi.org/10.1016/j.vetpar.2016.08.014)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

*Veterinary Parasitology*

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.





## Short communication

## Canine and ovine tick-borne pathogens in camels, Nigeria



Vincenzo Lorusso<sup>a</sup>, Michiel Wijnveld<sup>b</sup>, Maria S. Latrofa<sup>c</sup>, Akinyemi Fajinmi<sup>d</sup>,  
Ayodele O. Majekodunmi<sup>a</sup>, Abraham G. Dogo<sup>d</sup>, Augustine C. Igweh<sup>e</sup>, Domenico Otranto<sup>c</sup>,  
Frans Jongejan<sup>b,f</sup>, Susan C. Welburn<sup>a</sup>, Kim Picozzi (Dr.)<sup>a,\*</sup>

<sup>a</sup> Division of Infection and Pathway Medicine, Edinburgh Medical School: Biomedical Sciences, The University of Edinburgh, Edinburgh, United Kingdom

<sup>b</sup> Utrecht Centre for Tick-borne Diseases, Utrecht University, Utrecht, The Netherlands

<sup>c</sup> Department of Veterinary Medicine, The University of Bari, Bari, Italy

<sup>d</sup> Nigerian Institute for Trypanosomiasis Research, VOM, Jos, Plateau State, Nigeria

<sup>e</sup> National Veterinary Research Institute, VOM, Jos, Plateau State, Nigeria

<sup>f</sup> Department of Veterinary Tropical Diseases, Faculty of Veterinary Medicine, University of Pretoria, South Africa

## ARTICLE INFO

## Article history:

Received 28 June 2016

Received in revised form 15 August 2016

Accepted 19 August 2016

## Keywords:

*Anaplasma platys*

*Theileria ovis*

*Camelus dromedarius*

Camel

Tick-borne diseases

Nigeria

## ABSTRACT

In April 2008, whole blood samples were collected from 36 dromedary camels in Sokoto, North-western Nigeria. Following PCR and reverse line blotting, twenty-two samples (61%) resulted positive for *Ehrlichia/Anaplasma* spp. and three (8%) for *Theileria/Babesia* spp., with three (8%) cases of co-infections being found. Both sequence and BLAST analyses identified *Ehrlichia/Anaplasma* spp. and *Theileria/Babesia* spp. positive cases as *Anaplasma platys* and *Theileria ovis*, respectively.

This is the first report of the detection of *A. platys* and *T. ovis* in camels from sub-Saharan Africa. The epidemiological relevance of this finding is enhanced by the close living of these animals with both dogs and small ruminants. The high prevalence detected for *A. platys* suggests a possible role of camels as carriers of this infection.

© 2016 Published by Elsevier B.V.

Nigeria hosts a population of approximately 20,000 dromedary camels (i.e. *Camelus dromedarius*), most of which are found in its northern States (i.e. Sokoto State) (Mohammed and Hoffmann, 2006). Camels in Nigeria are reared for milk, meat and wool production; transport; traction in agriculture; recreation and beauty pageants (Mohammed and Hoffmann, 2006). In this country, ticks and tick-borne diseases (TBDs) represent a major constraint to its livestock health and productivity (Rabana et al., 2011). In camels, heavy tick infestation is associated with anaemia, rough hair coat, retarded growth, reduction in milk production and calf mortality (Rabana et al., 2011). However, little information is currently available on the occurrence of TBDs in camels in Nigeria, with all studies published to date relying on pathogen detection through cytological examination of blood smears (Mohammed et al., 2007; Bamaïyi et al., 2011; Rabana et al., 2011). Therefore, this study aimed to assess, by molecular tools, the occurrence of tick-borne microorganisms of veterinary and zoonotic importance in dromedary

camels from an area of Nigeria where camel rearing is of great economic relevance.

In April 2008, whole blood samples were collected, by jugular venipuncture, from 36 randomly selected dromedary camels (i.e. 13 males, 23 females) reared in the surroundings of the city of Sokoto (i.e. local government areas of Sokoto North, Central and South; Sokoto State), in North-western Nigeria. All sampled animals were restrained with the help of their owners and handled humanely. For each sampled animal, collected blood was applied to FTA<sup>TM</sup> cards (Whatman, BioScience, Cambridge, UK) and prepared for downstream analysis according to Ahmed et al. (2013). Based on physical examination (e.g. general somatic development and dentition) and on information provided by their owners, all sampled animals were identified as adults, being at least 5 year-old (FAO, 1990).

Reverse line blotting (RLB) targeting *Ehrlichia/Anaplasma* spp. and *Rickettsia* spp. 16S and *Theileria/Babesia* spp. 18S partial genes were carried out as described elsewhere (Lorusso et al., 2016). Following RLB, amplicons were sequenced to confirm their identity; selected sequences amongst those obtained were deposited in GenBank on May the 15th, 2014.

Positive samples were compared according to the sex of the animals using the Fisher's exact test with the WinPepi software.

\* Corresponding author at: Division of Infection and Pathway Medicine Edinburgh Medical School: Biomedical Sciences, The University of Edinburgh, 49 Little France Crescent, EH16 4SB, Edinburgh, United Kingdom.

E-mail address: [kim.picozzi@ed.ac.uk](mailto:kim.picozzi@ed.ac.uk) (K. Picozzi).

**Table 1**  
Sampled camels and results of the screening.

Camels	Sampled	Positive/Total Sampled			
		Total +	<i>Anaplasma platys</i> +	<i>Theileria ovis</i> +	<i>Anaplasma platys</i> + and <i>Theileria ovis</i> +
Males	13	7/13	7/13	0/13	0/13
Females	23	15/23	15/23	3/23	3/23
Total	36	22/36	22/36	3/36	3/36

P values lower than 0.05 were considered as indicative of significance.

At RLB, 22 samples (61%) resulted positive for *Ehrlichia/Anaplasma* spp. 'catch all' probe, three (8%) for *Theileria/Babesia* spp., with three (8%) cases of co-infections being found. Both sequence and BLAST analyses identified *Ehrlichia/Anaplasma* spp. and *Theileria/Babesia* spp. positive samples as *Anaplasma platys* (99–100% identity with GenBank accession no. JQ894779.2 and KJ659045.1) and *Theileria ovis* (99% identity with GenBank accession no. KJ452336.1), respectively. Sequences were deposited in GenBank (GenBank accession no. KJ832066 and KJ832067 for *A. platys*; KJ832064 and KJ832065 for *T. ovis*).

Camels positive for *A. platys* infection included 7/13 (54%) males and 15/23 (65%) females, with no statistically significant difference between the two sex groups ( $p=0.72$ ). However, animals positive also for *T. ovis* infection included only female camels (i.e. 3/23) (Table 1).

These findings are of novelty for camels in sub-Saharan Africa (SSA).

The high positive rate (61%) of *A. platys* in camels in SSA is unexpected in that the pathogen was initially considered to be restricted to dogs, but more recently reported in cats (Lima et al., 2010), sheep (Djiba et al., 2013), cattle (Lorusso et al., 2016) and also in humans (Arraga-Alvarado et al., 2014). Moreover, DNA of anaplasmataceae closely related to *A. platys* was recorded in the spleen of dromedary camels from Saudi Arabia (Bastos et al., 2015). This suggests that the host range of this canine pathogen may be broader than initially considered. Nevertheless, no information is available on the pathogenicity of this microorganism in animal hosts other than dogs. Evidence indicates the role of *Rhipicephalus sanguineus sensu lato* (s.l.) ticks in the transmission of this microorganism (Ramos et al., 2014). Though preferably feeding on dogs (Dantas-Torres, 2010), *R. sanguineus* s.l. can also be found infesting camels as well as other livestock species (Walker et al., 2003; Lorusso et al., 2013). The high prevalence of *A. platys* infection found in this study (i.e. 61%) is seemingly attributable to the high sensitivity of the RLB method employed (Lorusso et al., 2016). It is also possible that the rather small sample size contributed to some extent to the obtainment of this prevalence. Nevertheless, the adult age of all sampled animals, as well as their provenance from different localities, suggests the possible implication of camels as 'carriers' of *A. platys* infection in this area.

*Theileria ovis* was previously detected in camels in Egypt by cytological examination of blood smears, with a prevalence of 12.6% ( $n=24/190$ ) (Mazyad and Khalaf, 2002). Interestingly, DNA from a *Theileria* sp. showing 98% of identity with *T. ovis* was previously detected in one dog from North-Central Nigeria (GenBank accession no. GU726904) (Kamani et al., 2013). *T. ovis* can be transmitted by ticks of the genera *Hyalomma* and *Rhipicephalus* (Walker et al., 2003), both possibly infesting livestock and camels in North-western Nigeria (Lorusso et al., 2013). All camel keepers from the area of sampling also owned dogs and small ruminants.

The lack of significant difference ( $p=0.72$ ) in *A. platys* infection rates between male and female camels suggests that sex-related differences may not influence the establishment of this infection in these animals (Kamani et al., 2008; Bamaiyi et al., 2011; Ben Said et al., 2013). Furthermore, the fact that *T. ovis* positive cases

were recorded only in female camels, all of which were also positive for *A. platys*, could be related to the higher proportion of female animals compared to males sampled in the study (1.7:1) as well as to the fact that female camels, usually harboring more ticks than males (Elghali and Hassan, 2009), could have been infested by a larger number of tick specimens (i.e. *Hyalomma* spp.) responsible for the transmission of this microorganism in the study area. Due to impracticalities (i.e. lack of appropriate sampling and storing tools), this study did not include a collection or an estimation of the tick burden on the sampled animals. Nonetheless, it was noted that all camels sampled in this study were infested by ixodid ticks, including *Hyalomma* spp. and *Rhipicephalus* spp. specimens.

Future studies would therefore be advisable in order to ascertain i) the pathogenicity of *A. platys* and *T. ovis* in dromedary camels, ii) the tick species involved in their transmission to these vertebrate hosts, iii) the occurrence of these microorganisms in dogs and sheep from the study area, and thus iv) the role played by dromedary camels in the epidemiology of the canine and ovine infection respectively.

## Acknowledgments

The research leading to these results has received funding from the UK Department for International Development (DFID) under the umbrella of the 'Stamp Out Sleeping Sickness (SOS) Programme', the UK's Biotechnology and Biological Sciences Research Council (BBSRC) under the 'Combating Infectious Diseases in Livestock for International Development' (CIDLID) scheme, and the European Union's Seventh Framework Program (FP7/2007-2013) under grant agreement no 221948, Integrated Control of Neglected Zoonoses (ICONZ). We would like to thank Dr. Pwaveno H. Bamaiyi (University Malaysia Kelantan) for helping retrieving some of the literature on Nigerian camels. All farmers from the study area are acknowledged for their kind cooperation during sampling. This research has been approved by The University of Edinburgh Ethical Committee for the review of animal research conducted outside the UK.

## References

- Ahmed, H.A., Picozzi, K., Welburn, S.C., MacLeod, E.T., 2013. A comparative evaluation of PCR based methods for species specific determination of African animal trypanosomes in Ugandan cattle. *Parasit. Vectors.* 6, 316.
- Arraga-Alvarado, C.M., Qurollo, B., Parra, O.C., Berrueta, M.A., Hegarty, B.C., Breitschwerdt, E.B., 2014. Case report: molecular evidence of *Anaplasma platys* infection in two women from Venezuela. *Am. J. Trop. Med. Hyg.* 91, 1161–1165.
- Bamaiyi, P.H., Kalu, A.U., Ali, M., 2011. Haemoparasites of the trade camel (*Camelus dromedarius*) arriving for slaughter at Maiduguri Borno State, Nigeria. *Cont. J. Vet. Sci.* 5, 18–21.
- Bastos, A.D.S., Mohamed, O.B., Bennet, N.C., Petevinos, C., Alagaili, A.N., 2015. Molecular detection of novel Anaplasmataceae closely related to *Anaplasma platys* and *Ehrlichia canis* in the dromedary camel (*Camelus dromedarius*). *Vet. Microbiol.* 179, 310–314.
- Ben Said, M., Belkahia, H., Sayahi, L., Aloui, M., Jemli, M.H., Hadj Mohamed, B., Sassi, L., Darghouth, M.A., Djaïem, A.A., Bayouhd, M., Messadi, L., 2013. First serological study of the prevalence of *Anaplasma phagocytophilum* in dromedary (*Camelus dromedarius*) in Tunisia. *Bull. Soc. Pathol. Exot.* 107, 1–6.
- Dantas-Torres, F., 2010. Biology and ecology of the brown dog tick, *Rhipicephalus sanguineus*. *Parasit. Vectors.* 3, 26.
- Djiba, M.L., Mediannikov, O., Mbengue, M., Thiongane, Y., Molez, J.F., Seck, M.T., Fenollar, F., Raoult, D., Ndiaye, M., 2013. Survey of Anaplasmataceae bacteria in sheep from Senegal. *Trop. Anim. Health Prod.* 45, 1557–1561.

- Elghali, A., Hassan, S.M., 2009. Ticks (Acari: Ixodidae) infesting camels (*Camelus dromedarius*) in northern Sudan. *Onderstepoort J. Vet. Res.* 76, 177–185.
- FAO. 1990. Reproduction in camels – a review. By: Merkt, H., Rath, D., Musa, B., El-Naggar, M.A., FAO Animal Production and Health paper, 82, p. 25.
- Kamani, J., Turaki, A.U., Egwu, G.O., Mani, A.U., Saidu, M.K., Abdullhai, J.G., Kumshe, H.A., 2008. Haemoparasites of camels (*Camelus dromedaries*) in Maiduguri. *Nigeria Anim. Res. Int.* 5, 838–839.
- Kamani, J., Baneth, G., Mumcuoglu, K.Y., Waziri, N.E., Eyal, O., Guthmann, Y., Harrus, S., 2013. Molecular detection and characterization of tick-borne pathogens in dogs and ticks from Nigeria. *PLoS Negl. Trop. Dis.* 7, e2108.
- Lima, M.L., Soares, P.T., Ramos, C.A., Araújo, F.R., Ramos, R.A., Souza, I.I., Faustino, M.A., Alves, L.C., 2010. Molecular detection of *Anaplasma platys* in a naturally-infected cat in Brazil. *Braz. J. Microbiol.* 41, 381–385.
- Lorusso, V., Picozzi, K., de Bronsvort, B.M., Majekodunmi, A., Dongkum, C., Balak, G., Igweh, A., Welburn, S.C., 2013. Ixodid ticks of traditionally managed cattle in central Nigeria: where *Rhipicephalus (Boophilus) microplus* does not dare (yet?). *Parasites Vectors* 6, 171.
- Lorusso, V., Wijnveld, M., Majekodunmi, A.O., Dongkum, C., Fajinmi, A., Dogo, A.G., Thrusfield, M., Mugenyi, A., Vaumourin, E., Igweh, A.C., Jongejan, F., Welburn, S.C., Picozzi, K., 2016. Tick-borne pathogens of zoonotic and veterinary importance in Nigerian cattle. *Parasites Vectors* 9, 217.
- Mazyad, S.A., Khalaf, S.A., 2002. Studies on *Theileria* and *Babesia* infecting live and slaughtered animals in Al Arish and El Hasanah North Sinai Governorate, Egypt. *J. Egypt. Soc. Parasitol.* 32, 601–610.
- Mohammed, I., Hoffmann, I., 2006. Management of draught camels (*Camelus dromedarius*) in crop-livestock production systems in northwest Nigeria. *Livestock Res. Rural Dev.* 18.
- Mohammed, A.K., Sackey, A.K.B., Tekdek, L.B., Gefu, J.O., 2007. Common health problems of the one humped camel (*Camelus dromedarius*) introduced into sub-humid climate in Zaria, Nigeria. *Res. J. Anim. Sci.* 1, 1–5.
- Rabana, J.L., Kumshe, H.A., Kamani, J., Hafsat, G., Turaki, U.A., Dilli, H.K., 2011. Effects of parasitic infections on erythrocyte indices of camels in Nigeria. *Vet. Res. Forum* 2, 59–63.
- Ramos, R.A., Latrofa, M.S., Giannelli, A., Lacasella, V., Campbell, B.E., Dantas-Torres, F., Otranto, D., 2014. Detection of *Anaplasma platys* in dogs and *Rhipicephalus sanguineus* group ticks by a quantitative real-time PCR. *Vet. Parasitol.* 205, 285–288.
- Walker, A.R., Bouattour, A., Camicas, J.L., Estrada-Peña, A., Horak, I.G., Latif, A., Pegram, R.G., Preston, P.M., 2003. Ticks of domestic animals in Africa. In: *A Guide to Identification of Species Bioscience Reports*. Edinburgh, U. K, pp. 1–221.