



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Universal properties of partial quantum maps

Citation for published version:

Andres-Martinez, P, Heunen, C & Kaarsgaard, R 2023, Universal properties of partial quantum maps. in *Proceedings 19th International Conference on Quantum Physics and Logic*. vol. 394, EPTCS, pp. 192–207, Quantum Physics and Logic 2022, Oxford, United Kingdom, 27/06/22.
<https://doi.org/10.48550/arXiv.2206.04814>, <https://doi.org/10.4204/EPTCS.394.11>

Digital Object Identifier (DOI):

[10.48550/arXiv.2206.04814](https://doi.org/10.48550/arXiv.2206.04814)
[10.4204/EPTCS.394.11](https://doi.org/10.4204/EPTCS.394.11)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Proceedings 19th International Conference on Quantum Physics and Logic

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Universal Properties of Partial Quantum Maps

Pablo Andrés-Martínez*
University of Edinburgh

Chris Heunen†
University of Edinburgh

Robin Kaarsgaard‡
University of Edinburgh

We provide a universal construction of the category of finite-dimensional C^* -algebras and completely positive trace-nonincreasing maps from the rig category of finite-dimensional Hilbert spaces and unitaries. This construction, which can be applied to any dagger rig category, is described in three steps, each associated with their own universal property, and draws on results from dilation theory in finite dimension. In this way, we explicitly construct the category that captures hybrid quantum/classical computation with possible nontermination from the category of its reversible foundations. We discuss how this construction can be used in the design and semantics of quantum programming languages.

1 Introduction

The account of quantum measurement offered by *decoherence* establishes that the irreversible nature of mixed-state evolution occurs when a system is considered in isolation from its environment. When the environment is brought back into view, via mathematical techniques such as *quantum state purification* and *Stinespring dilation*, the reversible underpinnings of mixed-state evolution are exposed.

This perspective has in recent years led to the study of quantum theory through categorical completions of its reversible foundations, the category of finite-dimensional Hilbert spaces and unitaries, demonstrating connections between *universal* constructions and effectful quantum programming [10]. This article constructs in a universal way the category of finite-dimensional C^* -algebras and *partial quantum channels* (completely positive trace-nonincreasing maps) from the rig category of finite-dimensional Hilbert spaces and unitaries. The construction has three stages, each with a universal property of its own.

- Freely allowing *partiality respecting the dagger structure* (by making the additive unit a *zero object*) allows contractive maps to be described by unitaries through *Halmos dilation* [7, 24, 19].
- Freely allowing the *hiding of states* in a way that *respects partiality* (by making the multiplicative unit *terminal* for *total* maps) allows completely positive trace-nonincreasing maps to be described through contractions, using a variant of *Stinespring dilation* [28]. This construction has an interesting universal property as a pushout of monoidal categories.
- Freely splitting measurement maps between finite-dimensional Hilbert spaces yields finite-dimensional C^* -algebras, which describe *hybrid quantum/classical* computation.

All three universal constructions are abstract and apply to any suitably structured category. They show that the traditional model of C^* -algebras inevitably arises from the mere concepts of quantum circuits, partiality, hiding, and classical communication, without any concept of *e.g.* norm. Thus they inform the design of quantum programming languages [10], as part of a highly effective broader approach to program semantics from universal properties [29, 15, 26].

*Supported by EPSRC grant EP/L01503X/1 via the CDT in Pervasive Parallelism

†Supported by EPSRC Fellowship EP/R044759/1

‡Supported by DFF | *Natural Sciences* International Postdoctoral Fellowship 0131-00025B

Related work The role of universal properties and categorical completions in quantum theory, in particular the *monoidal indeterminates* construction [8], has been studied in recent years [23, 13, 14, 3, 9, 10] (see also [27] for a related approach in the probabilistic case, and [30, 21] for other accounts of dilations as universal properties). The role of partiality in effectus theory was studied in [2]. We deepen the connections between dilations and universal properties of functors between categories of quantum systems first observed in [13, 14]. A direct connection between universal constructions and the design and semantics of quantum programming languages with effects was demonstrated in [10]. The idempotent splitting of the category of finite-dimensional Hilbert spaces and completely positive maps, in particular the fact that it contains all finite-dimensional C*-algebras, has been the subject of study and discussion in [11, 23].

Overview Section 2 recalls some facts about rig categories and their additive affine completion and relation to partiality, which we extend in Section 3 to the biaffine completion and dagger partiality, and show that the biaffine completion of the category **Unitary** of (finite-dimensional Hilbert spaces and) unitaries is precisely **Contraction** of contractive maps. Section 4 describes a construction that completes **Contraction** to the category **FHil**_{CPTN} of *partial quantum channels* (completely positive trace-nonincreasing maps) using a variant of Stinespring dilation, and we show that this construction satisfies a universal property as a pushout in the category of monoidal categories. Finally, Section 5 shows that splitting along a particular class of idempotents, corresponding in **FHil**_{CPTN} to *measurements*, completes **FHil**_{CPTN} to the category **FCstar**_{CPTN} of finite-dimensional C*-algebras and partial quantum channels. We end in Section 6 with a discussion of the applications of these constructions to the design and semantics of quantum programming languages.

2 The additive affine completion of rig categories

We enter the story assuming that the reader is familiar with monoidal categories and dagger categories [12], and proceed with preliminaries about rig categories and their additive affine completion.

A *rig category* is a category which is symmetric monoidal in two different ways, such that one monoidal product distributes over the other, subject to a large amount of coherence equations [18]. In analogy with the situation in Hilbert spaces, we usually write these monoidal products as (\otimes, I) (the “tensor product”) and (\oplus, O) (the “direct sum”) with \otimes distributing (up to natural isomorphism) over \oplus via distributors $\delta^L: A \otimes (B \oplus C) \rightarrow (A \otimes B) \oplus (A \otimes C)$ and $\delta^R: (A \oplus B) \otimes C \rightarrow (A \otimes C) \oplus (B \otimes C)$ and annihilators (“nullary distributors”) $\delta_0^L: O \otimes A \rightarrow O$ and $\delta_0^R: A \otimes O \rightarrow O$. A *dagger rig category* is a dagger category with a rig structure such that all coherence isomorphisms are unitary (*i.e.*, satisfy $f^{-1} = f^\dagger$). Natural examples of rig categories are *distributive categories* where \otimes is a categorical product with a terminal object as its monoidal unit and \oplus is a categorical coproduct with an initial unit. However, not all rig categories are of this form: the category **Unitary** of finite-dimensional Hilbert spaces and unitaries (with tensor product and direct sum) and the category **FinBij** of finite sets and bijections (with cartesian product and disjoint union) are both (dagger) rig categories, but neither has products or coproducts.

2.1 Partiality and the additive affine completion

What is the appropriate notion of *partiality* for a given rig category? If coproducts and a terminal object are available the *lift monad* $(-)+1$ can answer this question, but not every rig category has these. Instead, we can think of a partial map $A \rightarrow B$ as a map $A \rightarrow B \oplus E$, *i.e.*, extend the output state space with an extra

part E to receive all the inputs we wish to be undefined. Any map $f: A \rightarrow B$ can be lifted to a total map $\rho_{\oplus}^{-1} \circ f: A \rightarrow B \oplus O$ using the inverse right unitor, and partial maps $f: A \rightarrow B \oplus E$ and $g: B \rightarrow C \oplus E'$ can be composed by composing their defined parts, i.e., $\alpha_{\oplus} \circ g \otimes \text{id}_E \circ f: A \rightarrow C \oplus (E' \oplus E)$. This is an information-preserving variant of Kleisli-composition for the lift monad.

This describes the *additive affine completion* of a rig category, barring one detail: given that E describes where the map is undefined, it shouldn't actually matter how we represent this particular part. For example, given some partial map $f: A \rightarrow B \oplus E$ and some manipulation $m: E \rightarrow E'$ of the undefined part, f and $(\text{id} \oplus m) \circ f$ morally describe the same partial map. Therefore, given morphisms $f: A \rightarrow B \oplus E$ and $f': A \rightarrow B \oplus E'$, we write $f \leq_L f'$ if and only if there exists some *mediator* $m: E \rightarrow E'$ such that

$$\begin{array}{ccc} & A & \\ f \swarrow & & \searrow f' \\ B \oplus E & \xrightarrow{\text{id} \oplus m} & B \oplus E' \end{array}$$

commutes. This straightforwardly gives a preorder, though not (necessarily) an equivalence relation since mediators need not be invertible. However, since we would like momentarily to treat it as an equivalence, we consider instead its equivalence closure \sim_L , i.e., the least equivalence relation containing \leq_L .

Definition 1. Given a rig category \mathbf{C} , its *additive affine completion* $L_{\oplus}(\mathbf{C})$ is the category whose

- objects are those of \mathbf{C} ,
- morphisms $[f, E]: A \rightarrow B$ are pairs of an object E and an equivalence class of morphisms $f: A \rightarrow B \oplus E$ of \mathbf{C} under \sim_L ,
- identities $A \rightarrow A$ are $[\rho_{\oplus}^{-1}, O]$ (with $\rho_{\oplus}: A \oplus O \rightarrow A$ the right unitor), and
- composition of $[f, E]: A \rightarrow B$ and $[g, E']: B \rightarrow C$ is $[\alpha_{\oplus} \circ g \otimes \text{id}_E \circ f, E' \oplus E]$.

There is a dual to this construction, the *additive coaffine completion* $R_{\oplus}(\mathbf{C})$, defined as $L_{\oplus}(\mathbf{C}^{\text{op}})^{\text{op}}$. Explicitly, morphisms $A \rightarrow B$ in $R_{\oplus}(\mathbf{C})$ are equivalence classes of morphisms $A \oplus E \rightarrow B$, and so hide part of their *source* space rather than their *target* space. We summarise some features of these categories.

Proposition 2. When \mathbf{C} is a rig category, so are $L_{\oplus}(\mathbf{C})$ and $R_{\oplus}(\mathbf{C})$.

Proof. That $R_{\oplus}(\mathbf{C})$ is a rig category was shown in [10, Lemma 12]; that $L_{\oplus}(\mathbf{C})$ is also a rig category follows by $L_{\oplus}(\mathbf{C}) \cong R_{\oplus}(\mathbf{C}^{\text{op}})^{\text{op}}$ and the fact that \mathbf{C} is a rig category iff \mathbf{C}^{op} is. \square

Proposition 3. The additive unit O is terminal in $L_{\oplus}(\mathbf{C})$ and initial in $R_{\oplus}(\mathbf{C})$.

Proof. The inverse left unitor $\lambda_{\oplus}^{-1}: A \rightarrow O \oplus A$ of \mathbf{C} represents a morphism $A \rightarrow O$ in $L_{\oplus}(\mathbf{C})$; this satisfies the universal property of the terminal object by definition of \sim_L . Dually, O is initial in $R_{\oplus}(\mathbf{C})$. \square

We call a rig category *additively coaffine* when the additive unit is initial, and *additively affine* when it is terminal.

Proposition 4. There are strict rig functors $\mathcal{D}: \mathbf{C} \rightarrow L_{\oplus}(\mathbf{C})$ and $\mathcal{E}: \mathbf{C} \rightarrow R_{\oplus}(\mathbf{C})$.

Proof. Define $\mathcal{D}(A) = A$ on objects, and $\mathcal{D}(f) = [\rho_{\oplus}^{-1} \circ f, O]$ on morphisms, and \mathcal{E} dually. Straightforward calculations show that this defines strict rig functors. \square

As the name suggests, these are, indeed, completions on rig categories.

Proposition 5. $L_{\oplus}(\mathbf{C})$ is the additive affine completion of \mathbf{C} in the following sense: given any additively affine rig category \mathbf{D} and strong rig functor $F: \mathbf{C} \rightarrow \mathbf{D}$, there is a unique strong rig functor $\hat{F}: L_{\oplus}(\mathbf{C}) \rightarrow \mathbf{D}$ making the diagram below commute.

$$\begin{array}{ccc} \mathbf{C} & \xrightarrow{\mathcal{G}} & L_{\oplus}(\mathbf{C}) \\ & \searrow F & \downarrow \hat{F} \\ & & \mathbf{D} \end{array}$$

Proof. By [10, Theorem 19] and duality. □

Dualising this result exhibits $R_{\oplus}(\mathbf{C})$ as the additive coaffine completion of \mathbf{C} .

3 Dagger partiality and the additive biaffine completion

As the (co)affine completion explicitly involves hiding a part of the source or target space of a morphism, we cannot expect to lift either construction to a completion of *dagger* rig categories. A great example of this fact is demonstrated by considering $R_{\oplus}(\mathbf{Unitary})$. Write **Isometry** for the category of finite-dimensional Hilbert spaces and morphisms satisfying $f^{\dagger} \circ f = \text{id}$, and **coIsometry** for its dual.

Proposition 6. *There are rig equivalences $R_{\oplus}(\mathbf{Unitary}) \simeq \mathbf{Isometry}$ and $L_{\oplus}(\mathbf{Unitary}) \simeq \mathbf{coIsometry}$.*

Proof. That $R_{\oplus}(\mathbf{Unitary}) \simeq \mathbf{Isometry}$ was shown in [13]. The other statement follows from duality: $L_{\oplus}(\mathbf{Unitary}) \simeq R_{\oplus}(\mathbf{Unitary}^{\text{op}})^{\text{op}} \simeq R_{\oplus}(\mathbf{Unitary})^{\text{op}} \simeq \mathbf{Isometry}^{\text{op}} \simeq \mathbf{coIsometry}$. □

However, though **Unitary** is a dagger rig category, **Isometry** and **coIsometry** are mere rig categories. Intuitively, this must be the case because dagger categories are self-dual (*i.e.*, satisfy $\mathbf{C} \cong \mathbf{C}^{\text{op}}$) so limits and colimits coincide, but the affine completion only adds (certain) limits without the corresponding colimits. However, this also suggests that if we seek a notion of partiality that respects daggers, we would need the additive unit to be both initial and terminal, *i.e.*, a *zero object*. Fortunately, we can ensure this by applying L_{\oplus} after R_{\oplus} or vice versa.

Proposition 7. *The additive unit O is a zero object in both $L_{\oplus}(R_{\oplus}(\mathbf{C}))$ and $R_{\oplus}(L_{\oplus}(\mathbf{C}))$.*

Proof. By [10, Lemma 11], $R_{\oplus}(-)$ preserves terminal objects, and by duality, $L_{\oplus}(-)$ preserves initial objects. Thus O is both initial and terminal (*i.e.*, a zero object) in both $L_{\oplus}(R_{\oplus}(\mathbf{C}))$ and $R_{\oplus}(L_{\oplus}(\mathbf{C}))$. □

The situation is interesting: neither $L_{\oplus}(-)$ nor $R_{\oplus}(-)$ on their own preserve dagger rig categories, but as we will see, their combination does. Hence it is advantageous to consider them together for dagger rig categories, which also leads to a slightly simpler presentation. In a rig category, define $\sim_{LR_{\oplus}}$ as the least equivalence relation containing the three relations $\sim_{\text{id}_{\oplus}}$, $\sim_{L_{\oplus}}$, and $\sim_{R_{\oplus}}$ defined as follows, for all $f: A \oplus H \rightarrow B \oplus G$:

- $f \sim_{L_{\oplus}} (\text{id} \oplus m) \circ f$ for all $m: G \rightarrow G'$;
- $f \sim_{R_{\oplus}} f \circ (\text{id} \oplus n)$ for all $n: H' \rightarrow H$;
- $f \sim_{\text{id}_{\oplus}} \alpha_{\oplus} \circ (f \oplus \text{id}_X) \circ \alpha_{\oplus}^{-1}$ for all identities id_X .

Definition 8. Given a rig category \mathbf{C} , its *biaffine completion* $LR_{\oplus}(\mathbf{C})$ is the category whose

- objects are those of \mathbf{C} ,

- morphisms $[H, f, G]: A \rightarrow B$ are triples consisting of two objects H and G and a morphism $A \oplus H \rightarrow B \oplus G$ of \mathbf{C} quotiented by $\sim_{LR_{\oplus}}$,
- identities $A \rightarrow A$ are $[O, \text{id}_{A \oplus O}, O]$, and
- the composition of $f: A \oplus H \rightarrow B \oplus G$ and $g: B \oplus H' \rightarrow C \oplus G'$ is given by

$$A \oplus (H \oplus H') \xrightarrow{\alpha_{\oplus}^{-1}} (A \oplus H) \oplus H' \xrightarrow{f \oplus \text{id}} (B \oplus G) \oplus H' \xrightarrow{\cong} (B \oplus H') \oplus G \xrightarrow{g \oplus \text{id}} (C \oplus G') \oplus G \xrightarrow{\alpha_{\oplus}} C \oplus (G' \oplus G).$$

We state some properties of the LR_{\oplus} -construction, the proofs of which can be found in the appendix.

Proposition 9. *The additive unit O is a zero object in $LR_{\oplus}(\mathbf{C})$.*

Proposition 10. *When \mathbf{C} is a dagger rig category, so is $LR_{\oplus}(\mathbf{C})$.*

As before, there is a functor $\mathcal{F}: \mathbf{C} \rightarrow LR_{\oplus}(\mathbf{C})$ given on objects by $\mathcal{F}(A) = A$ and on morphisms by $F(f) = [0, f \oplus \text{id}_0, 0]$, making $LR_{\oplus}(\mathbf{C})$ a completion in the formal sense. We say that a rig category is *additively biaffine* if the unit of the sum is a zero object.

Theorem 11. *$LR_{\oplus}(\mathbf{C})$ is the additive biaffine completion of \mathbf{C} in the following sense: given any additively biaffine rig category \mathbf{D} and strong rig functor $F: \mathbf{C} \rightarrow \mathbf{D}$, there is a unique strong rig functor $\widehat{F}: LR_{\oplus}(\mathbf{C}) \rightarrow \mathbf{D}$ making the diagram below commute.*

$$\begin{array}{ccc} \mathbf{C} & \xrightarrow{\mathcal{F}} & LR_{\oplus}(\mathbf{C}) \\ & \searrow F & \downarrow \widehat{F} \\ & & \mathbf{D} \end{array}$$

Proof. Assuming such a functor exists, we first prove its uniqueness. Imposing $F = \widehat{F} \circ \mathcal{F}$ implies that $\widehat{F}(A) = F(A)$ on objects and, on morphisms in the image of \mathcal{F} , we have that $\widehat{F}(\mathcal{F}(f)) = F(f)$. It is easy to check (for instance, diagrammatically) that any $[H, f, G]: A \rightarrow B$ in $LR_{\oplus}[\mathbf{C}]$ decomposes as follows:

$$[H, f, G] = A \xrightarrow{\rho^{-1}} A \oplus O \xrightarrow{\text{id} \oplus !} A \oplus H \xrightarrow{\mathcal{F}(f)} B \oplus G \xrightarrow{\text{id} \oplus !} B \oplus O \xrightarrow{\rho} B$$

where the objects and morphisms shown are in $LR_{\oplus}[\mathbf{C}]$, where O is a zero object, and $!$ refer to the corresponding unique morphisms. The image under \widehat{F} of each of these morphisms is uniquely determined because \widehat{F} is assumed to be monoidal, the fact that \mathbf{D} has $0 \cong F(O)$ as its zero object, and the equality $\widehat{F}(\mathcal{F}(f)) = F(f)$ discussed above. Thus it only remains to prove that \widehat{F} as defined above is indeed a strong rig functor. By definition, $\widehat{F}(\text{id}) = F(\text{id})$, and functoriality is easy to check using naturality of ρ and the fact that 0 is a zero object. The fact that \widehat{F} is a strong rig functor follows directly from the same property for F , since \mathcal{F} is a strict rig functor, $\widehat{F}(O) = F(O) \cong 0$, and $\widehat{F}(A \oplus B) = F(A \oplus B) \cong F(A) \oplus F(B) = \widehat{F}(A) \oplus \widehat{F}(B)$. \square

Using this characterisation, we can show commutativity of the L_{\oplus} and R_{\oplus} constructions, the proof of which is found in the appendix.

Proposition 12. *When \mathbf{C} is a rig category, $L_{\oplus}(R_{\oplus}(\mathbf{C})) \cong LR_{\oplus}(\mathbf{C}) \cong R_{\oplus}(L_{\oplus}(\mathbf{C}))$.*

3.1 Unitaries with dagger partiality are contractions

A linear map $f: A \rightarrow B$ between normed spaces A and B is (weakly) *contractive* iff $\|f(x)\| \leq \|x\|$ for all $x \in A$. So contractive maps include all isometries, coisometries, and unitaries. We now show that applying the LR_{\oplus} -construction to the category of finite-dimensional Hilbert spaces and unitaries is equivalent to the category **Contraction** of finite-dimensional Hilbert spaces and contractions. Another way of saying this is that extending the unitaries with dagger partiality gives precisely the contractions.

Theorem 13. $LR_{\oplus}(\mathbf{Unitary}) \cong \mathbf{Contraction}$.

Proof. Proposition 12 gives $LR_{\oplus}(\mathbf{Unitary}) \cong L_{\oplus}(R_{\oplus}(\mathbf{Unitary}))$, and Proposition 6 gives $R_{\oplus}(\mathbf{Unitary}) \cong \mathbf{Isometry}$, so it suffices to show $L_{\oplus}(\mathbf{Isometry}) \cong \mathbf{Contraction}$. The strategy is to define a functor $F: \mathbf{Contraction} \rightarrow L_{\oplus}(\mathbf{Isometry})$ and prove it is full, faithful and essentially surjective. On objects, $F(A) = A$ so F is essentially surjective. Let $T: A \rightarrow B$ be a contraction and let $f: A \rightarrow B \oplus G$ be an isometry such that $T = \pi_B \circ f$: for example, by [7, 24], an isometric dilation $A \rightarrow B \oplus A$ can be constructed as

$$f = \begin{pmatrix} T \\ (1 - T^{\dagger}T)^{\frac{1}{2}} \end{pmatrix}.$$

Let $F(T) = [f, G]$. We need to check this mapping is well-defined, *i.e.*, if $f': A \rightarrow B \oplus G'$ also satisfies $T = \pi_B \circ f'$, verify that $[f, G] = [f', G']$. To do this, first consider the Hilbert space $\text{im}(f) \subseteq B \oplus G$ and similarly $\text{im}(f')$ and define a function $g: \text{im}(f') \rightarrow \text{im}(f)$ by $f'(a) \mapsto f(a)$. This function is well-defined because f and f' are isometries and hence injective. It is easy to see that g is linear; also notice that

$$\langle g(f'a) | g(f'a) \rangle = \langle fa | fa \rangle = \langle a | a \rangle = \langle f'a | f'a \rangle$$

because f and f' are isometries. This applies to all vectors in $\text{im}(f')$, so g is an isometry; in fact, because $\dim(\text{im}(f)) = \dim(\text{im}(f'))$ is finite, g has an inverse that is also an isometry, so $g: \text{im}(f') \rightarrow \text{im}(f)$ is unitary. Whenever $f(a) \in B$ it is necessary that $f(a) = f'(a)$ for them to be dilations of T , which means that g acts as the identity on B and thus it has the shape

$$g = \begin{pmatrix} \text{id} & 0 \\ 0 & h \end{pmatrix}.$$

Next we lift h to a map $k: G' \rightarrow G$. Assume without loss of generality that $\dim(G') \leq \dim(G)$ and pick an isometry $k: G' \rightarrow G$ satisfying $(\text{id}_B \oplus k) \circ f' = f$. Such a function exists: let $k(x) = h(x)$ whenever $x \in \text{im}(f') \cap G'$ and, for each element $x \in G'$ not in the image of f' , choose an element $y \in G$ not in $\text{im}(f)$ and map x to y . Since $\dim(G') \leq \dim(G)$ by assumption, the latter choices can be made so that k is an isometry. Then, it is immediate that $f \sim_{L_{\oplus}} f'$ since $k: G' \rightarrow G$ is an isometry acting as their mediator, and we conclude that $[f, G] = [f', G']$. Moreover, if $F(T) = [f, G]$ and $F(S) = [g, G']$ then it's easy to check that $S \circ T = \pi \circ (g \oplus \text{id}_G) \circ f$, so F is indeed a functor. If $F(V) = F(V')$ then there is an isometry f such that $V = \pi \circ f = V'$ and, hence, F is faithful. Now $\pi \circ f$ is a contraction for all $[f, G]$, so F is full. \square

4 Hiding in a partial setting

The previous section considered (co)affine completions with respect to the direct sum. But rig categories have another monoidal structure, the tensor product. Just as we can form the *additive* affine completion

$L_{\oplus}(\mathbf{C})$ of a rig category \mathbf{C} , we can also form the *multiplicative* one $L_{\otimes}(\mathbf{C})$: objects are those of \mathbf{C} , morphisms $A \rightarrow B$ are equivalence classes of morphisms $A \rightarrow B \otimes G$ in \mathbf{C} , with identities and composition just as in Definition 1. Where the L_{\oplus} -construction captures *partiality* by allowing part of the *state space* to be hidden, the L_{\otimes} -construction models *discarding* by allowing any *state* to be hidden (partly or fully).

This construction was shown to capture Stinespring dilation in [13], where the authors argued that $L_{\otimes}(\mathbf{Isometry})$ is monoidally equivalent to the category $\mathbf{FHilb}_{\text{CPTP}}$ of finite-dimensional Hilbert spaces and completely positive trace-preserving (CPTP) maps (or *quantum channels*). A drawback is that this does not generally preserve the direct sum, so $L_{\otimes}(\mathbf{C})$ is generally only monoidal (under tensor product) when \mathbf{C} is a rig category (though remnants of the direct sum persist, see [10, Section 4.4]).

Unfortunately, we cannot hope to use the L_{\otimes} -construction to construct the category $\mathbf{FHilb}_{\text{CPTN}}$ of finite-dimensional Hilbert spaces and completely positive trace-nonincreasing (CPTN) maps (or *partial quantum channels*) for a simple reason: the unit I of the tensor product is terminal in $L_{\otimes}(\mathbf{C})$, but unlike $\mathbf{FHilb}_{\text{CPTP}}$, it is not terminal in $\mathbf{FHilb}_{\text{CPTN}}$. The unique map $A \rightarrow I$ in $\mathbf{FHilb}_{\text{CPTP}}$ is given by the trace of a density matrix on A , and it is unique because all density matrices have unit trace. On the other hand, the *subnormalised* density matrices found in $\mathbf{FHilb}_{\text{CPTN}}$ take their traces in the unit interval, so even though there is only one *trace-preserving* map $A \rightarrow I$, there are as many trace-nonincreasing ones as there are real numbers in $[0, 1]$. Another way to say this is that the L_{\otimes} -construction fails to respect *partiality* in $\mathbf{FHilb}_{\text{CPTN}}$ (as also discussed in [9]).

This section generalises the result from [13] to the partial case. To do this, we show a variant of Stinespring dilation for completely positive trace-nonincreasing maps, describe a construction $L'_{\otimes}(\mathbf{C})$ that extends \mathbf{C} with unique *total* deletion maps, and relate the two by showing that $L'_{\otimes}(\mathbf{Contraction}) \simeq \mathbf{FHilb}_{\text{CPTN}}$. We relate this to the total case by showing that $L'_{\otimes}(\mathbf{C})$ has a universal property as a certain pushout in the category of (locally small) monoidal categories.

4.1 Stinespring dilation for partial quantum channels

We begin with a small lemma, which turns out to be incredibly useful when working with contractions.

Proposition 14. *T is contractive if and only if $T^{\dagger}T \leq 1$.*

Proof. By definition, $T^{\dagger}T \leq 1$ iff $1 - T^{\dagger}T$ is positive semidefinite, which in turn is the case iff $\langle \phi | (1 - T^{\dagger}T) | \phi \rangle \geq 0$ for all $|\phi\rangle$. But since

$$\langle \phi | (1 - T^{\dagger}T) | \phi \rangle = \langle \phi | \phi \rangle - \langle \phi | T^{\dagger}T | \phi \rangle = \|\phi\|^2 - \|T|\phi\rangle\|^2,$$

$1 - T^{\dagger}T$ is positive semidefinite iff $\|\phi\|^2 - \|T|\phi\rangle\|^2 \geq 0$ for all $|\phi\rangle$, i.e., when T is contractive. \square

The previous proposition links handily to the following theorem about the Kraus representation of completely positive trace-nonincreasing maps.

Proposition 15 ([1]). *Any CPTN map Φ admits a representation $\Phi(\rho) = \sum_{i=1}^k M_i \rho M_i^{\dagger}$ with $\sum_i M_i^{\dagger} M_i \leq 1$.*

As in the trace-preserving case, we can construct a Stinespring dilation from the Kraus representation.

Proposition 16. *Every CPTN map Φ admits a Stinespring dilation $\Phi(\rho) = \text{tr}_E(T\rho T^{\dagger})$ for some contraction T and Hilbert space E .*

Proof. Let $\mathcal{B}(A) \xrightarrow{\Phi} \mathcal{B}(B)$ be a partial quantum channel with Kraus representation $\Phi(\rho) = \sum_{i=1}^k M_i \rho M_i^\dagger$. Define $E = \mathbb{C}^k$, let $A \otimes \mathbb{C} \xrightarrow{U_R} A$ denote the right unitor, and let $T = (\sum_{i=1}^k M_i \otimes |i\rangle) U_R^\dagger$. Now for any ρ :

$$\begin{aligned} \text{tr}_E(T\rho T^\dagger) &= \text{tr}_E \left(\left(\sum_{i=1}^k M_i \otimes |i\rangle \right) U_R^\dagger \rho U_R \left(\sum_{j=1}^k M_j^\dagger \otimes \langle j| \right) \right) = \text{tr}_E \left(\sum_{i=1}^k \sum_{j=1}^k M_i \rho M_j^\dagger \otimes |i\rangle \langle j| \right) \\ &= \sum_{i=1}^k \sum_{j=1}^k \text{tr}(|i\rangle \langle j|) M_i \rho M_j^\dagger = \sum_{i=1}^k M_i \rho M_i^\dagger = \Phi(\rho) . \end{aligned}$$

It remains to show that T is contractive:

$$\begin{aligned} T^\dagger T &= U_R \left(\sum_{j=1}^k M_j^\dagger \otimes \langle j| \right) \left(\sum_{i=1}^k M_i \otimes |i\rangle \right) U_R^\dagger = U_R \left(\sum_{j=1}^k \sum_{i=1}^k M_j^\dagger M_i \otimes \langle j|i\rangle \right) U_R^\dagger \\ &= U_R \left(\sum_{i=1}^k M_i^\dagger M_i \otimes \langle i|i\rangle \right) U_R^\dagger = \sum_{i=1}^k M_i^\dagger M_i \leq 1 \end{aligned}$$

which finishes the proof. \square

This representation is essentially unique, *i.e.*, unique up to an isometry applied to the ancilla.

Lemma 17 ([20, Theorem 8.2]). *Let Φ be a CP map with two Kraus representations $\Phi(\rho) = \sum_{i=1}^k E_i \rho E_i^\dagger$ and $\Phi(\rho) = \sum_{j=1}^{k'} F_j \rho F_j^\dagger$. Assume $k = k'$; if $k < k'$, add some $E_i = 0$ for all $k < i \leq k'$. There is a unitary k -by- k matrix $U = (u_{ij})$ such that $E_i = \sum_{j=1}^k u_{ij} F_j$ for all i .*

Proposition 18. *Let Φ be a CPTN map, and let $T_E: A \rightarrow B \otimes G$, $T_F: A \rightarrow B \otimes G'$ be two contractions such that $\Phi(\rho) = \text{tr}_G(T_E \rho T_E^\dagger) = \text{tr}_{G'}(T_F \rho T_F^\dagger)$. If $\dim(G) \leq \dim(G')$, then there is an isometry $W: G \rightarrow G'$ such that $T_F = (1 \otimes W)T_E$.*

Proof. Write U_R for the right unitor. Fix an orthonormal basis for G and define $E_i = U_R(1 \otimes \langle i|)T_E$; do the same for $F_j = U_R(1 \otimes \langle j|)T_F$. Notice that $T_E = (\sum_i E_i \otimes |i\rangle)U_R^\dagger$ thanks to $\sum_i |i\rangle \langle i| = 1_G$ for an orthonormal basis. Take V to be an injection of G into a Hilbert space with the same dimension as G' . Let $U = (u_{ij})$ be the unitary from the lemma above. Then:

$$T_E = \left(\sum_i E_i \otimes |i\rangle \right) U_R^\dagger = \left(\sum_i \left(\sum_j u_{ij} F_j \right) \otimes |i\rangle \right) U_R^\dagger = \left(\sum_i \left(\sum_j u_{ij} F_j \otimes |i\rangle \right) \right) U_R^\dagger = \left(\sum_j F_j \otimes \left(\sum_i u_{ij} |i\rangle \right) \right) U_R^\dagger.$$

Because U is unitary, $|\hat{j}\rangle = \sum_i u_{ij} |i\rangle$ form an orthonormal basis for G' . Finally, we obtain the isometry $W: G \rightarrow G'$ by composing $W = RUV$ where R is the unitary mapping $|\hat{j}\rangle \mapsto |j\rangle$. \square

Moreover, as in the trace-preserving case, contractions give rise to CPTN maps through conjugation.

Proposition 19. *Every contraction T gives rise to a CPTN map $\Phi(\rho) = T\rho T^\dagger$.*

Proof. Conjugation by any linear map is completely positive. That Φ is trace-nonincreasing follows by $\text{tr}(T\rho T^\dagger) = \text{tr}(\rho T^\dagger T) \leq \text{tr}(\rho 1) = \text{tr}(\rho)$. \square

Taken together, these results show that the situation between **Contraction** and **FHilb**_{CPTN} mirrors that between **Isometry** and **FHilb**_{CPTP}: there is a (strict monoidal) functor **Contraction** \rightarrow **FHilb**_{CPTN} that sends a contraction T to conjugation by T , and every CPTN map can be expressed this way (in an essentially unique way) if we allow ourselves an ancilla system.

4.2 Total eclipse of the state

We now formulate the new L_{\otimes}^t -construction, adding a notion of hiding that cooperates with the *total* maps in a monoidal dagger category. Consider a monoidal dagger category \mathbf{C} . We can think of its dagger monomorphisms (*i.e.*, morphisms f satisfying $f^\dagger \circ f = \text{id}$) as its *total maps*. These form a subcategory of \mathbf{C} , which we denote $\text{DagMon}(\mathbf{C})$. The significance of the total maps is that, instead of being able to mediate with *arbitrary* maps from \mathbf{C} (as in the L_{\otimes} -construction), we are only permitted to mediate with *total* maps (*i.e.*, morphisms in $\text{DagMon}(\mathbf{C})$) in the L_{\otimes}^t -construction. Explicitly, for $f: A \rightarrow B \otimes G$ and $f': A \rightarrow B \otimes G'$, write $f \leq_{L_{\otimes}^t} f'$ iff there exists a dagger monomorphism $m: G \rightarrow G'$ making

$$\begin{array}{ccc} & A & \\ f \swarrow & & \searrow f' \\ B \otimes G & \xrightarrow{\text{id} \otimes m} & B \otimes G' \end{array}$$

commute. Let $\sim_{L_{\otimes}^t}$ denote the equivalence closure of $\leq_{L_{\otimes}^t}$.

Definition 20. The *partial multiplicative affine completion* $L_{\otimes}^t(\mathbf{C})$ of a symmetric monoidal dagger category \mathbf{C} is the category whose

- objects are those of \mathbf{C} ,
- morphisms $[f, G]: A \rightarrow B$ are pairs of an object G and an equivalence class of morphisms $f: A \rightarrow B \otimes G$ of \mathbf{C} under $\sim_{L_{\otimes}^t}$,
- identities are $[\rho_{\otimes}^{-1}, I]$, and
- composition of $[f, G]: A \rightarrow B$ and $[g, G']: B \rightarrow C$ is $[\alpha_{\otimes} \circ g \otimes \text{id}_G \circ f, G' \otimes G]$.

Before proceeding with the universal property of this construction, we first establish that it succeeds in constructing $\mathbf{FHilb}_{\text{CPTN}}$.

Theorem 21. *There is an equivalence $L_{\otimes}^t(\mathbf{Contraction}) \simeq \mathbf{FHilb}_{\text{CPTN}}$ of monoidal categories.*

Proof. Construct a functor $L_{\otimes}^t(\mathbf{Contraction}) \rightarrow \mathbf{FHilb}_{\text{CPTN}}$ acting as the identity on objects, by sending $[f, G]: A \rightarrow B$ to the map $\rho \mapsto \text{tr}_G(f^\dagger \rho f)$, which is CPTN by Proposition 19 and since the partial trace is trace-preserving. This is well-defined since dagger monomorphisms in $\mathbf{Contraction}$ are precisely the isometries, and since Stinespring dilations are invariant under isometric manipulation of the ancilla system G . This functor is essentially surjective since it is identity on objects and $L_{\otimes}^t(\mathbf{Contraction})$ and $\mathbf{FHilb}_{\text{CPTN}}$ have the same objects; it is full since every CPTN map admits a Stinespring dilation (by Proposition 16); it is faithful since different Stinespring dilations of the same CPTN map are always connected by an isometry on the ancilla by Proposition 18; and it is (strict) monoidal since it preserves coherence isomorphisms. \square

An immediate consequence of the definition of $L_{\otimes}^t(\mathbf{C})$ is that each object comes equipped with a discarding map and chosen projections.

Proposition 22. *Every object A of $L_{\otimes}^t(\mathbf{C})$ has a discarding map $A \rightarrow I$, giving canonical projections $\pi_1: A \otimes B \rightarrow A$ and $\pi_2: A \otimes B \rightarrow B$.*

Proof. As in L_{\oplus} , construct the discarding map $A \rightarrow I$ as the equivalence class of the pair $[\lambda_{\otimes}^{-1}, A]$ where $\lambda_{\otimes}^{-1}: A \rightarrow I \otimes A$ is the inverse left unitor of \mathbf{C} . Projections are given by the equivalence class of $[\text{id}_{A \otimes B}, B]: A \otimes B \rightarrow A$ and that of $[\sigma_{\otimes}, A]: A \otimes B \rightarrow B$. \square

We state some basic properties of this construction, shown analogously to those for the L_{\oplus} -construction.

Proposition 23. *When \mathbf{C} is a symmetric monoidal dagger category, $L_{\otimes}^t(\mathbf{C})$ is symmetric monoidal.*

Proposition 24. *There is a strict monoidal functor $\mathcal{E}_t: \mathbf{C} \rightarrow L_{\otimes}^t(\mathbf{C})$ given by $\mathcal{E}_t(A) = A$ on objects and $\mathcal{E}_t(f) = [\rho_{\otimes}^{-1} \circ f, I]$ on morphisms.*

The connection between the L_{\otimes} and the L_{\otimes}^t construction is made clear by the following inclusion.

Proposition 25. *There is a strict monoidal inclusion functor $I_t: L_{\otimes}(\text{DagMon}(\mathbf{C})) \rightarrow L_{\otimes}^t(\mathbf{C})$.*

Indeed, as we will see momentarily, the L_{\otimes}^t -construction is characterised by this inclusion. Interestingly, this construction can be seen as a particular instance of the *monoidal indeterminates*-construction [8]. This gives a very useful factorisation lemma, which may be regarded as an instance of *purification*.

Lemma 26. *Let \mathbf{C} be a symmetric monoidal dagger category. Then*

- (i) *morphisms $[f, E]$ of $L_{\otimes}(\text{DagMon}(\mathbf{C}))$ factor uniquely as $\pi_1 \circ \mathcal{E}(f')$ for some f' in $\text{DagMon}(\mathbf{C})$,*
- (ii) *morphisms $[f, E]$ of $L_{\otimes}^t(\mathbf{C})$ factor uniquely as $\pi_1 \circ \mathcal{E}_t(f')$ for some f' in \mathbf{C} , and*
- (iii) *morphisms $[f, E]$ of $L_{\otimes}^t(\mathbf{C})$ factor uniquely as $I_t(\pi_1) \circ \mathcal{E}(f')$ for some f' in \mathbf{C} .*

Proof. (i,ii) are the expansion-raw factorisation of [8], and (iii) holds as $\pi_1 = I_t(\pi_1)$ in $L_{\otimes}^t(\mathbf{C})$. \square

We are now ready to establish the universal property of $L_{\otimes}^t(\mathbf{C})$.

Theorem 27. *$L_{\otimes}^t(\mathbf{C})$ is a pushout of $\mathcal{E}: \text{DagMon}(\mathbf{C}) \rightarrow L_{\otimes}(\text{DagMon}(\mathbf{C}))$ along the inclusion functor $\text{DagMon}(\mathbf{C}) \rightarrow \mathbf{C}$ in the category of locally small symmetric monoidal categories and strong monoidal functors.*

$$\begin{array}{ccc}
 \text{DagMon}(\mathbf{C}) & \xrightarrow{I} & \mathbf{C} \\
 \mathcal{E} \downarrow & & \downarrow \mathcal{E}_t \\
 L_{\otimes}(\text{DagMon}(\mathbf{C})) & \xrightarrow{I_t} & L_{\otimes}^t(\mathbf{C}) \\
 & \searrow F & \downarrow F_t \\
 & & \mathbf{D}
 \end{array}$$

(Note: A dashed arrow \hat{F} also points from $L_{\otimes}(\text{DagMon}(\mathbf{C}))$ to \mathbf{D} , and a solid arrow F points from $L_{\otimes}(\text{DagMon}(\mathbf{C}))$ to \mathbf{D} .)

Proof sketch. Notice first that the upper square commutes since all functors involved are strict monoidal, and I and I_t are merely inclusions behaving as the identity on objects and morphisms, while \mathcal{E} is precisely \mathcal{E}_t restricted to dagger monomorphisms of \mathbf{C} .

Next, since objects on \mathbf{C} , $\text{DagMon}(\mathbf{C})$, $L_{\otimes}^t(\mathbf{C})$, and $L_{\otimes}(\text{DagMon}(\mathbf{C}))$ all coincide, F and F_t must agree on objects, so we may define $\hat{F}(X) = F(X) = F_t(X)$ on objects, and $\hat{F} \circ I_t = F$ and $\hat{F} \circ \mathcal{E}_t = F_t$ on objects follows immediately. On morphisms we define $\hat{F}([f, E]) = F(\pi_1) \circ F_t(f)$. That this definition satisfies $\hat{F} \circ I_t = F$ and $\hat{F} \circ \mathcal{E}_t = F_t$ on morphisms as well, and that \hat{F} is unique, follows using the factorisation lemma above (full proof in the appendix). \square

Instantiating this property to the case of **Contraction**, where the functors **Contraction** \rightarrow **Isometry** and **FHilb**_{CPTP} \rightarrow **FHilb**_{CPTN} are inclusions, and the functors **Isometry** \rightarrow **FHilb**_{CPTP} and **Contraction** \rightarrow **FHilb**_{CPTN} conjugating by an isometry or contraction, we obtain the following characterisation.

Corollary 28. *The following square is a pushout of symmetric monoidal categories:*

$$\begin{array}{ccc}
 \mathbf{Isometry} & \longrightarrow & \mathbf{Contraction} \\
 \downarrow & & \downarrow \\
 \mathbf{FHilb}_{\text{CPTP}} & \longrightarrow & \mathbf{FHilb}_{\text{CPTN}}
 \end{array}$$

Proof. That $\text{DagMon}(\mathbf{Contraction}) \simeq \mathbf{Isometry}$ follows by the fact that every isometry f is contractive (it satisfies $\|f(x)\| = \|x\|$, so specifically $\|f(x)\| \leq \|x\|$) and satisfies $f^\dagger \circ f = \text{id}$. Finally, $L_\otimes(\mathbf{Isometry}) \simeq \mathbf{FHilb}_{\text{CPTN}}$ is known [13], and $L_\otimes(\mathbf{Contraction}) \simeq \mathbf{FHilb}_{\text{CPTN}}$ by Theorem 21. \square

5 Splitting measurements

The category $\mathbf{FHilb}_{\text{CPTN}}$ does not have coproducts, but it can nevertheless be instructive to see why a given candidate for cotupling fails. Working in $L_\otimes^t(\mathbf{Contraction})$, given two Stinespring dilations $f: A \rightarrow C \otimes G$ and $g: B \rightarrow C \otimes G'$, a candidate for the Stinespring dilation of their cotupling is the map $\delta_L^{-1} \circ f \oplus g: A \oplus B \rightarrow C \otimes (G \oplus G')$. In particular, if we were to try to cotuple the trivial Stinespring dilations of the canonical injections $i_1: A \rightarrow (A \oplus B) \otimes I$ and $i_2: B \rightarrow (A \oplus B) \otimes I$ in $\mathbf{Contraction}$, the result would be the map $\delta_L^{-1} \circ i_1 \oplus i_2: A \oplus B \rightarrow (A \oplus B) \otimes (I \oplus I)$. Notice the non-trivial nature of the ancilla of this dilation, which tracks whether the result came from i_1 or i_2 . This is not simply the identity, as it should be if this were an actual cotupling. Computing, we see that this map acts on block diagonal density matrices on $A \oplus B$ by measuring whether the result falls in A or in B , *i.e.*, by the mapping

$$\left(\begin{array}{c|c} X & Y \\ \hline Z & W \end{array} \right) \mapsto \left(\begin{array}{c|c} X & 0 \\ \hline 0 & W \end{array} \right).$$

Clearly, maps such as these are idempotent, but interestingly they do not split in $\mathbf{FHilb}_{\text{CPTN}}$. However, they *do* have a very natural splitting in the category $\mathbf{FCstar}_{\text{CPTN}}$ of finite-dimensional C^* -algebras and CPTN maps, via $m: \mathcal{B}(A \oplus B) \rightarrow \mathcal{B}(A) \oplus \mathcal{B}(B)$ and $p: \mathcal{B}(A) \oplus \mathcal{B}(B) \rightarrow \mathcal{B}(A \oplus B)$ given by

$$m \left(\begin{array}{c|c} X & Y \\ \hline Z & W \end{array} \right) = (X, W) \quad \text{and} \quad p(X, W) = \left(\begin{array}{c|c} X & 0 \\ \hline 0 & W \end{array} \right) \quad (1)$$

as we then have $m \circ p = \text{id}$ and $p \circ m$ is precisely the idempotent we started with. Indeed, this is a defining characteristic of finite-dimensional C^* -algebras compared to Hilbert spaces.

Definition 29. When \mathbf{C} is an additively coaffine rig category, define a *measurement* in $L_\otimes^t(\mathbf{C})$ to be the equivalence class of any n -ary ‘‘attempted cotupling’’ of injections

$$(\delta_L^{-1})^{n-1} \circ \bigoplus_{1 \leq j \leq n} i_j: \left(\bigoplus_{1 \leq j \leq n} A_j \right) \rightarrow \left(\bigoplus_{1 \leq j \leq n} A_j \right) \otimes (I^{\oplus n}).$$

Note that identities are measurements where $n = 1$. In $\mathbf{FHilb}_{\text{CPTN}}$, these correspond to projections onto arbitrary block diagonals, the images of which are precisely the C^* -algebras. We spell out Karoubi’s concrete characterisation of the idempotent splitting of measurements in \mathbf{C} .

Definition 30. The idempotent splitting of measurements in $L_\otimes^t(\mathbf{C})$ (where \mathbf{C} is an additively coaffine rig category) is the category $\mathbf{Split}_M(L_\otimes^t(\mathbf{C}))$ whose

- objects are pairs (A, e) of an object A of \mathbf{C} and a measurement $e: A \rightarrow A$,
- morphisms $f: (A, e) \rightarrow (B, e')$ are morphisms $f: A \rightarrow B$ of \mathbf{C} satisfying $e' \circ f \circ e = f$,
- each identity $(A, e) \rightarrow (A, e)$ is e , and
- composition is as in $L_\otimes^t(\mathbf{C})$.

There is an inclusion of $L_{\otimes}^t(\mathbf{C})$ in $\mathbf{Split}_M(L_{\otimes}^t(\mathbf{C}))$ sending objects A to (A, id) and leaving morphisms unchanged. This construction is well-known to be the free splitting of these idempotents; more abstractly, it is the completion of a category with regards to certain absolute colimits [22]. Now for the main result of this section, remembering that $\mathbf{FHilb}_{\text{CPTN}} \simeq L_{\otimes}^t(\mathbf{Contraction})$ (by Theorem 21) and that $\mathbf{Contraction}$ is an additively coaffine dagger rig category (by Proposition 10 and Theorems 11 and 13).

Theorem 31. *There is an equivalence $\mathbf{Split}_M(\mathbf{FHilb}_{\text{CPTN}}) \simeq \mathbf{FCstar}_{\text{CPTN}}$ of categories.*

Proof sketch. Define a functor $F: \mathbf{Split}_M(\mathbf{FHilb}_{\text{CPTN}}) \rightarrow \mathbf{FCstar}_{\text{CPTN}}$ as follows:

- on objects, set $F((A, e)) = \text{im}(e)$, the image of the idempotent e as a subspace of A ;
- on morphisms, $F(f: (A, e) \rightarrow (B, e')) = m_{e'} \circ f \circ p_e$.

Here, p_e is the inclusion of the subspace $\text{im}(e)$ in A , and $m_{e'}$ is the projection of B onto its subspace $\text{im}(e')$, as in (1). This is straightforwardly seen to be functorial. That this functor is essentially surjective follows by Artin-Wedderburn, and that it is full and faithful follows using the unique splitting of measurement idempotents in $\mathbf{FCstar}_{\text{CPTN}}$ (see appendix for full proof). \square

We noted about the L_{\otimes}^t -construction that it does not preserve direct sums, so even when acting on a rig category, it is only a completion of monoidal categories. However, when splitting its measurements using the construction above, the failed cotupling in $\mathbf{FHilb}_{\text{CPTN}}$ does become an actual cotupling of coproducts in $\mathbf{Split}_M(\mathbf{FHilb}_{\text{CPTN}})$. Likewise, though $L_{\otimes}^t(\mathbf{C})$ does contain enough remnants of the direct sum of a rig category \mathbf{C} to make it an *arrow with choice* [10] and so allows for a description of measurements even in the abstract, it is admittedly awkward. For this reason, though L_{\otimes}^t and \mathbf{Split}_M are described by universal properties separately, they are much better considered in combination, just like the LR_{\oplus} -construction.

6 Discussion

We have presented three universal constructions which, taken together, construct the category of finite-dimensional C^* -algebras and completely positive trace-nonincreasing maps from the category of finite-dimensional Hilbert spaces and unitaries. Though we have kept to the finite-dimensional case in this paper, there is reason to suspect that many of these results will generalise to infinite dimensions. For example, *Halmos dilation* has a far stronger statement as *Sz. Nagy dilation* in the infinite-dimensional case, and the usual Stinespring dilation theorem generalises to the infinite-dimensional case as well.

The key application that we envision for this work is in the design and semantics of quantum programming languages. One application of the LR_{\oplus} -construction is in the quantisation of reversible classical programs. It can be shown that applying the LR_{\oplus} -construction to the category \mathbf{FinBij} of finite sets and bijections yields the category $\mathbf{FinPInj}$ of finite sets and partial injective functions. In particular, this means that the quantisation functor $\mathbf{FinBij} \rightarrow \mathbf{Unitary}$ lifts uniquely to a functor $\mathbf{FinPInj} \rightarrow \mathbf{Contraction}$. This is interesting since $\mathbf{FinPInj}$ is the setting for (finite) reversible classical computing (see e.g. [17, 5, 6]), in particular *reversible (classical) flowcharts* [31, 4] (with a finite state space). This suggests that $\mathbf{Contraction}$ may be similarly considered as a setting for *reversible quantum flowcharts*, a kind of quantum flowcharts (see also [25]) which eschew measurements in favour of quantum control.

Another application concerns extending the quantum programming language $\mathcal{U}\Pi_a^{\chi}$ (“yuppie-chi-a”) [10] with *classical types*. $\mathcal{U}\Pi_a^{\chi}$ is an effectful extension of $\mathcal{U}\Pi$, a quantum extension to the strongly typed classical reversible programming language Π [16]. The effectful part of $\mathcal{U}\Pi_a^{\chi}$ is that it uses the L_{\otimes} and R_{\oplus} constructions in a very direct way to add measurement as an effect to an otherwise measurement-free language, taking its semantics from $\mathbf{Unitary}$ to $\mathbf{FHilb}_{\text{CPTN}}$. Since the measurement idempotents

described in Section 5 can all be described as $\mathcal{U}\Pi_a^\lambda$ programs, the \mathbf{Split}_M -construction can similarly be used very directly to add classical types to this language as a computational effect. This addresses a known shortcoming of the language by allowing a distinction between *e.g.* the type of *bits* and the type of *qubits* (as in *e.g.* [25]), but also allows for a much more fine-grained type-level separation of measurement maps in *how much* they measure.

References


- [1] V. Cappellini, H.-J. Sommers & K. Życzkowski (2007): *Subnormalized states and trace-nonincreasing maps*. *Journal of Mathematical Physics* 48(5), p. 052110.
- [2] K. Cho (2015): *Total and Partial Computation in Categorical Quantum Foundations*. In C. Heunen, P. Selinger & J. Vicary, editors: *Proceedings 12th International Workshop on Quantum Physics and Logic (QPL 2015)*, EPTCS 195, pp. 116–135.
- [3] C. Comfort (2021): *The ZX& calculus: A complete graphical calculus for classical circuits using spiders*. In B. Valiron, S. Mansfield, P. Arrighi & P. Panangaden, editors: *Proceedings 17th International Conference on Quantum Physics and Logic (QPL 2020)*, *Electronic Proceedings in Theoretical Computer Science* 340, OPA, pp. 60–90.
- [4] R. Glück & R. Kaarsgaard (2018): *A categorical foundation for structured reversible flowchart languages: Soundness and adequacy*. *Logical Methods in Computer Science* 14(3:16), pp. 1–38.
- [5] R. Glück, R. Kaarsgaard & T. Yokoyama (2019): *Reversible programs have reversible semantics*. In: *International Symposium on Formal Methods: FM 2019 International Workshops*, Springer, pp. 413–427.
- [6] R. Glück, R. Kaarsgaard & T. Yokoyama (2022): *From Reversible Programming Languages to Reversible Metalanguages*. *Theoretical Computer Science* 920, pp. 46–63.
- [7] P. R. Halmos (1950): *Normal Dilations and Extensions of Operators*. *Summa Brasiliensis Mathematica* 2, pp. 125–134.
- [8] C. Hermida & R. D. Tennent (2012): *Monoidal indeterminates and categories of possible worlds*. *Theoretical Computer Science* 430, pp. 3–22.
- [9] C. Heunen & R. Kaarsgaard (2021): *Bennett and Stinespring, Together at Last*. In C. Heunen & M. Backens, editors: *Proceedings 18th International Conference on Quantum Physics and Logic (QPL 2021)*, *Electronic Proceedings in Theoretical Computer Science* 343, OPA, pp. 102–118.
- [10] C. Heunen & R. Kaarsgaard (2022): *Quantum Information Effects*. *Proceedings of the ACM on Programming Languages* 6(POPL).
- [11] C. Heunen, A. Kissinger & P. Selinger (2014): *Completely positive projections and biproducts*. In: *Proceedings of the 10th International Workshop on Quantum Physics and Logic (QPL 2013)*, Open Publishing Association, pp. 71–83.
- [12] C. Heunen & J. Vicary (2019): *Categories for Quantum Theory*. Oxford University Press.
- [13] M. Huot & S. Staton (2018): *Universal properties in quantum theory*. In P. Selinger & G. Chiribella, editors: *Proceedings of the 15th International Conference on Quantum Physics and Logic (QPL 2018)*, *Electronic Proceedings in Theoretical Computer Science* 287, Open Publishing Association, pp. 213–224.
- [14] M. Huot & S. Staton (2019): *Quantum channels as a categorical completion*. In: *34th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS 2019)*, IEEE, pp. 1–13.
- [15] G. Hutton (1999): *A tutorial on the universality and expressiveness of fold*. *Journal of Functional Programming* 9(4), pp. 355–372.
- [16] R. P. James & A. Sabry (2012): *Information Effects*. In: *Proceedings of the 39th Annual ACM SIGPLAN-SIGACT Symposium on Principles of programming languages (POPL '12)*, ACM, pp. 73–84.

- [17] R. Kaarsgaard, H. B. Axelsen & R. Glück (2017): *Join inverse categories and reversible recursion*. *Journal of Logical and Algebraic Methods in Programming* 87, pp. 33–50.
- [18] M. Laplaza (1972): *Coherence for Distributivity*. *Lecture Notes in Mathematics* 281, pp. 29–72.
- [19] E. Levy & O. Shalit (2014): *Dilation theory in finite dimensions: The possible, the impossible, and the unknown*. *Rocky Mountain Journal of Mathematics* 44(1), pp. 203–221.
- [20] M. A. Nielsen & I. L. Chuang (2010): *Quantum Computation and Quantum Information: 10th Anniversary Edition*. Cambridge University Press, doi:10.1017/CB09780511976667.
- [21] A. J. Parzygnat (2019): *Stinespring’s construction as an adjunction*. *Compositionality* 1(2), pp. 1–42.
- [22] R. Paré (1971): *On absolute colimits*. *Journal of Algebra* 19(1), pp. 80–95.
- [23] M. Rennela, S. Staton & R. Furber (2017): *Infinite-dimensionality in quantum foundations: W^* -algebras as presheaves over matrix algebras*. In: *13th International Conference on Quantum Physics and Logic (QPL 2017)*, Open Publishing Association, pp. 161–173.
- [24] P. L. Robinson (2018): *Julia operators and Halmos dilations*. *arXiv preprint arXiv:1803.09329*.
- [25] P. Selinger (2004): *Towards a Quantum Programming Language*. *Mathematical Structures in Computer Science* 14(4), pp. 527–586.
- [26] S. Staton & P. B. Levy (2013): *Universal properties of impure programming languages*. In: *Proceedings of the 40th Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages (POPL ’13)*, ACM, pp. 179–192.
- [27] D. Stein & S. Staton (2021): *Compositional semantics for probabilistic programs with exact conditioning*. In: *36th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS 2021)*, IEEE, pp. 1–13.
- [28] W. F. Stinespring (1955): *Positive functions on C^* -algebras*. *Proceedings of the American Mathematical Society* 6(2), pp. 211–216.
- [29] P. Wadler (1989): *Theorems for free!* In: *Proceedings of the fourth international conference on Functional programming languages and computer architecture (FPCA ’89)*, ACM, pp. 347–359.
- [30] A. Westerbaan & B. Westerbaan (2016): *Paschke Dilations*. In R. Duncan & C. Heunen, editors: *Proceedings of the 13th International Conference on Quantum Physics and Logic (QPL 2016)*, *Electronic Proceedings in Theoretical Computer Science* 236, Open Publishing Association, pp. 229–244, doi:10.4204/EPTCS.236.15.
- [31] T. Yokoyama, H. B. Axelsen & R. Glück (2016): *Fundamentals of reversible flowchart languages*. *Theoretical Computer Science* 611, pp. 87–115.

A Deferred proofs

Proposition 9. The additive unit O is a zero object in $LR_{\oplus}(\mathbf{C})$.

Proof. Define maps $A \rightarrow O$ and $O \rightarrow A$ as the equivalence classes of the symmetry $\sigma_{\oplus}: A \oplus O \rightarrow O \oplus A$ and $\sigma_{\oplus}: O \oplus A \rightarrow A \oplus O$. To see that the map $A \rightarrow O$ is unique, let $f: A \oplus H \rightarrow O \oplus G$ be any other morphism of \mathbf{C} . Then $[O, \sigma, A] \sim_{LR_{\oplus}} [H, f, G]$ can be seen using the graphical language [12]:



That the map $O \rightarrow A$ is unique follows analogously, so O is both initial and terminal, *i.e.*, a zero object. \square

Proposition 10. When \mathbf{C} is a dagger rig category, so is $LR_{\oplus}(\mathbf{C})$.

Proof. For any morphism $[H, f, G]: A \rightarrow B$ in $LR_{\oplus}(\mathbf{C})$, its dagger is defined to be

$$[H, f, G]^{\dagger} = [G, f^{\dagger}, H].$$

First, we need to check this is well-defined, *i.e.*, given $[H, f, G] = [H', f', G']$, verify $[H, f, G]^{\dagger} = [H', f', G']^{\dagger}$. This is straightforward, as $f \sim_{LR_{\oplus}} g$ implies $f^{\dagger} \sim_{LR_{\oplus}} g^{\dagger}$; to see this, notice that

$$\begin{aligned} f \sim_{L_{\oplus}} g &\implies f^{\dagger} \sim_{R_{\oplus}} g^{\dagger} \\ f \sim_{R_{\oplus}} g &\implies f^{\dagger} \sim_{L_{\oplus}} g^{\dagger} \end{aligned}$$


by using the dagger of the mediators and, furthermore, $f \sim_{id_{\oplus}} g$ trivially implies $f^{\dagger} \sim_{id_{\oplus}} g^{\dagger}$. Clearly, this dagger is involutive and

$$id_A^{\dagger} = [O, id_{A \oplus O}, O]^{\dagger} = [O, id_{A \oplus O}^{\dagger}, O] = id_A.$$

Checking explicitly that $(g \circ f)^{\dagger} = f^{\dagger} \circ g^{\dagger}$ is more involved, but conceptually trivial: flip the diagram and use the fact that \mathbf{C} is a dagger category. This works because, in $LR_{\oplus}(\mathbf{C})$ there is symmetry between input and output: the dagger turns hidden parts of the input into a hidden part of the output, and vice versa. That coherence isomorphisms are unitary follows immediately by the fact that they are inherited from \mathbf{C} , where they are unitary by \mathbf{C} a dagger rig category. \square

Proposition 12. When \mathbf{C} is a rig category, $L_{\oplus}(R_{\oplus}(\mathbf{C})) \cong LR_{\oplus}(\mathbf{C}) \cong R_{\oplus}(L_{\oplus}(\mathbf{C}))$.

Proof. The fact that O is initial in $LR_{\oplus}(\mathbf{C})$ together with the universal property of R_{\oplus} implies that the top triangle of the diagram below commutes, for a unique functor Φ . Then, the fact that O is also terminal in $LR_{\oplus}(\mathbf{C})$ and the universal property of L_{\oplus} implies that the bottom triangle also commutes, for a unique functor $\widehat{\mathcal{F}}$.



Moreover, O is a zero object in $L_{\oplus}(R_{\oplus}(\mathbf{C}))$ and, thus, the universal property of LR_{\oplus} implies there is a unique strong monoidal functor $\widehat{\mathcal{E}} \circ \widehat{\mathcal{D}}: LR_{\oplus}(\mathbf{C}) \rightarrow L_{\oplus}(R_{\oplus}(\mathbf{C}))$ such that $\mathcal{E} \circ \mathcal{D} = \widehat{\mathcal{E}} \circ \widehat{\mathcal{D}} \circ \mathcal{F}$. This, together with the commuting diagram above implies:

$$\widehat{\mathcal{E}} \circ \widehat{\mathcal{D}} \circ \mathcal{F} \circ \mathcal{E} \circ \mathcal{D} = \widehat{\mathcal{E}} \circ \widehat{\mathcal{D}} \circ \Phi \circ \mathcal{D} = \widehat{\mathcal{E}} \circ \widehat{\mathcal{D}} \circ \mathcal{F} = \mathcal{E} \circ \mathcal{D}$$

which is captured in the outer triangle of the diagram below

$$\begin{array}{ccccc} \mathbf{C} & \xrightarrow{\mathcal{D}} & R_{\oplus}(\mathbf{C}) & \xrightarrow{\mathcal{E}} & L_{\oplus}(R_{\oplus}(\mathbf{C})) \\ & \searrow & \downarrow & \downarrow & \downarrow \\ & & & & L_{\oplus}(R_{\oplus}(\mathbf{C})) \end{array}$$

$\mathcal{E} \circ \mathcal{D}$ (diagonal solid line from \mathbf{C} to $L_{\oplus}(R_{\oplus}(\mathbf{C}))$)
 $\widehat{\mathcal{E}} \circ \widehat{\mathcal{D}} \circ \mathcal{F}$ (dashed line from $R_{\oplus}(\mathbf{C})$ to $L_{\oplus}(R_{\oplus}(\mathbf{C}))$)

Due to the universal property of R_{\oplus} , the functor shown above as a diagonal dashed line is unique, so it must be \mathcal{E} ; then, $\widehat{\mathcal{E}} \circ \widehat{\mathcal{D}} \circ \mathcal{F} = 1_{L_{\oplus}(R_{\oplus}(\mathbf{C}))}$ as both functors make the inner right triangle commute but, by the universal property of L_{\oplus} , there is only one such functor. By a similar argument $\widehat{\mathcal{F}} \circ \widehat{\mathcal{E}} \circ \widehat{\mathcal{D}} = 1_{LR_{\oplus}(\mathbf{C})}$ and hence $L_{\oplus}(R_{\oplus}(\mathbf{C})) \cong LR_{\oplus}(\mathbf{C})$. This is an equivalence of rig categories, as $\widehat{\mathcal{F}}$ and $\widehat{\mathcal{E}} \circ \widehat{\mathcal{D}}$ are rig functors by construction. The same strategy proves $R_{\oplus}(L_{\oplus}(\mathbf{C})) \cong LR_{\oplus}(\mathbf{C})$. \square

Theorem 27. $L'_{\otimes}(\mathbf{C})$ is a pushout of $\mathcal{E}: \text{DagMon}(\mathbf{C}) \rightarrow L_{\otimes}(\text{DagMon}(\mathbf{C}))$ along the inclusion functor $\text{DagMon}(\mathbf{C}) \hookrightarrow \mathbf{C}$ in the category of locally small symmetric monoidal categories and strong monoidal functors.

$$\begin{array}{ccc} \text{DagMon}(\mathbf{C}) & \xrightarrow{I} & \mathbf{C} \\ \mathcal{E} \downarrow & & \downarrow \mathcal{E}_t \\ L_{\otimes}(\text{DagMon}(\mathbf{C})) & \xrightarrow{I_t} & L'_{\otimes}(\mathbf{C}) \end{array}$$

F (solid line from $L_{\otimes}(\text{DagMon}(\mathbf{C}))$ to \mathbf{D})
 F_t (solid line from \mathbf{C} to \mathbf{D})
 \hat{F} (dotted line from $L'_{\otimes}(\mathbf{C})$ to \mathbf{D})

Proof. Notice first that the upper square commutes since all functors involved are strict monoidal, and I and I_t are merely inclusions behaving as the identity on objects and morphisms, while \mathcal{E} is precisely \mathcal{E}_t restricted to dagger monomorphisms of \mathbf{C} .

Next, since objects on \mathbf{C} , $\text{DagMon}(\mathbf{C})$, $L'_{\otimes}(\mathbf{C})$, and $L_{\otimes}(\text{DagMon}(\mathbf{C}))$ all coincide, F and F_t must agree on objects, so we may define $\hat{F}(X) = F(X) = F_t(X)$ on objects, and $\hat{F} \circ I_t = F$ and $\hat{F} \circ \mathcal{E}_t = F_t$ on objects follows immediately. On morphisms we define $\hat{F}([f, E]) = F(\pi_1) \circ F_t(f)$. Then

$$\begin{aligned} \hat{F}(I_t([f, E])) &= \hat{F}(I_t(\pi_1 \circ \mathcal{E}(f))) = \hat{F}(I_t(\pi_1) \circ I_t(\mathcal{E}(f))) = \hat{F}(I_t(\pi_1) \circ \mathcal{E}_t(I(f))) \\ &= \hat{F}([I(f), E]) = F(\pi_1) \circ F_t(I(f)) = F(\pi_1) \circ F(\mathcal{E}(f)) \\ &= F(\pi_1 \circ \mathcal{E}(f)) = F([f, E]) \end{aligned}$$

so $\hat{F} \circ I_t = F$ on morphisms as well. For the other triangle,

$$\begin{aligned} \hat{F}(\mathcal{E}_t(f)) &= \hat{F}(\pi_1 \circ \rho^{-1} \circ \mathcal{E}_t(f)) = \hat{F}(\pi_1 \circ \mathcal{E}_t(\rho^{-1}) \circ \mathcal{E}_t(f)) = \hat{F}(\pi_1 \circ \mathcal{E}_t(\rho^{-1} \circ f)) \\ &= \hat{F}([\rho^{-1} \circ f, I]) = F(\pi_1) \circ F_t(\rho^{-1} \circ f) = F(\pi_1) \circ F(\rho^{-1}) \circ F_t(f) \\ &= F(\pi_1) \circ F_t(I(\rho^{-1})) \circ F_t(f) = F(\pi_1) \circ F(\mathcal{E}(\rho^{-1})) \circ F_t(f) \\ &= F(\pi_1 \circ \mathcal{E}(\rho^{-1})) \circ F_t(f) = F([\rho^{-1}, I]) \circ F_t(f) = F(\text{id}) \circ F_t(f) = F_t(f) \end{aligned}$$

establishing $\hat{F} \circ \mathcal{E}_t = F_t$ on morphisms. Finally, suppose G satisfies $G \circ I_t = F$ and $G \circ \mathcal{E}_t = F_t$. Then

$$G([f, E]) = G(I_t(\pi_1) \circ \mathcal{E}_t(f)) = G(I_t(\pi_1)) \circ G(\mathcal{E}_t(f)) = F(\pi_1) \circ F_t(f) = \hat{F}([f, E])$$

on morphisms, and since G and \hat{F} must agree on objects as well (e.g. $\hat{F}(X) = F(X) = G(I_t(X)) = G(X)$), we get $G = \hat{F}$. Finally, \hat{F} is strong monoidal since \mathcal{E}_t is strict monoidal and F_t and \hat{F} agree on objects, so \hat{F} may reuse the coercions $I \cong F_t(I)$ and $F_t(A \otimes B) \cong F_t(A) \otimes F_t(B)$. \square

Theorem 31. There is an equivalence $\mathbf{Split}_M(\mathbf{FHilb}_{\text{CPTN}}) \simeq \mathbf{FCstar}_{\text{CPTN}}$ of categories.

Proof. Define a functor $F: \mathbf{Split}_M(\mathbf{FHilb}_{\text{CPTN}}) \rightarrow \mathbf{FCstar}_{\text{CPTN}}$ as follows:

- on objects, set $F((A, e)) = \text{im}(e)$, the image of the idempotent e as a subspace of A ;
- on morphisms, $F(f: (A, e) \rightarrow (B, e')) = m_{e'} \circ f \circ p_e$.

Here, p_e is the inclusion of the subspace $\text{im}(e)$ in A , and $m_{e'}$ is the projection of B onto its subspace $\text{im}(e')$, as in (1). This preserves identities since

$$F(\text{id}_{(H, e)}) = F(e) = m_e \circ e \circ p_e = m_e \circ p_e \circ m_e \circ p_e = \text{id} \circ \text{id} = \text{id}.$$

Likewise, if $f: (A, e) \rightarrow (B, e')$ and $g: (B, e') \rightarrow (C, e'')$, then

$$F(g \circ f) = m_{e''} \circ g \circ f \circ p_e = m_{e''} \circ g \circ e' \circ f \circ p_e = m_{e''} \circ g \circ p_{e'} \circ m_{e'} \circ f \circ p_e = F(g) \circ F(f)$$

using the fact that e' splits as $e' = p_{e'} \circ m_{e'}$. Thus F is functorial. That

Any finite-dimensional C^* -algebra is, up to isomorphism, of the form $\bigoplus \mathcal{B}(H_i)$, so the image of a block diagonal projection on $\mathcal{B}(\bigoplus H_i)$. Thus F is essentially surjective.

Given any CPTN map of C^* -algebras $f: \bigoplus \mathcal{B}(A_i) \rightarrow \bigoplus \mathcal{B}(B_j)$, let $m_A: \bigoplus \mathcal{B}(A_i) \rightarrow \mathcal{B}(\bigoplus A_i)$ and $m_B: \bigoplus \mathcal{B}(B_j) \rightarrow \mathcal{B}(\bigoplus B_j)$ denote the projections onto this subspace, and p_A and p_B their inclusions in the other direction. We see that $p_B \circ f \circ m_A: (\bigoplus A_i, p_A \circ m_A) \rightarrow (\bigoplus B_j, p_B \circ m_B)$ is a morphism of $\mathbf{Split}_M(\mathbf{FHilb}_{\text{CPTN}})$ since $m_A \circ p_A = \text{id}$ and $m_B \circ p_B = \text{id}$, and so

$$(p_B \circ m_B) \circ (p_B \circ f \circ m_A) \circ (p_A \circ m_A) = p_B \circ (m_B \circ p_B) \circ f \circ (m_A \circ p_A) \circ m_A = p_B \circ f \circ m_A.$$

Further,

$$F(p_B \circ f \circ m_A) = m_B \circ (p_B \circ f \circ m_A) \circ p_A = (m_B \circ p_B) \circ f \circ (m_A \circ p_A) = \text{id} \circ f \circ \text{id} = f$$

so F is full.

Finally, let $f: (A, e) \rightarrow (B, e')$ and $f': (A, e) \rightarrow (B, e')$ be morphisms of $\mathbf{Split}_M(\mathbf{FHilb}_{\text{CPTN}})$ such that $F(f) = F(f')$, i.e., $m_{e'} \circ f \circ p_e = m_{e'} \circ f' \circ p_e$. Since $f = e' \circ f \circ e$ and $f = e' \circ f' \circ e$ by definition of morphisms in $\mathbf{Split}_M(-)$, we have

$$f = e' \circ f \circ e = p_{e'} \circ m_{e'} \circ f \circ p_e \circ m_e = p_{e'} \circ m_{e'} \circ f' \circ p_e \circ m_e = e' \circ f' \circ e = f'$$

so F is faithful, and so an equivalence of categories. \square