



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Spatiotemporal profiling of luteinising hormone/human choriogonadotropin receptor in the human fetal testis

### Citation for published version:

Macdonald, J, Smith, LB, Mitchell, RT & Anderson, R 2016, 'Spatiotemporal profiling of luteinising hormone/human choriogonadotropin receptor in the human fetal testis', *The Lancet*, vol. 387, pp. S74. [https://doi.org/10.1016/S0140-6736\(16\)00461-X](https://doi.org/10.1016/S0140-6736(16)00461-X)

### Digital Object Identifier (DOI):

[10.1016/S0140-6736\(16\)00461-X](https://doi.org/10.1016/S0140-6736(16)00461-X)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Publisher's PDF, also known as Version of record

### Published In:

The Lancet

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Spatiotemporal profiling of luteinising hormone/human choriogonadotropin receptor in the human fetal testis

Joni Macdonald\*, Lee B Smith\*, Richard A Anderson, Rod T Mitchell

## Abstract

Published Online  
February 25, 2016

Poster 22

\*Contributed equally

University of Edinburgh,  
Edinburgh, UK  
(J Macdonald PhD,  
Prof L B Smith PhD,  
Prof R A Anderson PhD,  
R T Mitchell PhD)

Correspondence to:  
Dr Rod T Mitchell, MRC Centre  
for Reproductive Health,  
University of Edinburgh,  
Edinburgh EH16 4TJ, UK  
rod.mitchell@ed.ac.uk

**Background** Androgen signalling is essential for masculinisation of the male fetus. Failure of masculinisation during fetal life can result in disorders of sex development. Masculinisation is mediated partly by ligand binding to the luteinising hormone/human choriogonadotropin receptor (LHCGR). In rodents, the initial stages of masculinisation are gonadotropin independent, though whether this also occurs in human beings remains unknown because accurate timing and cell-type localisation of LHCGR protein expression in human fetal testis remains unresolved. We aimed to characterise the spatiotemporal expression of LHCGR in human fetal testis.

**Methods** Human fetal testis tissue (n=12, 8–20 weeks' gestation) was obtained from elective terminations after informed patient consent. We used a combination of RT-PCR for LHCGR and Leydig cells (marker 3 $\beta$  hydroxysteroid dehydrogenase [3 $\beta$ HSD]) and immunofluorescent co-localisation for LHCGR, Leydig cell steroidogenesis (3 $\beta$ HSD), and endothelial cells (markers CD31 and von Willebrand factor) to determine the exact timing and cellular location of LHCGR mRNA and protein in the human fetal testis. The study received ethics approval.

**Findings** LHCGR mRNA was detected from 9 weeks' gestation, but protein was not detectable until 12 weeks'. Expression of 3 $\beta$ HSD (indicating active steroidogenesis) was detected in the first trimester (10 weeks') preceding the expression of LHCGR. LHCGR did not co-localise with the Leydig cell marker 3 $\beta$ HSD during the early second trimester and only became consistently co-localised at 20 weeks' gestation. Before expression in fetal Leydig cells, LHCGR was restricted to endothelial cells (demonstrated by co-localisation with CD31 and von Willebrand factor).

**Interpretation** These results indicate that gonadotropin-mediated signalling through the LHCGR in human fetal Leydig cells is not established until later than previously described, suggesting a period of gonadotropin insensitivity during masculinisation in fetal life. Future functional studies will determine the mechanisms involved in androgen signalling in the human fetal testis. The present findings are important for understanding the pathogenesis of disorders of sex development in man.

**Funding** Wellcome Trust.

### Contributors

JM and LBS performed experiments and analysed the data. RAA analysed the data. RTM designed the study, analysed the data, and wrote the abstract.

### Declaration of interests

We declare no competing interests.