



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Radiologist opinions regarding reporting incidental coronary and cardiac calcification on thoracic CT

Citation for published version:

Williams, MC, Weir-McCall, J, Moss, AJ, Schmitt, M, Stirrup, J, Holloway, B, Gopalan, D, Deshpande, A, Hughes, GM, Agrawal, B, Nicol, E, Roditi, G, Shambrook, J & Bull, R 2022, 'Radiologist opinions regarding reporting incidental coronary and cardiac calcification on thoracic CT', *BJR|Open*, vol. 4, no. 1.
<https://doi.org/10.1259/bjro.20210057>

Digital Object Identifier (DOI):

[10.1259/bjro.20210057](https://doi.org/10.1259/bjro.20210057)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Published In:

BJR|Open

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Received:
17 September 2021

Revised:
07 January 2022

Accepted:
12 February 2022

Published online:
11 March 2022

Cite this article as:

Williams MC, Weir-McCall J, Moss AJ, Schmitt M, Stirrup J, Holloway B, et al. Radiologist opinions regarding reporting incidental coronary and cardiac calcification on thoracic CT. *BJR Open* (2022) 10.1259/bjro.20210057.

ORIGINAL RESEARCH

Radiologist opinions regarding reporting incidental coronary and cardiac calcification on thoracic CT

¹MICHELLE C WILLIAMS, ²JONATHAN WEIR-MCCALL, ³ALASTAIR J MOSS, ⁴MATTHIAS SCHMITT, ⁵JAMES STIRRUP, ⁶BEN HOLLOWAY, ⁷DEEPA GOPALAN, ⁸APARNA DESHPANDE, ⁹GARETH MORGAN HUGHES, ¹⁰BOBBY AGRAWAL, ¹¹EDWARD NICOL, ¹²GILES RODITI, ¹³JAMES SHAMBROOK and ¹⁴RUSSELL BULL

¹BHF Centre for Cardiovascular Science and Edinburgh Imaging, University of Edinburgh, Edinburgh, UK

²University of Cambridge School of Clinical Medicine, Cambridge, UK

³British Heart Foundation Cardiovascular Research Centre, University of Leicester, Leicester, UK

⁴North West Heart Centre, Manchester University NHS Foundation Trust, Manchester, UK

⁵Royal Berkshire NHS Foundation Trust, Reading, UK

⁶Queen Elizabeth Hospital Birmingham, Birmingham, UK

⁷Imperial College London, London, UK

⁸Glenfield Hospital, University Hospitals of Leicester, Leicester, UK

⁹Plymouth Hospitals NHS Trust, Plymouth, UK

¹⁰Royal Papworth Hospital, Cambridge, UK

¹¹Royal Brompton and Harefield NHS Foundation Trust Departments of Cardiology and Radiology, UK; National Heart and Lung Institute, Faculty of Medicine, Imperial College, London, London, UK

¹²Dept. of Radiology, Glasgow Royal Infirmary, NHS Greater Glasgow & Clyde, Glasgow, UK; Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK

¹³Southampton General Hospital, Southampton, UK

¹⁴Royal Bournemouth Hospital, Bournemouth, UK

Address correspondence to: Dr Michelle C Williams
E-mail: michelle.williams@ed.ac.uk

Objectives: Coronary and cardiac calcification are frequent incidental findings on non-gated thoracic computed tomography (CT). However, radiologist opinions and practices regarding the reporting of incidental calcification are poorly understood.

Methods: UK radiologists were invited to complete this online survey, organised by the British Society of Cardiovascular Imaging (BSCI). Questions included anonymous information on subspecialty, level of training and reporting practices for incidental coronary artery, aortic valve, mitral and thoracic aorta calcification.

Results: The survey was completed by 200 respondents: 10% trainees and 90% consultants. Calcification was not reported by 11% for the coronary arteries, 22% for the aortic valve, 35% for the mitral valve and 37% for the thoracic aorta. Those who did not subspecialise in cardiac imaging were less likely to report coronary

artery calcification ($p = 0.005$), aortic valve calcification ($p = 0.001$) or mitral valve calcification ($p = 0.008$), but there was no difference in the reporting of thoracic aorta calcification. Those who did not subspecialise in cardiac imaging were also less likely to provide management recommendations for coronary artery calcification ($p < 0.001$) or recommend echocardiography for aortic valve calcification ($p < 0.001$), but there was no difference for mitral valve or thoracic aorta recommendations.

Conclusion: Incidental coronary artery, valvular and aorta calcification are frequently not reported on thoracic CT and there are differences in reporting practices based on subspecialty.


Advances in knowledge: On routine thoracic CT, 11% of radiologists do not report coronary artery calcification. Radiologist reporting practices vary depending on subspecialty but not level of training.

INTRODUCTION

Calcification in the coronary arteries, cardiac valves and thoracic aorta are frequent incidental findings (Figure 1) on routine non-gated thoracic computed tomography (CT)^{1–11} and may indicate important underlying pathologies. However, little is known about the reporting practices of radiologists regarding these incidental findings.

Cardiac and vascular calcification share overlapping risk factors and pathogenesis. Coronary artery calcification is a marker of atherosclerosis and its presence and extent are associated with increased cardiovascular and all-cause mortality.^{12–21} Similarly, aortic valve calcification is a marker of aortic sclerosis or stenosis and its severity is associated with the degree of aortic

Figure 1. Examples of aortic valve calcification (blue arrow), mitral valve calcification (yellow arrow), coronary artery (green arrow) and thoracic aorta calcification (orange arrow).



valve dysfunction.^{22–24} The clinical significance of mitral calcification is less certain, as although mitral valve calcification is associated with mitral valve dysfunction mitral annular calcification is a frequent finding that is rarely associated with mitral valve dysfunction. Calcification in the thoracic aorta is also a frequent finding in asymptomatic patients, and its severity may be associated with the severity of coronary artery disease and cardiovascular mortality.^{11,25} Although the reporting of coronary artery calcification on non-gated thoracic CT is supported by national and international guidelines, it is frequently not reported.^{1–3,26}

This survey aims to investigate the current practices and opinions of UK radiologists regarding the reporting of cardiac and vascular calcification on routine non-gated thoracic CT.

METHODS

Survey design

This survey was organised by the British Society of Cardiovascular Imaging/British Society of Cardiac Computed Tomography (BSCI/BSCCT). The executive committee of the BSCI/BSCCT acted as the steering committee for the survey. Survey questions were designed and refined by the BSCI/BSCCT executive committee. An electronic platform (Survey Monkey) was used to collect anonymous survey data. When more than one option was provided as an answer to a question, these were provided in a random order. Invitations to participate were sent to all BSCI/BSCCT members and they were asked to forward the survey to all radiologists working in their hospitals.

Survey questions

Survey questions included information about the respondent's subspecialty and level of training. For each of coronary artery calcification, aortic valve calcification, mitral calcification (valve and/or annulus), and thoracic aorta calcification they were asked whether they routinely report its presence, what factors affected whether they would report its presence and if management recommendations were provided. For coronary artery calcification, they were asked what method they use to report it (visual assessment on a per patient level, visual assessment on a per vessel level, semi-quantitative score, or Agatston score). For aortic valve and mitral calcification, they were asked whether

they would recommend echocardiography. Respondents were free to choose not to answer certain questions. Additional comments were solicited at the end of the survey.

Statistical analysis

Statistical analysis was performed using R v. 4.0.1 (R Foundation for Statistical Computing, Vienna, Austria). Survey responses which only included information on subspecialty and job description questions were deemed incomplete and excluded from the analysis. Number and percentage are presented for categorical data with statistical significance assessed using a Pearson's chi-square test. Binomial logistic regression analysis was performed to assess the impact of level of training and subspecialisation on reporting practices. A statistically significant difference was defined as a two-sided p -value < 0.05 .

RESULTS

Demographic information

After excluding incomplete survey responses ($n = 12$), there were 200 survey responses that were included in this analysis. This included 20 (10%) trainees and 180 (90%) consultants. Of the consultants, 25% ($n = 45$) were less than 5-year post-training, 23% ($n = 41$) were 5 to 10 years post-training, and 52% ($n = 94$) were more than 10 years post-training. Subspecialty practice/training was reported by 50% ($n = 100$) for cardiac imaging, 57% ($n = 114$) for thoracic imaging, and 16% ($n = 33$) for vascular imaging. 30% ($n = 59$) of respondents did not subspecialise in cardiac, thoracic, or vascular imaging.

Coronary artery calcification

Coronary artery calcification was reported for all cases by 23% of radiologists, for most cases by 38% of radiologists, and some cases by 29% of radiologists (Table 1). Coronary artery calcification was not reported by 11% of all survey respondents (Figure 2), and those who were not subspecialised in cardiac imaging were less likely to report coronary artery calcification on routine thoracic CT (17% of non-cardiac specialists did not report coronary artery calcification, $n = 17/100$ vs 4% of cardiac specialists did not report coronary artery calcification, $n = 4/100$; $p = 0.005$). Trainees were more likely to report all, most or some coronary artery calcification (95%, $n = 19/20$), compared to consultants with less than 5 years of experience (93%, $n = 42/45$), consultants with 5 to 10 years of experience (93%, $n =$

Table 1. Frequency of reporting of cardiac and vascular calcification

	Coronary artery calcification	Aortic valve calcification	Mitral valve calcification	Thoracic aorta calcification
Number	200	198	196	195
Yes - for all cases	46 (23%)	47 (24%)	33 (17%)	12 (6%)
Yes - for most cases (>50%)	75 (38%)	45 (23%)	-	24 (12%)
Yes - for some cases (<50%)	58 (30%)	63 (32%)	93 (47%)	85 (43%)
No	21 (11%)	43 (22%)	70 (35%)	74 (37%)

38/41) and consultants with more than 10 years of experience (85%, $n = 80/94$, Table 2), but these differences did not reach statistical significance ($p = 0.293$).

Visual assessment on a per patient basis was the most frequent method used to report coronary artery calcification (Figure 3). Age and indication for imaging were the most frequent factors to influence whether coronary artery calcification was reported, influencing 55 and 42% of respondents, respectively (Figure 4).

Management recommendations for coronary artery calcification findings were provided for all cases by 7% ($n = 14$), for most cases 8% ($n = 15$) and for some cases by 22% ($n = 45$). Management recommendations were not provided by 63% ($n = 126$) and were less likely to be provided by those who did not subspecialise in cardiac imaging (79%, $p < 0.001$). Free text comments highlighted concerns regarding

the potential clinical benefit of reporting these findings, or the potential to increase unnecessary further investigations.

Aortic valve calcification

Aortic valve calcification was reported for all cases by 24% of radiologists, for most cases by 23% of radiologists, and some cases by 1% of radiologists (Table 1). Aortic valve calcification was not reported by 22% of all survey respondents, and those who did not subspecialise in cardiac imaging were less likely to report aortic valve calcification (31% of non-cardiac specialists did not report coronary artery calcification, $n = 31/100$ vs 12% of cardiac specialists did not report coronary artery calcification, $n = 12/100$; $p = 0.001$). Level of training did not impact reporting of aortic valve calcification (Table 2, $p = 0.293$). Age and previous cardiac intervention were the most frequent factors influencing reporting of aortic valve calcification, influencing 35 and

Figure 2. Reporting of coronary artery calcification by (A) all survey respondents and (B) those who did not subspecialise in cardiac imaging.




Table 2. Impact of level of training on reporting of cardiac and vascular calcification

	Trainee	Consultant		
		<5 years	5–10 years	>10 years
Coronary artery calcification ^a	19 (95%)	42 (93%)	39 (93%)	80 (85%)
Aortic valve calcification ^a	15 (75%)	34 (76%)	34 (83%)	72 (77%)
Mitral valve calcification ^a	12 (60%)	29 (64%)	23 (56%)	62 (66%)
Thoracic aorta calcification ^a	12 (65%)	28 (62%)	29 (71%)	51 (54%)

^aCombined “Yes - for all cases”, “Yes - for most cases” and “Yes - for some cases”

30% of respondents, respectively. Echocardiogram was recommended following the identification of aortic valve calcification by 4% (*n* = 8) for all cases, 12% (*n* = 24) for most cases and 20% (*n* = 40) for some cases. Recommendations for echocardiography were not provided by 63% (*n* = 126) and were less likely to be provided by those who did not subspecialise in cardiac imaging (77% of non-cardiac specialists did not provide recommendations for echocardiography, *n* = 77/100

vs 49% of cardiac specialists did not provide recommendations for echocardiography, *n* = 49/100; *p* < 0.001)

Mitral calcification

Mitral calcification (valve and/or annulus) was reported for all cases by 17% of radiologists and for some cases by 47% of radiologists (Table 1). Mitral calcification was not reported by 35% of all survey

Figure 3. Reporting methods used by respondents who report coronary artery calcification.





Figure 4. Factors that influence reporting of coronary artery calcification (CAC), aortic valve calcification (AVC), mitral valve calcification (MVC) and thoracic aorta calcification (TAC).



respondents, and those who did not subspecialise in cardiac imaging were less likely to report mitral calcification (41% of non-cardiac specialists did not report mitral calcification, $n = 41/100$ vs 29% of cardiac specialists did not report mitral calcification, $n = 29/100$; $p = 0.0996$), but this difference did not reach statistical significance. Level of training did not impact reporting of mitral calcification ($p = 0.599$, Table 2). Age and previous cardiac intervention were the most frequent factors influencing reporting of mitral calcification, influencing 25 and 21% of respondents, respectively. Echocardiogram was recommended following the identification of mitral calcification by 2% ($n = 4$) for all cases, 3% ($n = 5$) for most cases and 14% ($n = 28$) for some cases. Recommendations for echocardiography were not provided by 80% ($n = 159$) and were less likely to be provided by those who did not subspecialise in cardiac imaging, but this difference did not reach statistical significance (84% of non-cardiac specialists did not provide recommendations for echocardiography, $n = 84/100$ vs 75% of cardiac specialists did not provide recommendations for echocardiography, $n = 77/100$; $p = 0.092$).

Thoracic aorta

Thoracic aortic calcification was reported for all cases by 6% of radiologists, for most cases by 12% of radiologists, and for some cases by 43% of radiologists (Table 1). Thoracic aorta calcification was not reported by 37% of all survey respondents, this was similar for those who did not subspecialise in cardiac imaging and those who did (41% of non-cardiac specialists did not report thoracic aorta calcification, $n = 41/100$ vs 33% of cardiac specialists did not report thoracic aorta calcification, $n = 33/100$; $p = 0.420$). Level of training did not impact reporting of thoracic

aortic calcification ($p = 0.491$, Table 2). Age, previous cardiac intervention, and presence of comorbidities were the most frequent factors influencing reporting of thoracic aortic calcification, influencing 34%, 37 and 23% of respondents respectively (Figure 3).

Free text comments

Free text comments were provided by 33 (17%) respondents. Of these, 7 (21%) were regarding the importance of the topic of this survey. Two of the comments (6%) were concerned about the potential for reporting incidental calcification to cause patient anxiety. While 5 (15%) comments were concerned that reporting incidental calcification would cause unnecessary investigations and increased workload, 2 (6%) thought that reporting incidental calcification would be ignored by referrers. One (3%) comment was concerned about potential medicolegal implications. Seven (21%) of the comments suggested that clear guidelines would be useful for their clinical practice, including reporting templates and clear pathways for referrers. The remaining comments provided further detail on their current reporting practices.

DISCUSSION

This survey found that incidental calcification is frequently not reported on routine thoracic CT and that subspecialisation, but not level of training, influenced reporting practices. Subspecialisation in cardiac imaging influenced reporting of incidental calcification in the coronary arteries, aortic valve, and mitral valve, but not in the thoracic aorta. The primary factors which influenced reporting

of calcification were age and indication for imaging. Management recommendations were provided infrequently and were less likely to be provided by those who did not subspecialise in cardiac imaging.

Extensive research has shown that coronary artery calcification is an effective marker of the presence of coronary artery disease and is associated with both cardiovascular and all-cause mortality in patients undergoing dedicated cardiac imaging or routine thoracic imaging.^{4,12–21,27–31} Nevertheless, our survey showed that coronary artery calcification was not reported by 11% of survey respondents. At present, there remains debate in the radiology community regarding the reporting of cardiac calcification on routine thoracic CT. In our survey comments included concerns around patient anxiety, unnecessary investigations and increased workload for radiology and cardiology departments. Other concerns that have been raised include cost-effectiveness, ethical and medicolegal implications, additional reporting time and unknown clinical utility. Randomised controlled trials of management changes based on incidental calcification have not yet been completed. The Risk Or Benefit IN Screening for Cardiovascular diseases (ROBINSCA) randomised controlled trial is currently assessing the clinical impact of screening asymptomatic individuals with coronary artery calcium scoring compared to either screening based on traditional cardiovascular risk factors or no screening.³² Information on cardiovascular outcomes in this trial are awaited, but early publications show that basing medication choices on calcium scoring reduced the number of patients receiving preventative medication compared to screening using traditional cardiovascular risk factors.³² The results of this and other ongoing research will shape future guidelines in this area.

The recently published BSCI/BSCCT consensus statement on reporting incidental coronary, aortic valve and cardiac calcification on non-gated thoracic CT aims to address some of the concerns raised in this survey.³³ For coronary artery calcification, it suggests a simple patient-based score for coronary artery calcification (none, mild, moderate, severe) and highlights that many patients with incidental calcification do not require further investigation.³³ Repeating this survey after the publication of this consensus statement will be important to assess future changes in the opinions of radiologists throughout the UK. The Society of Cardiovascular Computed tomography have also published international guidelines which recommend that coronary artery calcification should be reported on all non-gated thoracic CT.³⁴ There are currently no reporting guidelines that address the issue of thoracic aortic calcification on

non-gated thoracic CT, highlighting on-going uncertainty in this area.

We found that on thoracic CT radiologists were less likely to report non-coronary cardiovascular calcification compared to coronary calcification. The degree of aortic valve calcification on electrocardiogram-gated CT is associated with the severity of aortic valve disease and reporting this is part of international guidelines for the management of valvular heart disease.³⁵ However, for mitral calcification and thoracic aortic calcification their impact is more debated. Although mitral valve leaflet calcification will indicate valvular dysfunction, mitral annular calcification is more frequently an asymptomatic incidental finding (although in some severe cases is associated with significant valvular dysfunction). The severity of thoracic aortic calcification correlates with coronary artery calcium scores, but it is not independent of coronary artery calcification in the assessment of cardiac events.^{36–38} Further research in these areas will help guide reporting of these incidental findings.

Limitations of this survey include the number of participants, the UK only participants and the unequal proportion of trainees and consultants. Those who completed this survey are more likely to be interested in this topic (selection bias), and therefore the true frequency of reporting is likely to be lower than that presented by this survey. As this was a survey of reporting practices rather than directly assessing CT reports there is the potential for recall bias to impact our results. We did not ask participants whether they reported findings differently on non-contrast compared to contrast enhanced scans and we did not discriminate between mitral valve and mitral annular calcification. We also did not enquire whether respondents worked in district general, tertiary care or academic departments, which may also influence reporting practices. Participants were also free not to answer all of the questions in the survey, meaning some of the data is incomplete. The survey was extensively reviewed by members of the BSCI/BSCCT executive committee, which includes consultant and trainee radiologists and cardiologists, but as this was not an externally validated survey the way that the questions were asked could have influenced the responses. Nevertheless, this survey provides an interesting insight into the practices of radiologists regarding the reporting of incidental calcification.

In conclusion, this survey found that coronary, valvular and aortic calcification is frequently not reported by UK radiologists. Subspecialisation but not level of training impacted reporting practices.

REFERENCES

- Williams KA, Kim JT, Holohan KM. Frequency of unrecognized, unreported, or underreported coronary artery and cardiovascular calcification on noncardiac chest ct. *J Cardiovasc Comput Tomogr* 2013; 7: S1934–5925(13)00109-3: 167–72. . <https://doi.org/10.1016/j.jcct.2013.05.003>
- Uretsky S, Chokshi N, Kobriniski T, Agarwal SK, Po JR, et al. The interplay of physician awareness and reporting of incidentally found coronary artery calcium on the clinical management of patients who underwent noncontrast chest computed tomography. *Am J Cardiol* 2015; 115: S0002-9149(15)00895-4: 1513–17. . <https://doi.org/10.1016/j.amjcard.2015.02.051>
- Balakrishnan R, Nguyen B, Raad R, Donnino R, Naidich DP, et al. Coronary artery calcification is common on nongated chest computed tomography imaging. *Clin Cardiol* 2017; 40: 498–502. <https://doi.org/10.1002/clc.22685>
- Phillips WJ, Johnson C, Law A, Turek M, Small AR, et al. Comparison of framingham risk score and chest-ct identified coronary artery calcification in breast cancer patients to predict cardiovascular events. *Int J Cardiol* 2019; 289: S0167-5273(18)35694-8: 138–43. . <https://doi.org/10.1016/j.ijcard.2019.01.056>

5. Rodriguez-Granillo GA, Reynoso E, Capunay C, Garcia-Garcia HM, Carrascosa P. Impact on mortality of coronary and non-coronary cardiovascular findings in non-gated thoracic ct by malignancy status. *Eur J Radiol* 2017; **93**: S0720-048X(17)30209-7: 169–77. . <https://doi.org/10.1016/j.ejrad.2017.05.030>
6. van de Wiel JCM, Wang Y, Xu DM, van der Zaag-Loonen HJ, van der Jagt EJ, et al. Neglectable benefit of searching for incidental findings in the dutch-belgian lung cancer screening trial (nelson) using low-dose multidetector ct. *Eur Radiol* 2007; **17**: 1474–82. <https://doi.org/10.1007/s00330-006-0532-7>
7. Raju P, Sallomi D, George B, Patel H, Patel N, et al. Aortic valve calcification - a commonly observed but frequently ignored finding during ct scanning of the chest. *Int J Clin Pract* 2012; **66**: 552–55. <https://doi.org/10.1111/j.1742-1241.2012.02916.x>
8. Lippert JA, White CS, Mason AC, Plotnick GD. Calcification of aortic valve detected incidentally on ct scans: prevalence and clinical significance. *AJR Am J Roentgenol* 1995; **164**: 73–77. <https://doi.org/10.2214/ajr.164.1.7998572>
9. Koos R, Kühl HP, Mühlenbruch G, Wildberger JE, Günther RW, et al. Prevalence and clinical importance of aortic valve calcification detected incidentally on ct scans: comparison with echocardiography. *Radiology* 2006; **241**: 76–82. <https://doi.org/10.1148/radiol.2411051163>
10. Mahnken AH, Mühlenbruch G, Das M, Wildberger JE, Kühl HP, et al. MDCT detection of mitral valve calcification: prevalence and clinical relevance compared with echocardiography. *AJR Am J Roentgenol* 2007; **188**: 1264–69. <https://doi.org/10.2214/AJR.06.1002>
11. Takasu J, Katz R, Nasir K, Carr JJ, Wong N, et al. Relationships of thoracic aortic wall calcification to cardiovascular risk factors: the multi-ethnic study of atherosclerosis (mesa). *Am Heart J* 2008; **155**: 765–71. <https://doi.org/10.1016/j.ahj.2007.11.019>
12. Rumberger JA, Simons DB, Fitzpatrick LA, Sheedy PF, Schwartz RS. Coronary artery calcium area by electron-beam computed tomography and coronary atherosclerotic plaque area. a histopathologic correlative study. *Circulation* 1995; **92**: 2157–62. <https://doi.org/10.1161/01.cir.92.8.2157>
13. Sangiorgi G, Rumberger JA, Severson A, Edwards WD, Gregoire J, et al. Arterial calcification and not lumen stenosis is highly correlated with atherosclerotic plaque burden in humans: a histologic study of 723 coronary artery segments using nondecalcifying methodology. *J Am Coll Cardiol* 1998; **31**: 126–33. [https://doi.org/10.1016/s0735-1097\(97\)00443-9](https://doi.org/10.1016/s0735-1097(97)00443-9)
14. Raggi P, Cooil B, Callister TQ. Use of electron beam tomography data to develop models for prediction of hard coronary events. *Am Heart J* 2001; **141**: 375–82. <https://doi.org/10.1067/mhj.2001.113220>
15. Detrano R, Guerci AD, Carr JJ, Bild DE, Burke G, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008; **358**: 1336–45. <https://doi.org/10.1056/NEJMoa072100>
16. Shaw LJ, Raggi P, Schisterman E, Berman DS, Callister TQ. Prognostic value of cardiac risk factors and coronary artery calcium screening for all-cause mortality. *Radiology* 2003; **228**: 826–33. <https://doi.org/10.1148/radiol.2283021006>
17. McClelland RL, Jorgensen NW, Budoff M, Blaha MJ, Post WS, et al. 10-year coronary heart disease risk prediction using coronary artery calcium and traditional risk factors: derivation in the mesa (multi-ethnic study of atherosclerosis) with validation in the hnr (heinz nixdorf recall) study and the dhs (dallas heart study). *J Am Coll Cardiol* 2015; **66**: S0735-1097(15)04976-1: 1643–53. . <https://doi.org/10.1016/j.jacc.2015.08.035>
18. Williams MC, Moss A, Dweck M, Hunter A, Pawade T, et al. Standardized reporting systems for computed tomography coronary angiography and calcium scoring: a real-world validation of cad-rads and cac-drs in patients with stable chest pain. *J Cardiovasc Comput Tomogr* 2020; **14**: S1934-5925(19)30209-6: 3–11. . <https://doi.org/10.1016/j.jcct.2019.07.010>
19. Budoff MJ, Mayrhofer T, Ferencik M, Bittner D, Lee KL, et al. Prognostic value of coronary artery calcium in the promise study (prospective multicenter imaging study for evaluation of chest pain). *Circulation* 2017; **136**: 1993–2005. <https://doi.org/10.1161/CIRCULATIONAHA.117.030578>
20. Lee JH, Rizvi A, Hartaigh BÓ, Han D, Park MW, et al. The predictive value of coronary artery calcium scoring for major adverse cardiac events according to renal function (from the coronary computed tomography angiography evaluation for clinical outcomes: an international multicenter [confirm] registry). [CONFIRM]. *Am J Cardiol* 2019; **123**: S0002-9149(19)30188-2: 1435–42. . <https://doi.org/10.1016/j.amjcard.2019.01.055>
21. Whelton SP, Al Rifai M, Dardari Z, Shaw LJ, Al-Mallah MH, et al. Coronary artery calcium and the competing long-term risk of cardiovascular vs. cancer mortality: the cac consortium. *Eur Heart J Cardiovasc Imaging* 2019; **20**: 389–95. <https://doi.org/10.1093/ehjci/jey176>
22. Messika-Zeitoun D, Aubry M-C, Detaint D, Bielak LF, Peyser PA, et al. Evaluation and clinical implications of aortic valve calcification measured by electron-beam computed tomography. *Circulation* 2004; **110**: 356–62. <https://doi.org/10.1161/01.CIR.0000135469.82545.D0>
23. Cowell SJ, Newby DE, Burton J, White A, Northridge DB, et al. Aortic valve calcification on computed tomography predicts the severity of aortic stenosis. *Clin Radiol* 2003; **58**: 712–16. [https://doi.org/10.1016/s0009-9260\(03\)00184-3](https://doi.org/10.1016/s0009-9260(03)00184-3)
24. Cueff C, Serfaty J-M, Cimadevilla C, Laissy J-P, Himbert D, et al. Measurement of aortic valve calcification using multislice computed tomography: correlation with haemodynamic severity of aortic stenosis and clinical implication for patients with low ejection fraction. *Heart* 2011; **97**: 721–26. <https://doi.org/10.1136/hrt.2010.198853>
25. Kälsch H, Lehmann N, Möhlenkamp S, Hammer C, Mahabadi AA, et al. Prevalence of thoracic aortic calcification and its relationship to cardiovascular risk factors and coronary calcification in an unselected population-based cohort: the heinz nixdorf recall study. *Int J Cardiovasc Imaging* 2013; **29**: 207–16. <https://doi.org/10.1007/s10554-012-0051-3>
26. Sverzellati N, Arcadi T, Salvolini L, Dore R, Zompatori M, Mereu M, et al. Under-reporting of cardiovascular findings on chest CT. *Radiol Med* 2016; **121**: 190–99. <https://doi.org/10.1007/s11547-015-0595-0>
27. Chiles C, Duan F, Gladish GW, Ravenel JG, Baginski SG, Snyder BS, et al. Association of Coronary Artery Calcification and Mortality in the National Lung Screening Trial: A Comparison of Three Scoring Methods. *Radiology*. 2015;276(1):82-90.
28. Shemesh J, Henschke CL, Farooqi A, Yip R, Yankelevitz DF, Shaham D, et al. Frequency of coronary artery calcification on low-dose computed tomography screening for lung cancer. *Clin Imaging*. 2006;30(3):181-5.
29. Williams MC, Murchison JT, Edwards LD, Agusti A, Bakke P, Calverley PM, et al. Coronary artery calcification is increased in patients with COPD and associated with increased morbidity and mortality. *Thorax*. 2014;69(8):718-23.
30. Williams MC, Morley NCD, Muir KC, Reid JH, van Beek EJR, et al. Coronary artery calcification is associated with mortality independent of pulmonary embolism severity: a retrospective cohort study. *Clin Radiol* 2019; **74**: S0009-9260(19)30549-5: 973. . <https://doi.org/10.1016/j.crad.2019.08.023>
31. Shao L, Yan AT, Lebovic G, Wong HH, Kirpalani A, et al. Prognostic value of visually detected coronary artery calcification

- on unenhanced non-gated thoracic computed tomography for prediction of non-fatal myocardial infarction and all-cause mortality. *J Cardiovasc Comput Tomogr* 2017; **11**: S1934-5925(17)30085-0: 196–202. . <https://doi.org/10.1016/j.jcct.2017.03.004>
32. van der Aalst CM, Denissen SJAM, Vonder M, Gratama JWC, Adriaansen HJ, et al. Screening for cardiovascular disease risk using traditional risk factor assessment or coronary artery calcium scoring: the robinsca trial. *Eur Heart J Cardiovasc Imaging* 2020; **21**: 1216–24. <https://doi.org/10.1093/ehjci/jeaa168>
33. Williams MC, Abbas A, Tirr E, Alam S, Nicol E, et al. Reporting incidental coronary, aortic valve and cardiac calcification on non-gated thoracic computed tomography, a consensus statement from the bsci/bsct and bsti. *Br J Radiol* 2021; **94**: 20200894: 1117: . <https://doi.org/10.1259/bjr.20200894>
34. Hecht HS, Cronin P, Blaha MJ, Budoff MJ, Kazerooni EA, et al. 2016 scct/str guidelines for coronary artery calcium scoring of noncontrast noncardiac chest ct scans: a report of the society of cardiovascular computed tomography and society of thoracic radiology. *J Cardiovasc Comput Tomogr* 2017; **11**: S1934-5925(16)30285-4: 74–84: . <https://doi.org/10.1016/j.jcct.2016.11.003>
35. Falk V, Baumgartner H, Bax JJ, De Bonis M, Hamm C, et al. 2017 esc/eacts guidelines for the management of valvular heart disease. *Eur J Cardiothorac Surg* 2017; **52**: 616–64. <https://doi.org/10.1093/ejcts/ezx324>
36. Budoff MJ, Nasir K, Katz R, Takasu J, Carr JJ, et al. Thoracic aortic calcification and coronary heart disease events: the multi-ethnic study of atherosclerosis (mesa). *Atherosclerosis* 2011; **215**: 196–202. <https://doi.org/10.1016/j.atherosclerosis.2010.11.017>
37. Hoffmann U, Massaro JM, D'Agostino RB, Kathiresan S, Fox CS, et al. Cardiovascular event prediction and risk reclassification by coronary, aortic, and valvular calcification in the framingham heart study. *J Am Heart Assoc* 2016; **5**(2): e003144. <https://doi.org/10.1161/JAHA.115.003144>
38. Mahabadi AA, Lehmann N, Möhlenkamp S, Pundt N, Dykun I, et al. Noncoronary measures enhance the predictive value of cardiac ct above traditional risk factors and cac score in the general population. *JACC Cardiovasc Imaging* 2016; **9**: S1936-878X(16)30408-9: 1177–85: . <https://doi.org/10.1016/j.jcmg.2015.12.024>