



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

On Engel groups, nilpotent groups, rings, braces and the Yang-Baxter equation

Citation for published version:

Smoktunowicz, A 2018, 'On Engel groups, nilpotent groups, rings, braces and the Yang-Baxter equation', *Transactions of the American Mathematical Society*, vol. 370, no. 9, pp. 6535-6564.
<https://doi.org/10.1090/tran/7179>

Digital Object Identifier (DOI):

[10.1090/tran/7179](https://doi.org/10.1090/tran/7179)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Transactions of the American Mathematical Society

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



ON ENGEL GROUPS, NILPOTENT GROUPS, RINGS, BRACES AND THE YANG-BAXTER EQUATION

AGATA SMOKTUNOWICZ

ABSTRACT. It is shown that over an arbitrary field there exists a nil algebra R whose adjoint group R° is not an Engel group. This answers a question by Amberg and Sysak from 1997 [5] and answers related questions from [3, 44]. The case of an uncountable field also answers a recent question by Zelmanov.

In [37], Rump introduced braces and radical chains $A^{n+1} = A \cdot A^n$ and $A^{(n+1)} = A^{(n)} \cdot A$ of a brace A . We show that the adjoint group A° of a finite right brace is a nilpotent group if and only if $A^{(n)} = 0$ for some n . We also show that the adjoint group of A° of a finite left brace A is a nilpotent group if and only if $A^n = 0$ for some n . Moreover, if A is a finite brace whose adjoint group A° is nilpotent then A is the direct sum of braces whose cardinalities are powers of prime numbers. Notice that A° is sometimes called the multiplicative group of a brace A (for example in [13]). We also introduce a chain of ideals $A^{[n]}$ of a left brace A and then use it to investigate braces which satisfy $A^n = 0$ and $A^{(m)} = 0$ for some m, n (Theorems 2, 3).

In Section 2 we describe connections between our results and braided groups and the non-degenerate involutive set-theoretic Yang-Baxter equation. It is worth noticing that by a result by Gateva-Ivanova [17] braces are in one-to-one correspondence with braided groups with involutive braided operators.

Keyword: Engel group, nil ring, Jacobson-radical ring, braces, braided groups, Yang-Baxter equation, nilpotent braces, braided groups.

1. INTRODUCTION

In [37], Rump introduced braces as a generalisation of Jacobson radical rings and as a tool for describing solutions of the Yang-Baxter equation. In the same paper, he introduced the following two series of subsets A^n and $A^{(n)}$ of a right brace A , defined inductively as $A^{n+1} = A \cdot A^n$ and $A^{(n+1)} = A^{(n)} \cdot A$, where $A = A^1 = A^{(1)}$. Let A be a finitely generated Jacobson radical ring. It is known that the adjoint group A° of A is a nilpotent group if and only if A is a nilpotent

Date: November 19, 2015.

2010 Mathematics Subject Classification. Primary 16N80, 16P90, 16N40 .

Key words and phrases. Engel group, nilpotent group, adjoint group of a ring , braces, nil rings, nil algebras, the Yang-Baxter equation.

ring, i.e., $A^n = 0$ for some n [3]. In this paper, we show that a similar result holds for finite braces.

Theorem 1. *Let A be a finite left brace. Then the adjoint group of A is nilpotent if and only if $A^n = 0$ for some n . Moreover, such a brace is the direct sum of braces whose cardinalities are powers of prime numbers.*

Recall that the direct sum $A = \bigoplus_{i=0}^n A_i$ of braces is defined in the same way as for rings; namely if $a = (a_1, \dots, a_n) \in A$ and $b = (b_1, \dots, b_n) \in A$, then $a + b = (a_1 + b_1, \dots, a_n + b_n)$ and $a \cdot b = (a_1 \cdot b_1, \dots, a_n \cdot b_n)$. Recall that a result of Rump shows that if A is a left brace whose adjoint group A° is a finite p -group then $A^n = 0$ for some n (Corollary after Proposition 8, [37]). Notice that if A is a left brace, then by using the opposite multiplication we get a right brace; therefore if A is a right brace, then the group A° is nilpotent if and only if $A^{(n)} = 0$.

Observe that by writing Example 3 of Rump from [37] in the language of left braces, we see that there is a left brace A of cardinality 6 such that $A^{(3)} = 0$ and $A^n \neq 0$ for every n , the adjoint group A° is not a nilpotent group. This shows that the adjoint group of a finite brace need not be an Engel group, and that the assumptions of Theorem 1 are necessary. Notice that by writing Example 2 of Rump from [37] in the language of left braces we get that there is a finite left brace A such that $A^4 = 0$ and $A^{(n)} \neq 0$ for every n , whose adjoint group A° is a nilpotent group ([37], Example 2).

Recall that in [37] Rump introduced the following two series of subsets A^n and $A^{(n)}$ of a right brace A , defined inductively as $A^{n+1} = A \cdot A^n$ and $A^{(n+1)} = A^{(n)} \cdot A$, where $A = A^1 = A^{(1)}$. We introduce the following chain $A^{[n]}$ of ideals of a brace:

$$A^{[n+1]} = \sum_{i=1}^n A^{[i]} \cdot A^{[n+1-i]},$$

where $A^{[1]} = A$. It is clear that $A^{[n]} \subseteq A^{[n-1]} \subseteq \dots \subseteq A^{[1]} = A$, and that for every i , $A^{[i]}$ is a two-sided ideal of A . Recall that for subsets $C, D \subseteq A$ we use notation $C \cdot D = \sum_{i=1}^{\infty} c_i d_i$ with $c_i \in C, d_i \in D$ where almost all c_i, d_i are zero (so the sums $\sum_{i=1}^{\infty} c_i d_i$ are finite). Our next results follow.

Theorem 2. *Let A be a left brace such that $A^{[s]} = 0$ for some s . If $a \in A^{[i]}$, $b \in A^{[j]}$, $c \in A^{[k]}$ then*

$$(a + b)c - ac - bc \in A^{[i+j+k]}.$$

Let $P \subseteq A$ and let S be the set of all products of elements from P . If R is the additive subgroup of A generated by elements from S , then R is a brace.

We obtain that the following result holds for both finite and infinite braces.

Theorem 3. *Let A be a left brace and let n be a natural number. Then the following assertions are equivalent:*

1. $A^{(n)} = 0$ and $A^m = 0$ for some natural numbers m, n .
2. $A^{[n]} = 0$ for some natural number n .
3. $A^{(n)} = 0$ for some n and the group A° is nilpotent.
4. The adjoint group A° is nilpotent, and the solution of the Yang-Baxter equation associated to A is a multipermutation solution (A° is also called the multiplicative group of the brace A in [13]).

Recall that in [16], Etingof, Shedler and Soloviev introduced a retraction of a solution of the Yang-Baxter equation. A solution (X, r) is called a multipermutation solution of level m if m is the smallest nonnegative integer that, after applying the operation of retraction m times, the obtained solution has cardinality 1. If such m exists the solution is also called retractable (see [16] or [14] page 3 for a more detailed definition). Such a solution is also called a *multipermutation solution*, that is a solution which has a finite multipermutation level (for a detailed definition see [17], [13]). There are many interesting results in this area [16, 14, 13, 22, 47]. Proposition 5.16 from [17] and the above Theorem 3 motivated the following related result:

Remark 4. [23] *Let A be a left brace, and let (A, r) be the solution to the Yang-Baxter equation associated to A (as at the beginning of the Section 2). Then (A, r) is a solution of multipermutation level $m < \infty$ if and only if $A^{(m+1)} = 0$ and $A^{(m)} \neq 0$.*

The proof of Remark 4 is very similar to the proof of Proposition 5.16 [17] and can be found in [23]; it is also possible to prove it by applying Proposition 7 [37] several times translated to left braces.

Our next result concerns adjoint groups of radical rings and nil rings. Recall that nil rings have been used by many authors to construct examples of groups; for example triply factorized groups, SN -groups, torsion groups, Engel groups and p -groups. Therefore, it might be useful to describe new methods for constructing and investigating such rings. This is one of the aims which motivated our next result.

Recall that if R is any ring then the adjoint semigroup of R is constructed according to the following rule: $a \circ b = ab + a + b$. It is also denoted $1 + R$, and it is a group if and only if R is a Jacobson radical ring. Amberg, Catino, Dickenschied,

Kazarin, Plotkin, Shalev, Sysak and others proved many interesting results on the adjoint group of a radical ring [7, 44, 6, 4, 10, 33, 8, 25, 13, 39]. Amongst many other interesting results, Amberg, Dickenschied and Sysak showed that the adjoint group R° of any Jacobson radical ring is an SN-group in which every finite subgroup is nilpotent [3]. As mentioned in their paper, by using Zelmanov's theorem on the restricted Burnside problem (see [49, 50, 51]) and properties of SN-groups they were able to deduce the following: If R is a finitely generated Jacobson radical ring, then the following are equivalent: (a) R is an n -Engel ring for some $n \geq 1$ (b) R is a nilpotent ring (c) R° is an n -Engel group for some $n \geq 1$. Recall that the aforementioned result of Zelmanov asserts that an n -Engel Lie algebra over an arbitrary field is locally nilpotent, and that any torsion free n -Engel Lie ring is nilpotent [49, 50, 51]. A surprisingly short proof by Shalev assures that if a radical ring R is an n -Engel algebra over a field of prime characteristic then the adjoint group R° of R is m -Engel for some m [39]. A natural question arises then whether an analogy of any of these results would hold for Engel groups and Engel Lie rings. Notice that every nil ring is an Engel Lie ring (for some interesting related results see [2, 32, 40]). Golod has constructed a nil ring whose adjoint group is an Engel group. In 1997 in [3], Amberg and Sysak asked the following question: *If R is a nil ring, is the adjoint group R° an Engel group?* Similar questions were also asked in [3, 44]. At the conference in Porto Cesareo in July 2015, after one of the talks Zelmanov asked the following question *If R is a nil algebra over an uncountable field, is the adjoint group R° an Engel group?* Our result answers these questions in the negative:

Theorem 5. *There is a nil ring R such that the adjoint group of R° is not an Engel group. Moreover, R can be taken to be an algebra over an arbitrary field.*

The paper is organized thus: in Section 2 we mention connections with the Yang-Baxter equation and braided groups. In Sections 6 – 12 we prove Theorem 5. In Sections 3-5 we prove Theorems 1, 2, 3. Sections 6-12 and Sections 3-5 can be read independently.

2. NOTATION AND APPLICATIONS FOR THE YANG-BAXTER EQUATIONS AND FOR BRAIDED GROUPS

Around 2005, Rump introduced braces as a generalisation of Jacobson radical rings. He also showed that braces correspond to solutions of the Yang-Baxter equation [37]. In [30] Lu, Yan and Zhu proposed a general way of constructing set-theoretical solutions of the Yang-Baxter equation using braiding operators on

groups. In this paper, by a solution of the Yang-Baxter equation we will mean non-degenerate involutive set-theoretic solution of the Yang-Baxter equation, as in [13].

Let R be a Jacobson radical ring; then R yields a solution to the Yang-Baxter equation with the Yang-Baxter operator $r(x, y) = (u, v)$, where $u = x \cdot y + y$ and $v = z(xy + x + y) + xy + x + y + z$ where z is the inverse of $u = xy + y$ in the adjoint group R° of R , for $x, y \in R$. The same holds when $(R, +, \cdot)$ is a left brace, and this solution is called the solution *associated to left brace* R , and will be denoted as (R, r) . This follows from the property $x \circ y = u \circ v$ which implies $zoxoy = v$ where $a \circ b = a \cdot b + a + b$.

In [35] pp 128, Rump gave the following definition of a right brace: “Let A be an abelian group together with a right distributive multiplication, that is,

$$(a + b)c = ac + bc$$

for all $a, b, c \in A$. We call A a *brace* if the *circle operation*

$$a \circ b = ab + a + b$$

makes A into a group. This group A° will be called the *adjoint group* of a brace A .” In [13] Cedó, Jespers and Okninski wrote the definition of a brace in terms of operation \circ ; in their paper the adjoint group A° is called the multiplicative group of brace A .

Similarly, a left brace is an abelian group $(A, +)$ together with a left distributive multiplication; that is, $a(b + c) = ab + ac$ such that the circle operation $a \circ b = ab + a + b$ makes A into group. For a left brace A , the associativity of A° is easily seen to be equivalent to the equation $(ab + a + b)c = a(bc) + ac + bc$. A right brace which is also a left brace is called a two-sided brace; Rump has shown that two-sided braces are exactly Jacobson radical rings.

In [16] Etingof, Shedler and Soloviev introduced a retraction of a solution of the Yang-Baxter equation, and described some classes of the solutions. They also introduced retractable solutions, which are now also called multipermutation solutions (see [16, 13, 17]). Theorem 2 [13] by Cedó, Jespers and Okninski assures that *If G is a left brace then there exists a solution X, r' of the Yang-Baxter equation such that the solution $\text{Ret}(X, r')$ is isomorphic to the solution associated to the left brace G , and moreover, $\mathcal{G}(X, r)$ is isomorphic to the multiplicative group of the left brace G . Furthermore, if G is finite then X can be taken a finite set.*

By writing Example 3 of Rump from [37] in the language of left braces, we get that there is a finite left brace A whose adjoint group A° is the symmetric group S_3

which is not a nilpotent group ([37], Example 3); moreover $A^{(3)} = 0$ and $A^n \neq 0$ for every n . By Remark 4 the solution to the Yang-Baxter equation associated to A is a multipermutation solution. Observe that by applying the aforementioned Theorem 2 from [13] to this example we obtain the following remark.

Remark 6. *(related to Example 3, [37]) There is a finite multipermutation solution (X, r) of the Yang-Baxter solution whose permutation group $\mathcal{G}(X, r)$ of left actions associated with (X, r) is not a nilpotent group.*

Recall that permutation group $\mathcal{G}(X, r)$ of left actions associated with (X, r) was introduced by Tatyana-Ivanova in [18] (see also [17]). By writing Example 2 from [37] in the language of a left brace we get that there is a finite left brace A such that $A^4 = 0$, $A^{(n)} \neq 0$ for every n , whose multiplicative group is a nilpotent group ([37], Example 2). By Remark 4 the solution associated to A is not a multipermutation solution. This implies, together with Theorem 2 from [13], the following remark.

Remark 7. *(related to Example 2, [37]) There is a finite solution (X, r) to the Yang-Baxter equation, which is not a multipermutation solution, and whose permutation group $\mathcal{G}(X, r)$ of left actions associated with (X, r) is a nilpotent group.*

We get a following related result for (possibly infinite) braces.

Proposition 8. *Let A be a left brace such that the solution of the Yang-Baxter equation associated to A is a multipermutation solution. Then the adjoint group A° of A is a nilpotent if and only if $A^n = 0$ for some natural number n .*

Gateva-Ivanova and Van den Bergh [24] and independently Etingof, Schedler and Soloviev [16] gave a group theoretical interpretation of the set theoretic involutive non-degenerate solutions of the Yang-Baxter equation. Cedó, Jespers and Okninski [13, 12] asked which groups are multiplicative groups of braces. A similar question in the language of ring theory was asked in [4, 5]. In this paper we obtain the following corollary of Theorem 5, which is related to this question.

Corollary 9. *There is a finitely-generated, two-sided brace whose multiplicative group is a torsion group but is not an Engel group.*

By a result of Gateva-Ivanova (see Theorem 3.7, [17]), every brace G can be considered as a braided group with the involutive braided operator. Moreover, by Proposition 6.2, [17], G is a two-sided brace if and only if the corresponding braided group satisfies the following identity:

$$c^{(abc)^{-1}}c = (b^{-1}c)^{((a^b)^{(b^{-1}c))^{-1}}c),$$

for every $a, b, c \in G$. By combining the Gateva-Ivanova result with Corollary 9, we obtain that:

Corollary 10. *There is a countable, braided group (G, σ) with an involutive braided operator σ which is a torsion-group and not an Engel group. Moreover, G satisfies a non-trivial identity*

$$c^{(abc)^{-1}}c = (b^{-1}c)^{((a^b)^{(b^{-1}c))^{-1}}c),$$

for all $a, b, c \in G$. We use notation $\sigma(a, b) = (a^b, a^b)$.

This shows that infinite braided groups satisfying non-trivial identities can be quite complicated.

We also get the following result for finite braided groups.

Proposition 11. *Let G be a finite nilpotent group and let (G, σ) be symmetric group (in the sense of Takeuchi). Let $(G, +, o)$ be a left brace associated to (G, σ) as in Theorem 3.8 in [17]. Then $(G, +, o)$ is a direct sum of left braces whose cardinalities are powers of prime numbers. These braces correspond to Sylow subgroups of G .*

3. BRACES WITH $A^n = 0$ AND $A^{(n)} = 0$

In [37] Rump introduced the following two series of subsets of a brace. One of the series introduced by Rump is $\dots \subseteq A^{(2)} \subseteq A^{(1)} = A$, where $A^{(n+1)} = A^{(n)} \cdot A$. The other series introduced by Rump is $\dots \subseteq A^2 \subseteq A^1 = A$, where $A^{n+1} = A \cdot A^n$. Rump has proved that the series A^n of every right brace consists of two-sided ideals. Similarly, for a left brace A , the series $A^{(n)}$ consists of two-sided ideals. Recall that I is an ideal in a brace A if for $i, j \in I$ and $a \in A$ we have $i + j \in I$ and $ai \in I, ia \in I$, see [37].

We propose another series, defined for any left or right brace. This series consists of two-sided ideals in A . We define the series $\dots \subseteq A^{[2]} \subseteq A^{[1]} = A$, where

$$A^{[n+1]} = \sum_{i=1}^n A^{[i]} \cdot A^{[n+1-i]}.$$

Then it is clear that $A^{[n]}$ is an ideal in A for every n , and $A^{[n+1]} \subseteq A^{[n]}$.

Recall that for subsets $C, D \subseteq A$ we use notation

$$C \cdot D = \sum_{i=1}^{\infty} c_i d_i$$

with $c_i \in C, d_i \in D$, and almost all c_i, d_i are zero (so the sums $\sum_{i=1}^{\infty} c_i d_i$ are finite).

Theorem 12. *Let $(A, \cdot, +)$ be a left or right brace. If m, n are natural numbers and $A^n = A^{(m)} = 0$ then $A^{[s]} = 0$ for some number s .*

Proof. We will prove the result in the case when A is a right brace, the case when A is a left brace is done by considering the brace with the opposite multiplication. We will proceed by induction on n . If $n = 2$ then $0 = A^2 = A \cdot A = A^{(2)} = A^{[2]}$, so the result holds. Suppose that there is a natural number $s_{n,m}$ such that any right brace satisfying $A^n = 0$ and $A^{(m)} = 0$ satisfies $A^{[s_{n,m}]} = 0$.

Assume now that our brace satisfies $A^{n+1} = 0$ and $A^{(m)} = 0$. Let $p > s_{n,m} \cdot m$, and suppose that $a \in A^{[p]}$. Then $a = \sum_i a_i b_i$ for some $a_i, b_i \in A$ with $a_i \in A^{[p-q_i]}$, $b_i \in A^{[q_i]}$, for some numbers q_i . Observe that if $q_i > s_{n,m}$ then $b_i \in A^n$ (by the inductive assumption applied to the brace A/A^n ; this brace is well defined as Rump proved that A^n is an ideal in A). In this case we get $a_i b_i \in A \cdot A^n = A^{n+1} = 0$. Therefore $q_i \leq s_{n,m}$, as otherwise $a_i b_i = 0$. Consequently we can assume that all $q_i \leq s_{n,m}$. For each i , we can now write $a_i = \sum_j a_{i,j} b_{i,j}$, and by the same argument as before, we get that each $b_{i,j} \in A^{[r_i]}$ for some $r_i \leq s_{n,m}$ (as otherwise $b_{i,j} \in A^n$ by the inductive assumption applied to A/A^n , and so $a_{i,j} b_{i,j} \in A^{n+1} = 0$). Observe now that since A is a right brace then

$$\sum_i a_i b_i = \sum_i \left(\sum_j a_{i,j} b_{i,j} \right) b_i = \sum_{i,j} (a_{i,j} b_{i,j}) b_i.$$

Continuing in this way we get that $a \in \sum_{c_1, \dots, c_m \in A} (((A \cdot c_1) \cdot c_2) \dots \cdot c_{m-1}) \cdot c_m$, and since $A^{(m)} = 0$ we get that each $a = 0$, so $A^{[p]} = 0$. \square

Theorem 13. *Let $(A, \cdot, +)$ be either a left brace or a right brace. If $A^n = A^{(m)} = 0$ for some natural numbers m, n , then the multiplicative group of A is a nilpotent group.*

Proof. Let $a, b \in A$, then $[a, b] = a \circ b \circ a^{-1} \circ b^{-1}$ where a^{-1} and b^{-1} are inverses of a and b respectively in the adjoint group A° . We will construct a finite lower central series of A° . We proceed by induction on S . If $A^{[2]} = 0$ then A is commutative so the result holds. Suppose that the result holds for all numbers smaller than s ; by the inductive assumption applied to $A' = A/A^{[s-1]}$ we get $[[[[A, A]A] \dots]A] \in A^{[s-1]}$ (m brackets for some m). Since $A^{[s-1]}$ is in the center of A we get that $[[[[A, A]A] \dots]A] = 0$ ($m+1$ brackets), hence A has a finite lower central series. \square

Theorem 14. *Let A be a left brace such that $A^{(n)} = 0$ for some n . If the multiplicative group of A is nilpotent then $A^m = 0$ for some m , and hence $A^{[s]} = 0$ for some s .*

Proof. By assumption $A^{(n)} = 0$ for some n . We can assume that n is minimal possible. Let $b \in A^{(n-1)}$, $a \in A$ and let a^{-1} and b^{-1} be the inverses of respectively a and b in the adjoint group A° . Recall that A° is the group under the circle operation $a \circ b = ab + a + b$. We will show that

$$a \circ b \circ a^{-1} \circ b^{-1} = ab.$$

Note that $A^{(n-1)} \subseteq \text{Soc}(A) = \{x \in A \mid x \circ a = x + a\}$. By [38, Corollary after Proposition 6], $A^{(n-1)}$ is an ideal. Hence $A^{(n-1)}$ is a normal subgroup of the multiplicative group of the left brace A . Let $b \in A^{(n-1)}$ and $a \in A$. Since $0 = b \circ b^{-1} = b + b^{-1}$, we have that $b^{-1} = -b$.

$$\begin{aligned} [a, b] &= a \circ b \circ a^{-1} \circ b^{-1} \\ &= a \circ b \circ a^{-1} + b^{-1} \quad (\text{since } a \circ b \circ a^{-1} \in A^{(n-1)}) \\ &= a \circ (b + a^{-1}) - b \\ &= a \circ b + a \circ a^{-1} - a - b \\ &= a \circ b - a - b \\ &= ab \end{aligned}$$

Therefore $[a, b] = a \circ b \circ a^{-1} \circ b^{-1} = ab$.

Since the multiplicative group A° of A is nilpotent we get

$$[a_m[\dots[a_2[a_1, b_1]]]] = 0$$

for some m . Therefore $[a_m[\dots[a_2[a_1, b_1]]]] = a_m(a_{m-1}(\dots(a_2(a_1b))))$. Consequently $A(A(\dots A(A^{(n-1)}))) = 0$ (m brackets).

We will now apply this result to prove our theorem; we will use induction on n (recall that n is such that $A^{(n)} = 0$). For $n = 2$ the result holds since $A^{(2)} = A^2 = A^{[2]}$. Suppose now that the result holds for all numbers smaller than n , so if B is a left brace and $B^{(n-1)} = 0$ and the adjoint group of B° is nilpotent then $B^{(n')} = 0 = B^{[n']}$ for some n' .

Recall that $A^{(n-1)}$ is an ideal in A , and hence a normal subgroup of A° [37, 13], hence the adjoint group of brace $A/A^{(n-1)}$ is nilpotent. We can apply the inductive assumption for brace $B' = A/A^{(n-1)}$ and we get that $B'^{n'} = 0$ hence $A^{n'} = A(A(\dots A)) \subseteq A^{(n-1)}$. Therefore $A^{m+n'} \subseteq A(A(\dots A(A^{(n-1)}))) = 0$. By Theorem 12 we get that $A^{[s]} = 0$ for some s . \square

Let us remark that the first part of the above proof was provided by Ferran Cedo after reading the original proof in the first version in this manuscript. The original proof, which is much longer, can be found in [43].

4. STRUCTURE OF LEFT BRACES WITH $A^n = 0$

In this section we observe some connections between nilpotent braces and nilpotent rings. We start with the following lemma.

Lemma 15. *Let s be a natural number and let A be a left brace such that $A^s = 0$ for some s . Let $a, b \in A$. Define inductively elements $d_i = d_i(a, b)$, $d'_i = d'_i(a, b)$ as follows: $d_0 = a$, $d'_0 = b$, and for $i \leq 1$ define $d_{i+1} = d_i + d'_i$ and $d'_{i+1} = d_i d'_i$. Then for every $c \in A$ we have*

$$(a + b)c = ac + bc + \sum_{i=0}^{2s} (-1)^{i+1} ((d_i d'_i)c - d_i(d'_i c)).$$

Proof. Observe first that by an inductive argument $d'_i \in A^i$ for each i . Observe that for $i \geq 1$ we have

$$d_{i+1} \cdot c = (d_i + d'_i) \cdot c = ((d_{i-1} + d'_{i-1}) + d_{i-1} d'_{i-1}) \cdot c.$$

Recall that since A is a left brace then

$$d_{i+1}c = (d_{i-1} + d'_{i-1} + d_{i-1}d'_{i-1}) \cdot c = d_{i-1}c + d'_{i-1}c + d_{i-1}(d'_{i-1}c).$$

The same holds when we increase i by 1, hence $d_{i+2}c = d_i c + d'_i c + d_i(d'_i c)$. Subtracting the above equation from the previous one we get

$$d_{i+1}c - d_{i+2}c = (d_{i-1}c - d_i c) + e_i$$

where $e_i = d'_{i-1}c - d'_i c + d_{i-1}(d'_{i-1}c) - d_i(d'_i c)$. Observe that

$$e_i = (d_{i-2}d'_{i-2})c - (d_{i-1}d'_{i-1})c + d_{i-1}(d'_{i-1}c) - d_i(d'_i c).$$

Therefore,

$$\sum_{i=1}^s e_{2i} = \sum_{i=1}^s (d_{2i-2}d'_{2i-2})c - (d_{2i-1}d'_{2i-1})c + d_{2i-1}(d'_{2i-1}c) - d_{2i}(d'_{2i}c)$$

Notice that if $i \geq s$ then $d'_i \in A^s = 0$. Therefore $\sum_{i=1}^s e_{2i} = (d_0 d'_0)c + q$ where

$$q = \sum_{i=1}^s d_{2i-1}(d'_{2i-1}c) - (d_{2i-1}d'_{2i-1})c - \sum_{i=1}^s d_{2i}(d'_{2i}c) - (d_{2i}d'_{2i})c.$$

Observe now that $d_{i+1}c - d_{i+2}c = (d_{i-1}c - d_i c) + e_i$ implies

$$\sum_{i=1}^s (d_{2i+1}c - d_{2i+2}c) = \sum_{i=1}^s (d_{2i-1}c - d_{2i}c) + \sum_{i=1}^s e_{2i},$$

therefore

$$d_{2s+1}c - d_{2s+2}c = d_1c - d_2c + \sum_{i=1}^s e_{2i}.$$

Observe that $d_{2s+2} = d_{2s+1} + d'_{2s+1} = d_{s+1}$ since $d'_{2s+1} \in A^s = 0$. Consequently $d_2c - d_1c = \sum_{i=1}^s e_{2i} = (d_0d'_0)c + q$. Recall that $d_0 = a$, $d'_0 = b$, $d_1 = a + b$, $d'_1 = ab$ and $d_2 = a + b + ab$. Therefore $d_1c = (a + b)c$ and $d_2c = (a + b + ab)c = ac + bc + a(bc)$. Consequently $d_1c = d_2c - (d_0d'_0)c + q$. It follows that $(a + b)c = ac + bc + a(bc) - (d_0d'_0)c - q = ac + bc + d_0(d'_0c) - (d_0d'_0)c - q$. Notice that $d_0(d'_0c) - (d_0d'_0)c - q = \sum_{i=0}^{2s} (-1)^{i+1} ((d_i d'_i)c - d_i(d'_i c))$, which finishes the proof. \square

Lemma 16. *Let the assumptions and notation be as in Lemma 15. Suppose that there is a natural number n such that $m \cdot a = m \cdot b = 0$. Let d_i, d'_i be as in Lemma 15; then $m \cdot d_i = m \cdot d_{i'} = 0$ for every $i \geq 1$.*

Proof. We will first show that $m \cdot d'_t = 0$ for every $t \geq 0$. For $t = 0$ we have $m' \cdot d'_0 = m \cdot b = 0$. Suppose the result holds for some $t \geq 0$, then $m \cdot d'_{t+1} = m \cdot (d_t d'_t) = d_t(m \cdot d'_t) = 0$.

We will now that $m \cdot d_t = 0$ for all $t \geq 0$. For $t = 0$ we have $m \cdot d_0 = m \cdot a = 0$. Suppose the result holds for some $t \geq 0$, then $m \cdot d_{t+1} = m \cdot d_t + m \cdot d'_t = 0$ by the inductive assumption. \square

Let A be a left brace and let $S, Q \subseteq A$ be additive subgroups of A . Then we denote $S + Q = \{s + q : s \in S, q \in Q\}$.

For an element $a \in A$ and a natural number i by

$$i \cdot a$$

we will denote the sum of i copies of element a (hence $0 \cdot a = 0$).

Lemma 17. *Let $(A, +, \cdot)$ be a finite left brace of cardinality $p_1^{\alpha_1} \dots p_k^{\alpha_k}$ for some prime pairwise distinct numbers p_1, \dots, p_k and natural numbers $\alpha_1, \dots, \alpha_k$. Then*

$$A = A_1 + A_2 + \dots + A_k$$

where A_i is the additive subgroup of the additive group $(A, +)$ of cardinality $p_i^{\alpha_i}$ for every $i \leq k$. Moreover, $(A_i, +, \cdot)$ is a brace for each $i \leq k$.

Proof. Since the additive group of A is a finite abelian group, then using the primary decomposition theorem we can decompose the additive group $(A, +)$ into a sum of additive subgroups of A ; we can call them A_1, \dots, A_k , where A_i is an additive subgroup of A of cardinality $p_i^{\alpha_i}$ and $A_i \cap A_j = 0$. Observe that if $x, y \in A$

and $p \cdot y = 0$ for some natural number p then $p \cdot (xy) = x \cdot (py) = 0$. Therefore if $a, a' \in A_i$ then $a \cdot a' \in A_i$, hence A_i is closed under the multiplication. We know that A_i is closed under the addition, hence it is also closed under the operation \circ , where $a \circ b = a \cdot b + a + b$ for $a, b \in A$. Observe that since A is a finite group, then the inverse of $a \in A$ in the adjoint group A° is of the form $a \circ a \circ \dots \circ a$, hence it belongs to A_i . It follows that A_i is a left brace. \square

Theorem 18. *Let A be a finite left brace such that $A^n = 0$ for some n . Then A is the direct sum of braces whose cardinalities are powers of prime numbers. In particular, the adjoint group A° of A is a nilpotent group.*

Proof. Let notation be as in Lemma 17. We will first show that if $a, b \in A$ and $m \cdot a = m' \cdot c = 0$ for some coprime natural numbers m, m' then $a \cdot c = 0$. Let $a \in A$ by $\deg(a)$ we will denote the largest number $i \leq n$ such that $a \in A^i$. We will proceed by induction on $i = 2n - \deg(a) - \deg(c)$. If $2n - \deg(a) - \deg(c) = 0$ then $a, c \in A^n = 0$, so the result holds. Suppose now that the $i > 0$ and that result holds when $2n - \deg(a) - \deg(c) < i$. We will show that the results also holds for $2n - \deg(a) - \deg(c) = i$.

Let j be a natural number. Let $q, d_1, d'_1, \dots, d_n, d_{n'}$ be as in Lemma 15, applied for a and for $b = ja$ and for $s = n$. Denote $q_j = q$ then by Lemma 15 we have

$$(a + ja)c = ac + (ja)c + q_j.$$

By Lemma 16, $m \cdot d_i = m \cdot d'_i$ for all $i \geq 0$. Observe now that for any i , the order of element $d'_i c$ is a divisor of m' and hence is coprime with m . This follows because, by the assumption at the beginning of the proof the order of c is m' and m' is coprime with m . Observe that $m' \cdot (d'_i c) = d'_i c + \dots + d'_i c = d'_i \cdot (m' \cdot c) = 0$.

Observe $d_i(d'_i c) = 0$ by the inductive assumption on, as $2n - \deg(d_i) - \deg(d'_i c) \leq 2n - \deg(d_i) - (\deg(c) + 1) < 2n - \deg(a) - \deg(c)$. Similarly $d_i(d'_i c) = 0$ by the inductive assumption since $2n - \deg(d_i d'_i) - \deg(c) \leq 2n - (\deg(d'_i) + 1) - \deg(c) < 2n - \deg(a) - \deg(c)$. Therefore $q_j = 0$. Consequently for every natural number j ,

$$(a + ja)c = ac + (ja)c.$$

Recall that m, m' are coprime numbers; therefore there are natural numbers ξ, β such that $\beta m' - \xi \cdot m = 1$. Denote $e = \xi \cdot m + 1 = \beta \cdot m'$. Observe now that by the above $ac = (ea)c = ((e-1)a)c + ac = ((e-2)a)c + ac + ac = \dots = e(ac) = a(ec) = 0$. We have proved that $ac = 0$. Therefore if $a \in A_i$ and $c \in A_j$ then $ac = 0$, provided that $i \neq j$.

Let $a_i \in A^i$ for $i = 1, \dots, k$ and $b \in A$. By the property of a left brace

$$b \cdot \left(\sum_{i=1}^k a_i \right) = \sum_{i=1}^k ba_i.$$

Let $c_i \in A_i$. To show that A is the direct sum of braces A_i it remains to show that $(\sum_{j=1}^k a_j)c_i = a_i c_i$. We will show that for every $l \leq k$, $(\sum_{j=1}^l a_j)c_i = a_i c_i$ if $i \leq l$ and $(\sum_{j=1}^l a_j)c_i = 0$ if $i > l$. We will proceed by induction on l . The result is true for $l = 1$. Let $l > 1$ and suppose that the result holds $l - 1$.

Observe first that $a_l \cdot (\sum_{j=1}^{l-1} a_j) = \sum_{j=1}^{l-1} a_l a_j = 0$ by the first part of the proof. Hence $(\sum_{j=1}^l a_j)c_i = (a_l + (\sum_{j=1}^{l-1} a_j) + a_l(\sum_{j=1}^{l-1} a_j))c_i = a_l c_i + (\sum_{j=1}^{l-1} a_j)c_i + a_l((\sum_{j=1}^{l-1} a_j)c_i)$. By the inductive assumption $(\sum_{j=1}^{l-1} a_j)c_i = a_i c_i$ if $i \leq l$ and $(\sum_{j=1}^{l-1} a_j)c_i = 0$ otherwise. Suppose that $i > l$ then $(\sum_{j=1}^{l-1} a_j)c_i = 0$ and $a_l c_i = 0$ hence $(\sum_{j=1}^l a_j)c_i = 0$, as required. If $i = l$ then $(\sum_{j=1}^{l-1} a_j)c_i = 0$ so $(\sum_{j=1}^l a_j)c_i = a_l c_i = a_i c_i$ as required. If $i < l$ then $(\sum_{j=1}^{l-1} a_j)c_i = a_i c_i$, $a_l c_i = 0$ and $(a_l(\sum_{j=1}^{l-1} a_j)c_i) = a_l(a_i c_i) = 0$ as $a_l \in A_l$ and $a_i c_i \in A_i$ and $l \neq i$. Hence $(\sum_{j=1}^l a_j)c_i = a_l c_i = a_i c_i$ as required. Therefore, A is the direct sum of braces A_i .

We will now show the nilpotency of A . Observe first that for every i , A_i is a p -group and hence is nilpotent. Observe then that if $a \in A_i$ and $b \in A_j$ for $i \neq j$ then $a \circ b = b \circ a$ since $a \circ b = a + b + ab = a + b$ and $b \circ a = b + a + ba = b + a$ by the above. Therefore, A° is the direct product of groups A_i for $i = 1, \dots, k$, and hence it is a nilpotent group. \square

5. BRACES WHOSE ADJOINT GROUP IS NILPOTENT

In this section we will investigate the structure of braces whose adjoint groups are nilpotent. For the following result we use a short proof which was provided by Ferran Cedó after reading the first version of this manuscript (see [43]). The original and much longer proof can be found in [43].

Theorem 19. *Let A be a left brace such that the adjoint group A° is a nilpotent group. Then A is a direct sum of braces whose cardinalities are powers of prime numbers. Assume that A has cardinality $p_1^{\alpha_1} \cdots p_k^{\alpha_k}$, for some prime pairwise distinct numbers p_1, \dots, p_k and some natural numbers $\alpha_1, \dots, \alpha_k$. Then $A^n = 0$ where n is the largest number from among $\alpha_1 + 1, \alpha_2 + 1, \dots, \alpha_k + 1$.*

Proof. (Provided by Ferran Cedó.) The first part is easier to prove using the equivalent definition of left brace. [[13], Definition 1]: A left brace is a set B with two binary operations: a sum $+$ and a multiplication \circ , such that $(B, +)$ is an abelian group, (B, \circ) is a group and $a \circ (b + c) + a = a \circ b + a \circ c$ for all $a, b, c \in B$.

Suppose that B is a finite left brace such that its multiplicative group is nilpotent. Let P be a Sylow p -subgroup of the additive group of the left brace B . By [[13], Lemma 1], $\lambda_a(P) = P$ for all $a \in B$, where $\lambda_a(b) = a \circ b - a$. In particular P is closed by the multiplication and hence it is a subgroup of the multiplicative group of the left brace B . Thus P is a Sylow p -subgroup of the multiplicative group of B . Since the multiplicative group of B is nilpotent, P is a normal subgroup in (B, \circ) . Hence P is an ideal of the left brace B (see [[13], Definition 3]). Therefore, if P_1, \dots, P_r are the Sylow subgroups of the additive group of B , then they are also the Sylow subgroups of the multiplicative group of B , in fact they are ideals of B and $B = P_1 \circ \dots \circ P_r = P_1 + \dots + P_r$ is the inner direct product of the subbraces P_1, \dots, P_r .

The second part of Theorem 19 is a consequence of [[37], Corollary after Proposition 8]. \square

Proof of Theorem 1 If $A^n = 0$ for some n , then by Theorem 48 the group A° is nilpotent, and A is the direct sum of braces whose cardinalities are prime numbers. On the other hand if A is a left brace and A° is nilpotent then $A^n = 0$ for some n , by Theorem 19.

Proof of Theorem 2 This follows from Lemma 15 applied several times, taking into account that $A^{[s]} = 0$.

Proof of Theorem 3 Notice that 1 and 2 are equivalent by Theorem 12. Notice that by Remark 4, 3 and 4 are equivalent. By Theorems 13 and 14 properties 3 and 1 are equivalent.

Proof of Proposition 8 This follows from Remark 4 and Theorems 13, 14.

6. JACOBSON RADICAL

In this chapter we give some preliminary results on Jacobson radical rings.

Lemma 20. *Let F be a field. Let n be a natural number. Let R be an F -algebra generated by elements a, b , and suppose that $a^2 = 0$ and $b^n = 0$ for some n . Let S be the F -linear space spanned by elements $a \cdot b^i$ for $1 < i < n$. If all finite matrices with entries from S are nilpotent, then R is a Jacobson radical ring.*

Proof. We will use the well-known fact that a one-sided ideal in which every element is quasi-regular generates a two-sided ideal which is Jacobson radical [29]. Let R' be a subring of R generated by elements from S . Since all matrices with entries from S are nilpotent, then by Theorem 1.2 from [42] R' is a Jacobson radical ring. Consider ring S' generated by elements from S and from Sa and by

element a . Recall that $a^2 = 0$, and so $SaS = 0$. Therefore Sa is a two sided-ideal in S' which is nilpotent; also S'/Sa is Jacobson radical, since R' is nilpotent. It follows that S' is Jacobson radical.

Observe that $S'R \subseteq S + Sa = S'$, hence S' is a right ideal in R . Therefore the two sided ideal generated by S' in R is Jacobson radical; we will call this ideal I . Observe now that the ring R/I is nilpotent, as it is generated by powers of b . It follows that R is Jacobson radical. \square

Lemma 21. *Let F be a field. Let n be a natural number. Let R be an F -algebra generated by elements a, b , and suppose that $a^2 = 0$ and $b^n = 0$ for some n . Let $R[x]$ be the polynomial ring in one variable x over R . Let Q be the F -linear space spanned by elements $a \cdot b^i x^j$ for $0 < i < n$, $0 \leq j$. If all finite matrices with entries from Q are nilpotent, then $R[x]$ is a Jacobson radical ring, and hence R is a nil ring.*

Proof. Amitsur's theorem assures that if R is a ring such that $R[x]$ is Jacobson radical then R is a nil ring. Therefore it suffices to show that $R[x]$ is Jacobson radical. Observe that by Theorem 1.2 from [42], if R' is a subring of $R[x]$ generated by elements from Q then R' is Jacobson radical. Let $S' = R' + R'a + F \cdot a$; then similarly as in Lemma 20 we get that S' is Jacobson radical. It then follows that the two-sided ideal I generated by S' in R is Jacobson radical, and moreover $R[x]/I$ is nil. Therefore $R[x]$ is Jacobson radical. \square

By R^1 we denote the usual extension of a ring R by the identity element.

Lemma 22. *Let F be a field, and $R = F[a, b]$ be the free algebra (without identity) generated by elements a, b . Let S be the linear F -subspace of $F[a, b]$ spanned by elements ab and $a \cdot b^2$. Let I be the ideal of $F[a, b]$ generated by a^2, b^3 and by elements from sets F_1, F_2, \dots such that $F_i \subseteq S^i$ for every i .*

1. *If $p + q + t + t' \in I$ and $p \in ab \cdot R^1, q \in bR^1, t \in F[a], t' \in F[b]$ then $p, q, t, t' \in I$.*
2. *If $p + q \in I$ and $p \in R^1 \cdot ba, q \in R \cdot b$ then $p, q \in I$.*
3. *If $p = e_1 + e_2 + \dots + e_n$ with $e_i \in F \cdot S^i$ and $p \in I$ then $e_i \in I$ for all $i \leq n$.*

Proof. **1.** By specialising at $b = 0$ we get that $t = 0 \in I$, and by specialising at $a = 0$ we get $t' = 0$; hence $p + q \in I$. Denote $Z = \bigcup_{i=1}^{\infty} F_i$. Observe that $Z \subseteq aR$ and $I \subseteq Z \cdot R^1 + b^3 \cdot R^1 + a^2 \cdot R^1 + bI + aI$. It follows that $I \subseteq aR + (bI \cap baR^1) + b^2R^1$. Notice that $p + q \in I$ and so $p + q - s \in b^2R^1$ for some $s \in aR + (bI \cap baR^1)$. Since s and $p + q$ have not terms from b^2R^1 , then $p + q - s = 0$. It follows that $p + q \in aR + (bI \cap baR^1)$. It follows that $q \in bI$, hence $p = (p + q) - q \in I$.

2. Observe now that $I \subseteq Ia + Ib + R^1 \cdot Z + R^1 \cdot a^2 + R^1 \cdot b^3$, hence $p + q \in (Ia \cap R^1ba) + Rb + R^1 \cdot a^2$. Notice that $p \in R^1ba$ and $q \in Rb$; it follows that $p \in Ia \cap R^1ba$, and hence $p \in I$ and so $q = (p + q) - p \in I$.

3. Let J be the ideal of R generated by elements from sets F_i , and let $\langle a^2 \rangle, \langle b^3 \rangle$ denote ideals generated by a^2 and b^3 respectively; then $p - j \in \langle a^2 \rangle + \langle b^3 \rangle$ for some $j \in J$. As j and p have no terms from $\langle a^2 \rangle + \langle b^3 \rangle$ then $p = j$, so $p \in J$. Consequently $p = e_1 + e_2 + \dots + e_n \in J$ with $e_i \in S^i$ implies that $p \in I' + bI' + I'a + bI'a$, where I' is the ideal of E generated by elements from sets F_i and F'_i , where E is the F -algebra generated by elements from S and $F'_i \in (F_i b + F_i b^2 + Rb^3) \cap E$. Hence $p - i \in bI' + I'a + bI'a$ for some $i \in I'$. Since the left hand side belongs to $aRb + ab \cdot F$ and the right hand side to $Ra + bR$ it follows that $p - i = 0$, so $p \in I' \subseteq E$. Therefore in the factor ring E/I' we have that $p + I'$ is the zero element.

Observe that I' is an homogeneous ideal in E when we assign gradation of elements from S to have gradation 1, from S^2 gradation 2 etc. Now $p + I' = 0$ in E/I' , so $\sum_{j=1}^n (e_j + I') = 0$ in E/I' , and since E/I' is a graded ring and each $e_j + I'$ has gradation j it follows that $e_i + I' = 0$, hence $e_i \in I$, for every I . \square

7. IDEALS GENERATED BY POWERS OF MATRICES ARE ‘SMALL’

Let R be a ring and $R[x]$ be the polynomial ring over R . Given a matrix M with entries from $R[x]$, let $P(M)$ denote the linear space spanned by coefficients of polynomials which are entries of matrix M .

We will say that a ring R and a linear space S satisfy **Assumption 1** when

1. R is the free algebra (without identity) generated by two elements a, b over a field F .
2. S is the linear F -subspace of $F[a, b]$ spanned by elements ab and $a \cdot b^2$.

Lemma 23. *Let R, S satisfy assumption 1, and let $R[x]$ be the polynomial ring over R in one variable x . Let m be a natural number and let M be a matrix with entries from $S^m \cdot F[x]$. Let $C = \{c_1, c_2, \dots, c_j\}$, where c_1, \dots, c_j are nonzero elements from $F \cdot S^m$. Let $r = r_1 r_2 r_3$ where r_i is a product of n_i elements from set C , for $i = 1, 2, 3$, with $n_i \geq 0$.*

If $r \in P(M^{n_1+n_2+n_3})$ then $r_i \in P(M^{n_i})$ for $i = 1, 2, 3$.

Proof. We can write $M^n = M^{n_1} \cdot M^{n_2} \cdot M^{n_3}$. Therefore $P(M^n) \subseteq P(M^{n_1}) \cdot P(M^{n_2}) \cdot P(M^{n_3})$. Hence $r = r_1 \cdot r_2 \cdot r_3 \in P(M^{n_1})P(M^{n_2})P(M^{n_3})$ and $r_i \in F \cdot S^{m \cdot n_i}, P(M^{n_i}) \subseteq F \cdot S^{m \cdot n_i}$; it follows that $r_i \in P(M^{n_i})$ for $i = 1, 2, 3$.

Indeed, if $r_j \notin P(M^{n_j})$ for some j , then we would find a linear mapping $f : F \cdot S^{m \cdot n_j} \rightarrow F \cdot S^{m \cdot n_j}$ such that $f(P(M^{n_j})) = 0$ and $f(r_j) \neq 0$, and we can apply this mapping to the above inclusion at appropriate places, obtaining a contradiction. \square

Definition 24. Let F be an infinite field. Let R, S satisfy Assumption 1. Let $f : F \cdot S^m \rightarrow F$ be a F -linear mapping. For every i we can extend the mapping f to the mapping $f : F \cdot S^{m \cdot i} \rightarrow F$ by defining $f(w_1 \cdots w_i) = f(w_1) \cdots f(w_i)$ for $w_1, \dots, w_i \in S^m$, and then extending it by linearity to all elements from $F \cdot S^{m \cdot i}$.

Let M be a matrix with entries $m_{i,j}$; by $f(M)$ we will denote the matrix with corresponding entries equal to $f(m_{i,j})$.

Similarly, if $g : F \cdot S^m \rightarrow F \cdot S^m$ is a linear mapping then for every i we can extend the mapping g to the mapping $g : F \cdot S^{m \cdot i} \rightarrow F \cdot S^{m \cdot i}$ by defining $g(w_1 \cdots w_i) = g(w_1) \cdots g(w_i)$ for $w_1, \dots, w_i \in S^m$ and then extending it by linearity to all elements from $F \cdot S^{m \cdot i}$.

Lemma 25. Let notation be as in Lemma 23. Assume that $f(c_i) \neq 0$ for all $i \leq j$, where $c_i \in S^m$ are as in Lemma 23. Let $f : F \cdot S^m \rightarrow F$ be a linear mapping, and $f : F \cdot S^{m \cdot n_2} \rightarrow F$ be as in Definition 24. Let F be an infinite field, and let $n = n_1 + n_2 + n_3$ be natural numbers. If $r = r_1 r_2 r_3 \in P(M^n)$ then

$$r_1 f(r_2) r_3 \in P(M^{n_1} f(M^{n_2}) M^{n_3}).$$

Proof. Let M be a d by d matrix and let $a_{i,j}(x)$ be the polynomial which is at the i, j entry of M . Notice that $a_{i,j}(x) = \sum_{k,l \leq d} b_{i,k}(x) c_{k,l}(x) d_{l,j}(x)$, where $b_{i,k}$ is the i, k entry of matrix M^{n_1} , $c_{k,l}(x)$ is the k, l entry of M^{n_2} and $d_{l,j}$ is the l, j entry of matrix M^{n_3} . Similarly $n_{i,j}(x) = \sum_{k,l \leq d} b_{i,k}(x) f(c_{k,l}(x)) d_{l,j}(x)$ is the i, j -th entry of matrix $M^{n_1} f(M^{n_2}) M^{n_3}$.

Notice that since F is infinite, then by a Vandermonde matrix argument we get that $P(M^n) = \sum_{i,j \leq d, p \in F} F \cdot a_{i,j}(p)$ and

$$P(M^{n_1} f(M^{n_2}) M^{n_3}) = \sum_{i,j \leq d, p \in F} F \cdot n_{i,j}(p)$$

If $r = r_1 r_2 r_3 \in P(M^n)$ then $r \in \sum_{i,j \leq d, p \in F} F \cdot a_{i,j}(p)$ hence

$$r_1 r_2 r_3 \in \text{span}_{p \in F, i, j \leq d} \sum_{k, l \leq d} b_{i,k}(p) c_{k,l}(p) d_{l,j}(p).$$

If we apply the mapping f as in Definition 24 at appropriate places we get that

$$r_1 f(r_2) r_3 \in \text{span}_{p \in F, i, j \leq d} \sum_{k, l \leq d} b_{i,k}(p) f(c_{k,l}(p)) d_{l,j}(p).$$

Recall that $n_{i,j}(x) = \sum_{k,l \leq d} b_{i,k}(x)f(c_{k,l}(x))d_{l,j}(x)$ is the i, j -th entry of matrix $M^{n_1}f(M^{n_2})M^{n_3}$. Therefore the linear space spanned by elements

$$\sum_{k,l \leq d} b_{i,k}(p)f(c_{k,l}(p))d_{l,j}(p)$$

for $p \in F$ equals the space spanned by $n_{i,j}(p)$ for $p \in P$. We have shown at the beginning of this proof that the latter space equals $P(M^{n_1}f(M^{n_2})M^{n_3})$. Therefore $r_1f(r_2)r_3 \in \text{span}_{i,j \leq d, p \in F} n_{i,j}(p) \subseteq P(M^{n_1}f(M^{n_2})M^{n_3})$. \square

Lemma 26. *Let R be an F -algebra. Let P be a linear space spanned by coefficients of polynomials $h_i(x) \in R[x]$ for $i = 1, 2, \dots$. Then for arbitrary non-zero polynomial $g(x)$ from $F[x]$ the linear space Q spanned by polynomials $g(x)h_i(x)$ equals P .*

Proof. Clearly $P \subseteq Q$. Let P_i denote the space spanned by coefficients by x^i of polynomials $h_1(x), h_2(x), \dots$

By calculating the coefficient by the smallest power of x in polynomials $g(x)h_i(x)$ we get that $P_0 \subseteq Q$. By then calculating the coefficient by the second-smallest power of x in $g(x)h_i(x)$ we get that $P_1 \in Q + P_0 \subseteq Q$. Continuing in this way we get $P_i \subseteq Q + P_0 + \dots + P_{i-1}$, so $P_i \subseteq Q$ for every i . It follows that $P \subseteq Q$. \square

Lemma 27. *Let notation be as in Definition 24 and Lemma 25. Let t be a natural number and let M be a d by d matrix. Let $n_1, n_2, n_3 \geq 0$. Then for all $i \geq 1$ we have*

$$P(M^{n_1}f(M^{n_2})M^{n_3}) \subseteq \sum_{i=1}^{d+1} P(M^{n_1}f(M)^iM^{n_3}).$$

Proof. Let $b_{i,k}$ denote the i, k entry of matrix M^{n_1} , $c_{k,l}(x)$ denote the k, l entry of M^{n_2} and $d_{l,j}$ denote the l, j entry of matrix M^{n_3} .

Let $n_{i,j}(x)$ be the i, j -th entry of matrix $M^{n_1}f(M^{n_2})M^{n_3}$, then $n_{i,j}(x) = \sum_{k,l \leq d} b_{i,k}(x)f(c_{k,l}(x))d_{l,j}(x)$.

Notice that $f(M)$ is a matrix with coefficients from $F[x]$. Every matrix with entries from the field of rational functions $F\{x\}$ in variable x satisfies its characteristic polynomial. It follows that there are polynomials $f_i(x)$ such that

$$\sum_{i=1}^{d+1} f_i(x)f(M)^i = 0$$

with $f_{d+1}(x)$ nonzero.

Therefore for every n there is a polynomial $g_n(x) \in F[x]$ such that $g_n(x)f(M)^n \in \sum_{i=1}^{d+1} F[x] \cdot f(M)^i$.

By Lemma 26, $P(M^{n_1} f(M^{n_2}) M^{n_3}) = P(M^{n_1} g(x) f(M^{n_2}) M^{n_3})$. We know that $g_n(x) f(M^{n_2}) \subseteq \sum_{i=1}^{d+1} F[x] f(M)^i$, hence

$$P(M^{n_1} g(x) f(M^{n_2}) M^{n_3}) \subseteq \sum_{i=1}^{d+1} P(F[x] \cdot M^{n_1} f(M)^i M^{n_3}).$$

By Lemma 26 we get $P(M^{n_1} f(M^{n_2}) M^{n_3}) \subseteq \sum_{i=1}^{d+1} P(M^{n_1} f(M)^i M^{n_3})$. □

We will say that M, R, S, m, d, α satisfy **Assumption 2** if

1. R, S satisfy Assumption 1 and m, d, α are natural numbers.
2. M is a d by d matrix with entries from $S^m \cdot F[x]$. Moreover,

$$M \subseteq R + Rx + Rx^2 + \dots Rx^\alpha.$$

Let c_1, \dots, c_j be linearly independent elements from $F \cdot S^m$ and denote $C = \{c_1, \dots, c_j\}$. Let $v = c_{i_1} \dots c_{i_j}$ and $v' = c_{k_1} \dots c_{k_j}$ for some i_1, \dots, i_j , and v, v' some k_1, \dots, k_j . We will say that words v and v' are distinct if $i_l \neq j_l$ for some $l \leq j$.

Let r be a product of elements from the set C . We say that w is a subword of degree n in r if w is a product of n elements from C and $r = vwv'$ for some v, v' which are also products of elements from C .

Lemma 28. *Let F be an infinite field. M, R, S, m, d, α satisfy Assumption 2. Let q be natural number. Let c_1, \dots, c_j be linearly independent elements from $F \cdot S^m$, and let r, r' be products of q elements from the set $C = \{c_1, \dots, c_j\}$. If $n \geq 8d^3 \cdot (\alpha + 1)$ and r has at least n pairwise distinct subwords of length n , and r' has at least n pairwise distinct subwords of length n then $r \cdot r' \notin P(M^t)$, for any t .*

Proof. Suppose on the contrary that $rr' \in P(M^t)$. Let p_1, \dots, p_n be subwords of r of degree n , and q_1, q_2, \dots, q_n be subwords of r' of length n . Then there are $s_{i,k}$ such that $p_i s_{i,k} q_k$ is a subword of $r \cdot r'$ for all $i, k \leq n$. By Lemma 23, $r \cdot r' \in P(M^t)$ implies $p_i s_{i,k} q_k \in P(M^{m_{i,k}})$ for some $m_{i,k}$. Let $f : F \cdot S^m \rightarrow F$ be a linear mapping, and $f : F \cdot S^{m \cdot n_2} \rightarrow F$ be as in Definition 24. We can choose f that $f(c_i) \neq 0$ for $i = 1, 2, \dots, j$ and hence $f(s_{i,k}) \neq 0$ for every $i, k \leq n$. By Lemma 25,

$$p_i f(s_{i,k}) q_k \in P(M^n f(M^{m_{i,k}-2n}) M^n).$$

By Lemma 27, $p_i q_k \in \sum_{l=1}^{d+1} P(M^n f(M^l) M^n)$. Notice that the linear space $\sum_{l=1}^{d+1} P(M^n f(M^l) M^n)$ has dimension smaller than $d^2(d+1) \cdot (2n\alpha + 2)$.

Observe now that since p_i and q_i are products of n elements c_i , and c_i 's are linearly independent over F , then elements $p_i q_k$ are linearly independent over F . Therefore elements $p_i q_k$ span a linear space over field F of dimension at least n^2 . Hence $n^2 \leq d^2(d+1) \cdot (2n\alpha + 2) < 8d^3 \cdot (\alpha + 1) \cdot n$, a contradiction. □

8. SUBSPACES E_i AND E'_i

Let F be a countable field and let R, S satisfy Assumption 1. Since F is countable, we can enumerate finite matrices with entries in $S \cdot F[x]$ as X_1, X_2, \dots . We can assume that the matrix X_i is of dimension at most i and X_i had entries in $F \cdot (S + Sx + S \cdot x^2 + \dots + S \cdot x^i)$ for every i , if necessary taking $X_i = 0$ for some i .

The following is similar to Theorem 5 from [41].

Theorem 29. *Let F be a countable field, let R, S satisfy Assumption 1 and let matrices X_1, X_2, \dots be as above. Let $0 < m_1 < m_2 < \dots$ be a sequence of natural numbers such that m_i is a power of two and $2^{2^{m_i}}$ divides m_{i+1} for all $i \geq 1$. Denote $R(m) = F \cdot S^m$ for every m . Let E'_i be the linear space spanned by all coefficients of polynomials which are entries of the matrix $X_i^{2^{2^{m_i}}}$ and let*

$$E_i = \sum_{j=0}^{\infty} R(j \cdot m_{i+1}) E'_i S R.$$

Then there is an ideal I in R contained in $\sum_{i=1}^{\infty} E_i + bE_i + b^2E_i + \langle a^2 \rangle + \langle b^3 \rangle$ and such that R/I is a nil ring, where $\langle a^2 \rangle, \langle b^3 \rangle$ denote ideals in R generated by elements a^2 and b^3 .

Proof. Observe first that the ideal I_k of R generated by coefficients of polynomials which are entries of the matrices $X_k^{2^{m_{k+1}+2}}$ is contained in the subspace $E_k + bE_k + b^2E_k$. It follows because entries of every matrix X_k have degree one in the subring generated by S with elements of S of degree one. In general if $n > m_{k+1} + 2^{2^{m_k}} + 1$ then coefficients of polynomials which are entries of matrix X_k^n belong to $R(i)E'_k R(1)R$ for every $0 \leq i < n - m_{k+1} - 1$.

Define $I = \sum_{i=1}^{\infty} I_k + \langle a^2 \rangle + \langle b^3 \rangle$; then $I \subseteq \sum_{i=1}^{\infty} E_i + bE_i + b^2E_i + \langle a^2 \rangle + \langle b^3 \rangle$. Observe also that, by Lemma 21, R/I is a nil ring. \square

Lemma 30. *Let F be an infinite field and let $T \subseteq L$ be finitely dimensional F -linear spaces. Let $c_1, c_2, \dots, c_j \in L$ and $c_1, c_2, \dots, c_j \notin T$. Then there is a linear mapping $f : L \rightarrow F$ such that T is contained in the kernel of f and c_1, c_2, \dots, c_j are not contained in the kernel of f .*

Moreover there is a linear mapping $g : L \rightarrow L$ such that T is contained in the kernel of g and c_1, c_2, \dots, c_j are not contained in the kernel of g and the image of g has co-dimension 1 in L .

Proof. Let Q be a maximal linear space such that $c_1, c_2, \dots, c_m \notin Q$ and $T \subseteq Q$. We will show that L/Q is a one dimensional linear space. Suppose on the contrary, then there are two elements $x + Q, y + Q$ in L/Q which are linearly independent

over F . By maximality of Q , we get that there are $\alpha \neq \beta$ such that linear spaces $Q + F \cdot (x + \alpha \cdot y)$ and $Q + F \cdot (x + \beta \cdot y)$ both contain some element c_i . Then $c_i - t_1 \cdot (x + \alpha \cdot y) \in Q$ and $c_i - t_2 \cdot (x + \beta \cdot y) \in Q$ for some $t_1, t_2 \in F$. It follows that $t_1 \cdot (x + \alpha \cdot y) - t_2 \cdot (x + \beta \cdot y) \in Q$, a contradiction since $x + Q$ and $y + Q$ are linearly independent in L/Q . We can now take $g : L \rightarrow L$ to be such that $f(q) = 0$ for every $q \in Q$, and the image of g has co-dimension one in L . Observe that the natural linear mapping $f : L \rightarrow L/Q$; then L/Q has dimension 1, so L/Q is isomorphic as a linear space to F . In this way we can define the mapping f . \square

9. WORDS w_i

In this chapter we give some supporting results on some monomials related to Engel elements.

Definition 31. *Let M be the free monoid generated by elements A, B, A', B' . Define inductively a sequence of infinite monomials W_i as follows.*

$$W_1 = A, W_2 = ABA', W_3 = W_2B\bar{W}_2$$

and

$$W_{n+1} = W_nB\bar{W}_n,$$

where $\bar{W} = (\alpha(W))^{op}$ where $\alpha : M \rightarrow M$ is a homomorphism of monoids such that $\alpha(A) = A'$, $\alpha(A') = A$, $\alpha(B) = B'$ and $\alpha(B') = B$. Recall that if $x_i \in \{A, A', B, B'\}$ then $(x_1x_2 \dots x_n)^{op} = x_nx_{n-1} \dots x_1$. We will sometimes refer to monomials from M as words. Observe also that for every $n > 0$,

$$\bar{W}_{n+1} = W_nB'\bar{W}_n.$$

Lemma 32. *For every i , W_i has length $2^i - 1$.*

Proof. By induction on i . \square

Lemma 33. *Let notation be as in Lemma 32. Let $\alpha(c, v)$ denote the number of occurrences of c in a word $v \in M$. Then for every $n \geq 1$ $\alpha(A, W_{n+1}) = \alpha(A', W_{n+1}) = \alpha(B, W_{n+1}) = 2^{n-1}$ and $\alpha(B', W_{n+1}) = 2^{n-1} - 1$. Moreover in the word W_{n+1} after elements A, A' appears either B or B' ; and after elements B, B' appears either A or A' .*

Proof. Observe that for $v = W_nB'$ we have $\alpha(A, v) + \alpha(A', v) = \alpha(B, v) + \alpha(B', v)$, as in v after A and A' always appears either B or B' and vice-versa. We will now proceed with the proof of our theorem by induction on n . For $n = 1$ we have

$W_2 = ABA'$, so the result holds in this case. Suppose that the result holds for some $n \geq 1$; we will show that it holds for $n + 1$.

Observe that for every word v $\alpha(A, v) = \alpha(A', \bar{v})$ and $\alpha(B, v) = \alpha(B', \bar{v})$. Recall that $W_{n+1} = W_n B \bar{W}_n$, hence $\alpha(A, W_{n+1}) = \alpha(A', W_{n+1})$, $\alpha(B, W_{n+1}) = \alpha(B', W_{n+1}) + 1$. Let $v = W_{n+1} B'$; then by the above $\alpha(A, v) = \alpha(A', v)$ and $\alpha(B, v) = \alpha(B', v)$.

By Lemma 32, $\alpha(A, v) + \alpha(A', v) + \alpha(B, v) + \alpha(B', v) = 2^{n+1}$. Hence $\alpha(A, v) = \alpha(A', v) = \alpha(B, v) = \alpha(B', v) = 2^{n-1}$. The result follows. \square

Definition 34. Let M be a free monoid generated by elements A, A', B, B' . Let $v \in M$. Let R be a ring, and let $x, y, z, t \in R$. By $v(x, y, z, t)$ we will denote an element of R obtained by substituting $A = x$, $B = y$, $A' = z$ and $B' = t$ in the word v .

Let R be a ring; by R^1 we will denote the ring with the identity element which is the usual extension of R .

Definition 35. Let F be a field, and let R be a ring generated by elements a, b , such that $a^2 = 0, b^3 = 0$. Then $1 + a$ and $1 + b$ are invertible elements in R^1 . Denote

$$\begin{aligned} v_1 &= [1 + a, 1 + b] = (1 + a)(1 + b)(1 + a)^{-1}(1 + b)^{-1}, \\ v_2 &= [v_1, 1 + b] = v_1(1 + b)v_1^{-1}(1 + b)^{-1}, \\ v_{n+1} &= [v_n, 1 + b] = v_n(1 + b)v_n^{-1}(1 + b)^{-1}. \end{aligned}$$

Lemma 36. Let notation be as in Definition 35. Denote $z_{n+1} = v_n \cdot (1 + b)$. Then $z_2 = (1 + a)(1 + b)(1 + a)^{-1} = (1 + a)(1 + b)(1 - a)$ and

$$z_{n+1} = z_n \cdot (1 + b) \cdot z_n^{-1}$$

for $n = 2, 3, \dots$. Moreover,

$$(z_{n+1})^{-1} = z_n \cdot (1 - b + b^2) \cdot z_n^{-1}.$$

Proof. It is clear that $z_2 = v_1(1 + a) = (1 + a)(1 + b)(1 + a)^{-1} = (1 + a)(1 + b)(1 - a)$. By the definition of v_{n+1} we get $z_{n+1} = v_{n+2}(1 + b) = v_{n+1} \cdot (1 + b) \cdot v_{n+1}^{-1} = [v_n(1 + b)] \cdot (1 + b) \cdot [(1 + b)^{-1}v_n^{-1}] = z_n \cdot (1 + b) \cdot z_n^{-1}$. This implies $z_{n+1}^{-1} = z_n(1 - b + b^2)z_n^{-1}$. \square

Notation. Let notation be as in Definition 31. Let R be a ring generated by elements a, b such that $a^2 = 0$ and $b^3 = 0$. In what follows we will use the following notation.

$$w_n = W_n(a, b, -a, b^2 - b), \bar{w}_n = \bar{W}_n(a, b, -a, b^2 - b).$$

Lemma 37. *Let notation be as above and as in Definition 31. Then $w_1 = a$, $w_2 = -aba$, $\bar{w}_1 = -a$ and $\bar{w}_2 = -a(b^2 - b)a$. Moreover, for every n , $w_{n+1} = w_n \cdot b \cdot \bar{w}_n$ and $\bar{w}_{n+1} = w_n \cdot (b^2 - b) \cdot \bar{w}_n$.*

Proof. It follows from Definition 31 by induction on n . □

Lemma 38. *Let notation be as in Lemmas 37, 36. Let $T(j)$ be the linear space spanned by all monomials $x_1x_2 \dots x_n$ such that $x_i \in \{a, b\}$ and the cardinality of the set $\{1 \leq i \leq n-1 : x_i x_{i+1} \in \{ab, ba\}\}$ is at most j (n is arbitrary).*

Then for every $n \geq 2$,

$$z_n - w_n - 1, z_n^{-1} - \bar{w}_n - 1 \in T(2^n - 3)$$

and

$$w_n, \bar{w}_n \in F \cdot S^{2^{n-1}-1}a \subseteq T(2^n - 2).$$

Proof. We will proceed by induction. We will use Lemmas 37 and 36.

For $n = 2$ we have $w_2 = -aba \subseteq F \cdot Sa \subseteq T(2) = T(2^2 - 2)$, and $z_2 = (1+a)(1+b)(1-a) = -aba + (ab + b - a^2 - ba) + 1$. Therefore, $z_2 - w_2 - 1 = ab + b - a^2 - ba \in T(1) = T(2^2 - 3)$, as required.

Recall that $\bar{w}_2 = -a(b^2 - b)a \in F \cdot Sa \subseteq T(2)$. Observe also that $z_2^{-1} = (1+a)(1-b+b^2)(1-a) = -a(b^2 - b)a + a(b^2 - b) + (b^2 - b) - a^2 - (b^2 - b)a$ hence $z_2^{-1} - \bar{w}_2 - 1 \in T(1)$, as required.

Suppose now that the result holds for some number $n \geq 2$; we will prove it for $n+1$. Observe that for all i, j , we have $T(i)T(j) \subseteq T(i+j+1)$, as we have only one more place when the words from $T(i)$ and $T(j)$ meet where a change from a to b or from b to a can appear.

Observe that $w_{n+1} = w_n b \bar{w}_n$, hence by the inductive assumption $w_{i+1} \in T(2^n - 2)T(0)T(2^n - 2) \subseteq T(2^n - 2 + 0 + 2^n - 2 + 2) = T(2^{n+1} - 2)$.

We have $z_{n+1} = z_n(1+b)z_n^{-1}$, hence for some $q, q' \in T(2^n - 3) + F$

$$z_{n+1} = (w_n + q)(1+b)(\bar{w}_n + q').$$

By the inductive assumption $q(1+b)q' \subseteq T(2^n - 3)T(0)T(2^n - 3) \subseteq T(2^n - 3 + 0 + 2^n - 3 + 2) = T(2^{n+1} - 4)$. Similarly $q(1+b)\bar{w}_n \subseteq T(2^n - 3)T(0)T(2^n - 2) \subseteq T(2^{n+1} - 3)$ and $w_n(1+b)q' \subseteq T(2^{n+1} - 3)$. Consequently, $z_{n+1} - w_{n+1} - 1 \in T(2^{n+1} - 3)$. The proof that $w_{n+1}^{-1} \in T(2^{n+1} - 2)$ and $z_{n+1}^{-1} - \bar{w}_{n+1} \in T(2^{n+1} - 3)$ is analogous.

Notice also that by Lemma 33 and by Notation before Lemma 37

$$w_n, \bar{w}_n \in S^{2^{n-1}-1}a \subseteq T(2^n - 2).$$

□

Lemma 39. *Let F be an infinite field. Let R, S satisfy Assumption 1. Let I be the ideal of R generated by elements from sets F_1, F_2, \dots , where $F_i \subseteq F \cdot S^i$ for every i and by elements a^2 and b^3 . Let $w'_{n+1} = W_{n+1}(a, b, -a, b^2 - b)$, $w'_{n+2} = W_{n+2}(a, b, -a, b^2 - b)$. Let v_n, z_{n+1} be as in Definition 35 and Lemma 36 applied for ring $\bar{R} = R / \langle a^2, b^2 \rangle$. Let I' be the ideal of \bar{R} which is generated by images in $R / \langle a^2, b^3 \rangle$ of elements from sets F_1, F_2, \dots .*

If $v'_n - 1 \in I'$ for some $n > 1$ then $w'_{n+1}b \in I$, and hence $w'_{n+2} \in I$.

Proof. Let $z'_{n+1} \in R$ be such that the image of z'_{n+1} in $R / \langle a^2, b^2 \rangle$ is z_{n+1} . Observe that $v_n - 1 \in I'$ is equivalent to $z_{n+1} - 1 \in I'$ and hence $z'_{n+1} - (b+1) \in I$. Observe that $R/I' = \bar{R}/I$. We can write $z'_{n+1} - (b+1) = p + q + t + t' + t'' \in I$ for some $p \in ab \cdot R^1$, $q \in bR^1$, $t \in F[a]$, $t' \in F[b]$, $t'' \in a^2R^1$. Notice that $t'' \in I$; hence $p + q + t + t' \in I$. By Lemma 22 [1.] we get $p, q, t, t' \in I$. By Lemma 38 we get $p = w'_{n+1} + v$ for some $v \in T(2^{n+1} - 3) \cap ab \cdot R^1$. We can write $v = v' + v'' + v'''$ where $v' \in Rba$, $v'' \in Rb$ and $v''' \in Ra^2$. Then $v''' \in I$ and $w_{n+1} + v' + v'' \in I$. By Lemma 22 [2.] we get that $w_{n+1} + v' \in I$. Then $w_{n+1}b + v'b \in I$ and $v'b \in T(2^{n+1} - 2)$. Notice that $vb \in ab \cdot R^1 \cap R^1 \cdot ab$. It follows that $v'b \in F + F \cdot S + F \cdot S^2 + \dots + F \cdot S^{2^n - 1}$. By Lemma 33 $w'_{n+1}b \subseteq F \cdot S^{2^n}$. By Lemma 22 [3.] $w'_{n+1}b \in I$. Observe also that $w'_{n+1}b \in I$ implies $w'_{n+2} = w'_{n+1}bw'_{n+1} \in I$. \square

10. COMBINATORICS OF WORDS

The following lemma immediately follows from the proof of Theorem 1.3.13, pp.22, [31]. We repeat a slightly modified proof from Theorem 1.3.13, pp.22, [31].

Lemma 40. *Let n be a natural number. Let w be an infinite word which has less than n subwords of degree n ; then $u = cddd\dots$ for some words c, d such that c has length smaller than $n!$ and d has length $n!$.*

Moreover if u is a finite word which has less than n subwords of degree n then $u = cddd\dots d$ for some words c, d such that c has length smaller than $n! + n!$ and d has length $n!$.

Proof. Notice that for some $m \leq n$, w has the same number of words of length m and $m + 1$. Hence for every subword v_1 in w of length m we have exactly one possibility of a subword u_1 of w which has length $m + 1$ and which contains v_1 as the beginning. Let v_2 be the ending of length m in u_1 . We can apply the same reasoning to v_2 instead of v_1 and find word u_2 containing v_2 as the beginning. After at most n steps we get $v_i = v_j$ for some $i \neq j$ and then from the step j $v_{k+j} = v_k$ for each k . Therefore $w = v_1 \dots v_{i-1} v v v v v v \dots$ where v is some word of length $t < n + 1$. Because t divides $n!$ we get the result for w .

If u is a finite word then we can apply similar reasoning. \square

We can then get the following.

Lemma 41. *Let R, S satisfy assumption 1. Let $v' = c_1 \dots c_m$ with each $c_i \in F \cdot S^m$ for some m . We say that w is a subword of v' of length n if $v' = uwu'$ where for some k , $u = c_1 \dots c_k$, $w = c_{k+1}c_{k+2} \dots c_{k+n}$, $u' = c_{k+n+1} \dots c_m$ (u and u' may be trivial words). Let n be a natural number. Assume that v' has less than n subwords of degree n then $u = cddd \dots d$ for some subwords c, d such that c has length smaller than $n! + n!$ and d has length $n!$.*

Proof. It follows from Lemma 40. \square

Theorem 42. *Let $0 < m_1 < m_2 < \dots$ be such that each m_i is a power of two, $2^{2^{m_i}} < m_{i+1}$ and $2^{2^{m_i}}$ divides m_{i+1} for every i . Let $R(i), S, E_1$ be as in Theorem 29. Assume that $2^{m_1} > 17! \cdot 10$. Then $w_n b \notin E_1$ for any n , where $w_n = W_n(a, b, -a, b^2 - b)$.*

Proof. Suppose, on the contrary, that $w_n = W_n(a, b, -a, b^2 - b)b \in E_1$. We can assume that $n > m_2$, since $w_i b \in E_1$ implies $w_{i+1} b \in E_1$, by the definition of W_i and F_1 . Recall that $R(2^{n-1}) = F \cdot S^{2^{n-1}}$. Since $w_n b \in R(2^{n-1})$, by Lemma 32 and m_2 is a power of two, we can assume that 2^{n-1} is divisible by m_2 (because we can take larger n if needed by the argument from the first lines of this proof).

Denote $w = w_n b$. We can write $w = w'_1 \dots w'_t$ with each $w'_i \in R(m_2)$. Since $w \in E_1$ it follows that some $w'_k \in E_1 \cap R(m_2)$. Then $w'_k = v_1 \dots v_{m_2}$ for $v_i \in R(m_1)$. Denote $\alpha = 2^{2^{m_1}}$, then $v_1 \dots v_\alpha \in E'_1$, where E'_1 is the linear space spanned by all coefficients which are entries of the matrix $X_1^{2^{2^{m_1}}}$, by Theorem 29.

Write $v_1 \dots v_\alpha = vv'$, where $v, v' \in R(\alpha/2)$ and $\alpha = 2^{2^{m_1}}$. Notice that $vv' \in P(X_1^j)$ for $j = 2^{2^{m_1}}$. By Lemma 28 we get that for every $n > 8d^3 \cdot (\alpha + 1)$ either v or v' has less than n subwords of length n ; without restraining generality we can assume that it happens for v' (by a subword we mean a product $v_i v_{i+1} \dots v_j$ for some $i \leq j$).

In our case $\alpha = 1$, $d = 1$ because of assumptions on matrix $M = X_1$ (before Theorem 29), so we can take $n = 17$. By Lemma 41 $v' = cddd \dots d$ for some words c, d such that c has length smaller than $17! + 17!$ and d has length $17!$.

By assumption $2^{2^{m_1}} = 2^k$ for some k . By the definition of words W_i we get that $vv' = w_{k+1}b$ and $vv' = w_k b \bar{w}_k b$. It follows that $v = w_k b$ and $v' = \bar{w}_k b$.

Observe now that $v' = \bar{w}_k b$ implies $v' = w_{k-1}(b^2 - b)\bar{w}_{k-1}b$.

By Lemma 33 we can write $w_{k-1} = ab_{2^{k-2}-1} \cdot b_{2^{k-2}-1}a \cdots ab_2 \cdot ab_1a$ where $b_i \in \{b, b^2 - b\}$. Then $v' = w_{k-1}(b^2 - b)\bar{w}_{k-1}b$ yields

$$v' = (ab_{2^{k-2}-1} \cdot ab_{2^{k-2}-2} \cdots ab_2 \cdot ab_1 \cdot a(b^2 - b))(a\bar{b}_1 \cdot a\bar{b}_2 \cdots a\bar{b}_{2^{k-2}-1} \cdot ab),$$

where $\bar{b} = b^2 - b$ and $b^2 - b = b$. By Lemma 41 applied to matrix v' and to $m = 1$ we get that $v' = cddd \dots d$ for some words $c \in R(\alpha)$, $d \in R(17!)$ where c has length smaller than $17! + 17!$, so $\alpha < 17! + 17!$.

By the assumptions of our theorem $k > 17! \cdot 10$. We can write

$$v' = w_{k-1}(b^2 - b)\bar{w}_{k-1}b = s \cdot p \cdot q \cdot r \cdot t$$

for some $s, t \in R$ and some $p, q, r \in R(17!)$ with $spq = w_{k-1}(b^2 - b)$. Notice that then $s \in R(q)$ for some $q > 17! \cdot 2$. Then because $v' = cddd \dots$ and d has length $17!$ we get $p = q = r$. Recall that $spq = w_{k-1}(b^2 - b)$, and by the above

$$s \cdot p \cdot q = (ab_{2^{k-2}-1} \cdot ab_{2^{k-2}-2} \cdots ab_2 \cdot ab_1 \cdot a(b^2 - b)).$$

Consequently, $q = ab_{17!-1}a \dots b_2ab_1a(b^2 - b)$, $r = a\bar{b}_1a\bar{b}_2a \dots a\bar{b}_{17!}$, and $p = p'ab_{17!}$, for some p' . Recall that $p = r$; it follows that $\bar{b}_{17!} = b_{17!}$, which is impossible, as $b_{17!} \in \{b, b^2 - b\}$ and $\bar{b} = b^2 - b$ and $b^2 - b = b$. We have obtained a contradiction. \square

11. MAPPING T

In this section we will use the following notation

$$w_t = W_t(a, b, -a, b^2 - b), \bar{w}_t = \bar{W}_t(a, b, -a, b^2 - b).$$

Recall that by Lemma 32 we have

$$w_k b, w_k(b^2 - b) \in R(2^{k-1}).$$

Recall that, given matrix M with entries in $R[x]$, by $P(M)$ we denote the linear space spanned by coefficients of polynomials which are entries of matrix M . The linear spaces E_i and E'_i are as in Theorem 29.

Let R, S satisfy Assumption 1, by S -monomial we will mean a product of elements from $S = \{ab, ab^2\}$.

Lemma 43. *Let F be a field and let R, S satisfy Assumption 1. Let n, j be natural numbers. Let d_1, \dots, d_j be S -monomials from $R_{m_{n+1}}$ such that $d_1, \dots, d_j \notin \sum_{k=1}^n E_k$. Then there is a linear mapping $T' : R(m_{n+1}) \rightarrow R(m_{n+1})$ such that*

$$T' \left(\sum_{k=1}^n E_k \cap R(m_{n+1}) \right) = 0.$$

Moreover there are nonzero S -monomials $m_1 \dots m_j$ and nonzero $\alpha_1, \dots, \alpha_j \in F$ such that $T'(d_k) = \alpha_k \cdot m_k$ for all $k \leq j$. Moreover, there is a linear mapping $L : R(m_{n+1} - 1) \rightarrow R(m_{n+1} - 1)$ such that $T'(uv) = L(u)v$ for all S -monomians u, v with $v \in R(1)$ and $u \in R(m_{n+1} - 1)$.

Proof. By the definition of sets E_k , $R(m_{i+1}) \cap \sum_{k=1}^i E_k = P' \cdot R(1)$ for some linear space P' . Each d_k can be written as $d_k = c_k e_k$ where c_k, e_k are S -monomials, and where $e_k \in R(1)$. It follows that $c_k \notin P$ for each k . We apply Lemma 30 for $L = R(m_{i+1} - 1)$ and $T = P'$ to get linear mapping $f : R(m_{i+1} - 1) \rightarrow R(m_{i+1} - 1)$ such that $f(c_k) \neq 0$ for every k , and the image of f has codimension 1 in $R(m_{i+1} - 1)$. There is a monomial $m \neq 0$ such that $m \in \text{Im}(f)$, and since f has codimension one, then the image of f is $F \cdot m$. Hence $f(c_k)$ is a non-zero multiply of a monomial for every k . Moreover by Lemma 30 we have $f(P') = 0$. Therefore mapping $T'(uv) = L(u)v$ defined for monomials u, v with $v \in R(1)$, $u \in R(m_{i+1} - 1)$ and extended by linearity to all elements of $R(m_{i+1})$ satisfies the thesis of our theorem. \square

Definition 44. Let $T' : R(m_{n+1}) \rightarrow R(m_{n+1})$ be a mapping as in Lemma 43. For every j , we can extend the mapping T' to the mapping $T : R(j \cdot m_{n+1}) \rightarrow R(j \cdot m_{n+1})$ by defining $T(w_1 \dots w_j) = T'(w_1) \dots T'(w_j)$ for $w_1, \dots, w_j \in R(m_{n+1})$, and then extending it by linearity to all elements from $R(j \cdot m_{n+1})$.

Let M be a matrix with entries $m_{i,j}$, by $T(M)$ we will denote the matrix with corresponding entries equal to $T(m_{i,j})$.

Lemma 45. Let $n \geq 1$ be a natural number. Let $0 < m_1 < m_2 < \dots$ be such that each m_i is a power of two, $2^{2^{m_i}} < m_{i+1}$ and $2^{2^{m_i}}$ divides m_{i+1} for every i . Let T be a mapping satisfying assertions of Definition 44 and let

$$M = T(X_{n+1}^{m_{n+1}})$$

where X_{n+1} is as in Theorem 29. Denote $\beta = 2^{2^{m_{n+1}}} / m_{n+1}$ and $k = 2^{m_{n+1}} - 1$. Suppose that $w_j b \notin \sum_{i=1}^n E_i$ for every j , and $w_t b \in \sum_{i=1}^{n+1} E_i$ for some t . Then there is $w \in \{w_k b, w_k(b^2 - b), \bar{w}_k b, \bar{w}_k(b^2 - b)\}$ such that $T(w) \in P(M^\beta)$.

Proof. Observe that we can assume that t is arbitrarily large, since $w_i b \in \sum_{i=1}^\infty E_i$ implies $w_{i+1} b \in \sum_{i=1}^\infty E_i$ by the definition of words w_i . Therefore we can assume that $t > m_{n+2}$. Observe that since $w_t b \in \sum_{i=1}^{n+1} E_i$ then $w_t b \in (\sum_{i=1}^{n+1} E_i \cap R(m_{n+2}))^l$ for some l . By the definition of words $w_i b = W_i(a, b, -a, b^2 - b)b$ and by Lemma 32 we see that $w_t b = u_1 \dots u_l$ with each $u_i \in R(m_{n+2})$ (since length of

$w_t b$ is 2^t we can do it since $2^t \geq m_{n+2}$). It follows that for some ξ we have

$$u_\xi \in \sum_{i=1}^{n+1} E_i \cap R_{m_{j+2}}.$$

By the definition of words $w_i = W_i(a, b, -a, b^2 - b)$ it follows that $u_\xi \in Z_\gamma$ where

$$Z_\gamma = \{w_\gamma b, w_\gamma(b^2 - b), \bar{w}_\gamma b, \bar{w}_\gamma(b^2 - b)\}$$

where $2^{\gamma-1} = m_{n+2}$. By Theorem 29

$$u_\xi = v_1 \dots v_{m_{n+2}/m_{n+1}}$$

for $v_i \in R(m_{n+1})$. Observe that $v_1 \in E'_{n+1} + \sum_{i=1}^n E_i$, since otherwise $u_\xi \in \sum_{i=1}^n E_i \cap R_{m_{j+2}}$, and hence $w_t b \in \sum_{i=1}^n E_i \cap R_{m_{j+2}}$ contradicting the inductive assumption (where E'_{n+1} is the linear space spanned by all coefficients which are entries of the matrix $X_{n+1}^{2^{2^{m_{n+1}}}}$). Observe that $T(\sum_{i=1}^n E_i \cap R(m_{n+2})) = 0$ since by Lemma 43, $T'(\sum_{i=1}^n E_i \cap R(m_{n+1})) = 0$. It follows that

$$T(u_\xi) = T(v_1 v_2 \dots v_{m_{n+2}/m_{n+1}}) \in T(E_{n+1}).$$

Notice that by the definition of the mapping T we get

$$T(v_1)T(v_2) \dots T(v_{m_{n+2}/m_{n+1}}) = T(v_1 v_2 \dots v_{m_{n+2}/m_{n+1}}) \in T(E_{n+1}).$$

By the definition of E_{n+1} it follows that

$$T(v_1 \dots v_\beta) \in T(E'_{n+1}),$$

where $\beta = 2^{2^{m_{n+1}}}/m_{n+1}$. Notice that $T(E'_{n+1})$ is the linear space spanned by coefficients of matrix $T(X_{n+1}^{m_{n+1} \cdot \beta})$. Observe that $T(X_{n+1}^{m_{n+1} \cdot \beta}) = T(X_{n+1}^{m_{n+1}})^\beta$.

Therefore

$$T(v_1) \dots T(v_\beta) = T(v_1 \dots v_\beta) \in P(M^\beta)$$

where $M = T(X_{n+1}^{m_{n+1}})$. Therefore $w = u_\xi$ satisfies thesis of our theorem. \square

Theorem 46. *Let $0 < m_1 < m_2 < \dots$ be such that each m_i is a power of two, $2^{2^{m_i}} < m_{i+1}$ and $2^{2^{m_i}}$ divides m_{i+1} for every i . Assume that $2^{m_1} > 17! \cdot 10$. Let R, I satisfy assumptions of Theorem 29 for these m_i . Let $R(i), S$ be as in Theorem 29. Let n be a natural number. Then $w_t b \notin \sum_{i=1}^n E_i$ for any t .*

Proof. We proceed by induction on n . For $n = 1$ the result holds by Theorem 42. Suppose now that $w_j b \notin \sum_{i=1}^n E_i$ for any j . We will show that $w_j b \notin \sum_{i=1}^{n+1} E_i$ for some j . Suppose on the contrary that $w_t b \in \sum_{i=1}^{n+1} E_i$ for some t , then by the definition of words $w_i, w_{t+j} b \in \sum_{i=1}^n E_i$ for every $j \geq 0$.

Let w be as in Lemma 46. Without restricting the generality we can assume that $w = w_k b$, the proof in the other cases is similar. By the definition of words W_i we get that $w = v_1 \cdots v_\beta$ for some $v_1, \dots, v_\beta \in R(m_{n+1})$, moreover each v_i is an S -monomial. Observe that for each i , $v_i \notin \sum_{i=1}^n E_i$ as otherwise $w_t b \in \sum_{i=1}^n E_i$ contradicting the inductive assumption.

Let T' be a linear mapping as in Lemma 43 applied for S -monomials $d_i = v_i$ for $i = 1, 2, \dots, \beta$ and let T be a mapping constructed using T' as in Definition 44. By Lemma 46 we know that $T(w) \in P(M^\beta)$.

Let $c_1, c_2, \dots, c_j \in R(m_{n+1})$ be distinct S -monomials which form a basis of the linear space $F \cdot v_1 + \dots + F \cdot v_\beta$. Then every $v_k = c_{i_k} \alpha_k$ for some i_k and some $0 \neq \alpha \in F$ and v_i are S -monomials. Then $w = w' \cdot \alpha$ where $w' = c_{i_1} \dots c_{i_\beta} \cdot \alpha$ (since c_i, v_i are S -monomials). It follows that $T(w') \in P(M^\beta)$. Notice that by Definition 44, $T(c_i) = T'(c_i)$, for every i .

Denote

$$v = T'(v_1) \dots T'(v_{\beta/2}), v' = T(v_{\beta/2+1}) \dots T(v_\beta)$$

and

$$u = T'(c_{i_1}) \dots T'(c_{i_{\beta/2}}), u' = T(c_{i_{\beta/2+1}}) \dots T(c_{i_\beta}),$$

then $T(w') = T(c_{i_1} \dots c_{i_\beta}) = uu'$ and $u, u' \in R(2^{2^{m_{n+1}}}/2) = R(m_{n+1})^{\beta/2}$. Moreover there are $0 \neq \beta, \beta' \in F$ such that $v = \beta \cdot u$ and $v' = \beta' \cdot u'$.

By Lemma 28 we get that for $n' \geq 8d^3 \cdot (\alpha + 1)$ either u or u' has less than m subwords of length n' ; without restraining generality we can assume that it happens for u' . In our case $\alpha = (n + 1) \cdot m_{n+1}$, $d = n + 1$ because of assumptions on matrix $M = T(X_{n+1}^{m_{n+1}})$ (before Lemma 28), so we can take $n' = 8(n+2)^4 m_{n+1}$, so v' has less than n' subwords of degree n' . By Lemma 41 we get that

$$u' = cddd \dots d$$

for some words c, d such that c has length smaller than $n'! + n'!$ and d has length $n'!$, so $d \in R(n'! \cdot m_{n+1})$ and $c \in R(l \cdot m_{n+1})$ for some $l < n'! + n'!$. It follows that

$$v' = \beta' \cdot cddd \dots d.$$

Clearly $2^{2^{m_{n+1}}} = 2^k$ for some k . By the definition of words W_i we get that $w \in \{w_{k+1}b, w_{k+1}(b^2 - b), \bar{w}_{k+1}b, \bar{w}_{k+1}(b^2 - b)\}$. Therefore

$$vv' \in \{T(w_{k+1}b), T(w_{k+1}(b^2 - b)), T(\bar{w}_{k+1}b), T(\bar{w}_{k+1}(b^2 - b))\}.$$

Without restricting the generality we can assume that $vv' = T(w_{k+1}b)$, the proof in the other cases is similar. Therefore $vv' = T(w_k b \bar{w}_k b)$. It follows that $v = T(w_k b)$

and $v' = T(\bar{w}_k b)$. Observe now that $v' = T(\bar{w}_k b)$ implies

$$v' = T(w_{k-1}(b^2 - b)\bar{w}_{k-1}b).$$

By Lemma 33 we can write $w_{k-1} = ab_{2^{k-2}-1} \cdot b_{2^{k-2}-2}a \cdots ab_2 \cdot ab_1a$ where $b_i \in \{b, b^2 - b\}$. Then $v' = T(w_{k-1}(b^2 - b)\bar{w}_{k-2}b)$ gives

$$v' = T(ab_{2^{k-2}-1} \cdot ab_{2^{k-2}-2} \cdots ab_2 \cdot ab_1 \cdot a(b^2 - b)(\bar{a}b_1 \cdot \bar{a}b_2 \cdots \bar{a}b_{2^{k-2}-1} \cdot ab)),$$

where $\bar{b} = b^2 - b$ and $b^2 \bar{b} = b$.

Recall that $v' = \beta' \cdot cddd \dots d$ for some words c, d such that c has length smaller than $n'! + n'$ and d has length $n'!$ where $n' = 8(n+2)^4 m_{n+1}$. By the assumptions $v' \in R(2^{2^{m_{n+1}}}/2) = R(m_{n+1})^{\beta/2}$. Observe that $\beta/2 > n'! \cdot 10 \cdot m_{n+1}$. Therefore we can write

$$\bar{w}_{k-1}(b^2 - b)w_{k-1}b = s \cdot p \cdot q \cdot r \cdot t$$

for some $s, t \in R$ and some $p, q, r \in R(n'! \cdot m_{n+1})$ with $s \cdot p \cdot q = w_{k-1}(b^2 - b)$.

Notice that then $s \in R(q)$ for some $q > n'! \cdot 2$. Then

$$v' = T(w_{k-1}(b^2 - b)\bar{w}_{k-1}b) = T(s \cdot p \cdot q \cdot r \cdot t).$$

Notice that $spq = w_{k-1}(b^2 - b)$ implies $spq \in R(\gamma)$ for some γ divisible by m_{n+1} , as m_{n+1} is a power of two and $w_{k-1}(b^2 - b) \in R(2^{2^{m_{n+1}}}/4)$. By the definition of T , $v' = T(spq)T(qr)$. Because $p, q, r \in R(m! \cdot m_{j+1})$ we get

$$v' = T(s)T(p)T(q)T(r)T(t).$$

Then because $v' = cddd \dots \beta'$ and d has length $m!$ we get $\gamma \cdot T(p) = \gamma' \cdot T(q) = \gamma'' \cdot T(r) = m$ for some $0 \neq \gamma, \gamma', \gamma'' \in F$ and some S -monomial m . Denote $\xi = n'! \cdot m_{j+1}$, then $q = ab_{\xi-1}a \dots b_2 ab_1 a(b^2 - b)$, $r = \bar{a}b_1 \bar{a}b_2 a \dots \bar{a}b_\xi$, and $p = p' ab_\xi$, for some p' (for some $b_i \in \{b, b^2 - b\}$ where $\bar{b} = b^2 - b$ and $b^2 \bar{b} = b$). Recall that $\gamma \cdot T(p) = \gamma'' \cdot T(r)$. Recall also that by Lemma 43 we have $T(p) = s \cdot ab_\xi$ and $T(r) = s' \cdot ab\bar{b}_\xi$ for some s, s' ; it follows that $\bar{b}_\xi = b_\xi$, which is impossible, as $b_\xi \in \{b, b^2 - b\}$ and $\bar{b} = b^2 - b$ and $b^2 \bar{b} = b$. We have obtained a contradiction. \square

Theorem 47. *There is a nil ring R such that the adjoint group of R^o is not an Engel group. Moreover R can be taken to be an algebra over an arbitrary countable field.*

Proof. Suppose first that F is an infinite field. Let $0 < m_1 < m_2 < \dots$ be such that each m_i is a power of two, $2^{2^{m_i}} < m_{i+1}$ and $2^{2^{m_i}}$ divides m_{i+1} for every i . Assume moreover that $2^{m_1} > 17! \cdot 10$. Let R, I satisfy assumptions of

Theorem 29 for these m_i . By Theorem 46 we have $w_nb \notin \sum_{i=1}^j E_i$ for any n, j , where $w_n = W_n(a, b, -a, b^2 - b)$. By Theorem 29 we see that $I \subseteq \sum_{i=1}^{\infty} E_i + bE_i + b^2E_i + \langle a^2 \rangle + \langle b^3 \rangle$. Suppose that $w_nb \in I$ for some n , then there is $w \in \sum_{i=1}^{\infty} E_i$ such that $w_nb - w \in bR + \langle a^2 \rangle + \langle b^3 \rangle$. Observe that $(bR + \langle a^2 \rangle + \langle b^3 \rangle) \cap (\sum_{i=1}^{\infty} E_i) = 0$ and so $w_nb - w = 0$ so $w_nb \in \sum_{i=1}^{\infty} E_i$, a contradiction. It follows that $w_nb \notin I$ for every n .

By Lemma 39 we have $v_n - 1 \notin I'$ for every n , and hence $(R/I)^\circ$ is not an Engel group (I' is as in Lemma 39). By Theorem 29, R/I is nil. Therefore R/I is a nil algebra over field F such that the adjoint group R/I° of this algebra is not an Engel group.

If F is a finite field then we proceed in the following way: Let \bar{F} be the algebraic closure of F , then \bar{F} is infinite. Hence there is a nil algebra A over \bar{F} such that the adjoint group A° is not nil. Let $x, y \in A$ be such that $[x[\dots[x[x, y]]]] \neq 1$ (n copies of x) for every n . Let A' be the smallest subring of A containing x, y and such that such that if $r \in A'$ then $f \cdot r \in A'$ for every $f \in F$. Then A' is an F -algebra which is generated as F -algebra by x, y and which is nil. Since $x, y \in A'$ then the adjoint group A'° is not an Engel group. \square

Proof of Corollary 9. Let $(R, +, \cdot)$ be a ring such that R is nil and R° is not an Engel group. Assume moreover that R is an algebra over a finite field of cardinality p for some prime number p . Let $(R, +, \circ)$ be the associated brace, so $a \circ b = a \cdot b + a + b$ for all $a, b \in R$, and the addition is the same as in the ring R . It follows that $(R, +, \circ)$ satisfies the thesis of Corollary 9.

12. ZELMANOV'S QUESTION

Observe that in the case of algebras over uncountable fields we have the following result analogous to Lemma 20.

Lemma 48. *Let F be an uncountable field and let F' be a countable subfield of F . Let R be an F -algebra generated by elements a, b , and suppose that $a^2 = 0$ and $b^3 = 0$. Let $R[x_1, x_2, \dots]$ be the polynomial ring over R in infinitely many commuting variables x_1, x_2, \dots . Let $F'[x_1, x_2, \dots]$ be the polynomial ring over F' in infinitely many commuting variables x_1, x_2, \dots .*

Let S' be the F' -linear space spanned by elements abx_i and ab^2x_i for $0 \leq i$.

If all finite matrices with entries from S' are nilpotent then R is a Jacobson radical ring.

Proof. Observe that if all finite matrices with entries from S are nilpotent, then after substituting arbitrary elements $\alpha_1, \alpha_2, \dots \in F$ for variables x_1, x_2, \dots we get that all matrices with entries from F' -linear space spanned by elements $ab^i\alpha_j$ are nilpotent. Therefore every matrix with entries in the F -linear space spanned by elements ab and ab^2 is nilpotent. By Lemma 20, R is a Jacobson radical F -algebra. \square

We recall Amitsur's theorem.

Theorem 49. *Let R be a finitely generated algebra over an uncountable field. If R is a Jacobson radical algebra then R is nil.*

We will now use Lemma 48 to give an analogon of Theorem 29.

Let F be a field and let F' be a countable subfield of F . Let R be as in Lemma 48 and let S be the F' -linear space spanned by elements ab and ab^2 . Let $X = \{x_1, x_2, \dots\}$ and let $F'[X]$ denote the polynomial ring over F' in infinitely many variables x_1, x_2, \dots

We can enumerate all finite matrices with entries from $S \cdot F'[X]$ as X_1, X_2, \dots . We can assume that the matrix X_i is of dimension at most i and X_i has entries in $S \cdot y_1 + S \cdot y_2 + \dots S \cdot y_i$ for some $y_1, y_2, \dots, y_i \in F'[X]$ (if necessary taking $X_i = 0$ for some i). The following theorem has the same proof as Theorem 29.

Theorem 50. *Let F be an uncountable field, and let R, S and the matrices X_1, X_2, \dots be as above. Let $0 < m_1 < m_2 < \dots$ be a sequence of natural numbers such that $2^{2^{m_i}}$ divides m_{i+1} for every $i \geq 1$. Denote $R(m) = F \cdot S^m$ for every m . Let E'_i be the linear space spanned by all coefficients of polynomials which are entries of the matrix $X_i^{2^{2^{m_i}}}$ and let*

$$E_i = \sum_{j=0}^{\infty} R(j \cdot m_{i+1}) E'_i S R.$$

Then there is an ideal I in R contained in $\sum_{i=1}^{\infty} E_i + bE_i + b^2E_i + \langle a^2 \rangle + \langle b^3 \rangle$ and such that R/I is a nil ring, where $\langle a^2 \rangle, \langle b^3 \rangle$ denote ideals in R generated by elements a^2 and b^3 .

Proof. Observe first that the ideal I_k of R generated by coefficients of polynomials which are entries of the matrices $X_k^{2^{2^{m_{k+1}}}}$ is contained in the subspace $E_k + bE_k + b^2E_k$. It follows because entries of every matrix X_k have degree one in the subring generated by S with elements of S of degree one. In general if $n > m_{k+1} + 2^{2^{m_k}} + 1$ then every entry of matrix X_k^n belongs to $R(i)E'_k R(1)R$ for every $0 \leq i < n - m_{k+1} - 1$.

Define $I = \sum_{i=1}^{\infty} I_k + \langle a^2 \rangle + \langle b^3 \rangle$, then $I \subseteq \sum_{i=1}^{\infty} E_i + bE_i + b^2E_i + \langle a^2 \rangle + \langle b^3 \rangle$. Observe also that, by Lemma 48 and Theorem 49, R/I is a nil ring. \square

We will say that $M, R, F', S, r_1, r_2, m, d, \alpha$ satisfy **Assumption 3** if

1. R, F' are as in Lemma 48 and S is the F' -linear space spanned by elements ab and ab^2 , and m, d, α are natural numbers.
2. M is a d by d matrix with entries from $S^m \cdot F[X]$. Moreover,

$$M \subseteq R + R \cdot y_1 + R \cdot y_2 + \dots R \cdot y_{\alpha},$$

for some $y_1, y_2, \dots, y_{\alpha} \in F[X]$, where $X = \{x_1, x_2, \dots\}$ is an infinite set.

We now propose a generalisation of Lemma 28.

Lemma 51. *Let F be an infinite field. M, R, S, m, d, α satisfy Assumption 3. Let q be natural number. Let c_1, \dots, c_k be linearly independent elements from $F \cdot S^m$, and let $r_1, r_2, \dots, r_{\alpha+1}$ be products of q elements from the set $C = \{c_1, \dots, c_k\}$. If $n > d^{\alpha+2} \cdot (\alpha + 1)^{\alpha}$ and for each i , r_i has more than n subwords of length n then $r_1 \cdot r_2 \cdots r_{\alpha+1} \notin P(M^j)$, for any j .*

We say that w is a subword of degree n in r , if w is a product of n elements from C , and $r = vwv'$ for some v, v' which are also products of elements from C .

Proof. Denote $r = r_1 \cdots r_{\alpha+1}$. Suppose on the contrary that $r \in P(M^j)$. Let $p_{1,i}, \dots, p_{n,i}$ be subwords of degree n in r_i for all $i \leq \alpha + 1$. Fix numbers $\beta(1), \dots, \beta(\alpha + 1) \leq \alpha + 1$, then there are q_1, \dots, q_{α} such that

$$s = \left(\prod_{i=1}^{\alpha} p_{\beta(i),i} q_i \right) p_{\beta(\alpha+1),\alpha+1}$$

is a subword of r .

By Lemma 23, $r \in P(M^j)$ implies $s \in P(M^{j'})$, for some j' .

Let $f : F \cdot S^m \rightarrow F$ be a linear map such that $f(c_i) \neq 0$, and let $f : F \cdot S^{m \cdot n'} \rightarrow F$ be as in Definition 24. Then $f(c_{i_1} c_{i_2} \dots c_{i_{n'}}) = f(c_1) \dots f(c_{n'}) \neq 0$ for every choice of $i_1, i_2, \dots, i_{n'} \leq k$. By Lemma 25 applied several times we get that

$$\left(\prod_{i=1}^{\alpha} p_{\beta(i),i} f(q_i) \right) p_{\beta(\alpha+1),\alpha+1} \in P\left(\left(\prod_{i=1}^{\alpha} M^n f(M^{\deg q_i}) M^n \right) \right).$$

By an analogous argument to Lemma 27,

$$p_{\beta(1),1} p_{\beta(2),2} \cdots p_{\beta(\alpha+1),\alpha+1} \in P\left(\left(\prod_{i=1}^{\alpha} M^n Q_i \right) M^n \right),$$

where $Q_i = \sum_{j=1}^{d+1} F \cdot f(M^j)$.

Notice that the linear space $(\prod_{i=1}^{\alpha} M^n Q_i)M^n$ has dimension smaller than d^α . Therefore $P((\prod_{i=1}^{\alpha} M^n Q_i)M^n)$ has dimension at most $d^\alpha \cdot d^2 \cdot ((\alpha + 1) \cdot n)^\alpha$.

Observe now that since each $p_{i,j}$ is a product of n elements from the set C , then elements

$$p_{\beta(1),1} p_{\beta(2),2} \cdots p_{\beta(\alpha+1),\alpha+1}$$

span linear space over field F of dimension at least $n^{\alpha+1}$. Hence $d^\alpha \cdot d^2 \cdot ((\alpha + 1) \cdot n)^\alpha < n^{\alpha+1}$, a contradiction with the assumptions on n . \square

Proof of Theorem 5. The proof is the same as the proof of Theorem 47 but instead of Lemma 28 we use Lemma 51; and instead of Theorem 29 we use Theorem 50.

ACKNOWLEDGEMENTS

I would like to thank Efim Zelmanov and Gunnar Traustason for their helpful comments on Engel groups, and Tatyana Gateva-Ivanova for introducing me to research into braided groups, braces and solutions to the Yang-Baxter equation. I am also grateful to Jan Krempa and Jan Okniński for their helpful suggestions. I am very grateful to Ferran Cedó for reading the manuscript and for his very useful comments. This research was supported with ERC grant 320974 and I would like to thank them for their support.

REFERENCES

- [1] Alireza Abdollahi, *Engel elements in groups*, Groups St Andrews 2009 in Bath, London Mathematical Society Lecture Notes Series, 94-117.
- [2] Adel Alahmadi, Hamed Alsulami, S.K. Jain, Efim Zelmanov, *Finite generation of Lie algebras associated with associative algebras*, Journal of Algebra 426 (2015) 69–78.
- [3] B. Amberg, O. Dickenschied, Ya.P. Sysak Subgroups of the adjoint group of a radical ring, *Canad. J. Math.* 50 (1998), 3–15.
- [4] B. Amberg, L. Kazarin, *Nilpotent p -algebras and factorized p -groups*, Proceedings of "Groups St. Andrews 2005", London Math. Society Lecture Note Series 339, Vol 1, Cambridge Univ. Press (2007), 130–147.
- [5] B. Amberg, Ya.P. Sysak, *Radical rings and products of groups*, Groups St Andrews 1997 in Bath: Edited by C. M. Campbell, 1–19.
- [6] B. Amberg, Ya. P. Sysak, *Radical rings with soluble adjoint group*, *J. Algebra* 247 (2002), 692-702
- [7] Bernhard Amberg, Yaroslav P. Sysak, *Radical Rings with Engel Conditions*, Journal of Algebra Volume 231, Issue 1, 1 September 2000, Pages 36–373.
- [8] J.C. Ault, J.F. Watters, *Circle groups of nilpotent rings*, *Am. Math. Mon.* 80, (1973), 48–52.

- [9] David Bachiller, *Counterexample to a conjecture about braces*, arXiv:1507.02137v1 [math.GR] 8 July 2015.
- [10] F. Catino, M. M. Miccolia, Ya. P. Sysak, *On the Adjoint Group of Semiprime Rings*, Communications in Algebra, Volume 35, Issue 1, 2006, 265–270.
- [11] F. Catino, R. Rizzo, *Regular subgroups of the affine group and radical circle algebras*, Bull. Aust. Math. Soc. 79 (2009), 103–107.
- [12] Ferran Cedó, *Braces and the Yang-Baxter equation*, (joint work with Eric Jespers and Jan Okniński), slides from the talk in Stuttgart, June 2012.
- [13] Ferran Cedó, Eric Jespers, Jan Okniński, *Braces and the Yang-Baxter Equation*, Communications in Mathematical Physics April 2014, Volume 327, Issue 1, pp 101–116.
- [14] Ferran Cedó, Eric Jespers, Jan Okniński, *Retractability of set theoretic solutions of the Yang-Baxter equation*, Adv. Math. 224 (2010), 2472–2484.
- [15] F. Cedó, E. Jespers and A. del Rio, *Involutive Yang-Baxter Groups*, Trans. Amer. Math. Soc. 362 (2010), 2541–2558.
- [16] P. Etingof, T. Schedler, T. Solovov, *A set theoretical solutions to the quantum Yang-Baxter equation*, Duke. Math. J. 100 (1999), 169–209.
- [17] Tatiana Gateva-Ivanova, *Set-theoretic solutions of the Yang-Baxter equation, Braces, and Symmetric groups*, arXiv:1507.02602 [math.QA].
- [18] T. Gateva-Ivanova, *A combinatorial approach to the set-theoretic solutions of the Yang-Baxter equation*, J.Math.Phys. 45 (2004), 3828–3858.
- [19] T. Gateva-Ivanova, *A combinatorial approach to the set-theoretic solutions of the Yang-Baxter equation*, J. Math. Phys. 45 (2004), 3828–3858.
- [20] T. Gateva-Ivanova and S. Majid, *Matched pairs approach to set-theoretic solutions of the Yang-Baxter equation*, J. Algebra 319 (2008), 1462–1529.
- [21] T. Gateva-Ivanova and S. Majid, *Quantum spaces associated to multipermutation solutions of level two*, arXiv: 0806.2928.
- [22] T. Gateva-Ivanova, Peter Cameron, *Multipermutation solutions of the Yang-Baxter equation*, Comm. Math. Phys. 309 (2012), 583–621.
- [23] T. Gateva-Ivanova, and A. Smoktunowicz *Multipermutation solutions of the Yang-Baxter equation, Braces and Symmetric groups*, Preprint (2015).
- [24] T. Gateva-Ivanova and M. Van den Bergh, *Semigroups of I-type*, J. Algebra 206 (1998), 97112.
- [25] Hales, A.W., Passi, I.B.S.: *The second augmentation quotient of an integral group ring*. Arch. Math. (Basel) 31, (1978), 259–265
- [26] Natalia Iyudu, Stanislav Shkarin, *Finite dimensional semigroup quadratic algebras with the minimal number of relations*, Monatshefte für Mathematik, November 2012, Volume 168, Issue 2, pp 239–252.
- [27] E. Jespers and J. Okniński, *Monoids and groups of I-type*, Algebras Repres. Theory 8 (2005), 709729.
- [28] E. Jespers and J. Okniński, *Noetherian Semigroup Algebras*, Springer, Dordrecht, 2007.
- [29] T.Y. Lam, *A First Course in Noncommutative Rings*, Springer-Verlag, New York, 1991.
- [30] J. Lu, M. Yan and Y. Zhu, *On the set-theoretical Yang-Baxter equation*, Duke. Math. J. 104, 2000, 1–18.

- [31] M. Lothaire, Algebraic Combinatorics on Words, E-Book, DOI: <http://dx.doi.org/10.1017/CBO9781107326019>.
- [32] V. M. Petrogradsky, I.P. Shestakov, E. Zelmanov, *Nil graded self-similar algebras*, Groups Geom. Dyn. 4, No. 4, 873-900 (2010).
- [33] Boris Plotkin, *Notes on Engel groups and Engel elements in groups. Some generalizations*, arXiv:math/0406100 [math.GR].
- [34] D. J. S. Robinson, *A course on group theory*, Springer-Verlag, GTM Vol. 80 (1996).
- [35] W. Rump, *Modules over braces*, Algebra Discrete Math., 2006, Issue 2, 127-137.
- [36] W. Rump, *The brace of a classical group*, Note Mat. **34** no. 1, 115-14.
- [37] Wolfgang Rump, *Braces, radical rings, and the quantum YangBaxter equation*, Journal of Algebra Volume 307, Issue 1, 1 January 2007, Pages 153–170.
- [38] W. Rump, *A decomposition theorem for square-free unitary solutions of the Yang-Baxter equation*, Adv. Math. 101 (1990), no. 3, 583-591.
- [39] A. Shalev, *On associative algebras satisfying the Engel condition*, Israel J.Math. 67, (1989), 287–290.
- [40] I. P. Shestakov and E. Zelmanov, *Some examples of nil Lie algebras*, J. Eur. Math. Soc. (JEMS) 10 (2008), no. 2, 391-398.
- [41] Agata Smoktunowicz, *How far can we go with Amitsur's theorem*, arXiv:1504.01341v3 [math.RA].
- [42] Agata Smoktunowicz, *The Jacobson radical of rings with nilpotent homogeneous elements*, Bull. London Math. Soc. (2008) 40 (6): 917–928
- [43] Agata Smoktunowicz, *On Engel groups, nilpotent groups, rings, braces and the Yang-Baxter equation*, arXiv:1509.00420 [math.RA].
- [44] Ya. P. Sysak, *The adjoint group of radical rings and related questions*, Ischia Group Theory 2010: Proceedings of the Conference. Edited by Patrizia Longobardi, Mariagrazia Bianchi, Mercede Maj, 344–365.
- [45] Ya. P. Sysak, *Products of groups and local nearrings*, Note. Mat. 28 (2008), suppl. n. 2. 177-214.
- [46] G. Traustason, *Engel groups*, in Groups St Andrews 2009 in Bath, Volume II, 520-550. London Math. Soc. Lecture Note Ser. 388, Cambridge University Press, Cambridge 2011.
- [47] L. Vendramin, *Extensions of set-theoretic solutions of the Yang-Baxter equation and a conjecture of Gateva-Ivanova*, arXiv:1502.00790v3 [math.QA] 20 April 2015.
- [48] I. Wisliceny, *Konstruktion nilpotenter associativer*, Algebren mit wenig Relationen. Math. Nachr. 147, 7582 (1990).
- [49] E. I. Zelmanov, *On some problems of group theory and Lie algebra*, Mat. Sb. 66, No. 1 (1990), 159-167.
- [50] E. I. Zelmanov, *The solution of the restricted Burnside problem for groups of odd exponent*, Math. USSR. Izu. 36, No. 1 (1991), 41-60.
- [51] E. I. Zelmanov, *The solution of the restricted Burnside problem for 2- groups*, Mat. Sb. 182, No. 4 (1991), 568–592.

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, JAMES CLERK MAXWELL BUILDING, THE KINGS BUILDINGS, MAYFIELD ROAD EH9 3JZ, EDINBURGH

E-mail address: A.Smoktunowicz@ed.ac.uk