



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Association of polygenic risk for major psychiatric illness with subcortical volumes and white matter integrity in UK Biobank

Citation for published version:

Reus, L, Shen, X, Gibson, J, Wigmore, E, Ligthart, L, Adams, M, Davies, G, Cox, S, Hagenaars, S, Bastin, M, Deary, I, Sibley, H & McIntosh, A 2017, 'Association of polygenic risk for major psychiatric illness with subcortical volumes and white matter integrity in UK Biobank', *Scientific Reports*, vol. 7, 42140. <https://doi.org/10.1038/srep42140>

Digital Object Identifier (DOI):

[10.1038/srep42140](https://doi.org/10.1038/srep42140)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Scientific Reports

Publisher Rights Statement:

© The Author(s) 2017

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



SCIENTIFIC REPORTS



OPEN

Association of polygenic risk for major psychiatric illness with subcortical volumes and white matter integrity in UK Biobank

L. M. Reus¹, X. Shen¹, J. Gibson¹, E. Wigmore¹, L. Ligthart², M. J. Adams¹, G. Davies³, S. R. Cox^{3,4}, S. P. Hagenaars^{1,3,4}, M. E. Bastin³, I. J. Deary^{3,4}, H. C. Whalley¹ & A. M. McIntosh^{1,3}

Major depressive disorder (MDD), schizophrenia (SCZ) and bipolar disorder (BP) are common, disabling and heritable psychiatric diseases with a complex overlapping polygenic architecture. Individuals with these disorders, as well as their unaffected relatives, show widespread structural differences in corticostriatal and limbic networks. Structural variation in many of these brain regions is also heritable and polygenic but whether their genetic architecture overlaps with that of major psychiatric disorders is unknown. We sought to address this issue by examining the impact of polygenic risk of MDD, SCZ, and BP on subcortical brain volumes and white matter (WM) microstructure in a large single sample of neuroimaging data; the UK Biobank Imaging study. The first release of UK Biobank imaging data comprised participants with overlapping genetic data and subcortical volumes (N = 978) and WM measures (N = 816). The calculation of polygenic risk scores was based on genome-wide association study results generated by the Psychiatric Genomics Consortium. Our findings indicated no statistically significant associations between either subcortical volumes or WM microstructure, and polygenic risk for MDD, SCZ or BP. These findings suggest that subcortical brain volumes and WM microstructure may not be closely linked to the genetic mechanisms of major psychiatric disorders.

Major depressive disorder (MDD), schizophrenia (SCZ) and bipolar disorder (BP) are major psychiatric disorders affecting between 1% and 13% of the general population^{1,2}. Twin and family studies find a strong genetic contribution to all three of these disorders^{3,4} and genome-wide association studies (GWAS) suggest that these effects are conferred by the cumulative impact of many loci, each of small effect^{5–8}. Whilst progress in identifying the specific risk-conferring loci in SCZ has proved highly productive⁵, progress in BD and MDD has been slower^{6,7}. Several strategies may accelerate the discovery of risk-loci, including the use of quantitative traits such as brain structure and connectivity.

Brain structure and connectivity measures have been shown to be heritable quantitative traits, with high heritabilities reported for both subcortical volumes (44–88%)⁹ and white matter (WM) integrity (53–90%)¹⁰. Furthermore, several genetic variants that influence brain structure have been identified^{11,12}. Subcortical brain volume and WM microstructure abnormalities in corticostriatal and limbic networks have also been shown to distinguish individuals with SCZ, BD and MDD from controls in several studies, albeit with some inconsistencies^{13–18}. Differences between patients and controls in global measures of WM microstructure, diffusion tensor imaging (DTI) biomarkers, fractional anisotropy (FA), and mean diffusivity (MD), have been more consistently reported in these disorders^{19–24}, and could suggest wide-spread WM integrity reductions.

Together, these volumetric and WM brain differences potentially describe a network of abnormality associated with major psychiatric illness. This network may have an important mechanistic role in the development of these disorders. This possibility is supported by several studies that show an association between brain structure and genetic liability for psychiatric disorders, though due to the cost of imaging these samples are typically an order

¹Division of Psychiatry, University of Edinburgh, Royal Edinburgh Hospital, Edinburgh, EH10 5HF, United Kingdom.

²Department of Biological Psychology, Vrije Universiteit, Amsterdam, the Netherlands. ³Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, EH8 9JZ, United Kingdom. ⁴Department of Psychology, University of Edinburgh, Edinburgh, EH8 9JZ, United Kingdom. Correspondence and requests for materials should be addressed to H.C.W. (email: heather.whalley@ed.ac.uk)

of magnitude smaller in size than the GWAS that preceded them. Notably, Van Scheltinga *et al.*²⁵ ($N_{\text{SCZ}} = 152$, $N_{\text{healthy}} = 142$) found higher polygenic risk for SCZ to be modestly ($R^2 = 0.048$) associated with smaller total brain volume, while Caseras *et al.*²⁶ ($N_{\text{healthy}} = 274$) reported a negative association between polygenic risk scores (PGRS) for SCZ and BP, and globus pallidus and amygdala grey matter (GM) volumes. Similar results have been found for the association between less optimal WM microstructure and greater genetic liability for SCZ^{25,27,28}.

These associations have not been consistently reported; specifically, several studies have found no association between subcortical volumes and polygenic risk for MDD and SCZ^{29–31}. However, these studies have used differing methodologies (*i.e.*, multi-scanner study³¹ $N = 11,840$, from 22 studies) with relatively small sample sizes ($N_{\text{healthy}} = 438$, $N_{\text{healthy}} = 122$), which may contribute to inconsistencies in findings.

The current study sought to investigate the genetic relationship between major mental disorders and structural brain measures in a single large quality-controlled dataset using the same scanner and the same analysis pipelines. We used newly acquired data from the first release of the UK Biobank Imaging study to examine whether structural imaging measures, including subcortical volumes ($N = 978$) and WM microstructure measures ($N = 816$), were associated with genetic risk of MDD, SCZ, and BP³² (<http://www.ukbiobank.ac.uk>). We hypothesized that higher polygenic risk for these disorders would be associated with decreased volumetric and WM integrity measures in specifically corticostriatal and limbic networks.

Methods

Study population. The first release of UK Biobank imaging data consisted of 4,446 subjects ($N_{\text{females}}/N_{\text{males}} = 2,342/2,104$; mean age \pm s.d. = 55.52 ± 7.62 years; age range = 40–70 years) with quality-checked volumetric and DTI data for the analysis (conducted by UK Biobank, Brain Imaging Documentation V1.1, <http://www.ukbiobank.ac.uk>)³³. In the current study, subjects were excluded if they did not provide genetic data, or if they were known to have participated in studies from the Psychiatric Genomics Consortium (PGC) GWAS dataset as this was used to construct the polygenic risk scores (PGRS). In total, the volumetric analysis included 978 subjects ($N_{\text{females}}/N_{\text{males}} = 499/479$; mean age \pm s.d. = 55.32 ± 7.38 years; age range = 40–70 years) and the DTI analysis included 816 subjects ($N_{\text{females}}/N_{\text{males}} = 413/403$; mean age \pm s.d. = 55.49 ± 7.26 years; age range = 40–70 years). Sample overlap (individuals included in both the DTI and volumetric analysis) included $N = 816$ participants and the volumetric analysis included $N = 162$ extra participants. In addition, a sensitivity analysis was performed where we imposed internal imaging quality checks, excluding individuals whose subcortical volume or FA values were greater than three standard deviations above or below the mean for that measure. The sensitivity analysis, further referred to as the sample excluding outliers, included 892 (exclusion of 86 subjects; $N_{\text{females}}/N_{\text{males}} = 477/415$; mean age \pm s.d. = 55.29 ± 7.31 years; age range = 40–70 years) and 733 (exclusion of 83 subjects; $N_{\text{females}}/N_{\text{males}} = 373/360$; mean age \pm s.d. = 55.31 ± 7.22 years; age range = 40–70 years) subjects for the volumetric and DTI analysis respectively. Sample overlap for the sample excluding outliers included $N = 679$, and the volumetric analysis included 213 unique observations, whereas the DTI analysis included 54 unique observations.

This study has been approved by the National Health Service (NHS) Research Ethics Service (approval letter dated 17th June 2011, reference: 11/NW/0382), and by the UK Biobank Access Committee (Project #4844). All experiments were performed in accordance with guidelines and regulations from these committees. Written informed consent was obtained from each subject.

Genotyping and derivation of polygenic risk scores. Procedures for DNA collection and genotyping in UK Biobank have been described previously by Hagenaars *et al.*³⁴. Subjects were excluded from the PGRS analysis if they had a non-British ancestry, gender mismatch, relatedness ($r > 0.044$), and genotype missingness bigger than 2%.

PGRS for the three traits of interest (*i.e.*, MDD, SCZ, BP) were created using PRSice³⁵. For each subject PGRS were calculated by summing the number of risk alleles weighted by the strength of its association with the trait of interest. The strength of this association has been calculated previously by the PGC (PGC2 for SCZ)^{5,7,8}. Prior to calculating the PGRS, single nucleotide polymorphisms (SNPs) were excluded if they had a minor allele frequency less than 1%, deviated significantly from Hardy-Weinberg equilibrium ($p < 1 \times 10^{-6}$) in the total sample of founder individuals, or had a call rate of less than 99%. The remaining SNPs were used to calculate 15 multidimensional scaling (MDS) ancestry components (to account for population structure). For the main results, the SNP inclusion threshold was set as $p \leq 0.5$, as Purcell *et al.*³⁶ and studies following this work, have shown that this threshold is generally the most predictive of case-control status in subsequent studies^{5–8}. Using the $p \leq 0.5$ SNP inclusion threshold, 631,763, 5,422,836 and 1,283,324 SNPs (after LD pruning) were included for the calculation of MDD-PGRS, SCZ-PGRS and BP-PGRS, respectively. Results generated by a SNP inclusion threshold of $p \leq 0.01$, $p \leq 0.05$, $p \leq 0.1$ and $p \leq 1.0$ are reported in supplementary materials.

Image acquisition and Neuroimaging data pre-processing. Procedures for image acquisition and pre-processing have been performed by UK Biobank, and are available on the UK Biobank website (<http://www.ukbiobank.ac.uk/>), and have been documented previously^{33,37}. In short, images were collected in Cheadle Manchester on a Siemens Skyra 3.0 T scanner (Siemens Medical Solutions, Germany) with a 32-channel head coil (<https://www.healthcare.siemens.com/magnetic-resonance-imaging>). T1-weighted scans were acquired using a 3D MPRAGE (resolution 1 mm³ isotropic voxels) sequence, and diffusion-weighted (DW) scans were acquired with a monopolar Steejskal-Tanner pulse sequence and multi-shell acquisition ($b_0 = 0$ s/mm², $b = 1,000$ s/mm², $b = 2,000$ s/mm²). Images were preprocessed and analysed with the FMRIB Software Library (FSL) (<http://fsl.fmrib.ox.ac.uk/fsl>). More detailed information is reported in the supplementary materials.

Statistical analysis. All analyses were performed using R (version 3.2.3) in the Linux environment (R Development Core Team, 2010). To examine whether higher PGRS were associated with globally poor WM

microstructure, general components for FA (gFA) and MD (gMD) across the 27 WM tracts were calculated³⁷. Latent measures, explaining a portion of variance WM structure, were calculated for both FA and MD by extracting the scores on the first unrotated components, using a principal component analysis (PCA) on the 27 WM tracts. The first components accounted for a mean of 44.82% and 41.73% of total variance in FA and, 47.45% and 45.35% in MD, in the sample including and excluding outliers respectively (loadings of tracts on the first latent component >0.15). To examine whether higher PGRS were selectively associated with poor WM structure in a certain classification, this PCA was also performed separately for groups of different categories of WM tracts, including association (inferior fronto-occipital fasciculus, inferior longitudinal fasciculus, superior longitudinal fasciculus, uncinate fasciculus, cingulum), projection (forceps minor, forceps major, corticospinal tracts, acoustic radiation, medial lemniscus, middle cerebellar peduncle) and thalamic radiation fibers, separately³⁷ (sample including outliers: variance explained = 54.52%, 38.68% and 61.92% respectively for each subset, and loading of first component >0.33; sample excluding outliers: variance explained = 60.05%, 32.05% and 72.05% respectively for each subset and loading of first component >0.27).

For bilateral brain regions, associations between the three PGRS and each imaging measure were examined using a repeated measure linear mixed-effects analysis, modelling hemisphere as a random factor (where the PGRS * hemisphere interaction itself was found to be non-significant). For unilateral structures (*e.g.* total GM volume, forceps major, forceps minor and middle cerebellar peduncle), and those where there was a significant interaction between PGRS and hemisphere, associations were analysed as single structures without repeated measurements and without hemisphere as a separate term in the model.

For all models, additional fixed effects included age, age², gender, genotype batch, genotype array and 15 ancestry-informative MDS components. Intracranial volume, calculated as the sum of WM, GM and ventricular CSF, was also modelled as a fixed effect in the analysis of subcortical volumes in order to adjust for differences in overall brain size. PGRS and imaging measures were scaled to zero mean and unitary standard deviation. Standardised Beta values are reported throughout results. Results were corrected for multiple comparisons using a false discovery rate (FDR) correction³⁸. Nagelkerke's R² was used to estimate the proportion of variation explained by PGRS.

Post-hoc positive control analyses, with age as an independent variable and each of the brain measures as dependent variable, were performed in order to test the sensitivity of UK Biobank imaging data, by verifying that we could replicate the widely reported negative associations between many structural brain measures and age (including those reported previously in an overlapping sample in UK Biobank)³⁷. Fixed effects included gender for all structures, and side of hemisphere for lateralized structures only.

Results

PGRS and total brain and subcortical volumes. No significant associations were observed between PGRS (MDD, SCZ, or BP) at the threshold of $p \leq 0.5$ and total GM, WM or CSF volume, in samples including or excluding outliers (Table 1). Furthermore, no significant associations were found between any of the three sets of PGRS and total GM, WM and CSF volume for any of the other PGRS p -value thresholds (all $\beta < 0.054$; all $p_{\text{uncorrected}}$ and $p_{\text{FDR}} > 0.05$) (Table S1).

A modest negative association was observed between thalamus volumes and the polygenic risk for SCZ at the threshold of $p \leq 0.05$, $p \leq 0.1$, $p \leq 0.5$ and $p \leq 1.0$ (β range = -0.043 – -0.039 ; $p_{\text{uncorrected}} = 0.052$, $p_{\text{uncorrected}} = 0.030$, $p_{\text{uncorrected}} = 0.054$, $p_{\text{uncorrected}} = 0.048$ for all PGRS thresholds respectively) (Table 1 and S3). However, following FDR-correction no significant associations were found between PGRS (MDD, SCZ, or BP) at the threshold of $p \leq 0.5$ and all subcortical volumes, before and after exclusion of outliers (Table 1). These results were unchanged at the other p -value thresholds (Tables S2, S3 and S4). For subcortical volumetric measures, no significant PGRS * hemisphere interactions were found; therefore, all analyses of bilateral structures could be conducted using a repeated measures design with hemisphere as a repeated factor. Mean values of the subcortical volumes are presented in supplementary Table S15.

PGRS and general and tract-specific diffusion measures. Associations between PGRS (MDD, SCZ, BP) and measures of general WM microstructure (gFA and gMD) were not significant (gFA β range = -0.048 – -0.017 ; p range = 0.216 – 0.771 , gMD β range = -0.047 – -0.031 ; p range = 0.181 – 0.985) (Tables 2 and 3). Moreover, when gFA and gMD were calculated separately for projection, association, and thalamic radiation fibres, there remained no significant association between these measures and PGRS for MDD, SCZ and BP at $p \leq 0.5$ threshold (Tables 2 and 3). These findings were similar at other p -value thresholds (Tables S5 and S6).

Trend-wise associations were observed between some polygenic MDD risk scores and MD values in the anterior thalamic radiation, a group of WM fibers connecting frontal regions to anterior and middle nuclear groups of the thalamus ($\beta = 0.084$; p -FDR = 0.085 , $\beta = 0.088$; p -FDR = 0.061 , $\beta = 0.086$; p -FDR = 0.065 for PGRS threshold $p \leq 0.01$, $p \leq 0.05$ and $p \leq 0.5$ respectively) (Tables S9 and S10). Polygenic MDD, SCZ or BP risk was also not significantly associated with other individual tract-specific water diffusion measures in the sample including ($N = 816$) and excluding outliers ($N = 733$) (Tables S7 and S9). These results were again similar for PGRS at the other thresholds (Tables S8 and S10).

No significant PGRS * hemisphere interactions were found for the majority of individual tract water diffusion values, and therefore analyses on bilateral structures were conducted as a repeated measure. However, the SCZ-PGRS * hemisphere interaction was found to be significant for FA values in the parahippocampal portion of the cingulum (connecting orbitofrontal regions to the hippocampus) in the sample including outliers ($\beta = -0.052$; $p_{\text{corrected}} = 0.032$). We therefore conducted tests of this association with SCZ-PGRS separately for each lateralised structure for this region. However, these were also not significant for FA in either the left ($\beta = -0.040$;

	Including outliers (N = 978)					Excluding outliers (N = 892)				
	Beta: z ratio (S.D.)	t statistic	p -uncorr.	p-FDR	R ²	Beta: z ratio (S.D.)	t statistic	p -uncorr.	p-FDR	R ²
MDD-PGRS										
GM volume	0.001 (0.027)	0.049	0.961	1.000	0.000	-0.001 (0.029)	-0.051	0.959	1.000	0.000
WM volume	-0.007 (0.027)	-0.262	0.793	1.000	0.005	-0.026 (0.028)	-0.913	0.362	1.000	0.067
CSF volume	0.042 (0.030)	1.392	0.164	0.492	0.174	0.054 (0.031)	1.748	0.081	0.243	0.289
Caudate	0.026 (0.026)	0.970	0.332	1.000	0.066	0.031 (0.028)	1.119	0.264	1.000	0.099
Hippocampus	-0.002 (0.026)	-0.082	0.935	1.000	0.000	-0.003 (0.026)	-0.130	0.896	1.000	0.001
Pallidum	0.003 (0.027)	0.107	0.915	1.000	0.001	-0.013 (0.027)	-0.466	0.641	1.000	0.016
Thalamus	-0.012 (0.020)	-0.611	0.541	1.000	0.015	-0.025 (0.020)	-1.218	0.223	1.000	0.060
Amygdala	-0.000 (0.027)	-0.010	0.992	1.000	0.000	0.015 (0.027)	0.551	0.582	1.000	0.023
Nucleus accumbens	-0.019 (0.026)	-0.742	0.458	1.000	0.038	-0.030 (0.027)	-1.126	0.260	1.000	0.093
Putamen	0.007 (0.022)	0.309	0.757	1.000	0.005	-0.010 (0.023)	-0.451	0.652	1.000	0.011
SCZ-PGRS										
GM volume	-0.012 (0.028)	-0.444	0.657	1.000	0.001	-0.018 (0.029)	-0.613	0.540	1.000	0.033
WM volume	0.000 (0.027)	0.032	0.974	1.000	0.000	-0.004 (0.029)	-0.147	0.883	1.000	0.002
CSF volume	0.049 (0.030)	1.616	0.106	0.319	0.236	0.063 (0.031)	2.029	0.043	0.128	0.397
Caudate	-0.003 (0.026)	-0.101	0.919	1.000	0.001	-0.016 (0.028)	-0.559	0.577	1.000	0.025
Hippocampus	-0.001 (0.025)	-0.044	0.965	1.000	0.000	0.015 (0.026)	0.558	0.577	1.000	0.021
Pallidum	-0.015 (0.027)	-0.578	0.564	1.000	0.024	-0.022 (0.027)	-0.795	0.427	1.000	0.047
Thalamus	-0.039 (0.020)	-1.929	0.054	0.378	0.149	-0.039 (0.020)	-1.947	0.052	0.363	0.155
Amygdala	0.015 (0.026)	0.558	0.577	1.000	0.022	0.025 (0.028)	0.912	0.362	1.000	0.063
Nucleus accumbens	-0.024 (0.026)	-0.924	0.356	1.000	0.058	-0.023 (0.027)	-0.843	0.400	1.000	0.053
Putamen	0.005 (0.022)	0.211	0.833	1.000	0.002	-0.016 (0.023)	-0.672	0.502	1.000	0.025
BP-PGRS										
GM volume	0.003 (0.027)	0.093	0.926	1.000	0.001	-0.012 (0.029)	-0.405	0.686	1.000	0.013
WM volume	0.000 (0.026)	0.019	0.985	1.000	0.000	0.006 (0.028)	0.229	0.819	1.000	0.004
CSF volume	-0.007 (0.029)	-0.246	0.805	1.000	0.005	0.007 (0.030)	0.216	0.829	1.000	0.004
Caudate	0.017 (0.026)	0.653	0.514	1.000	0.029	0.007 (0.028)	0.245	0.806	1.000	0.005
Hippocampus	0.024 (0.025)	0.963	0.336	1.000	0.058	0.011 (0.026)	0.421	0.674	1.000	0.012
Pallidum	-0.000 (0.026)	-0.009	0.993	1.000	0.000	-0.023 (0.027)	-0.865	0.387	1.000	0.054
Thalamus	0.010 (0.020)	0.517	0.605	1.000	0.010	0.005 (0.020)	0.249	0.803	1.000	0.002
Amygdala	0.030 (0.026)	1.158	0.247	1.000	0.091	0.032 (0.027)	1.186	0.236	1.000	0.103
Nucleus accumbens	-0.008 (0.026)	-0.296	0.767	1.000	0.006	-0.017 (0.027)	-0.626	0.532	1.000	0.028
Putamen	0.025 (0.022)	1.133	0.257	1.000	0.061	0.015 (0.023)	0.659	0.510	1.000	0.023

Table 1. Association of PGRS (MDD, SCZ or BP) at ≤ 0.5 with total grey matter, white matter, cerebrospinal fluid, and subcortical volumes, in sample including and excluding outliers. MDD: major depressive disorder, SCZ: schizophrenia, BP: bipolar disorder, PGRS: polygenic risk scores, uncorr.: uncorrected, FDR: false discovery rate, GM: grey matter, WM: white matter, CSF: cerebrospinal fluid, S.D.: standard deviation. Controlled for age, age², gender, genotype batch and array, and 15 MDS components. Additional fixed effects for subcortical volumes include intracranial volume and side of hemisphere. R² = estimate of variance explained by PGRS in %.

$p_{\text{uncorrected}} = 0.262$) or right ($\beta = -0.079$; $p_{\text{uncorrected}} = 0.029$) parahippocampal portion of the cingulum, following FDR correction. Mean values of the tract-specific water diffusion measures are presented in supplementary Table S16.

Age-related effects in structural brain measures. We additionally performed a post-hoc validation analysis in order to verify that despite our lack of polygenic associations we could indeed replicate the widely reported negative associations between structural brain measures and age. We conducted our analyses using age as an independent variable and each of the brain measures as dependent variable. As expected, total GM and WM volumes showed a significant negative association with age, whereas this association was positive for total CSF volume (Table S11). Additionally, highly significant ($\beta = -0.392$ – -0.166 ; p -FDR < 0.001) negative associations were reported between age and the majority of subcortical volumes. Amygdala volumes were not significant associated with age (Table S12).

The association between age and WM microstructure has been examined in a larger sample (N = 3,513) of the UK Biobank dataset³⁷. Age effects on WM in this sample with overlapping MRI and genetic data are broadly in agreement with this previous study, as indicated by a negative association between age and WM microstructure quality (Tables S13 and S14).

	Including outliers (N = 816)				Excluding outliers (N = 733)			
	Beta: z ratio (S.D.)	t statistic	p	R ²	Beta: z ratio (S.D.)	t statistic	p	R ²
MDD-PGRS								
gFA	0.017 (0.036)	0.476	0.634	0.030	-0.011 (0.038)	-0.291	0.771	0.012
gFA Association	0.037 (0.036)	1.026	0.305	0.139	0.011 (0.039)	0.296	0.767	0.013
gFA Projection	-0.020 (0.036)	-0.560	0.576	0.041	-0.049 (0.038)	-1.297	0.195	0.244
gFA Thalamic	0.002 (0.037)	-0.043	0.966	0.000	-0.025 (0.039)	-0.650	0.516	0.063
SCZ-PGRS								
gFA	-0.028 (0.036)	-0.768	0.443	0.078	-0.048 (0.039)	-1.237	0.216	0.230
gFA Association	-0.020 (0.036)	-0.559	0.576	0.041	-0.040 (0.039)	-1.034	0.302	0.161
gFA Projection	-0.034 (0.036)	-0.947	0.344	0.116	-0.036 (0.038)	-0.947	0.344	0.132
gFA Thalamic	-0.032 (0.037)	0.885	0.376	0.105	-0.059 (0.039)	-1.521	0.129	0.348
BP-PGRS								
gFA	-0.013 (0.036)	-0.367	0.714	0.017	-0.014 (0.038)	-0.370	0.711	0.019
gFA Association	-0.016 (0.036)	-0.442	0.659	0.025	-0.017 (0.038)	-0.445	0.657	0.028
gFA Projection	-0.023 (0.035)	-0.665	0.506	0.055	-0.014 (0.037)	-0.369	0.712	0.019
gFA Thalamic	-0.000 (0.036)	0.0223	0.982	0.000	-0.008 (0.038)	-0.214	0.831	0.008

Table 2. Association of PGRS (MDD, SCZ or BP) at $p \leq 0.5$ with gFA and gFA for association, projection and thalamic WM fibers. FA: fractional anisotropy, WM: white matter, g: general component, MDD: major depressive disorder, SCZ: schizophrenia, BP: bipolar disorder, PGRS: polygenic risk scores, S.D.: standard deviation. Controlled for age, age² and gender, genotype batch and array, and 15 MDS components. R² = estimate of variance explained by PGRS in %.

	Including outliers (N = 816)				Excluding outliers (N = 733)			
	Beta: z ratio (S.D.)	t statistic	p	R ²	Beta: z ratio (S.D.)	t statistic	p	R ²
MDD-PGRS								
gMD	-0.047 (0.035)	-1.340	0.181	0.224	-0.028 (0.037)	-0.756	0.450	0.079
gMD Association	-0.053 (0.036)	-1.484	0.138	0.281	-0.030 (0.038)	-0.810	0.418	0.093
gMD Projection	-0.053 (0.037)	-1.417	0.157	0.277	-0.042 (0.039)	-1.075	0.283	0.179
gMD Thalamic	-0.031 (0.034)	-0.907	0.365	0.094	-0.017 (0.035)	-0.479	0.632	0.029
SCZ-PGRS								
gMD	-0.018 (0.035)	-0.503	0.615	0.031	0.001 (0.037)	0.019	0.985	0.000
gMD Association	-0.006 (0.036)	-0.179	0.858	0.004	0.005 (0.038)	0.121	0.903	0.002
gMD Projection	-0.035 (0.037)	-0.951	0.342	0.124	-0.021 (0.040)	-0.517	0.606	0.042
gMD Thalamic	-0.026 (0.034)	-0.777	0.438	0.068	0.003 (0.036)	0.084	0.933	0.001
BP-PGRS								
gMD	0.014 (0.034)	0.408	0.684	0.020	0.031 (0.036)	0.865	0.387	0.099
gMD Association	0.017 (0.035)	0.500	0.617	0.030	0.032 (0.037)	0.874	0.383	0.104
gMD Projection	0.003 (0.036)	0.082	0.935	0.000	-0.009 (0.039)	0.232	0.817	0.008
gFMDThalamic	0.011 (0.033)	0.328	0.743	0.012	0.032 (0.035)	0.933	0.351	0.104

Table 3. Association of PGRS (MDD, SCZ or BP) at $p \leq 0.5$ with gMD and gMD for association, projection and thalamic WM fibers. MD: mean diffusivity, WM: white matter, g: general factor, MDD: major depressive disorder, SCZ: schizophrenia, BP: bipolar disorder, PGRS: polygenic risk scores, S.D.: standard deviation. Controlled for age, age² and gender, genotype batch and array, and 15 MDS components. R² = estimate of variance explained by PGRS in %.

Discussion

Structural brain abnormalities in corticostriatal and limbic structures are thought to play an important role in the pathophysiology of common disabling psychiatric disorders, such as MDD, SCZ and BP. Since these structural brain abnormalities have been observed in unaffected relatives of patients and have been shown to be heritable quantitative traits, the current study sought to test the impact of the polygenic liability for MDD, SCZ, or BP on subcortical brain volumes and WM microstructure in individuals from a population-based study. Contrary to our predictions, we found no evidence for an association between polygenic risk for MDD, SCZ and BP and either subcortical brain regions or WM microstructure. These findings provide no support for the hypothesis that the polygenic liability for MDD, SCZ and BP is linked to changes in subcortical brain volumes and WM microstructure in major psychiatric illness.

Our findings are in contrast to many published studies that report a relationship between subcortical GM volume and PGRS for psychiatric disorders^{25,26}. However, these studies have generally been smaller than UK

Biobank ($N^{25,26} < 294$), increasing the risk of false positive results. In a complementary approach to our study, schizophrenia polygenic risk scores and linkage disequilibrium (LD) score regression have been used to test for shared genetic architecture between subcortical brain volumes and schizophrenia ($N = 11,840$)³¹. As in the current study, this study also reported a lack of overlapping genetic architecture between SCZ and subcortical volumes³¹. Moreover, Van Scheltinga *et al.*'s²⁵ result of a significant association between polygenic liability for SCZ and total brain volume could not be replicated in another study³⁰. Holmes *et al.*²⁹ ($N = 438$) also reported no association between MDD-PGRS and amygdala volume²⁹. Here we extend these findings to SCZ, BD and MDD and across a range of subcortical measures (Table 1 and S1–S4) using a large, single scanner sample.

Only a few studies have been published regarding the association between genetic liability for psychiatric illness and WM integrity. No associations between WM integrity and polygenic liability have been reported for BP, whereas polygenic liability for MDD and SCZ have been reported to be associated with decreased WM integrity^{27,28}. Furthermore, individuals at high genetic risk for psychiatric disorders, such as relatives of patients, have shown reduced WM integrity^{39–41}. Conversely, our study found no relation between genetic liability for psychiatric disorders and WM integrity (Tables 2 and 3 and S5–S10).

Our results provide no evidence for an association between PGRS and subcortical volumes and WM microstructure, however there are a number of limitations to this study that should be taken into account. Firstly, it is possible that any shared genetic architecture with brain structure was too small to detect in the current sample. Although our sample sizes ($N_{\text{volumetric}}/N_{\text{DTI}} = 978/816$) are large for imaging research, they are relatively small when compared to a typical PGRS study. A general limitation of PGRS is that they account for only a small proportion of total phenotypic variance, far short of their narrow sense heritability. The addition of further discovery GWA samples for MDD, SCZ and BD will improve the predictive accuracy of PGRS and alongside developments in imaging consortia, such as ENIGMA (<http://enigma.ini.usc.edu/>), will make it possible to test for smaller effects in future studies⁴². The current results may indicate that polygenic risk score associations with brain structure should be interpreted cautiously in smaller studies. Our findings should, however, be replicated to exclude the presence of a small genetic correlation between psychiatric disorder and brain structure and connectivity measures. Moreover, UK Biobank is a mixed community population, including both healthy individuals, patients, and individuals without clinical records. Although some previous studies did find associations in both healthy individuals and patients separately^{25,26}, analysing these groups together could have resulted in differences between the current and previous findings. To exclude the possibility of different genetic associations across clinical groups, future research should analyse cases and controls separately. Additionally, previous studies included relatively young participants (age range = 18–35 years)^{25,26,29–31}, whereas UK Biobank is a middle-older age study (age range = 40–70 years)³². Genetic associations between brain structure and major psychiatric disorders might be more difficult to detect in older populations, as older age groups might engender greater time for differential environmental exposure on the brain. Our results are, therefore, limited to the studied age range and should be replicated in independent studies. Furthermore, future work should also examine whether the genetic architecture of other *in vivo* brain parameters, such as measures of cortical morphology⁴³ and brain function, overlap with major mental illness.

In summary, the current study reports no evidence of shared genetic architecture between psychiatric disorders and either subcortical brain volumes or WM integrity. Our results do not replicate the findings of several published studies with small sample sizes, but are supported by similar recent work, particularly in SCZ.

References

- McGrath, J., Saha, S., Chant, D. & Welham, J. Schizophrenia: A Concise Overview of Incidence, Prevalence, and Mortality. *Epidemiol. Rev.* **30**, 67–76 (2008).
- Fernandez-Pujals, A. M. *et al.* Epidemiology and Heritability of Major Depressive Disorder, Stratified by Age of Onset, Sex, and Illness Course in Generation Scotland: Scottish Family Health Study (GS:SFHS). *PLoS One* **10**, e0142197 (2015).
- Shih, R. A., Belmonte, P. L. & Zandi, P. P. A review of the evidence from family, twin and adoption studies for a genetic contribution to adult psychiatric disorders. *Int. Rev. psychiatry* **16**, 260–283 (2004).
- Gejman, P. V., Sanders, A. R. & Duan, J. The Role of Genetics in the Etiology of Schizophrenia. *Psychiatr. Clin. North Am.* **33**, 35–66 (2010).
- Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature* **511**, 421–427 (2014).
- CONVERGE consortium. Sparse whole-genome sequencing identifies two loci for major depressive disorder. *Nature* **523**, 588–591 (2015).
- Sklar, P. *et al.* Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. *Nat. Genet.* **43**, 977–983 (2011).
- Ripke, S. *et al.* A mega-analysis of genome-wide association studies for major depressive disorder. *Mol. Psychiatry* **18**, 497–511 (2013).
- Braber, A. den, Bohlken, M., Brouwer, R. & Ent, D. van't. Heritability of subcortical brain measures: a perspective for future genome-wide association studies. *Neuroimage* **83**, 98–102 (2013).
- Kochunov, P., Jahanshad, N., Marcus, D. & Winkler, A. Heritability of fractional anisotropy in human white matter: a comparison of Human Connectome Project and ENIGMA-DTI data. *Neuroimage* **111**, 300–311 (2015).
- Hibar, D., Stein, J., Renteria, M. & Arias-Vasquez, A. Common genetic variants influence human subcortical brain structures. *Nature* **520**, 224–229 (2015).
- Zuliani, R. *et al.* Genetic variants in the ErbB4 gene are associated with white matter integrity. *Psychiatry Res. - Neuroimaging* **191**, 133–137 (2011).
- Arnone, D. *et al.* Magnetic resonance imaging studies in bipolar disorder and schizophrenia: meta-analysis. *Br. J. Psychiatry* **195**, 194–201 (2009).
- Haijma, S., Haren, N. Van, Cahn, W. & Koolschijn, P. Brain Volumes in Schizophrenia: A Meta-Analysis in Over 18 000 Subjects. *Schizophr. Bull.* **39**, 1129–1138 (2012).
- Hibar, D., Westlye, L., Thompson, P. & Andreassen, O. ENIGMA Bipolar disorder working group findings from 1,747 cases and 2,615 controls. *Hum. Brain Mapp* (2014).
- van Erp, T. G. M. *et al.* Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol. Psychiatry* **21**, 547–553 (2016).

17. Schmaal, L. *et al.* Subcortical brain alterations in major depressive disorder: findings from the ENIGMA Major Depressive Disorder working group. *Mol. Psychiatry* 1–7 doi: 10.1038/mp.2015.69 (2015).
18. Zhang, K. *et al.* Molecular, Functional, and Structural Imaging of Major Depressive Disorder. *Neuroscience* (2016).
19. Cullen, K. R. *et al.* Altered White Matter Microstructure in Adolescents With Major Depression: A Preliminary Study. *J. Am. Acad. Child Adolesc. Psychiatry* 49, 173–183.e1 (2010).
20. Korgaonkar, M., Grieve, S., Koslow, S. & Gabrieli, J. Loss of White Matter Integrity in Major Depressive Disorder: Evidence Using Tract-Based Spatial Statistical Analysis of Diffusion Tensor Imaging. *Hum. Brain Mapp.* 32, 2161–2171 (2011).
21. Zhang, A. *et al.* Quantitative tract-specific measures of uncinate and cingulum in major depression using diffusion tensor imaging. *Neuropsychopharmacology* 37, 959–67 (2012).
22. Anderson, D., Ardekani, B. & Burdick, K. Overlapping and distinct gray and white matter abnormalities in schizophrenia and bipolar I disorder. *Bipolar* 15, 680–693 (2013).
23. Lener, M., Wong, E., Tang, C., Byne, W. & Goldstein, K. White Matter Abnormalities in Schizophrenia and Schizotypal Personality Disorder. *Schizophr. Bull.* (2014).
24. Roalf, D., Gur, R., Verma, R. & Parker, W. White matter microstructure in schizophrenia: Associations to neurocognition and clinical symptomatology. *Schizophrenia* 161, 42–49 (2015).
25. Terwisscha Van Scheltinga, A. F. *et al.* Genetic schizophrenia risk variants jointly modulate total brain and white matter volume. *Biol. Psychiatry* 73, 525–531 (2013).
26. Caseras, X., Tansey, K. E., Foley, S. & Linden, D. Association between genetic risk scoring for schizophrenia and bipolar disorder with regional subcortical volumes. *Transl. Psychiatry* 5, 1–5 (2015).
27. Bohlken, M. M. *et al.* Structural Brain Connectivity as a Genetic Marker for Schizophrenia. *JAMA Psychiatry* 73, 11–19 (2016).
28. Whalley, H. C. *et al.* Polygenic risk and white matter integrity in individuals at high risk of mood disorder. *Biol. Psychiatry* 74, 280–286 (2014).
29. Holmes, A. J. *et al.* Individual differences in amygdala-medial prefrontal anatomy link negative affect, impaired social functioning, and polygenic depression risk. *J. Neurosci.* 32, 18087–18100 (2012).
30. Papiol, S., Mitjans, M., Assogna, F. & Piras, F. Polygenic determinants of white matter volume derived from GWAS lack reproducibility in a replicate sample. *Transl. Psychiatry* 4, 1–4 (2014).
31. Franke, B. *et al.* Genetic influences on schizophrenia and subcortical brain volumes: large-scale proof of concept. *Nat. Neurosci.* 19, 420–431 (2016).
32. Collins, R. What makes UK Biobank special? *Lancet* 379, 1173–1174 (2012).
33. Miller, K. L. *et al.* Multimodal population brain imaging in the UK Biobank prospective epidemiological study. *Nat. Neurosci.* 19 (2016).
34. Hagenaars, S. P. *et al.* Shared genetic aetiology between cognitive functions and physical and mental health in UK Biobank (N = 112 151) and 24 GWAS consortia. *Mol. Psychiatry* 1–4, doi: 10.1038/mp.2015.225 (2016).
35. Euesden, J., Lewis, C. M. & Reilly, P. F. O. Genome analysis PRSice: Polygenic Risk Score software. 31, 1466–1468 (2015).
36. Purcell, S. M. *et al.* Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 460, 748–52 (2009).
37. Cox, S. R., Ritchie, S. J., Tucker-Drob, E. M., Liewald, D. C., Hagenaars, S. P., Davies, G., Wardlaw, J. M., Gale, C. R., Bastin, M. E. & D. I. J. Ageing and brain white matter structure in 3,513 UK Biobank participants. *Nature Communications* 7, 13629, doi:10.1038/ncomms13629 (2016).
38. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: a Practical and Powerful Approach to Multiple Testing. *J.R. Stat. Soc. B* 57, 289–300 (1995).
39. Sprooten, E. *et al.* White matter integrity in individuals at high genetic risk of bipolar disorder. *Biol. Psychiatry* 70, 350–356 (2011).
40. Huang, H., Fan, X., Williamson, D. E. & Rao, U. White matter changes in healthy adolescents at familial risk for unipolar depression: a diffusion tensor imaging study. *Neuropsychopharmacology* 36, 684–91 (2011).
41. Munoz Maniega, S. *et al.* A diffusion tensor MRI study of white matter integrity in subjects at high genetic risk of schizophrenia. *Schizophr. Res.* 106, 132–139 (2008).
42. Thompson, P. M. *et al.* The ENIGMA Consortium: large-scale collaborative analyses of neuroimaging and genetic data. *Brain Imaging Behav.* 8, 153–182 (2014).
43. Liu, B. *et al.* Polygenic Risk for Schizophrenia Influences Cortical Gyrfication in 2 Independent General Populations. *Schizophr. Bull.* sbw051, doi: 10.1093/schbul/sbw051 (2016).

Acknowledgements

This work was supported by a grant of the Wellcome Trust Strategic Award “Stratifying Resilience and Depression Longitudinally” (STRADL) (reference 104036/Z/14/Z). We thank the UK Biobank participants for their participation, and the UK Biobank team for their work in collecting and providing these data for analysis. Part of the work was undertaken in The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology (CCACE), part of the cross council Lifelong Health and Wellbeing Initiative (MR/K026992/1); funding from the Biotechnology and Biological Sciences Research Council (BBSRC) and Medical Research Council (MRC) is gratefully acknowledged. Age UK (The Disconnected Mind project) also provided support for the work undertaken at CCACE. LMR was supported by an Erasmus Traineeship Grant. XS receives support from China Scholarship Council. HCW is supported by a JMAS SIM fellowship from the Royal College of Physicians of Edinburgh and by an ESAT College Fellowship from the University of Edinburgh. SRC was supported by MRC grant MR/M013111/1. IJD is supported by the Medical Research Council award to CCACE (MR/K026992/1) and by the Dementias Platform UK (MR/L015382/1). IJD and SRC are additionally supported by the Age UK-funded Disconnected Mind project (<http://www.disconnectedmind.ed.ac.uk>). A.M.M. has previously received grant support from Pfizer, Lilly and Janssen.

Author Contributions

A.M.M., H.C.W., S.R.C., X.S. and L.M.R. contributed to the design of the study, analysis of the data, and writing the manuscript. M.J.A. was involved in curating the data. I.J.D. and M.E.B. were involved in the conception of the study and overseeing analysis methodology. J.G., E.W. and S.P.H. were involved in the preprocessing and calculation of the polygenic risk scores. UK Biobank collected all data and was involved in the preprocessing of imaging data. All authors discussed and commented on the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Reus, L. M. *et al.* Association of polygenic risk for major psychiatric illness with subcortical volumes and white matter integrity in UK Biobank. *Sci. Rep.* 7, 42140; doi: 10.1038/srep42140 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017