



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Intelligence and neuroticism in relation to depression and psychological distress: Evidence from two large population cohorts

Citation for published version:

Navrady, L, Ritchie, SJ, Chan, SWY, Kerr, D, Adams, M, Hawkins, E, Porteous, D, Deary, I, Gale, C, Batty, GD & McIntosh, A 2017, 'Intelligence and neuroticism in relation to depression and psychological distress: Evidence from two large population cohorts', *European Psychiatry*, vol. 43, pp. 58-65.
<https://doi.org/10.1016/j.eurpsy.2016.12.012>

Digital Object Identifier (DOI):

[10.1016/j.eurpsy.2016.12.012](https://doi.org/10.1016/j.eurpsy.2016.12.012)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

European Psychiatry

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Contents lists available at ScienceDirect

European Psychiatry

journal homepage: <http://www.eurpsy-journal.com>



Original article

Intelligence and neuroticism in relation to depression and psychological distress: Evidence from two large population cohorts

Q1 L.B. Navrady^{a,*}, S.J. Ritchie^{b,c}, S.W.Y. Chan^d, D. Kerr^{a,b,c,d,e,f,g,h,i}, M.J. Adams^a, E. Hawkins^a,
D. Porteous^{b,e,f}, I.J. Deary^{b,c,f}, C.R. Gale^{b,g}, G.D. Batty^{b,h,i}, A.M. McIntosh^{a,b,f}

^a Division of Psychiatry, University of Edinburgh, Royal Edinburgh Hospital, Edinburgh, EH10 5HF, UK

^b Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, 7, George Square, Edinburgh, EH8 9JZ, UK

^c Department of Psychology, University of Edinburgh, 7, George Square, Edinburgh, EH8 9JZ, UK

^d Section of Clinical Psychology, University of Edinburgh, Medical Quad, Teviot Place, Edinburgh, EH8 9AG, UK

^e Medical Genetics Section, Centre for Genetics and Experimental Medicine, Institute for Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh, EH4 2XU, UK

^f Generation Scotland, Centre for Genetics and Experimental Medicine, Institute for Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh, EH4 2XU, UK

^g MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton, SO16 6YD, UK

^h Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London, WC1E 6BT, UK

ⁱ Alzheimer Scotland Dementia Research Centre, Department of Psychology, University of Edinburgh, 7, George Square, Edinburgh, EH8 9JZ, UK

ARTICLE INFO

Article history:

Received 24 October 2016

Received in revised form 23 December 2016

Accepted 28 December 2016

Available online xxx

Keywords:

Depression

Intelligence

Neuroticism

UK Biobank

Generation Scotland

Psychological distress

ABSTRACT

Background: Neuroticism is a risk factor for selected mental and physical illnesses and is inversely associated with intelligence. Intelligence appears to interact with neuroticism and mitigate its detrimental effects on physical health and mortality. However, the inter-relationships of neuroticism and intelligence for major depressive disorder (MDD) and psychological distress has not been well examined.

Methods: Associations and interactions between neuroticism and general intelligence (*g*) on MDD, self-reported depression, and psychological distress were examined in two population-based cohorts: Generation Scotland: Scottish Family Health Study (GS:SFHS, *n* = 19,200) and UK Biobank (*n* = 90,529). The Eysenck Personality Scale Short Form-Revised measured neuroticism and *g* was extracted from multiple cognitive ability tests in each cohort. Family structure was adjusted for in GS:SFHS.

Results: Neuroticism was strongly associated with increased risk for depression and higher psychological distress in both samples. Although intelligence conferred no consistent independent effects on depression, it did increase the risk for depression across samples once neuroticism was adjusted for. Results suggest that higher intelligence may ameliorate the association between neuroticism and self-reported depression although no significant interaction was found for clinical MDD. Intelligence was inversely associated with psychological distress across cohorts. A small interaction was found across samples such that lower psychological distress associates with higher intelligence and lower neuroticism, although effect sizes were small.

Conclusions: From two large cohort studies, our findings suggest intelligence acts a protective factor in mitigating the effects of neuroticism on psychological distress. Intelligence does not confer protection against diagnosis of depression in those high in neuroticism.

© 2017 The Authors. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Q2 1. Introduction

Major depressive disorder (MDD) is a leading cause of disease burden worldwide [1]. Although MDD aetiology remains elusive, a

large proportion of its genetic covariance is attributable to neuroticism [2,3], suggesting a causal relationship. Neuroticism is a partially-heritable personality trait representing high emotionality and stress sensitivity [4], which correlates highly with MDD [5]. Cross-sectional studies suggest a strong positive association between neuroticism and MDD [6–8], whilst higher neuroticism prospectively associates with depression longitudinally [2,9–12], even when controlling for overlapping criteria [13–15] and

* Corresponding author. Floor 7, Kennedy Tower, Royal Edinburgh Hospital, Edinburgh, EH10 5HF, UK.

E-mail address: s1467731@sms.ed.ac.uk (L.B. Navrady).

<http://dx.doi.org/10.1016/j.eurpsy.2016.12.012>

0924-9338/© 2017 The Authors. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

demographics [16,17]. Whilst the public health impacts of neuroticism are wide-ranging (for a comprehensive review see Lahey), neuroticism may be an indirect measure of later MDD risk, rather than the causative risk factor itself. Whilst MDD is often recurrent [18], neuroticism is a stable trait [19] suggesting that their correlation is unlikely to be substantially attributable to an effect of MDD on neuroticism.

General intelligence (*g*) is a latent construct theorized to explain the common observation that people who excel in one type of cognitive task tend to excel in others [20]. When reduced to a single factor (*g*) these correlations explain approximately 50% of the covariance between tests. Lower intelligence in early life has been found to be a risk factor for poor physical health [21] and early mortality in adulthood [22,23]. Although research specifically regarding MDD is relatively sparse [24], there is evidence to suggest that *g* is impaired in depression [25,26] with longitudinal studies suggesting lower *g* in childhood or adolescence confers vulnerability to psychopathology in adulthood [27–30].

Psychological distress represents a cluster of emotional symptoms linked to depression [31–33]. Although symptoms of distress are common in population samples [34,35], they indicate only subthreshold mental health problems. With self-report measures of distress [36,37] freely available in epidemiological research, their measurement provides greater detective power to make distinctions between syndrome and subthreshold symptoms. Longitudinal research suggests neuroticism has a strong, direct effect on psychological distress [38]. Low childhood intelligence strongly associates with increased psychological distress in adulthood [27,39], which may precede MDD onset [40]. However, this is not a universal observation, particularly in studies accounting for socioeconomic status (SES).

Intelligence and neuroticism may interact to influence indices of health. A longitudinal study of war veterans [41] found high neuroticism and low cognitive ability were separate risk factors for mortality. Specifically, a 1-standard deviation increase in neuroticism resulted in a 33% increase in mortality; a 1-standard deviation decrease in intelligence associated with a 27% increase in mortality. An interaction (hazards ratio of 0.89) suggested that high neuroticism with low cognitive ability associates with high risk of poor health and reduced lifespan. Furthermore, high cognitive ability moderates the adverse effects of neuroticism on adjustment [42]. Whether similar interactions exist with regard to their effects on depression remains unknown. No investigation has yet examined how intelligence and neuroticism influence risk for MDD and how they may moderate each other's associations in

depression and psychological distress. Such an analysis may serve to clarify the mechanisms underlying MDD.

In this study, two large population-based cohorts were examined – Generation Scotland: Scottish Family Health Study (GS: SFHS) [43,44] and UK Biobank [45,46]. As previous studies suggest strong associations of neuroticism with risk of MDD [2,5], the same effect was hypothesised here. We hypothesised that higher intelligence may reduce MDD risk by mitigating the adverse effects of neuroticism, similarly to the interaction identified for mortality [41]. This reasoning transfers to psychological distress, hypothesising a positive association between neuroticism and psychological distress would be ameliorated by higher intelligence.

2. Method

2.1. GS:SFHS Overview

GS:SFHS is a family and population-based cohort recruited throughout Scotland between 2006 and 2011 [43]. During clinic assessment, participants aged 18–98 ($n = 24,084$) provided clinical, cognitive and biological data. Full details are provided elsewhere [43,44]. The GS:SFHS sample is predominately female (59%), and generally healthier and wealthier than the Scottish population [43]. This study includes 19,200 individuals with complete data of interest. Demographic information from this cohort is provided in Table 1 and within the Supplementary materials.

Study assessments: during clinic assessment, participants were screened for lifetime history of MDD using a structured clinical interview [47]. Diagnosis of MDD follows DSM-IV criteria; if either symptoms of depressive mood or anhedonia are endorsed, a minimum of four further symptoms must also be endorsed. Clinical significance must be endorsed, too (ie., symptoms lasting nearly all day, every day for a minimum of two weeks). This study includes 2481 individuals meeting criteria for lifetime history of MDD (13%), and 16,719 non-MDD cases (87%).

Four cognitive tests measuring intelligence were administered during clinic assessment [43,44]. The Wechsler Digit Symbol Substitution Task [48] measured processing speed. One paragraph from The Wechsler Logical Memory Test I & II [49] measured verbal declarative memory. The Verbal Fluency Test measured executive function [48] using phonemic lists of C, F and L. Vocabulary was measured with The Mill-Hill Vocabulary Test [50], using combined junior and senior synonyms. General intelligence (*g*) was extracted from these tests, as the first un-rotated principal component [51], explaining 41% of the variance. Loadings for processing speed,

Table 1
Demographic, clinical, and cognitive characteristics of GS:SFHS and UK Biobank individuals in the current study.

	GS:SFHS			UK Biobank		
	Total ($n = 19,200$)	Control ($n = 16,719$)	Lifetime MDD ($n = 2481$)	Total ($n = 90,529$)	Control ($n = 60,402$)	Lifetime MDD ($n = 30,127$)
Age	47.16 (14.97)	47.23 (15.27)	46.39 (12.89)	56.64 (8.13)	57.15 (8.16)	55.60 (7.98) [*]
Sex (% female)	59	57	72 [*]	52	46	65 [*]
Neuroticism	3.84 (3.16)	3.45 (2.94)	6.45 (3.32) [*]	3.46 (2.86)	2.65 (2.43)	5.09 (2.96) [*]
GHQ score	15.93 (8.81)	14.93 (7.56)	22.70 (12.77) [*]	–	–	–
PHQ score	–	–	–	1.36 (1.91)	0.89 (1.33)	2.30 (2.47) [*]
Wechsler Digit Symbol Substitution Task	72.31 (17.09)	72.45 (17.23)	71.44 (16.06) [*]	–	–	–
Mill-Hill Vocabulary Test	30.06 (4.76)	30.05 (4.75)	30.15 (4.84)	–	–	–
Wechsler Logical Memory Test I & II	31.01 (8.04)	30.99 (8.09)	31.02 (7.68) [*]	–	–	–
Verbal Fluency Test	25.68 (8.10)	25.60 (8.11)	26.21 (8.05) [*]	–	–	–
Reaction time	–	–	–	564.00 (119.87)	564.70 (119.98)	562.58 (119.66) [*]
Visual memory	–	–	–	4.04 (3.21)	4.04 (3.23)	4.04 (3.17)
Verbal-numerical reasoning	–	–	–	6.09 (2.14)	6.07 (2.16)	6.12 (2.11) [*]
SIMD	3903.82 (1851.91)	3957.58 (1832.28)	3541.51 (1941.03) [*]	–	–	–
Townsend Score	–	–	–	–1.37 (2.84)	–1.47 (2.77)	–1.06 (2.94) [*]

GS:SFHS: Generation Scotland: the Scottish Family Health Study; MDD: Major Depressive Disorder; GHQ: General Health Questionnaire; PHQ: Patient Health Questionnaire; SIMD: the Scottish Index of Multiple Deprivation. With the exception of sex, values represent Mean (SD).

^{*} Significantly different from controls at $P < 0.05$.

vocabulary, verbal declarative memory and executive function were 0.57, 0.68, 0.63 and 0.69 respectively.

The self-reported Eysenck Personality Questionnaire Short Form-Revised (EPQ-SF) [52] measured neuroticism. Twenty-four questions assessed neuroticism and extraversion, with total scores on each subscale ranging from 0–12. Higher scores indicate higher levels of each trait. This scale has been concurrently validated [53] with high reliability [54].

Psychological distress was self-reported using the General Health Questionnaire (GHQ-28) [36]. Twenty-eight items were scored from 0 (“not at all”) to 3 (“much more than usual”) with a total score ranging from 0–84. Higher scores indicate increased psychological distress.

The Scottish Index of Multiple Deprivation (SIMD) [55] is an official tool which identifies deprivation by combining different indicators (eg., income, crime) into a single index. The SIMD divides Scotland into 6505 small areas based on participant postcode, and assigns them a relative ranking from 1 (most deprived) to 6505 (least deprived).

2.2. UK Biobank Overview

UK Biobank is a population cohort recruited across the UK from 2006–2010. During an extensive baseline assessments [56] participants aged 40–69 ($n = 502,682$) provided biological, physical, and touch-screen questionnaire measures of socio-demographics (e.g., age, sex), psychosocial factors (e.g., mental health), and cognitive function. UK Biobank represents a wide range of exposures typical within the UK population [57], and has been described in detail elsewhere [45,46]. In this study, 147 individuals were removed from analysis due to participation in GS:SFHS. In total, 90,529 individuals with complete data of interest were included. Demographic information is provided in Table 1 and in the Supplementary materials.

Study assessments: between 2008–2010, a touch-screen questionnaire was added to the protocol to assess probable depression ($n = 172,751$) [58]. Although depression was not assessed using a precise diagnostic tool, the classification followed a self-report approach within the guidelines of the ICD-10 [59] and the DSM-IV [60]. Lifetime history of depression was assessed using items relating to the lifetime experience of depressive symptoms and help-seeking for mental health. A detailed description of how this phenotype was derived is provided elsewhere [56]. This study included 30,127 (33%) individuals self-reporting lifetime history of depression, and 60,402 (67%) non-depressed cases.

Three novel cognitive tests were administered via touch-screen questionnaire measuring reaction time, verbal-numerical reasoning, and visual memory [56]. A timed symbol matching test measured reaction time as the mean response time in ms over 12 trials; higher reaction times equate to poorer performance. Thirteen logic/reasoning-type questions assessed verbal-numerical reasoning - the total number of correct answers given within two-minutes was analysed. A visuo-spatial memory task measured the number of errors made when matching card pairs, higher scores reflect poorer cognitive function. From these tests, g was extracted as the first un-rotated principal component [51], explaining 44% of the variance in scores. Loadings onto g were: -0.61 (verbal-numeric reasoning), 0.57 (visual memory), and 0.55 (reaction time).

Neuroticism was assessed using 12 questions from the Eysenck Personality Questionnaire Short Form-Revised (EPQ-SF) [52], administered via a touch-screen questionnaire. A total score from 0–12 was produced, with higher scores reflecting increasing neuroticism.

The first four questions of the Patient Health Questionnaire–9 (PHQ9) [37] were administered by touch-screen questionnaire to

measure psychological distress. Responses on a scale from 0 (“Not at all”) to 3 (“Nearly every day”) were aggregated and a higher total score denoted higher levels of psychological distress.

The Townsend Deprivation Index [61] is a census-based measure of deprivation, incorporating unemployment, non-car ownership, non-home ownership and household overcrowding into a single index. Small geographical areas based on postcode information are allocated Townsend Scores. Higher scores represent greater deprivation.

2.3. Statistical analysis

In GS:SFHS, the MCMCglmm package was used. The Markov Chain Monte Carlo estimator produces generalised linear mixed models for binary outcomes (using the “threshold” family with a probit link function). The threshold link is unique to MCMCglmm, and although produces very similar results to a logit function, threshold links most closely match the underlying assumptions of latent normal errors in pedigree-based mixed effect models [62]. MCMCglmm was essential to control for genetic relatedness of the sample, which was fitted as a random effect using an inverse pedigree matrix. Due to limitations within MCMCglmm with missing predictor variables, only complete data can be used. An interaction was fitted to estimate the moderating effect of g on the contribution of neuroticism to MDD. Another model examined this interaction while conditioning on deprivation. Regression coefficients are reported as Odds-Ratios. In a second set of analyses, GHQ was modelled as a normally distributed outcome variable. Neuroticism and GHQ were standardised to have a mean of zero and a standard deviation of 1. Age (standardised) and sex were used as fixed effects throughout.

In UK Biobank, generalized linear regression analyses were conducted as kinship need not be accounted for. The main effects of neuroticism and g were examined as predictors for self-reported depression. The interaction between neuroticism and g on depression was modelled. Another model examined this interaction while adjusting for deprivation. Generalized linear regressions were fitted with a logit link function and Odds-Ratios reported. A second set of analyses examined psychological distress (PHQ) using linear regression models. Neuroticism and PHQ were standardised to have a mean of zero and a standard deviation of one. Reaction time was log transformed due to a significantly positive skew. Visual memory was transformed with a log + 1 transformation because it was significantly skewed and zero-inflated. All regression analyses co-varied for age, and sex.

3. Results

3.1. GS:SFHS

As seen in Table 1, MDD cases were younger, predominately female, and had higher GHQ and neuroticism scores. No group differences were found in general intelligence; ($t(3243.38) = -1.39$, $P = 0.17$, Cohen’s $d = 0.03$). Group differences were found in processing speed and executive function. MDD cases were from less deprived areas; ($t(3171.20) = 9.93$, $P = 2.20 \times 10^{-16}$, Cohen’s $d = 0.22$). Full statistical output can be found in the Supplementary materials.

3.1.1. Associations of neuroticism and g with MDD status

Higher neuroticism was strongly associated with increased risk for MDD. A 1SD-increase in neuroticism increased MDD risk by an odds-ratio of 3.61 (95% CIs = [3.28, 4.01], $P < 1.00 \times 10^{-4}$). Although no age effects were found, being female increased risk for MDD by an Odds-Ratio of 1.76 (95% CIs = [1.52, 2.03],




Fig. 1. Predicted risk for MDD and self-reported depression from the interaction of neuroticism and *g* in both GS:SFHS and UK Biobank. Regression lines reflect the interaction at mean *g* (black line) and 2SD above (blue line) and below mean *g* (pink line).

$P < 1.00 \times 10^{-4}$. *g* had no independent effect on risk for MDD (OR = 1.02, 95% CIs = [0.99, 1.07], $P = 0.53$).

3.1.2. Interaction between neuroticism and *g* on MDD

No interaction was found between neuroticism and *g* (OR = 1.03, 95% CI = [0.98, 1.08], $P = 0.32$), see Fig. 1 and Table 2, even after co-varying for SIMD. However, the main effect of neuroticism was strongly associated with MDD risk (OR = 3.71, 95% CI = [3.37, 4.12], $P < 1.00 \times 10^{-4}$) whilst *g* was associated with a small increase in MDD risk (OR = 1.14, 95% CIs = [1.07, 1.20], $P < 1.00 \times 10^{-4}$). A main effect was found whereby higher deprivation confers risk for MDD (OR = 0.80, 95% CIs = [0.75, 0.86], $P < 1.00 \times 10^{-4}$).

3.1.3. Associations of neuroticism and *g* with psychological distress

Neuroticism was associated with increased psychological distress; a 1SD increase in neuroticism was associated with an increase in GHQ of β 0.52 (95% CIs = [0.50, 0.53], $P < 1.00 \times 10^{-4}$). A small inverse relationship was found whereby higher *g* was associated with decreased levels of psychological distress ($\beta = -0.08$, 95% CIs = [-0.09, -0.07], $P < 1.00 \times 10^{-4}$).

3.1.4. Interaction between neuroticism and *g* on psychological distress

A small interaction suggested higher *g* interacts with neuroticism to mitigate neuroticism's detrimental association on GHQ ($\beta = -0.05$, 95% CIs = [-0.06, -0.04], $P < 1.00 \times 10^{-4}$), see Fig. 2 and Table 2. This interaction remained after co-varying for deprivation.

Table 2
Results of a MCMC generalized linear mixed model from GS:SFHS predicting Odds-Ratios of MDD status, beta-coefficients for psychological distress (GHQ), *P*-value, upper and lower 95% confidence intervals and the Deviance Information Criterion AND results of a logistic regression from UK Biobank predicting Odds-Ratios for MDD status, beta-coefficients for psychological distress (PHQ), *P*-value, upper and lower 95% confidence intervals, the Akaike Information Criterion and adjusted R^2 value for the model.

Sample	Outcome	Variables	Odds-Ratio	β	Lower 95% CIs	Upper 95% CIs	<i>P</i> -value	DIC	AIC	R^2
GS:SFHS	MDD	Age	1.00	-	0.99	1.01	9.71×10^{-2}	12,561.35	-	-
		Sex (F)	1.71	-	1.48	1.97	$< 1.00 \times 10^{-4}$			
		Neuroticism	3.71	-	3.37	4.12	$< 1.00 \times 10^{-4}$			
		<i>g</i>	1.14	-	1.07	1.20	$< 1.00 \times 10^{-4}$			
		Neuroticism* <i>g</i>	1.03	-	0.98	1.08	0.32			
UK Biobank	MDD	Age	0.98	-	0.99	0.99	$< 2.00 \times 10^{-16}$	-	98,785	-
		Sex (F)	1.34	-	1.32	1.36	$< 2.00 \times 10^{-16}$			
		Neuroticism	2.40	-	2.36	2.44	$< 2.00 \times 10^{-16}$			
		<i>g</i>	1.06	-	1.04	1.07	5.08×10^{-14}			
		Neuroticism* <i>g</i>	0.96	-	0.95	0.98	$< 1.09 \times 10^{-7}$			
GS:SFHS	GHQ	Age	-	0.00	-0.00	0.00	0.59	47,873.87	-	-
		Sex (F)	-	0.04	0.02	0.07	2.63×10^{-3}			
		Neuroticism	-	0.50	0.49	0.52	$< 1.00 \times 10^{-4}$			
		<i>g</i>	-	-0.04	-0.05	-0.03	$< 1.00 \times 10^{-4}$			
		Neuroticism* <i>g</i>	-	-0.05	-0.06	-0.04	$< 1.00 \times 10^{-4}$			
UK Biobank	PHQ	Age	-	-0.02	-0.02	-0.01	$< 2.00 \times 10^{-16}$	-	-	0.2976
		Sex (F)	-	-0.02	-0.02	-0.01	1.57×10^{-8}			
		Neuroticism	-	0.51	0.51	0.52	$< 2.00 \times 10^{-16}$			
		<i>g</i>	-	-0.05	-0.06	-0.05	$< 2.00 \times 10^{-16}$			
		Neuroticism* <i>g</i>	-	-0.02	-0.03	-0.02	$< 2.00 \times 10^{-16}$			

MCMC: Markov Chain Monte Carlo; GS:SFHS: Generation Scotland: the Scottish Family Health Study; GHQ: General Health Questionnaire; DIC: Deviance Information Criterion; *g*: General Intelligence; MDD: major depressive disorder; PHQ: Patient Health Questionnaire; AIC: Akaike Information Criterion.




Fig. 2. Psychological distress scores from the interaction of neuroticism and *g* in both GS:SFHS (GHQ) and UK Biobank. Regression lines reflect the interaction at mean *g* (black line) and 2SD above (blue line) and below mean *g* (pink line).

3.2. UK Biobank

As reported in Table 1, MDD cases were younger, predominately female, and had higher psychological distress (PHQ) and neuroticism scores than non-depressed cases. Significant differences were found in verbal-numerical reasoning (in which non-depressed cases performed better) and reaction time (in which depressed cases performed better). *g* was higher in depressed cases ($t(61357) = -2.65, P = 8.12 \times 10^{-3}$, Cohen's $d = 0.02$). Non-depressed cases had lower deprivation scores than depressed cases; ($t(57110) = -20.08, P = 2.2 \times 10^{-16}$, Cohen's $d = 0.14$), although this difference was small. See the Supplementary materials for full statistical output.

3.2.1. Associations of neuroticism and *g* with MDD status

Higher neuroticism was associated with increased likelihood of self-reported depression. For every 1SD increase in neuroticism, the odds for depression increased by 2.39 (95% CIs = [2.35, 2.43], $P < 2.00 \times 10^{-16}$). No main effects of *g* were found (OR = 1.00, 95% CIs = [0.99, 1.01], $P = 0.86$). Small effects of age and sex were found.

3.2.2. Interaction between neuroticism and *g* on MDD

A small interaction was found in which high levels of intelligence and neuroticism associate with reduced self-reported depression (OR = 0.96, 95% CIs = [0.95, 0.98], $P = 1.09 \times 10^{-7}$), see Table 2 and Fig. 1. This interaction remained after co-varying for deprivation.

3.2.3. Associations of neuroticism and *g* with psychological distress

Neuroticism was moderately associated with increased levels of psychological distress. For every 1SD increase in neuroticism, PHQ increased by β 0.52 (95% confidence intervals = [0.51, 0.52], $P < 2.00 \times 10^{-16}$). *g* was associated with a small reduction in PHQ ($\beta = -0.08$, 95% CIs = [-0.08, -0.07], $P < 2.00 \times 10^{-16}$).

3.2.4. Interaction between neuroticism and *g* on psychological distress

A small interaction was found in which *g* moderates the detrimental effects of neuroticism on psychological distress ($\beta = -0.02$, 95% CIs = [-0.03, -0.02], $P < 2.00 \times 10^{-16}$), see

Table 2 and Fig. 2. This interaction remained after co-varying for deprivation.

4. Discussion

The cross-sectional associations between neuroticism, general intelligence (*g*), MDD, self-reported depression, and psychological distress were examined in two large population based cohorts; GS:SFHS and UK Biobank. Neuroticism was strongly associated with increased risk for both MDD diagnosis and self-reported depression, replicating previous findings [6,7]. Intelligence conferred no consistent independent effects but associated with an increased risk for depression once neuroticism was adjusted for. UK Biobank data suggest an interaction whereby higher *g* has a small effect in reducing the impact of neuroticism on self-reported depression. This interaction was small, both absolutely, and in comparison to the main effects of neuroticism. No such interaction was found in GS:SFHS using a clinical measure of MDD. However, across samples, the risk conferred by neuroticism after co-varying for *g* appears to be increased in terms of the absolute OR value when compared to basic models. Overall, results demonstrate an association whereby intelligence provides modest protection against the risk-conferring effects of neuroticism on self-reported depression, but not clinical MDD.

Consistent and replicable findings were found suggesting higher neuroticism associates with increased psychological distress, whereas higher intelligence associates with reduced psychological distress. A small interaction was found across samples such that lower distress associates with higher intelligence and lower neuroticism. Although these results are of small magnitude, they suggest an important interaction whereby higher *g* lessens the strength of the neuroticism–distress association.

This is the first study of intelligence's potential protective influence on MDD [63], self-reported depression, and psychological distress in high neuroticism individuals. Consistent with previous research the strong link between neuroticism with increased risk for depression and psychological distress was replicated with moderate effect sizes. Although longitudinal work suggests intelligence provides protection to mental health

[24,28,29], we found *g* increased the risk for depression when adjusted for neuroticism. The magnitude of this risk was very small, however. Across cohorts, intelligence associated with decreased levels of psychological distress. A modest association of intelligence as a mitigating factor in reducing psychological distress in individuals with high neuroticism was found in both cohorts. Although this study suggests intelligence provides a protective function in self-reported depression and psychological distress (which mirrors previous research [23,41,42]), intelligence was not found to be protective against diagnosis of depression in those high in neuroticism.

It is unclear why intelligence associates with protection to risk for psychological distress, but not MDD. One supposition is that individuals with higher intelligence may be more likely to seek help, and therefore are more likely to receive a clinical diagnosis of depression. Another postulation could be that intelligence has an effect only during times of depressive episode. A state-dependent association of cognitive ability has been suggested in which variability in intelligence co-varies with depressive episode and remission (for a comprehensive review, see Sackeim and Steif [26]). As such, subsequent investigations may benefit from addressing the same hypotheses examining individuals with current MDD in comparison to individuals in remission, and controls. Increased psychological distress is an established symptom of depression and often used in clinical diagnosis [31,32]. Goldberg [33] described distress as representing the overall severity of depression and so it is likely that individuals scoring highly on measures of psychological distress may be more likely to self-report the disorder, irrespective of its clinical significance. However, we must be mindful of the complexities of causality; whilst it is likely that the neuroticism trait prospectively predicts later distress and self-reported depression, we cannot be certain that these factors are not manifestations of the same underlying risk.

Intelligence could be a marker of system integrity [64] in which increased intelligence circumvents negative mood biasing in individuals high in neuroticism that may lead to distress and disorder [65]. Alternatively, more intelligent individuals may be better able to employ successful coping mechanisms during times of distress: higher intelligence associates with increased resilience to adversity in children [66]. Research suggests that psychosocial factors are associated with resilience to mood disorders [67]. Proactive and psychosocial coping mechanisms may enable individuals decrease transient feelings of distress and to implement established, effective strategies learned from previous exposure to distress or depression [68]. This possibility is consistent with the finding that whereas *g* and neuroticism interacted to associate with reduced psychological distress, the same interaction was not found in clinical MDD. It would be interesting to explore intelligence's influences on coping style [69] and subsequent psychological distress and MDD diagnosis in future investigations. Intelligence may influence the adoption of specific coping strategies, and this could be a mediating factor in the 'depressogenic' process.

Some caveats merit comment. Different cognitive tasks were used to generate *g* across our samples. In GF:SFHS, pre-existing, standardized measures were used, whereas UK Biobank used bespoke cognitive tasks. Further replication utilising standardised measures would be beneficial. A second limitation is the differing MDD phenotypes used in each sample. In GS:SFHS, MDD was determined using a semi-structured interview [47], obtaining a robust MDD phenotype based on a standardised diagnostic tool. In UK Biobank, self-reported questionnaires were aggregated to form a depression phenotype; this data is not as comprehensive. Although it is of benefit to have conducted an independent replication within this study, the disparity in depression phenotypes may explain not

only the difference in prevalence rates across samples, but also why an interaction was found in UK Biobank and not GS:SFHS. Thirdly, this investigation only examined neuroticism. Personality represents stable individual dispositions in emotional reactivity, behavioural tendencies, and cognitive styles [22,70], which may be moderated by intelligence in predicting mental health outcomes. Examining such associations between all major dimensions of personality in subsequent research is advised. As neuroticism and MDD share genetic aetiology [2,3], causality cannot be inferred here, although the associations reported do make a significant contribution to the literature. Because neuroticism is a stable trait and MDD is a disease with a given age of onset, we can use neuroticism to predict an individual's risk for depression, without needing to infer causality.

In conclusion, this study fails to demonstrate that intelligence confers protection to clinical MDD in those with high neuroticism. However, in both samples, a modest interaction was found in which higher intelligence appears to ameliorate the detrimental association between neuroticism and psychological distress. It would be useful to determine this relationship prospectively in a sample where incident cases of MDD can be identified. An important corollary of this work may inform risk and resilience mechanisms in MDD. Future studies to disentangle the mechanisms driving depression are an important next step in further elucidating the aetiology of the disorder.

Author contributions

L.N. wrote the manuscript text and prepared all tables and figures. A.M. was the main supervisor for the project, with co-supervision provided by S.R. and S.C. M.J. aided in the statistical analysis. L.N. and E.H. contributed to the data entry for the project. D.K., D.P., I.D., C.G. and D.B. reviewed the manuscript.

Funding

The authors declare no competing financial interests.

Disclosure of interest

The authors have not supplied their declaration of competing interest.

Acknowledgements

GS:SFHS received ethical approval from the NHS Tayside Committee on Medical Research Ethics (REC Reference Number: 05/S1401/89). Methods cited are in accordance with the relevant guidelines and regulations. Written consent for the use of data was obtained from all participants. Genotyping of the GS:SFHS samples was carried out by the Genetics Core Laboratory at the Wellcome Trust Clinical Research Facility, Edinburgh, Scotland and was funded by the Medical Research Council UK and the Wellcome Trust (Wellcome Trust Strategic Award "Stratifying Resilience and Depression Longitudinally" (STRADL) Reference 104036/Z/14/Z). This investigation was supported by Dr Mortimer and Theresa Sackler Foundation in addition to core support being provided by the Chief Scientist Office of the Scottish Government Health Directorates [CZD/16/6] and the Scottish Funding Council [HR03006].

UK Biobank received ethical approval from the North West Multicentre Research Ethics Committee (REC Reference Number: 11/NW/0382), and all methods were conducted in accordance with the relevant guidelines. Written consent for the use of data was obtained from all participants. This study was conducted under UK Biobank application 4844 "Stratifying Resilience and Depression Longitudinally" (PI Andrew McIntosh).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.eurpsy.2016.12.012>.

References

- [1] Ferrari AJ, Charlson FJ, Norman RE, Patten SB, Freedman G, Murray CJ, et al. Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS Med* 2013;10(11):e1001547.
- [2] Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ. A longitudinal twin study of personality and major depression in women. *Arch Gen Psychiatry* 1993;50(11):853–62.
- [3] Jardine R, Martin NG, Henderson AS. Genetic covariation between neuroticism and the symptoms of anxiety and depression. *Genet Epidemiol* 1984;1(2): 89–107.
- [4] Matthews G, Deary I, Whiteman M. Personality traits. 3rd ed., Cambridge: Cambridge University Press; 2009.
- [5] Jylhä P, Isometsä E. The relationship of neuroticism and extraversion to symptoms of anxiety and depression in the general population. *Depress Anxiety* 2006;23(5):281–9.
- [6] Chan SW, Goodwin GM, Harmer CJ. Highly neurotic never-depressed students have negative biases in information processing. *Psychol Med* 2007;37(9): 1281–91.
- [7] Muris P, Roelofs J, Rassin E, Franken I, Mayer B. Mediating effects of rumination and worry on the links between neuroticism, anxiety and depression. *Pers Individual Differences* 2005;1105–11.
- [8] Roelofs J, Huibers M, Peeters F, Arntz A, van Os J. Rumination and worrying as possible mediators in the relation between neuroticism and symptoms of depression and anxiety in clinically depressed individuals. *Behav Res Ther* 2008;46(12):1283–9.
- [9] Farmer A, Korszun A, Owen MJ, Craddock N, Jones L, Jones I, et al. Medical disorders in people with recurrent depression. *Br J Psychiatry* 2008;192(5): 351–5.
- [10] Hirschfeld RM, Klerman GL, Lavori P, Keller MB, Griffith P, Coryell W. Pre-morbid personality assessments of first onset of major depression. *Arch Gen Psychiatry* 1989;46(4):345–50.
- [11] Kendler KS, Gatz M, Gardner CO, Pedersen NL. Personality and major depression: a Swedish longitudinal, population-based twin study. *Arch Gen Psychiatry* 2006;63(10):1113–20.
- [12] Fanous AH, Neale MC, Aggen SH, Kendler KS. A longitudinal study of personality and major depression in a population-based sample of male twins. *Psychol Med* 2007;37(8):1163–72.
- [13] Schmutte PS, Ryff CD. Personality and well-being: reexamining methods and meanings. *J Pers Soc Psychol* 1997;73(3):549–59.
- [14] Spijker J, de Graaf R, Oldehinkel AJ, Nolen WA, Ormel J. Are the vulnerability effects of personality and psychosocial functioning on depression accounted for by subthreshold symptoms? *Depress Anxiety* 2007;24(7):472–8.
- [15] Fergusson DM, Horwood LJ, Lawton JM. The relationships between neuroticism and depressive symptoms. *Soc Psychiatry Psychiatr Epidemiol* 1989;24(6):275–81.
- [16] Kendler KS, Kuhn J, Prescott CA. The interrelationship of neuroticism, sex, and stressful life events in the prediction of episodes of major depression. *Am J Psychiatry* 2004;161(4):631–6.
- [17] Neeleman J, Ormel J, Bijl RV. The distribution of psychiatric and somatic III health: associations with personality and socioeconomic status. *Psychosom Med* 2001;63(2):239–47.
- [18] Hardeveld F, Spijker J, De Graaf R, Nolen WA, Beekman AT. Recurrence of major depressive disorder and its predictors in the general population: results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Psychol Med* 2013;43(1):39–48.
- [19] Conley JJ. Longitudinal stability of personality traits: a multitrait-multimethod-multioccasion analysis. *J Pers Soc Psychol* 1985;49(5):1266–82.
- [20] Humphreys L. The construct of general intelligence. *Intelligence* 1979;105–20.
- [21] Wraw C, Deary IJ, Gale CR, Der G. Intelligence in youth and health at age 50. *Intelligence* 2015;53:23–32.
- [22] Deary IJ, Weiss A, Batty GD. Intelligence and personality as predictors of illness and death: how researchers in differential psychology and chronic disease epidemiology are collaborating to understand and address health inequalities. *Psychol Sci Public Interest* 2010;11(2):53–79.
- [23] Calvin CM, Deary IJ, Fenton C, Roberts BA, Der G, Leckenby N, et al. Intelligence in youth and all-cause-mortality: systematic review with meta-analysis. *Int J Epidemiol* 2011;40(3):626–44.
- [24] Gale CR, Deary IJ, Boyle SH, Barefoot J, Mortensen LH, Batty GD. Cognitive ability in early adulthood and risk of 5 specific psychiatric disorders in middle age: the Vietnam experience study. *Arch Gen Psychiatry* 2008;65(12):1410–8.
- [25] Marazziti D, Consoli G, Picchetti M, Carlini M, Faravelli L. Cognitive impairment in major depression. *Eur J Pharmacol* 2010;626(1):83–6.
- [26] Sackeim H, Steif B. The neuropsychology of depression and mania. In: Georgotas A, Cancro R, editors. *Depression and mania*. New York: Elsevier; 1988. p. 265–89.
- [27] Gale C, Hatch S, Batty G, Deary I. Intelligence in childhood and risk of psychological distress in adulthood: the 1958 National Child Development Survey and the 1970 British Cohort Study. *Intelligence* 2009;92–9.
- [28] Gale CR, Batty GD, Tynelius P, Deary IJ, Rasmussen F. Intelligence in early adulthood and subsequent hospitalization for mental disorders. *Epidemiology* 2010;21(1):70–7.
- [29] Maccabe JH. Population-based cohort studies on premorbid cognitive function in schizophrenia. *Epidemiol Rev* 2008;30:77–83.
- [30] Zammit S, Allebeck P, David AS, Dalman C, Hemmingsson T, Lundberg I, et al. A longitudinal study of premorbid IQ Score and risk of developing schizophrenia, bipolar disorder, severe depression, and other non-affective psychoses. *Arch Gen Psychiatry* 2004;61(4):354–60.
- [31] Beck A, Rush A, Shaw B. *Cognitive therapy of depression*. New York; 1979.
- [32] Snaith RP. The concepts of mild depression. *Br J Psychiatry* 1987;150: 387–93.
- [33] Goldberg DP, Bridges K, Duncan-Jones P, Grayson D. Dimensions of neuroses seen in primary-care settings. *Psychol Med* 1987;17(2):461–70.
- [34] Kessler RC, Wang PS. The descriptive epidemiology of commonly occurring mental disorders in the United States. *Annu Rev Public Health* 2008;29: 115–29.
- [35] Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H. Psychiatric morbidity among adults living in private households, 2000. *Int Rev Psychiatry* 2003;15(1–2):65–73.
- [36] Goldberg D, Hillier V. A scaled version of the General Health Questionnaire. *Psychol Med* 1979;9:139–45.
- [37] Kroencke K, Spitzer R, Williams J. The phq-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;606–13.
- [38] Ormel J, Wohlfarth T. How neuroticism, long-term difficulties, and life situation change influence psychological distress: a longitudinal model. *J Pers Soc Psychol* 1991;60(5):744–55.
- [39] Hatch SL, Jones PB, Kuh D, Hardy R, Wadsworth ME, Richards M. Childhood cognitive ability and adult mental health in the British 1946 birth cohort. *Soc Sci Med* 2007;64(11):2285–96.
- [40] Gulliver A, Griffiths KM, Christensen H, Brewer JL. A systematic review of help-seeking interventions for depression, anxiety and general psychological distress. *BMC Psychiatry* 2012;12:81.
- [41] Weiss A, Gale C, Batty D, Deary I. Emotionally stable, intelligent men live longer: the Vietnam Experience Study Cohort. *Psychosom Med* 2009; 385–94.
- [42] Leikas S, Mäkinen S, Lönnqvist JE, Verkasalo M. Cognitive ability × emotional stability interactions on adjustment. *Eur J Pers* 2009;329–42.
- [43] Smith BH, Campbell A, Linksted P, Fitzpatrick B, Jackson C, Kerr SM, et al. Cohort profile: Generation Scotland: Scottish Family Health Study (GS:SFHS). The study, its participants and their potential for genetic research on health and illness. *Int J Epidemiol* 2013;42(3):689–700.
- [44] Smith BH, Campbell H, Blackwood D, Connell J, Connor M, Deary IJ, et al. Generation Scotland: the Scottish Family Health Study; a new resource for researching genes and heritability. *BMC Med Genet* 2006;7:74.
- [45] Allen N, Sudlow C, Downey P, Peakman T, Danesh J, Elliot P, et al. UK Biobank: current status and what it means for epidemiology. *Health Policy Technol* 2012;123–6.
- [46] Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* 2015;12(3):e1001779.
- [47] First M, Spitzer R, Gibbon M, Williams J. *Structured Clinical Interview for DSM-IV Axis I Disorders, Clinician Version (SCID-CV)*. Washington, DC, USA: American Psychiatric Publishing, Inc; 1997.
- [48] Wechsler D. *The measurement and appraisal of adult intelligence*. Baltimore, Md: Williams & Wilkens; 1958.
- [49] Wechsler D. A standardized memory scale for clinical use. *J Psychol* 1945; 87–95.
- [50] Raven J. *Guide to using the Mill Hill Vocabulary Scale with the Progressive Matrices Scales*. Oxford, England: H.K. Lewis & Co; 1958.
- [51] Marioni R, Batty G, Hayward C, Kerr S, Campbell A, Hocking L, et al. Common genetic variants explain the majority of the correlation between height and intelligence: the Generation Scotland Study. *Behav Genet* 2014;91–6.
- [52] Eysenck H. Dimensions of personality: 16, 5 or 3 criteria for a taxonomic paradigm. *Pers Individual Differ* 1991;773–90.
- [53] Gow A, Whiteman M, Pattie A, Deary I. Goldberg's 'IPIP' Big-Five factor markers: internal consistency and concurrent validation in Scotland. *Pers Individual Differ* 2005;317–29.
- [54] Eysenck S, Eysenck H, Barrett P. A revised version of the psychoticism scale. *Pers Individual Differ* 1985;21–9.
- [55] Payne R, Abel G. UK indices of multiple deprivation – a way to make comparisons across constituent countries easier. *Health Stat Q* 2012.
- [56] Smith DJ, Nicholl BI, Cullen B, Martin D, Ul-Haq Z, Evans J, et al. Prevalence and characteristics of probable major depression and bipolar disorder within UK biobank: cross-sectional study of 172,751 participants. *PLoS One* 2013;8(11): e75362.
- [57] Biobank U. UK Biobank: protocol for a large-scale prospective epidemiological resource; 2011 [Available from: <http://www.ukbiobank.ac.uk/wp-content/uploads/2011/11/UK-Biobank-Protocol.pdf>]; cited 2011 24/10/2016].
- [58] Biobank U. Touchscreen questionnaire; 2012 [Available from: http://www.ukbiobank.ac.uk/wp-content/uploads/2011/06/Touchscreen_questionnaire.pdf?phpMyAdmin=trmkQjYdjjnQjg%2CfAzikMhEnx6].
- [59] WHO. *The ICD-10 classification of mental and behavioural disorders*. Geneva: WHO; 1992.
- [60] APA. *Washington, DC: American Psychiatric Association; 1994.*
- [61] Townsend P. Deprivation. *J Soc Policy* 1987;125–46.

- [62] Lynch M, Wasl B. Genetics and analysis of quantitative traits. Sunderland, Massachusetts: Sinauer Associates; 1998.
- [63] Robinson M, Oishi S. Trait self-reports as a 'fill-in' belief system: categorization speed moderates the extraversion/life satisfaction relation. *Self Identity* 2006;15–34.
- [64] Deary IJ. Looking for 'system integrity' in cognitive epidemiology. *Gerontology* 2012;58(6):545–53.
- [65] Hasler G, Drevets WC, Manji HK, Charney DS. Discovering endophenotypes for major depression. *Neuropsychopharmacology* 2004;29(10):1765–81.
- [66] Fergusson M, Horwood L, Ridder E. Show me the child at seven II: childhood intelligence and later outcomes in adolescence and young adulthood. *J Child Psychol Psychiatry* 2005;850–8.
- [67] Garmezy N, Masten AS, Tellegen A. The study of stress and competence in children: a building block for developmental psychopathology. *Child Dev* 1984;55(1):97–111.
- [68] LeDoux JE, Gorman JM. A call to action: overcoming anxiety through active coping. *Am J Psychiatry* 2001;158(12):1953–5.
- [69] Higgins J, Endler N. Coping, life stress, and psychological and somatic distress. *Eur J Pers* 2006;253–70.
- [70] Roberts BW, Kuncel NR, Shiner R, Caspi A, Goldberg LR. The power of personality: the comparative validity of personality traits, socioeconomic status, and cognitive ability for predicting important life outcomes. *Perspect Psychol Sci* 2007;2(4):313–45.

640
641
642
643
644
645
646
647
648
649
650

UNCORRECTED PROOF