



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Species-specific variation in RELA underlies differences in NF- κ B activity

a potential role in African swine fever pathogenesis

Citation for published version:

Palgrave, CJ, Gilmour, L, Lowden, CS, Lillico, SG, Mellencamp, MA & Whitelaw, CBA 2011, 'Species-specific variation in RELA underlies differences in NF- κ B activity: a potential role in African swine fever pathogenesis', *Journal of Virology*, vol. 85, no. 12, pp. 6008-14. <https://doi.org/10.1128/jvi.00331-11>

Digital Object Identifier (DOI):

[10.1128/jvi.00331-11](https://doi.org/10.1128/jvi.00331-11)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Virology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Species-Specific Variation in RELA Underlies Differences in NF- κ B Activity: a Potential Role in African Swine Fever Pathogenesis[∇]

Christopher J. Palgrave,^{1,2,‡} Linzi Gilmour,^{2,‡} C. Stewart Lowden,³ Simon G. Lillico,²
Martha A. Mellencamp,^{4,§} and C. Bruce A. Whitelaw^{2,*}

Veterinary Pathology Unit, Division of Veterinary Clinical Sciences, Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Veterinary Centre, Roslin, Midlothian EH25 9RG, United Kingdom¹; Division of Developmental Biology, The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Campus, Roslin, Midlothian EH25 9RG, United Kingdom²; Veterinary Health Research Pty Ltd., Trevenna Rd., West Armidale, NSW 2350, Australia³; and Genus/PIC, 100 Bluegrass Commons Blvd., Hendersonville, Tennessee 37075⁴

Received 17 February 2011/Accepted 15 March 2011

African swine fever virus (ASFV) is a highly infectious disease of domestic pigs, with virulent isolates causing a rapidly fatal hemorrhagic fever. In contrast, the porcine species endogenous to Africa tolerate infection. The ability of the virus to persist in one host while killing another genetically related host implies that disease severity may be, in part, modulated by host genetic variation. To complement transcription profiling approaches to identify the underlying genetic variation in the host response to ASFV, we have taken a candidate gene approach based on known signaling pathways that interact with the virus-encoded immunomodulatory protein A238L. We report the sequencing of these genes from different pig species and the identification and initial *in vitro* characterization of polymorphic variation in RELA (p65; v-rel reticuloendotheliosis viral oncogene homolog A), the major component of the NF- κ B transcription factor. Warthog RELA and domestic pig RELA differ at three amino acids. Transient cell transfection assays indicate that this variation is reflected in reduced NF- κ B activity *in vitro* for warthog RELA but not for domestic pig RELA. Induction assays indicate that warthog RELA and domestic pig RELA are elevated essentially to the same extent. Finally, mutational studies indicate that the S531P site conveys the majority of the functional variation between warthog RELA and domestic pig RELA. We propose that the variation in RELA identified between the warthog and domestic pig has the potential to underlie the difference between tolerance and rapid death upon ASFV infection.

African swine fever (ASF) virus (ASFV) is a pathogen of the Suidae (domestic and wild pig species), which may be transmitted directly or via an arthropod vector in the form of *Ornithodoros* ticks (35). ASFV is highly infectious, with virulent isolates causing an acute, rapidly fatal hemorrhagic fever in domestic pigs (*Sus scrofa*) (10, 34). This is thought, in part, to be the result of a proinflammatory cytokine storm driven by infected macrophages (9, 15, 16, 42–44, 56). Initiation of a systemic inflammatory response results in severe hematological and vascular perturbations, ultimately leading to cardiovascular collapse in a manner not dissimilar to septic shock (3, 17, 18, 20, 21, 40, 52, 53). In addition to hemorrhage, severe widespread apoptosis of infected macrophages and uninfected lymphocytes is a prominent feature of the disease; this is also likely related to markedly elevated proinflammatory cytokine levels (19, 33, 37, 43). In comparison to the severe disease which occurs in domestic pigs, in its natural hosts, warthogs (*Phacochoerus* sp.) and bushpigs (*Potamochoerus* sp.), ASF is subclinical and persistent (2, 32, 50, 51).

ASFV is notifiable to the World Organization for Animal

Health (OIE), placing it in the highest category of infectious animal pathogens. It exhibits remarkable potential for transboundary spread, and outbreaks in domestic pig populations have a serious socioeconomic impact worldwide. Furthermore, ASF is considered to be the major limiting factor to pig production in Africa (34). ASFV is a large, double-stranded DNA virus and the only member of the *Asfarviridae* family (12), suggesting that it may carry novel genes that are not carried by other virus families. Furthermore, the ability of the virus to persist in one host while killing another genetically related host alludes to the possibility that disease severity may, in part, be modulated by host genetic variation.

Several candidate ASFV-encoded immune modulatory factors have been identified, including homologues of CD2 (8-DR/CD2v) (5, 6, 41), IAP (A224L) (31, 39), Bcl-2 (A179L; 5-HL) (1, 7, 8, 30), and I κ B α (A238L; 5-EL) (36, 49). Of these, A238L shares 40% sequence homology and 20% identity with domestic pig I κ B α (NFKBIA) and substitutes for NFKBIA by binding to the RELA (p65; v-rel reticuloendotheliosis viral oncogene homolog A) subunit of NF- κ B. Thus, A238L reduces the ability of NF- κ B to be activated (36, 49). In addition to inhibiting host NF- κ B, A238L also suppresses calcineurin phosphatase activation of NFAT signaling by the following two mechanisms: direct binding to calcineurin phosphatase 3, β isoform (PPP3CB), and binding to the immunophilin carrier cyclophilin A (PPIA) in a manner similar to that of the immunosuppressive drug cyclosporine A (28, 29).

Various groups have initiated transcription profiling of host genes implicated in ASFV infection (15, 16, 42–44, 56). These

* Corresponding author. Mailing address: Division of Developmental Biology, The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Campus, Roslin, Midlothian EH25 9RG, United Kingdom. Phone: 44 (0)131 651 9175. Fax: 44 (0)131 651 9105. E-mail: bruce.whitelaw@roslin.ed.ac.uk.

‡ C.J.P. and L.G. contributed equally to this work.

§ Present address: Ralco Nutrition Inc., 1600 Hahn Road, Marshall, MN 56258.

[∇] Published ahead of print on 30 March 2011.

TABLE 1. Gene-specific PCR primers

Gene/domain	Primer sequence (5'–3')	
	Sense	Antisense
<i>PPIA</i> (cDNA)	ATC TNT CAG TGC TGC TCA GC	CAG AAG GAA TGG TCT GAT GG
<i>NFKBIA</i> (cDNA)	AAG GAG CGG CTA CTG GAC G	CAT GGT CTT TTA GAC ACT TTC CA
<i>NFKBIA</i> (5'-end DNA)	CTC ATC GCA GGG AGT TTC TC	TCC TCG TCC TTC ATG GAG TC
<i>NFATC1</i> (cDNA)	ATC TCA GCT GTT GGG TCA GC	AGT GAG GGT GAG TGG TCC AG
<i>PPP3CB</i> (cDNA)	CCC AAC ACA TCG TTT GAC AT	ATG TGA GAG TCC CTG TGA AG
<i>RELA</i> (cDNA)	GAC CTC TTC CCC CTC ATC TT	CCC CTT AGG AGC TGA TCT GA
<i>RELA</i> transactivation domains (DNA)	GGA AGG GAC ACTGAC AGA GG	TCA GAA GGG CTG AGA AGT CC

studies identified numerous upregulated host genes, but to date, all are limited to analysis in domestic pig cells. In this study, we take a complementary approach to this question by the testing of variation in targeted candidate genes. Clearly A238L represents a novel and versatile immunoregulatory mechanism by which ASFV can inhibit both the NF-κB and NFAT signaling pathways (11, 28, 29, 36, 49). We therefore consider the three A238L target proteins, RELA, PPP3CB, and PPIA, and the two proteins it mimics, NFKBIA and NFATC1, as candidates for the genetic variation between pig species which may contribute to species-specific responses to ASFV infection. We now report the sequencing of these genes from different pig species and identification and initial *in vitro* characterization of polymorphic variation in one of them.

MATERIALS AND METHODS

mRNA isolation, cDNA synthesis, and DNA sequencing. Whole blood (5 ml) was collected into EDTA from a domestic pig (*Sus scrofa*, commercial pig; United Kingdom), common warthog (*Phacochoerus africanus*; Rotterdam Zoo, Holland), and babirusa (*Babyrusa babyrussa*; Marwell Zoo, United Kingdom) and transferred immediately into DNA/RNA stabilization reagent for blood/bone marrow (Roche Diagnostics). This was processed using an mRNA isolation kit for white blood cells (Roche Diagnostics). Initially, cDNA libraries were synthesized using the SMART RACE (rapid amplification of cDNA ends) cDNA amplification kit (Clontech) to enable partial sequencing and design of species-specific primers (Table 1) for synthesis of individual cDNAs using proof-reading PCR using a 1-μl sample in a 25-μl PCR mixture consisting of 20 pmol of each primer in 2 mM MgCl₂ and 2 mM deoxynucleoside triphosphates (dNTPs) with 0.7 U High-Fidelity DNA polymerase (Roche Diagnostics). The PCR cycling conditions used were optimized for each gene (data not shown). PCR products were resolved on a 1% agarose gel, excised with a scalpel blade, and extracted using a QIAquick gel extraction kit (Qiagen). These were cloned into pGEM-T Easy (Promega) and sequenced in both directions. To achieve the full coding sequence for NFKBIA, 5'-end fragments were amplified directly from genomic DNA.

Genomic DNA isolation and sequencing. Samples of skeletal muscle were collected into 20% dimethyl sulfoxide (DMSO)-saturated salt (NaCl) solution and stored at -70°C. DNA was extracted from 0.2 g muscle using the BACC2 extraction kit for blood and cell cultures (Nucleon Biosciences). For PCR, 50 ng genomic DNA was used as template in a 25-μl PCR mixture consisting of 20 pmol of each primer in 2 mM MgCl₂ and 2 mM dNTPs, with 0.7 U High-Fidelity DNA polymerase (Roche Diagnostics).

Plasmid construction. Restriction sites were introduced into RELA products from the above-described sequencing study by nested PCR; this enabled insertion of domestic pig and warthog RELA genotypes into the multiple cloning site of the pFLAG-CMV-4 vector (Sigma-Aldrich). PCR products were diluted to 1:500 in sterile water, and 1 μl was used as a template in a 25-μl PCR mixture consisting of 20 pmol of each primer (forward HindIII [5'-CCA AGC TTG ACC TCT TCC CCC TCA TCT T-3'] and reverse NotI [5'-GCG CGG CCG CTT AGG AGC TGA TCT GA-3']) in 2 mM MgCl₂ and 2 mM dNTPs, with 0.7 U High-Fidelity DNA polymerase (Roche Diagnostics). Restriction sites are underlined. Each ~1.6-kbp PCR product was resolved on a 1% agarose gel, excised with a scalpel blade, and extracted using the QIAquick gel extraction kit (Qiagen). Following HindIII and NotI restriction digestion of the vector and RELA

PCR products, the products were ligated into the open plasmid. These constructs allowed constitutive expression of warthog RELA and domestic pig RELA driven by the cytomegalovirus (CMV) promoter. In addition, an N-terminal eight-amino-acid FLAG sequence is incorporated into the protein, which is recognized by an anti-FLAG monoclonal antibody. Site-directed mutagenesis was performed using a QuikChange II site-directed mutagenesis kit (Stratagene).

Plasmid transient transfection was performed using Lipofectamine2000 (Invitrogen) in ~1 × 10⁵ cells/well of a 12-well plate in culture medium at 37°C and 5% CO₂. COS-7 cells were cultured in Glasgow minimal essential medium (GMEM; Sigma), containing 10% fetal bovine serum (Invitrogen), 1% L-glutamine (Invitrogen), 1% sodium pyruvate (Invitrogen), 1% nonessential amino acids (Invitrogen), and 0.2% β-mercaptoethanol (Invitrogen). Mouse embryonic fibroblasts (MEFs) and RELA^{-/-} MEFs were cultured in Dulbecco's modified Eagle medium (DMEM; Sigma), containing 10% fetal bovine serum (Invitrogen). Luciferase activity was determined in triplicate using the dual reporter assay system (Promega) and analyzed with Excel (Microsoft), with total protein measured with the bicinchoninic acid (BCA) protein assay kit (Pierce). The data are presented as the means ± standard deviations; statistical significance was evaluated using the unpaired Student *t* test, with a difference between groups being considered statistically significant if the *P* value of the comparison was <0.05.

Western blotting. For Western blot analysis, denatured protein (10 μg) was run in Tris-glycine-SDS running buffer (National Diagnostics) on precast Nu-Page 12% Tris-glycine gels (Invitrogen) before transfer to a nitrocellulose membrane (National Diagnostics). Proteins were visualized using anti-FLAG M2-peroxidase (horseradish peroxidase [HRP]) (Sigma)-conjugated primary antibody and mouse β-actin primary antibody with goat anti-mouse IgG HRP (Sigma)-conjugated secondary antibody and detected with Immobilon Western (Millipore) chemiluminescent HRP substrate.

RESULTS

Limited sequence variation in candidate porcine genes. cDNAs were produced for *PPIA*, *NFKBIA*, *NFATC1* (regulatory domain), *PPP3CB*, and *RELA*. These were obtained from the following three pig species: the domestic pig (*Sus scrofa*), common warthog (*Phacochoerus africanus*), and babirusa (*Babyrusa babyrussa*). The domestic pig and warthog represent ASFV-susceptible and ASFV-tolerant species, respectively. The “outspecies” is the babirusa, which is considered to be the most ancient extant species of pig; its range is restricted to the island of Sulawesi in the Indonesian archipelago. It has no common ancestor with the domestic pig more recently than approximately 10 to 19 million years ago (38). It is not known how this pig species would respond regarding ASFV infection. We generated sequence data (deposited in the EMBL database) (Table 2) and aligned all sequences to those of the human homologues which we used as reference sequences.

Limited sequence variation was observed in this study (data not shown), with the following genes displaying complete homology at the translated protein level between the domestic pig and warthog: PPIA, NFATC1 regulatory domain, and

TABLE 2. Sequences deposited into the EMBL database

Gene	EMBL accession no.		
	Domestic pig	Warthog	Babirusa
<i>PPIA</i>	FN401368	FN401369	FN401370
<i>NFKBIA</i> (cDNA)	FN421467	FN421468	FN421469
<i>NFKBIA</i> (5'-end DNA)	FN421464	FN421465	FN421466
<i>NFATC1</i>	FN421470	FN421471	FN421472
<i>PPP3CB</i>	FN421473	FN421474	FN421475
<i>RELA</i>	FN999988	FN999989	FN999990

NFKBIA. In the warthog, *PPP3CB* contained two insertions with respect to the domestic pig. However, these were located outwith the known functional domains. Furthermore, these correlate with splice variants described in human *PPP3CB* (22, 27). Only in the *RELA* subunit of *NF-κB* were potentially significant coding differences between the domestic pig and warthog identified (Fig. 1).

Sequence of the porcine *RELA* gene. The *RELA* open reading frame (ORF) is 1,662 nucleotides in the domestic pig and warthog and is slightly smaller at 1,556 nucleotides in the babirusa. These encode proteins which are 554 and 552 amino acids in length, respectively. Human *RELA* ORF and protein are the same lengths as the babirusa sequences. Due to insufficient primer binding sites in the 5' untranslated region

(UTR), we sequenced all but the first 6 nucleotides for all three pig species (Fig. 1).

Although the babirusa and human sequences are both 6 nucleotides shorter than the other pig sequences, the nucleotide deletions occur at different locations. The babirusa sequence has a single 6-nucleotide deletion, in relation to those of the other pig species, located between the Rel homology domain and the transactivation 2 domain. Human *RELA* has two 3-nucleotide deletions, one 20 nucleotides upstream from the babirusa deletion and the other in the transactivation 2 domain.

Of the 13 nucleotide differences between the domestic pig and warthog, only three are nonsynonymous and result in codon changes, T448A, S485P, and S531P. The threonine at 448 in the domestic pig occurs as an alanine in the warthog and babirusa and is absent in human *RELA*. The serine at 485 in the domestic pig sequence is a proline in the other porcine sequences and in human *RELA*. The serine at 531 in the domestic pig is also found in babirusa and human but is a proline in warthog *RELA*. All three amino acid differences occur outside the Rel homology domain, with amino acids 448 and 485 located within transactivation domain 2 and amino acid 531 within transactivation domain 1. To confirm these sequence differences, a 268-nucleotide region was amplified directly from genomic DNA and sequenced for an additional 5




FIG. 1. *RELA* primary protein sequence minus the first two amino acids of the porcine sequences. The Rel homology domain is underlined, the transactivation 2 domain is dashed underlined, and the transactivation 1 domain is double underlined. The nuclear localization signal (KRKR) is dotted underlined. The T448A, S485P, and S531P variations between the domestic pig and warthog are highlighted. SS, *Sus scrofa*; PA, *Phacochoerus africanus*; BB, *Babyrousa babyrussa*; HS, *Homo sapiens* (M62399).




FIG. 2. Effect of polymorphic *RELA* on basal NF- κ B activity. (A) Western blot of COS-7 cells at 24 h after transfection with 1 μ g FLAG-tagged warthog (WH) or domestic pig (DP) *RELA* or empty FLAG vector (-ve) and established *RELA*-FLAG expressing COS-7 cells (+ve). (B) Fold difference between NF- κ B-luciferase activity in COS-7 cells after 24 h of transient transfection with 1 μ g warthog (WH) or 1 μ g domestic pig (DP) *RELA*. Activity is presented as the fold difference between *RELA*-induced NF- κ B-luciferase activity normalized to cotransfected TK-Renilla luciferase activity relative to that of the empty vector. Error bars = standard deviations from the means. $P < 0.03$ between WH and DP. (C) Fold difference between NF- κ B-luciferase activities in *RELA*^{-/-} MEFs after 24 h of transient transfection with warthog (WH) and domestic pig (DP) *RELA* genes. Activity is presented as the fold difference between *RELA*-induced NF- κ B-luciferase activity normalized to cotransfected TK-Renilla luciferase activity relative to that of the empty vector normalized to cotransfected TK-Renilla luciferase activity. Error bars = standard deviations from the means. $P < 0.05$ between WH and DP.

domestic pigs and 11 warthogs (domestic pig sequences, EMBL accession numbers FN424224 to FN424228; warthog sequences, EMBL accession numbers FN424229 to FN424239). All domestic pig sequences were identical to each other over this region, and similarly, all warthog sequences were identical to each other over this region.

Predicted structural changes as a consequence of *RELA* sequence variation. Given that even single amino acid changes can have significant effects on protein structure, we performed *in silico* analyses on the identified changes in *RELA* in an attempt to determine the structural consequences of the amino acid substitutions. At the three sites, the greater hydrophobicity was conferred by the threonine at 448 in the domestic pig sequence and the prolines at 385 and 531 in the warthog sequence. No difference in structural disorder was evident (Phyre analysis) (25), and the only site predicted to be highly likely (0.994) to undergo phosphorylation was the domestic pig serine at 531 (485S = 0.072; 448T = 0.0142) (NetPhos2.0) (4). However, the Phospho.ELM database infers that the threonine at 448 could also be a target for phosphorylation (23).

Basal activity of porcine *RELA* variants. To determine whether the identified sequence variation between domestic pig *RELA* and warthog *RELA* affects NF- κ B activity, we established a cell transfection assay. Cells were transiently transfected in duplicate using a triple plasmid cotransfection strategy involving the following: (i) 1 μ g expression vector for FLAG-tagged domestic pig or warthog *RELA* or an empty vector control (pFLAG-CMV4; Sigma Aldrich); (ii) 1 μ g NF- κ B reporter plasmid comprising 4 copies of the NF- κ B consensus binding sequence, driving expression of firefly luciferase (pNF κ B-Luc; BD Biosciences, Clontech); and (iii) 1 μ g transfection control vector expressing Renilla luciferase driven off the herpes simplex virus (HSV) thymidine kinase (TK) promoter (pRL-TK; Promega). Cells were harvested at 24 h posttransfection, and a dual luciferase assay (firefly luciferase activity relative to Renilla luciferase activity) was performed on quantified protein extracts. Luciferase activity values for each *RELA* type were averaged, and fold differences were calculated against background values for empty vector transfections. This strategy resulted in the delivery of an equivalent amount

of either domestic pig or warthog *RELA*, as determined by Western blotting (Fig. 2A). Initially, we established the assay in COS-7 cells, which have a functional endogenous *RELA* gene. Repeat transfection experiments ($n = 5$) demonstrated that domestic pig *RELA* is 50% more active than warthog *RELA* (Fig. 2B). A similar (44%) differential activity was also evident in *RELA*^{-/-} mouse embryonic fibroblasts (MEFs) that lacked endogenous *RELA* activity (derived from *RELA*^{-/-} murine embryos) using the same transfection regime ($n = 3$) (Fig. 2C).

We attempted to produce stably transfected *RELA*^{-/-} MEFs for the domestic pig and warthog *RELA* genes. Although we could generate numerous colonies for warthog *RELA*, we were reproducibly unsuccessful in generating those for domestic pig *RELA*. It is possible that the higher activity level of domestic pig *RELA* may not be compatible with survival in these cells; a similar scenario may have occurred during a study investigating phosphorylation at the equivalent site (S529) in human *RELA* (54).

Effect of *RELA* sequence variation on induced NF- κ B activity. To determine if the ability to induce NF- κ B activity was altered by the different pig *RELA* proteins, we stimulated transiently transfected *RELA*^{-/-} MEFs with known inducers of NF- κ B signaling (9, 14, 15, 16, 42–44, 56). Induction with tumor necrosis factor alpha (TNF- α) nearly doubled (90% increase) warthog *RELA* activity, whereas domestic pig *RELA* showed a 70% increase in activity (Fig. 3). We tested a further three known NF- κ B-inducing agents, lipopolysaccharide, phorbol-12-myristate-13-acetate, and hydrogen peroxide. For all three additional agents, the fold induction observed for warthog *RELA* was always greater than that of domestic pig (Fig. 3). FLAG immunostaining of COS-7 cells transfected with either domestic pig or warthog *RELA* and treated with phorbol-12-myristate-13-acetate did not identify any gross differences in cytoplasm-nuclear transit time (*RELA* of both species was in the nucleus within 1 min and back in the cytoplasm after 20 min) (data not shown).

Identification of functional mutations in porcine *RELA*. In our sequence data, we identified three amino acid differences between domestic pig *RELA* and warthog pig *RELA*. To determine if the *in vitro* difference in domestic pig and warthog




FIG. 3. Induction of *RELA* allelic variants. *RELA*^{-/-} MEFs were transiently cotransfected with 1 μ g warthog (clear box) or 1 μ g domestic pig (filled box) *RELA*. Cells were treated with TNF- α (30 ng/ml), lipopolysaccharide (LPS; 10 μ g/ml), phorbol-12-myristate 13-acetate (PMA; 20 mM), and hydrogen peroxide (10 μ M) immediately after and for the duration of the transfection. Cells were harvested at 24 h. *RELA*-induced NF- κ B-luciferase (1 μ g) activity was normalized to cotransfected TK-Renilla luciferase (1 μ g) activity and presented as the fold induction above nondrug-treated cells (relative value of 1 depicted by the line).

RELA activities was a cumulative effect of these three differences, or due to one or other individual mutations, we generated versions of *RELA* carrying the following single-base changes: T448A, S485P, and S531P. Transfection of these *RELA* variants into *RELA*^{-/-} MEFs demonstrated that the majority of the reduced basal activity observed for warthog *RELA* compared to that observed for domestic pig *RELA* was attributed to the S531P mutation (Fig. 4).

DISCUSSION

In this study, we have taken a candidate approach, identifying genes which may affect the severity of the host response to ASFV infection in the highly susceptible domestic pig and ASFV-tolerant warthog. The ASFV immunomodulatory factor A238L is known to interact with components of both NF- κ B and NFAT host signaling pathways (11, 28, 29, 36, 49). We have sequenced five key factors in these pathways, including three A238L-targeted proteins, *RELA*, *PPP3CB*, and *PPIA*, and the two proteins it mimics, *NFKBIA* and *NFATC1* (regulatory domain). Modest sequence differences have been identified at the cDNA (mRNA) level between the domestic pig and warthog; however, the majority of these are synonymous (silent) and do not alter the resulting amino acid sequences. We also sequenced the same genes from the ancient babirusa and observed a small number of amino acid differences between this species and the other two species of pig. These differences likely reflect the long evolutionary distance that exists between these species (38).

Despite the high degree of conservation observed between warthog and domestic pig *PPP3CB*-, *PPIA*-, *NFKBIA*-, and *NFATC1*-translated protein sequences, significant variation was detected in *RELA*. *RELA* is the predominant member of the heterodimeric transcription factor NF- κ B (47). Moreover, the sequence variation includes a phosphorylation site in transactivation domain 1 (position 531), which is highly conserved across mammals and has been demonstrated to modulate the activity of human *RELA* (equivalent to S529) (54, 55). The function of approximately one-third of all eukaryote proteins is controlled by phosphorylation; thus, the observed S531P sequence variation represents an intriguing candidate regulator




FIG. 4. Comparison of individual allelic *RELA* variation on NF- κ B activity. Comparison of NF- κ B-luciferase activity in *RELA*^{-/-} MEFs after 24 h transient transfection with 1 μ g warthog (WH) or 1 μ g domestic pig (DP) *RELA* or 1 μ g of *RELA* variants encoding the individual amino acid substitutions T448A, S485P, and S531P. Activity is presented as the fold difference between *RELA*-induced NF- κ B-luciferase activity normalized to cotransfected TK-Renilla luciferase activity relative to that of the empty vector normalized to cotransfected TK-Renilla luciferase activity. Error bars = standard deviations from the means, *, $P < 0.08$ versus DP.

for the reduced pathology observed in African pigs infected with ASFV. We demonstrate that the genetic variation in the *RELA* sequence between the domestic pig and warthog is reflected in NF- κ B activity *in vitro*, with warthog *RELA* displaying significantly reduced basal and induced NF- κ B activity. We discuss three (not mutually exclusive) scenarios of how the genetic variation we have identified between the domestic pig and warthog may underlie the dramatic phenotypic difference in how these two pig species respond to ASFV.

First, as suggested by our *in vitro* assays, warthog *RELA* is inherently less active than the domestic pig *RELA*, exhibiting lower basal and induced levels. In a study of macrophage transcription profiles following ASFV infection *in vitro*, several factors within the NF- κ B signaling pathway displayed elevated expression (i.e., *NFKB1*, *NFKBIA*); however, *RELA* expression was not observed to be altered (56). The impact of ASFV infection on the warthog *RELA* expression level is not known, since this study used only domestic pig macrophages. Therefore, although we have not determined whether the porcine *RELA* variants are differentially phosphorylated, it would appear that variation in domestic pig *RELA* activity may not be determined by altered expression levels. Alternatively, the warthog will have to function with reduced basal NF- κ B activity, which would presumably indicate an adapted NF- κ B-dependent transcriptome between the two species.

In the second scenario, phosphorylation of domestic pig *RELA* at S531 activates a set(s) of genes which are not activated by warthog *RELA*, as it lacks this phosphorylation site. Much is known about activation of the NF- κ B pathway, with data coming primarily from studies of human *RELA* (47). NF- κ B activity is regulated by two different mechanisms. The classical canonical pathway involves inhibitors (e.g., *NFKBIA*) that sequester this transcription factor in the cytoplasm until they are proteolytically degraded by the ubiquitin pathway (24). The alternative pathway revolves around posttranscriptional modifications, predominantly phosphorylation of *RELA* (46). At least 8 inducible phosphorylation sites have been identified in human *RELA*, which enable transcription of subsets of NF- κ B-dependent genes (26, 45). One such site is S529, which is equivalent to domestic pig S531, suggesting that this site may perform a similar role. Indeed, differential expression of subsets of immune and inflammatory proteins as a result of S531

phosphorylation could play a role in determining ASFV pathogenesis in the domestic pig. Furthermore, phosphorylation not only controls specific transcription profiles but also can underlie the developmental timing in gene activation (13). It is tempting to speculate that if a similar mechanism is applied through the domestic pig S531 site, then it could also play a role in the gross physical differences that characterize the various pig species.

In the third scenario, we consider whether the S531P variation in RELA results in different outcomes of interaction with A238L during ASFV infection. Phosphorylation of human S529 (equivalent to domestic pig S531) is inhibited by the interaction of RELA with NFKBIA; only upon activation and degradation of NFKBIA can S529 phosphorylation occur (55). During ASFV infection, NFKBIA is degraded and replaced by A238L, which mimics NFKBIA but is not susceptible to proteolytic degradation (36, 49). As a result, in domestic pig cells infected with ASFV, S531 may not be exposed for phosphorylation. In comparison, warthogs lack this phosphorylation site; therefore, the ability of A238L to block phosphorylation at this site is irrelevant. The role of NFKBIA is to prevent nuclear translocation of NF- κ B (24). We did not observe any gross temporal differences in nuclear transportation rates upon stimulation between cells expressing domestic pig RELA and warthog RELA; likewise, phosphorylation of human S529 also does not affect nuclear translocation (54). Furthermore, studies using recombinant ASFV lacking A238L indicate that neither nuclear import nor export of RELA is affected by this immune modulator (48). This suggests that the immunomodulatory functions of A238L are not the result of preventing nuclear translocation in a manner similar to that used by NFKBIA. Instead, A238L may inhibit NF- κ B-mediated transcription by other mechanisms, for example, by preventing phosphorylation of RELA in domestic pigs, as discussed above. Whether such differences in how domestic pig RELA and warthog RELA interact with A238L exist will require further investigation. Likewise, elucidating how different mechanisms governing RELA activity and NF- κ B-mediated transcription have evolved in these species, and determining the full extent of their functional implications on the immune system and wider transcriptome, will require additional study.

ASFV is highly infectious, with virulent isolates causing an acute, rapidly fatal hemorrhagic fever in domestic pigs (10, 34). In contrast, the porcine species endogenous to Africa tolerate the virus. Outbreaks have significant economic repercussions in addition to welfare concerns. In Africa, ASFV limits the use of the genetically improved breeding pig stock that is pervasive in the Eurasian landscape. Furthermore, ASF poses a constant threat to Europe and Asia, as documented by the list of outbreaks that have occurred over the last 30 to 40 years. As our world climate changes and the international movement of pork products continues to rise, this risk may well increase. No effective vaccine has been developed, so many look to genetic strategies to mitigate the geographical limits of pig breeding imposed by ASFV. We have demonstrated that a polymorphic RELA variant found in warthogs has the potential to underlie the difference between tolerance and rapid death upon infection with ASFV.

ACKNOWLEDGMENTS

This work was supported by the BBSRC (United Kingdom) and a Genus/PIC Postgraduate Studentship. C.J.P. was also funded by a Postgraduate Centenary Fellowship (University of Edinburgh). L.G. was funded through a Faraday-BBSRC CASE studentship.

RELA^{-/-} MEFs were kindly donated by Ron Hay (University of Dundee). The pcDNA-A238L expression vector was kindly donated by Linda Dixon (Institute for Animal Health, Pirbright). We are grateful to Susan Rhind, Liz Glass, Sarah Howie, and Paul Hopwood (University of Edinburgh) for their support of this work.

REFERENCES

- Afonso, C. L., J. G. Neilan, G. F. Kutish, and D. L. Rock. 1996. An African swine fever virus Bcl-2 homolog, 5-HL, suppresses apoptotic cell death. *J. Virol.* **70**:4858–4863.
- Anderson, E. C., G. H. Hutchings, N. Mukarati, and P. J. Wilkinson. 1998. African swine fever infection of the bushpig (*Potamochoerus porcus*) and its significance to the epidemiology of the disease. *Vet. Microbiol.* **62**:1–15.
- Anderson, E. C., S. M. Williams, S. P. Fisher-Hoch, and P. J. Wilkinson. 1987. Arachidonic acid metabolites in the pathophysiology of thrombocytopenia and haemorrhage in acute African swine fever. *Res. Vet. Sci.* **42**:387–394.
- Blom, N., S. Gammeltoft, and S. Brunak. 1999. Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *J. Mol. Biol.* **294**:1351–1362.
- Borca, M. V., et al. 1998. Deletion of a CD2-like gene, 8-DR, from African swine fever virus affects viral infection in domestic swine. *J. Virol.* **72**:2881–2889.
- Borca, M. V., et al. 1994. An African swine fever virus gene with similarity to the T-lymphocyte surface antigen CD2 mediates hemadsorption. *Virology* **199**:463–468.
- Brun, A., C. Rivas, M. Esteban, J. M. Escibano, and C. Alonso. 1996. African swine fever virus gene A179L, a viral homologue of bcl-2, protects cells from programmed cell death. *Virology* **225**:227–230.
- Brun, A., F. Rodriguez, J. M. Escibano, and C. Alonso. 1998. Functionality and cell anchorage dependence of the African swine fever virus gene A179L, a viral bcl-2 homolog, in insect cells. *J. Virol.* **72**:10227–10233.
- Carrasco, L., et al. 2002. African swine fever: expression of interleukin-1 alpha and tumour necrosis factor-alpha by pulmonary intravascular macrophages. *J. Comp. Pathol.* **126**:194–201.
- Colgrove, G. S., E. O. Haelterman, and L. Coggins. 1969. Pathogenesis of African swine fever in young pigs. *Am. J. Vet. Res.* **30**:1343–1359.
- Dixon, L. K., et al. 2004. African swine fever virus proteins involved in evading host defence systems. *Vet. Immunol. Immunopathol.* **100**:117–134.
- Dixon, L. K., et al. 2005. Family *Asfarviridae*, p. 135–143. *In* C. M. Fauquet, M. A. Mayo, J. Maniloff, U. Desselberger, and L. A. Ball (ed.), *Virus taxonomy*. Eighth report of the International Committee on Taxonomy of Viruses. Elsevier Academic Press, London, England.
- Gavalda, N., H. Gutierrez, and A. M. Davies. 2009. Developmental switch in NF-kappaB signalling required for neurite growth. *Development* **136**:3405–3412.
- Ghosh, S., M. J. May, and E. B. Kopp. 1998. NF- κ B and Rel proteins: evolutionary conserved mediators of immune responses. *Annu. Rev. Immunol.* **16**:225–260.
- Gil, S., et al. 2003. Expression at mRNA level of cytokines and A238L gene in porcine blood-derived macrophages infected in vitro with African swine fever virus (ASFV) isolates of different virulence. *Arch. Virol.* **148**:2077–2097.
- Gomez del Moral, M., et al. 1999. African swine fever virus infection induces tumor necrosis factor alpha production: implications in pathogenesis. *J. Virol.* **73**:2173–2180.
- Gomez-Villamandos, J. C., et al. 1997. African swine fever virus infection of bone marrow: lesions and pathogenesis. *Vet. Pathol.* **34**:97–107.
- Gomez-Villamandos, J. C., et al. 1998. Thrombocytopenia associated with apoptotic megakaryocytes in a viral haemorrhagic syndrome induced by a moderately virulent strain of African swine fever virus. *J. Comp. Pathol.* **118**:1–13.
- Gomez-Villamandos, J. C., et al. 1995. Experimental African swine fever: apoptosis of lymphocytes and virus replication in other cells. *J. Gen. Virol.* **76**(Pt. 9):2399–2405.
- Gomez-Villamandos, J. C., et al. 1995. A pathological study of the perisinusoidal unit of the liver in acute African swine fever. *Res. Vet. Sci.* **59**:146–151.
- Gomez-Villamandos, J. C., et al. 1995. Pathological changes in the renal interstitial capillaries of pigs inoculated with two different strains of African swine fever virus. *J. Comp. Pathol.* **112**:283–298.
- Guerini, D., and C. B. Klee. 1989. Cloning of human calcineurin A: evidence for two isozymes and identification of a proline structural domain. *Proc. Natl. Acad. Sci. U. S. A.* **86**:9183–9187.

23. Hammet, A., et al. 2003. FHA domains as phospho-threonine binding modules in cell signaling. *IUBMB Life* **55**:23–27.
24. Karin, M. 2006. Nuclear factor-kappaB in cancer development and progression. *Nature* **441**:431–436.
25. Kelley, L. A., and M. J. Sternberg. 2009. Protein structure prediction on the Web: a case study using the Phyre server. *Nat. Protoc.* **4**:363–371.
26. Law, M., P. Corsino, N. T. Parker, and B. K. Law. 2010. Identification of a small molecule inhibitor of serine 276 phosphorylation of the p65 subunit of NF-kappaB using in silico molecular docking. *Cancer Lett.* **291**:217–224.
27. McPartlin, A. E., H. M. Barker, and P. T. Cohen. 1991. Identification of a third alternatively spliced cDNA encoding the catalytic subunit of protein phosphatase 2B beta. *Biochim. Biophys. Acta* **1088**(2):308–310.
28. Miskin, J. E., C. C. Abrams, and L. K. Dixon. 2000. African swine fever virus protein A238L interacts with the cellular phosphatase calcineurin via a binding domain similar to that of NFAT. *J. Virol.* **74**:9412–9420.
29. Miskin, J. E., C. C. Abrams, L. C. Goatley, and L. K. Dixon. 1998. A viral mechanism for inhibition of the cellular phosphatase calcineurin. *Science* **281**:562–565.
30. Neilan, J. G., et al. 1993. An African swine fever virus gene with similarity to the proto-oncogene bcl-2 and the Epstein-Barr virus gene BHRF1. *J. Virol.* **67**:4391–4394.
31. Nogal, M. L., et al. 2001. African swine fever virus IAP homologue inhibits caspase activation and promotes cell survival in mammalian cells. *J. Virol.* **75**:2535–2543.
32. Oura, C. A., P. P. Powell, E. Anderson, and R. M. Parkhouse. 1998. The pathogenesis of African swine fever in the resistant bushpig. *J. Gen. Virol.* **79**:1439–1443.
33. Oura, C. A., P. P. Powell, and R. M. Parkhouse. 1998. African swine fever: a disease characterized by apoptosis. *J. Gen. Virol.* **79**:1427–1438.
34. Penrith, M.-L., G. R. Thomson, and A. D. S. Bastos. 2004. African swine fever, p. 1088–1119. In J. A. W. Coetzer and R. C. Tustin (ed.), *Infectious diseases of livestock*, 2nd ed., vol. 2. Oxford University Press, Oxford, United Kingdom.
35. Plowright, W., J. Parker, and M. A. Pierce. 1969. The epizootiology of African swine fever in Africa. *Vet. Rec.* **85**:668–674.
36. Powell, P. P., L. K. Dixon, and R. M. Parkhouse. 1996. An IkappaB homologue encoded by African swine fever virus provides a novel mechanism for down-regulation of proinflammatory cytokine responses in host macrophages. *J. Virol.* **70**:8527–8533.
37. Ramiro-Ibanez, F., A. Ortega, A. Brun, J. M. Escribano, and C. Alonso. 1996. Apoptosis: a mechanism of cell killing and lymphoid organ impairment during acute African swine fever virus infection. *J. Gen. Virol.* **77**(Pt. 9): 2209–2219.
38. Randi, E., V. Lucchini, and C. Hoong Dieng. 1996. Evolutionary genetics of the Suiformes as reconstructed using mtDNA sequencing. *J. Mamm. Evol.* **3**:163–194.
39. Rodriguez, C. I., et al. 2002. African swine fever virus IAP-like protein induces the activation of nuclear factor kappa B. *J. Virol.* **76**:3936–3942.
40. Rodriguez, F., A. Fernandez, J. P. Martin de las Mulas, M. A. Sierra, and A. Jover. 1996. African swine fever: morphopathology of a viral haemorrhagic disease. *Vet. Rec.* **139**:249–254.
41. Rodriguez, J. M., R. J. Yanez, F. Almazan, E. Vinuela, and J. F. Rodriguez. 1993. African swine fever virus encodes a CD2 homolog responsible for the adhesion of erythrocytes to infected cells. *J. Virol.* **67**:5312–5320.
42. Salguero, F. J., et al. 2002. Changes in macrophages in spleen and lymph nodes during acute African swine fever: expression of cytokines. *Vet. Immunol. Immunopathol.* **90**:11–22.
43. Salguero, F. J., P. J. Sanchez-Cordon, A. Nunez, M. Fernandez de Marco, and J. C. Gomez-Villamandos. 2005. Proinflammatory cytokines induce lymphocyte apoptosis in acute African swine fever infection. *J. Comp. Pathol.* **132**:289–302.
44. Salguero, F. J., et al. 2004. Apoptosis of thymocytes in experimental African swine fever virus infection. *Histol. Histopathol.* **19**:77–84.
45. Sasaki, C. Y., T. J. Barberi, P. Ghosh, and D. L. Longo. 2005. Phosphorylation of RelA/p65 on serine 536 defines an I{kappa}B{alpha}-independent NF-{kappa}B pathway. *J. Biol. Chem.* **280**:34538–34547.
46. Schmitz, M. L., S. Bacher, and M. Kracht. 2001. I{kappa}B-independent control of NF- κ B activity by modulatory phosphorylations. *Trends Biochem. Sci.* **26**(3): 186–190.
47. Schmitz, M. L., I. Mattioli, H. Buss, and M. Kracht. 2004. NF-kappaB: a multifaceted transcription factor regulated at several levels. *ChemBiochem* **5**:1348–1358.
48. Silk, R. N., G. C. Bowick, C. C. Abrams, and L. K. Dixon. 2007. African swine fever virus A238L inhibitor of NF-kappaB and of calcineurin phosphatase is imported actively into the nucleus and exported by a CRM1-mediated pathway. *J. Gen. Virol.* **88**:411–419.
49. Tait, S. W. G., E. B. Reid, D. R. Greaves, T. E. Wileman, and P. P. Powell. 2000. Mechanism of inactivation of NF- κ B by a viral homologue of I{kappa}B α . *J. Biol. Chem.* **275**:34656–34664.
50. Thomson, G. R. 1985. The epidemiology of African swine fever: the role of free-living hosts in Africa. *Onderstepoort J. Vet. Res.* **52**:201–209.
51. Thomson, G. R., M. D. Gainaru, and A. F. Van Dellen. 1980. Experimental infection of warthogs (*Phacochoerus aethiopicus*) with African swine fever virus. *Onderstepoort J. Vet. Res.* **47**:19–22.
52. Villeda, C. J., S. M. Williams, P. J. Wilkinson, and E. Vinuela. 1993. Consumption coagulopathy associated with shock in acute African swine fever. *Arch. Virol.* **133**:467–475.
53. Villeda, C. J., S. M. Williams, P. J. Wilkinson, and E. Vinuela. 1993. Haemostatic abnormalities in African swine fever a comparison of two virus strains of different virulence (Dominican Republic '78 and Malta '78). *Arch. Virol.* **130**:71–83.
54. Wang, D., and A. S. Baldwin, Jr. 1998. Activation of nuclear factor- κ B-dependent transcription by tumor necrosis factor- α is mediated through phosphorylation of RelA/p65 on serine 529. *J. Biol. Chem.* **273**:29411–29416.
55. Wang, D., S. D. Westerheide, J. L. Hanson, and A. S. Baldwin, Jr. 2000. Tumor necrosis factor α -induced phosphorylation of RelA/p65 on Ser⁵²⁹ is controlled by casein kinase II. *J. Biol. Chem.* **275**:32592–32597.
56. Zhang, F., et al. 2006. Macrophage transcriptional responses following in vitro infection with a highly virulent African swine fever virus isolate. *J. Virol.* **80**:10514–10521.