

**SINGLE AND COMBINED USE OF FALL RISK INCREASING DRUGS AND FRACTURE RISK:
A POPULATION-BASED CASE CONTROL STUDY**

Jonathan Hauff¹
M Rottenkolber¹
Patrick Oehler¹
S Fischer¹
J Gensichen¹
M Drey²
G. Caleb Alexander, MD, MS¹⁻³
B Guthrie⁴
Tobias Dreischulte, PhD^{1*}

Affiliations

1. Institute of General Practice and Family Medicine, University Hospital of Ludwig-Maximilians-University Munich, Nußbaumstr. 5, 80336, Munich, Germany.
2. Center for Drug Safety and Effectiveness, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD
3. Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD
4. Department of Medicine IV, Geriatrics, University Hospital, Ludwig-Maximilians-Universität Munich, Ziemssenstraße 5, Munich 80336, Germany.

Correspondence

Professor Dr Tobias Dreischulte, PhD, MSc MPharm
Nussbaumstr. 5
80336 München
Phone: ++49(0)89440044557
Email: tobias.dreischulte@med.uni-muenchen.de

Funding

This work was partially funded by 'Verein zur Förderung von Wissenschaft und Forschung (WiFoMed)' at the medical faculty of the Ludwig-Maximilians-University, Munich, Germany

Length: Abstract 250 words; text 2997 words; 34 references; 3 tables; 1 figure; x eTables; y eFigures

