



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Vascularizing the kidney in the embryo and organoid: questioning assumptions about renal vasculogenesis

Citation for published version:

Munro, D & Davies, J 2018, 'Vascularizing the kidney in the embryo and organoid: questioning assumptions about renal vasculogenesis', *Journal of the American Society of Nephrology*.
<https://doi.org/10.1681/ASN.2018020179>

Digital Object Identifier (DOI):

[10.1681/ASN.2018020179](https://doi.org/10.1681/ASN.2018020179)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of the American Society of Nephrology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Perspectives article for *JASN*:

Vascularizing the kidney in the embryo and organoid: questioning assumptions about renal vasculogenesis

Authors: David A. D. Munro^{1*} and Jamie A. Davies¹

Running title: Vascularizing the kidney

Affiliations: Deanery of Biomedical Sciences, University of Edinburgh

***Corresponding author address:** Room 339, Hugh Robson Building, 15 George Square, University of Edinburgh, Edinburgh, UK, EH8 9XD

***Email:** s1471287@sms.ed.ac.uk

Vascularizing the kidney in the embryo and organoid: questioning assumptions about renal vasculogenesis

Despite much debate, there remains no clear consensus on how the kidney becomes vascularized. In general, blood vessels form by one of two processes, vasculogenesis (assembly of vessels *de novo* from mesodermal precursors), and angiogenesis (branching of new vessels from existing ones). In principle, renal endothelia may assemble through *vasculogenesis-only* (improbable and not supported), *angiogenesis-only* (conceivable but overlooked), or a *combination* of both processes (the option favoured by most accounts; for a summary of pertinent reviews since 1995 see <http://dx.doi.org/10.7488/ds/2307>).

Though the *combination* option is favoured, opinions are inconsistent regarding the relative contribution of each mechanism. We challenge this prevailing viewpoint and argue that the renal vasculature (specifically, the endothelial component) instead develops through *angiogenesis-only*.

An early proponent for kidney vasculogenesis was Herring (1900)¹, who speculated that glomerular capillaries are generated from *in situ* precursors within the cleft of the S-shaped nephron. This concept persists, despite electron microscopy and immunohistochemistry images suggesting that these early glomerular capillaries instead stem from local, pre-existing vessels^{2,3}. Of course, still images only show cellular arrangement in the kidney at a single moment (at the point of fixation) and cannot prove or disprove dynamic processes such as angiogenesis or vasculogenesis. Here, time-lapse imaging of cultured embryonic kidneys has provided insights, showing fluorescence-tagged endothelia forming via angiogenesis⁴ and migrating from pre-existing vessels into the S-shaped cleft⁵ (Figure 1A).

More recent support for vasculogenesis comes from cross-transplantation experiments. In transplanted embryonic kidneys, both host- and graft-derived vessels usually developed, and observations of graft-derived vessels were taken as support for vasculogenesis. However, these studies assumed that the transplanted kidney rudiments were avascular, even when using embryonic day 12-12.5 (E12-E12.5) mouse kidneys. We find these to be highly vascularized (Figure 1B). In fact, even the earliest kidney contains blood vessels (E11; Figure 1C). More significantly, even if truly

avascular kidneys were transplanted and vasculogenesis occurred, this does not necessarily mean that it is the normal mechanism of kidney vascularisation. Transplanted kidneys are put into artificial environments that may encourage vasculogenesis – it is unclear how this relates to ‘natural’ development and we suggest that investigating kidney vascularization during normal development *in vivo* might be more revealing.

In vivo, mouse kidney vascularization starts by E11, when systemically connected vessels wrap around the ureteric bud⁶ (the precursor of collecting ducts and ureter). Over the next day, blood vessels continue to disperse throughout the kidney mesenchyme, in connection with extrinsic vessels⁶ (Figure 1B). Later, vessels at the border of the kidney form polygonal networks around populations of nephron progenitor cells⁶ (they organise this way in human embryos too⁷ - <https://transparent-human-embryo.com/?p=1001>; Figure 1D-E), while cortical and medullary vessels are simultaneously developing.

Based on our data, normal kidney vascularization relies on growth and remodelling of pre-existing vessels (angiogenesis) and does not depend on vasculogenesis at any point. When assessing the entire 3D vascular tree, isolated endothelia are never observed⁶. Even when immunostaining for stem cell leukemia (SCL), a marker of the most primitive endothelial progenitors⁸, staining is observed only within mature vessels that are traceable to the renal artery⁶. Other studies, however, suggest that a small population of renal endothelia are stroma-derived⁹, and the mechanisms by which these cells contribute to vascular development should be explored.

Uncovering the mechanisms of kidney vascularization may allow us to better recapitulate developmental processes to vascularize kidney organoids. Though organoids are generally improving, they lack anatomically realistic vascular systems. Based on the ‘angiogenesis-only’ hypothesis, we suggest that, rather than trying to induce vasculogenesis, methodological advances should promote invasion of renal organoids by exogenous vessels, both in culture and transplantation settings.

Illustrating the need for kidney organoid invasion by exogenous vessels, van den Berg et al¹⁰ generated pluripotent stem cell-derived kidney organoids and demonstrated that long periods of culture (where exogenous vessels are lacking) led to gradual loss

PERSPECTIVES

of endothelia, whereas transplantation into pre-vascularised host tissue (beneath the murine renal capsule) led to the development of a functional perfused vasculature.

van den Berg et al¹⁰ proposed that maturation of large kidney organoids in culture is limited due to hypoxia and metabolic deficiencies. To facilitate continued maturation in culture, it may be possible to generate perfused engineered vessels that can invade the organoid. Perfused vessels could supply oxygen and nutrients to the organoid, while also generating shear-forces to encourage vessel maturation. This could be achieved by connecting engineered vessels to a pumping system where the flow-input could be stringently controlled.

Improved *in vitro* vascularization and maturation prior to transplantation would allow for the engraftment of a functionally enhanced organoid. Upon transplantation, host-derived blood vessels can invade and connect with kidney organoid-derived vessels¹⁰. If a vascularized organoid with functioning glomeruli and nephrons can be transplanted, and its flow quickly re-established via anastomoses with the host's vasculature, the regenerative potential of kidney organoids may begin to be realised.

ACKNOWLEDGMENTS

Thanks to Karen Chapman and Peter Hohenstein for their useful comments.

DISCLOSURES

None to declare.

REFERENCES

1. Herring PT: The development of the Malpighian bodies of the kidney, and its relation to pathological changes which occur in them. *J Pathol* 6: 459–496, 1900.
2. Kazimierczak J: A study of scanning (SEM) and transmission (TEM) electron microscopy of the glomerular capillaries in developing rat kidney. *Cell Tissue Res* 212: 241-55, 1980
3. Naruse K, Fujieda M, Miyazaki E, Hayashi Y, Toi M, Fukui T, et al.: An immunohistochemical study of developing glomeruli in human fetal kidneys. *Kidney Int* 57:1836-46, 2000.
4. Halt KJ, Pärssinen HE, Junttila SM, Saarela U, Sims-Lucas S, Koivunen P, et al.: CD146⁺ cells are essential for kidney vasculature development. *Kidney Int* 90: 311–324, 2016.
5. Saarela U, Akram SU, Desgrange A, Rak-Raszewska A, Shan J, Cereghini S, et al.: Novel fixed z-direction (FiZD) kidney primordia and an organoid culture system for time-lapse confocal imaging. *Development*. 144:1113-1117, 2017
6. Munro DAD, Hohenstein P, Davies JA: Cycles of vascular plexus formation within the nephrogenic zone of the developing mouse kidney. *Sci Rep*. 7: 3273, 2017
7. Belle M, Godefroy D, Couly G, Malone SA, Collier F, Giacobini P, et al.: Tridimensional visualization and analysis of early human development. *Cell* 169: 161-173, 2017
8. Drake CJ, Brandt SJ, Trusk TC, Little CD: TAL1/SCL is expressed in endothelial progenitor cells/angioblasts and defines a dorsal-to-ventral gradient of vasculogenesis. *Dev Biol*. 192: 17-30, 1997
9. Sims-Lucas S, Schaefer C, Bushnell D, Ho J, Logar A, Prochownik E, et al.: Endothelial Progenitors Exist within the Kidney and Lung Mesenchyme. *PLoS ONE*. 8: e65993, 2013
10. van den Berg CW, Ritsma L, Avramut MC, Wiersma LE, van den Berg BM, Leuning DG, et al.: Renal Subcapsular Transplantation of PSC-Derived Kidney Organoids Induces Neo-vasculogenesis and Significant Glomerular and Tubular Maturation In Vivo. *Stem Cell Reports*. 13: 751-765. 2018.

Figure 1. Kidney vascularization. **(A)** Earliest glomerular endothelia forming into the cleft of an S-shaped nephron from pre-existing vessels in time-lapse culture (white arrowhead; adapted from⁵). **(B)** Model of early kidney vascularization *in vivo*. Some argue that the first vessels form via vasculogenesis, but our results suggest otherwise⁶. UB, ureteric bud; WD, Wolffian duct; PWM, peri-Wolffian mesenchyme. Blood vessels shown in red. **(C)** The E11 kidney is vascularized by systemically connected vessels (these vessels carry erythrocytes and connect to major arteries; black arrowhead), surrounding the ureteric bud (adapted from⁶), which calls into question whether avascular kidneys can be dissected at any age. Scale bar: 100 μm . **(D)** Representative image of the vascular network at the periphery of the E16.5 kidney. Some argue that these peripheral vessels form via vasculogenesis; however, they carry blood, are always enclosed by basement membrane, and connect with pre-existing vessels that can be traced to renal arteries⁶. Scale bar: 50 μm . **(E)** As with mouse embryonic kidneys, peripheral vessels in human embryonic kidneys arrange as polygonal networks around nephron progenitors (prepared from data accessed at <https://transparent-human-embryo.com>⁷).