



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Nonuniversality and finite dissipation in decaying magnetohydrodynamic turbulence

**Citation for published version:**

Linkmann, MF, Berera, A, McComb, WD & McKay, ME 2015, 'Nonuniversality and finite dissipation in decaying magnetohydrodynamic turbulence', *Physical Review Letters*, vol. 114, no. 23, 235001. <https://doi.org/10.1103/PhysRevLett.114.235001>

**Digital Object Identifier (DOI):**

[10.1103/PhysRevLett.114.235001](https://doi.org/10.1103/PhysRevLett.114.235001)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Physical Review Letters

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## Nonuniversality and Finite Dissipation in Decaying Magnetohydrodynamic Turbulence

M. F. Linkmann,<sup>\*</sup> A. Berera,<sup>†</sup> W. D. McComb, and M. E. McKay

*SUPA, School of Physics and Astronomy, University of Edinburgh, Peter Guthrie Tait Road, Edinburgh H9 3FD, United Kingdom*

(Received 2 March 2015; published 11 June 2015)

A model equation for the Reynolds number dependence of the dimensionless dissipation rate in freely decaying homogeneous magnetohydrodynamic turbulence in the absence of a mean magnetic field is derived from the real-space energy balance equation, leading to  $C_\varepsilon = C_{\varepsilon,\infty} + C/R_- + O(1/R_-^2)$ , where  $R_-$  is a generalized Reynolds number. The constant  $C_{\varepsilon,\infty}$  describes the total energy transfer flux. This flux depends on magnetic and cross helicities, because these affect the nonlinear transfer of energy, suggesting that the value of  $C_{\varepsilon,\infty}$  is not universal. Direct numerical simulations were conducted on up to  $2048^3$  grid points, showing good agreement between data and the model. The model suggests that the magnitude of cosmological-scale magnetic fields is controlled by the values of the vector field correlations. The ideas introduced here can be used to derive similar model equations for other turbulent systems.

DOI: 10.1103/PhysRevLett.114.235001

PACS numbers: 47.65.-d, 47.27.Gs, 47.27.Jv, 52.30.Cv

Magnetohydrodynamic (MHD) turbulence is present in many areas of physics, ranging from industrial applications such as liquid metal technology to nuclear fusion and plasma physics, geo-, astro-, and solar physics, and even cosmology. The numerous different MHD flow types that arise in different settings due to anisotropy, alignment, different values of the diffusivities, to name only a few, lead to the question of universality in MHD turbulence, which has been the subject of intensive research by many groups [1–12]. The behavior of the (dimensionless) dissipation rate is connected to this problem, in the sense that correlation (alignment) of the different vector fields could influence the energy transfer across the scales [2,13,14], and thus possibly the amount of energy that is eventually dissipated at the small scales.

For neutral fluids it has been known for a long time that the dimensionless dissipation rate in forced and freely decaying homogeneous isotropic turbulence tends to a constant with increasing Reynolds number. The first evidence for this was reported by Batchelor [15] in 1953, while the experimental results reviewed by Sreenivasan in 1984 [16], and subsequent experimental and numerical work by many groups, established the now well-known characteristic curve of the dimensionless dissipation rate against Reynolds number; see [17–20], and references therein. For statistically steady isotropic turbulence, the theoretical explanation of this curve was recently found to be connected to the energy balance equation for forced turbulent flows [19], where the asymptote describes the maximal inertial transfer flux in the limit of infinite Reynolds number.

For freely decaying MHD, recent results suggest that the temporal maximum of the total dissipation tends to a constant value with increasing Reynolds number. The first evidence for this behavior in MHD was put forward in 2009 by Mininni and Pouquet [21], using results from direct

numerical simulations (DNSs) of homogeneous MHD turbulence. The temporal maximum of the total dissipation rate  $\varepsilon(t)$  became independent of Reynolds number at a Taylor-scale Reynolds number  $R_\lambda$  [measured at the peak of  $\varepsilon(t)$ ] of about 200.

Dallas and Alexakis [22] measured the dimensionless dissipation rate  $C_\varepsilon$  from DNS data, where  $\varepsilon$  was non-dimensionalized with respect to the initial values of the rms velocity  $U(t)$  and the integral length scale  $L(t)$  (here defined with respect to the total energy), for random initial fields with strong correlations between the velocity field and the current density. The authors compared data with Ref. [21], and again it was found that  $C_\varepsilon \rightarrow \text{const}$  with increasing Reynolds number. Interestingly the approach to the asymptote was slower than for the data of Ref. [21].

In this Letter we propose a model for the Reynolds number dependence of the dimensionless dissipation rate derived from the energy balance equation for MHD turbulence in terms of Elsässer fields [23], which predicts nonuniversal values of the dimensionless dissipation rate in the infinite Reynolds number limit. In order to compare the predictions of the model against data, we carried out a series of DNSs of decaying MHD turbulence without a mean magnetic field. First we explain the derivation of the model equation, then proceed to a description of our numerical simulations and, subsequently, compare the model to DNS results. We conclude with a discussion of the results and suggestions for further research.

The equations describing incompressible decaying MHD flows are

$$\partial_t \mathbf{u} = -\frac{1}{\rho} \nabla P - (\mathbf{u} \cdot \nabla) \mathbf{u} + \frac{1}{\rho} (\nabla \times \mathbf{b}) \times \mathbf{b} + \nu \Delta \mathbf{u}, \quad (1)$$

$$\partial_t \mathbf{b} = (\mathbf{b} \cdot \nabla) \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{b} + \eta \Delta \mathbf{b}, \quad (2)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{and} \quad \nabla \cdot \mathbf{b} = 0, \quad (3)$$

where  $\mathbf{u}$  denotes the velocity field,  $\mathbf{b}$  the magnetic induction expressed in Alfvén units,  $\nu$  the kinematic viscosity,  $\eta$  the resistivity,  $P$  the pressure, and  $\rho = 1$  the density. For simplicity, and in order to compare to results in the literature, we consider the case of unit magnetic Prandtl number, that is  $Pm = \nu/\eta = 1$ .

For freely decaying MHD turbulence the decay rate of the total energy  $\varepsilon_D = -\partial_t E_{\text{tot}}$  equals the total dissipation rate  $\varepsilon$ , and the time evolution of the total energy is governed by the energy balance equation of MHD turbulence in real space, which is derived from the MHD equations (1)–(3). This suggests that the energy balance equation can be used in order to derive the Reynolds number dependence of the total dissipation rate.

Since we are interested in the total dissipation  $\varepsilon = \varepsilon_{\text{mag}} + \varepsilon_{\text{kin}}$ , where  $\varepsilon_{\text{mag}} = 2\eta \int_0^\infty dk k^2 E_{\text{mag}}(k)$ , and  $\varepsilon_{\text{kin}} = 2\nu \int_0^\infty dk k^2 E_{\text{kin}}(k)$  [ $E_{\text{mag}}(k)$  and  $E_{\text{kin}}(k)$  denoting magnetic and kinetic energy spectra], are the magnetic and kinetic dissipation rates, respectively, we could take two approaches, either formulating the energy balance in terms of the primary fields  $\mathbf{u}$  and  $\mathbf{b}$  or in terms of the Elsässer fields  $z^\pm = \mathbf{u} \pm \mathbf{b}$ . Since

$$\partial_t \langle |z^\pm|^2 \rangle = 2\partial_t E_{\text{tot}} \pm 2\partial_t H_c, \quad (4)$$

where  $H_c = \langle \mathbf{u} \cdot \mathbf{b} \rangle$  is the cross helicity, we can describe the total dissipation either by the energy balance equations for  $\langle |z^\pm|^2 \rangle$  [23] or by the sum of the energy balance equations for  $E_{\text{mag}}(t) = \int_0^\infty dk E_{\text{mag}}(k)$  and  $E_{\text{kin}}(t) = \int_0^\infty dk E_{\text{kin}}(k)$  [24,25].

This, however, is not the case if we are interested in the *dimensionless* dissipation rate. Unlike in hydrodynamics, there are several choices of scales with which to non-dimensionalize  $\varepsilon(t)$ , and thus with respect to which to define an MHD analogue to the Taylor surrogate expression [15,18]. For example  $U$  and  $L$  could be used, or the rms  $\mathbf{b}$  field  $B$  and  $L$  or  $U$  and  $L_{\text{kin}}$  etc., or scales defined with respect to  $z^\pm$ . The physical interpretation is different for the different scaling quantities. Since the total dissipation must equal the total flux of energy passed through the scales by the kinetic and magnetic energy transfer terms, a scaling with  $U$  will be appropriate only for hydrodynamic transfer as this transfer term scales as  $U^3/L_{\text{kin}}$ . All other transfer terms include  $\mathbf{b}$  and  $\mathbf{u}$  and thus should be scaled accordingly. This also precludes the most straightforward generalization of the Taylor surrogate, which would be a scaling of  $\varepsilon$  with  $L$  and  $\sqrt{U^2 + B^2}$ . A hydrodynamic transfer term would then be scaled partly with magnetic quantities, while the appropriate scaling should only involve kinetic quantities.

Instead, we propose to define the dimensionless dissipation rate for MHD turbulence with respect to the Elsässer variables

$$C_\varepsilon = \frac{C_\varepsilon^+ + C_\varepsilon^-}{2} \equiv \frac{1}{2} \left( \frac{\varepsilon L_+}{z^{+2} z^-} + \frac{\varepsilon L_-}{z^{-2} z^+} \right), \quad (5)$$

where  $L_\pm = (3\pi \int_0^\infty dk k^{-1} \langle |z^\pm|^2 \rangle) / (4 \int_0^\infty dk \langle |z^\pm|^2 \rangle)$  are the integral scales defined with respect to  $z^\pm$ , and  $z^\pm$  denote the rms values of  $z^\pm$  [26].

Using this definition we can now consistently non-dimensionalize the evolution equations of  $\langle |z^\pm|^2 \rangle$ . For conciseness we outline the arguments for the  $\langle |z^+|^2 \rangle$  case, since the  $\langle |z^-|^2 \rangle$  case proceeds analogously [27].

Following [23] the energy balance for  $\langle |z^+|^2 \rangle$  reads for the case  $Pm = 1$

$$-\frac{1}{2} \partial_t \langle |z^+|^2 \rangle = -\frac{3}{4} \partial_t B_{LL}^{++} - \frac{\partial_r}{r^4} \left( \frac{3r^4}{2} C_{LL,L}^{+++} \right) + \frac{3(\nu + \eta)}{2r^4} \partial_r (r^4 \partial_r B_{LL}^{++}), \quad (6)$$

where  $C_{LL,L}^{+++}(r)$  and  $B_{LL}^{++}(r)$  are the longitudinal third-order correlation function and the second-order longitudinal structure function of the Elsässer fields, respectively. The definitions of these functions can be found in the Supplemental Material [27]. Using Eq. (4) one can express the left-hand side of Eq. (6) in terms of  $\varepsilon(t)$  and  $\partial_t H_c$ .

If we now introduce the nondimensional variable  $\sigma = r/L_+$  [3] and non-dimensionalize Eq. (6) with respect to  $z^\pm$  and  $L_+$  as proposed in the definition of  $C_\varepsilon$  in Eq. (5), we obtain

$$C_\varepsilon^+ = -\frac{\partial_\sigma}{\sigma^4} \left( \frac{3\sigma^4 C_{LL,L}^{+++}}{2z^{+2} z^-} \right) - \frac{L_+}{z^{+2} z^-} \partial_t \frac{3B_{LL}^{++}}{4} + \frac{L_+}{z^{+2} z^-} \partial_t H_c + \frac{\nu + \eta}{L_+ z^-} \frac{3\partial_\sigma}{2\sigma^4} \left( \sigma^4 \partial_\sigma \frac{B_{LL}^{++}}{z^{+2}} \right). \quad (7)$$

In this way we arrive at a consistent scaling for the transfer term in Eq. (6) with the appropriate quantity, as the function  $C_{LL,L}^{+++}(r)$  scales with  $z^{+2} z^-$ .

Since the inverse of the coefficient in front of the dissipative term is similar to a Reynolds number, we introduce the generalized large-scale Reynolds number

$$R_- = \frac{2z^- L_+}{\nu + \eta}; \quad (8)$$

hence, Eq. (7) suggests a dependence of  $C_\varepsilon^+$  on  $1/R_-$ . However, the structure and correlation functions and the cross helicity flux also depend on the Reynolds number.

For conciseness we introduce dimensionless versions of all terms present on the right-hand side of Eq. (7), such that

$$C_{LL,L}^{+++}(r, t) = z^{+2} z^- g^{+++}(\sigma, t), \quad (9)$$

$$B_{LL}^{++}(r, t) = z^{+2} h_2^{++}(\sigma, t), \quad (10)$$

$$\partial_t B_{LL}^{++}(r, t) = \frac{(z^+)^2 z^-}{L_+} F^+(\sigma, t), \quad (11)$$

$$\partial_t H_c(t) = \frac{(z^+)^2 z^-}{L_+} G^+(t), \quad (12)$$

which leads to a dimensionless version of the  $\langle |z^+|^2 \rangle$  energy balance equation for freely decaying MHD turbulence

$$C_\varepsilon^+ = \frac{\varepsilon L_+}{z^{+2} z^-} = -\frac{\partial_\sigma}{\sigma^4} \left( \frac{3\sigma^4}{2} g^{++} \right) - \frac{3}{4} F^+ + G^+ + \frac{3}{R_-} \frac{\partial_\sigma}{\sigma^4} (\sigma^4 \partial_\sigma h_2^{++}). \quad (13)$$

After nondimensionalization, the highest derivative in the differential equation is multiplied with the small parameter  $1/R_-$ , suggesting that this can be viewed as a singular perturbation problem [28]; and thus we consider asymptotic expansions of the dimensionless functions in inverse powers of  $R_-$  [19,29].

The formal asymptotic series of a generic function  $f$  [used for conciseness in place of the functions on the right-hand side of Eq. (13)] up to second order in  $1/R_-$  reads

$$f = f_0 + \frac{1}{R_-} f_1 + \frac{1}{R_-^2} f_2 + O(R_-^{-3}). \quad (14)$$

After substitution of the expansions into Eq. (13) and following the same steps for the evolution equation for  $\langle |z^-|^2 \rangle$ , we arrive at model equations for  $C_\varepsilon^+$  and  $C_\varepsilon^-$

$$C_\varepsilon^\pm = C_{\varepsilon,\infty}^\pm + \frac{C^\pm}{R_\mp} + \frac{D^\pm}{R_\mp^2} + O(R_\mp^{-3}), \quad (15)$$

up to third order in  $1/R_\mp$ , where we defined the coefficients  $C_{\varepsilon,\infty}^\pm$ ,  $C^\pm$ , and  $D^\pm$ ,

$$C_{\varepsilon,\infty}^\pm = -\frac{\partial_\sigma}{\sigma^4} \left( \frac{3\sigma^4}{2} g_0^{\pm\mp\pm} \right) - \frac{3}{4} F_0^\pm \pm G_0^\pm, \quad (16)$$

$$C^\pm = \frac{3\partial_\sigma}{\sigma^4} \left[ \sigma^4 \left( \partial_\sigma h_{2,0}^{\pm\pm} - \frac{g_1^{\pm\mp\pm}}{2} \right) \right] - \frac{3}{4} F_1^\pm \pm G_1^\pm, \quad (17)$$

$$D^\pm = \frac{3\partial_\sigma}{\sigma^4} \left[ \sigma^4 \left( \partial_\sigma h_{2,1}^{\pm\pm} - \frac{g_2^{\pm\mp\pm}}{2} \right) \right] - \frac{3}{4} F_2^\pm \pm G_2^\pm, \quad (18)$$

in order to write Eq. (13) in a more concise way. Using  $R_+ = (L_-/L_+)(z^+/z^-)R_-$  to define

$$C = \frac{1}{2} \left( C^+ + \frac{L_- z^+}{L_+ z^-} C^- \right), \quad (19)$$

[ $D$  is defined analogously], finally one obtains for the dimensionless dissipation rate  $C_\varepsilon$ ,

$$C_\varepsilon = C_{\varepsilon,\infty} + \frac{C}{R_-} + \frac{D}{R_-^2} + O(R_-^{-3}). \quad (20)$$

Since the time dependence of the various quantities in this problem has been suppressed for conciseness, we stress that Eq. (20) is time dependent, including the Reynolds number  $R_-$ . A normalization using initial values of  $z^\pm$  and  $L^\pm$  would have resulted in a dependence of  $C_\varepsilon(t)$  on initial values of  $R_-$ , which only describe the initial conditions and not the evolved flow for which  $C_\varepsilon$  is measured.

At the peak of  $\varepsilon(t)$  the additional terms  $F_0^\pm$  should in fact vanish for constant flux of cross helicity (that is,  $\partial_t^2 H_c = 0$ ), since in the infinite Reynolds number limit the second-order structure function will have its inertial range form at all scales. By self-similarity the spatial and temporal dependences of, e.g.,  $B_{LL}^{++}$  should be separable in the inertial range, that is  $B_{LL}^{++}(r, t) \sim [e^+(t)r]^\alpha$  for some value  $\alpha$ , and  $\partial_t B_{LL}^{++} \sim \alpha e^+(t)^{\alpha-1} \partial_t e^+ r^\alpha$ . At the peak of dissipation  $\partial_t \varepsilon^+|_{t_{\text{peak}}} = \partial_t \varepsilon|_{t_{\text{peak}}} - \partial_t^2 H_c = \partial_t \varepsilon|_{t_{\text{peak}}} = 0$ , and we obtain  $F_0^+(t_{\text{peak}}) = 0$ . As the terms  $G_0^\pm$ , which describe the flux of cross helicity in the infinite Reynolds number limit, cancel the corresponding contribution from the transfer terms [27], the asymptotes  $C_{\varepsilon,\infty}^\pm$  describe the flux of total energy provided the model (15) is applied at  $t_{\text{peak}}$ .

Because of selective decay, that is the faster decay of the total energy compared to  $H_c$  and  $H_{\text{mag}}$  [14], one could perhaps expect  $\partial_t H_c$  to be small compared to  $\varepsilon$  in the infinite Reynolds number limit in most situations. In this case we obtain  $G_0^\pm \approx 0$  and

$$C_{\varepsilon,\infty}^\pm(t_{\text{peak}}) = -\frac{\partial_\sigma}{\sigma^4} \left( \frac{3\sigma^4}{2} g_0^{\pm\mp\pm} \right), \quad (21)$$

which recovers the inertial-range scaling results of Ref. [23] and reduces to Kolmogorov's 4/5th law for  $\mathbf{b} = 0$ .

Since  $C_{\varepsilon,\infty}$  is a measure of the flux of total energy across different scales in the inertial range, differences for the value of this asymptote should be expected for systems with different initial values for the ideal invariants  $H_c$  and magnetic helicity  $H_{\text{mag}} = \langle \mathbf{a} \cdot \mathbf{b} \rangle$ , where  $\mathbf{a}$  is the vector potential  $\mathbf{b} = \nabla \times \mathbf{a}$ . In the case of  $H_{\text{mag}} \neq 0$ , the value of  $C_{\varepsilon,\infty}$  should be *less* than for  $H_{\text{mag}} = 0$  due to a more pronounced reverse energy transfer in the helical case [13,30], the result of which is *less* forward transfer and thus a smaller value of the flux of total energy. For  $H_c \neq 0$  we expect  $C_{\varepsilon,\infty}$  to be smaller than for  $H_c = 0$ , since alignment of  $\mathbf{u}$  and  $\mathbf{b}$  weakens the coupling of the two fields in the induction equation, which leads to less transfer of magnetic energy across different scales and presumably

also less transfer of kinetic to magnetic energy. In short, one should expect nonuniversal values of  $C_{\varepsilon,\infty}$ .

Before we compare the model equation with DNS data and address this question of nonuniversality numerically, we briefly outline our numerical method. Equations (1)–(3) are solved numerically in a periodic box of length  $L_{\text{box}} = 2\pi$  using a fully de-aliased pseudospectral MHD code [33,34]. All simulations satisfy  $k_{\text{max}}\eta_{\text{mag,kin}} \geq 1$ , where  $\eta_{\text{mag,kin}}$  are the magnetic and kinetic Kolmogorov scales, respectively. We do not impose a background magnetic field, and both the initial magnetic and velocity fields are random Gaussian with zero mean, with initial magnetic and kinetic energy spectra of the form  $E_{\text{mag,kin}}(k) \sim k^4 \exp[-k^2/(2k_0)^2]$ , where  $k_0 \geq 5$  and further simulation details are specified in Table 1 of Ref. [27]. The initial relative magnetic helicity is  $\rho_{\text{mag}}(k) = kH_{\text{mag}}(k)/2E_{\text{mag}}(k) = 1$  for all runs of series H and zero for the runs labeled NH. The initial relative cross helicity was  $\rho_c(0) = H_c(0)/(|\mathbf{u}(0)||\mathbf{b}(0)|) = 0$  for runs of the H and NH series and  $\rho_c(0) = 0.6$  for series CH06H and CH06NH, while initial magnetic and kinetic energies were in equipartition. All spectral quantities have been shell and ensemble averaged, with ensemble sizes restricted by computational resources to up to 10 runs per ensemble. The total dissipation rate  $\varepsilon$  was measured at its maximum.

Figure 1 shows fits of the model equation to DNS data for data sets that differ in the initial value of  $H_{\text{mag}}$  and  $H_c$ . As can be seen, the model fits the data very well. For the series H runs and for  $R_- > 70$  it is sufficient to consider terms of first order in  $R_-$ , while for the series NH the first-order approximation is valid for  $R_- > 100$ . The cross-helical CH06H runs gave consistently lower values of  $C_\varepsilon$  compared to the series H runs, while little difference was observed between series CH06NH and NH. The




FIG. 1 (color online). The solid, dotted, and dash-dotted lines show Eq. (20) fitted to helical, nonhelical, and cross-helical DNS data, respectively. The red (gray) lines refer to fits using the first-order model equation, the black lines use the model equation up to second order in  $1/R_-$ . As can be seen, the respective asymptotes differ for the data sets.

asymptotes were  $C_{\varepsilon,\infty} = 0.241 \pm 0.008$  for the H series,  $C_{\varepsilon,\infty} = 0.265 \pm 0.013$  for the NH series,  $C_{\varepsilon,\infty} = 0.193 \pm 0.006$  for the CH06H series and  $C_{\varepsilon,\infty} = 0.268 \pm 0.005$  for the CH06NH series.

As predicted by the qualitative theoretical arguments outlined before, the measurements show that the asymptote calculated from the nonhelical runs is larger than for the helical case, as can be seen in Fig. 1. The asymptotes of the series H and NH do not lie within 1 standard error of one another. Simulations carried out with  $H_c \neq 0$  suggest little difference in  $C_\varepsilon$  for magnetic fields with initially zero magnetic helicity. For initially helical magnetic fields  $C_\varepsilon$  is further quenched if  $H_c \neq 0$ . In view of nonuniversality, an even larger variance of  $C_{\varepsilon,\infty}$  can be expected once other parameters such as external forcing, plasma  $\beta$ ,  $Pm$ , etc., are taken into account. Here we have restricted ourselves to nonuniversality caused by different values of vector field correlations.

In summary, a definition for the dimensionless dissipation rate  $C_\varepsilon$  for MHD turbulence has been proposed, where  $\varepsilon$  was nondimensionalized with respect to the Elsässer fields instead of the rms velocity. For this definition of  $C_\varepsilon$  and the case of the unit Prandtl number we derived a model for the dependence of  $C_\varepsilon$  on a generalized Reynolds number  $R_-$ . The model predicts that  $C_\varepsilon \rightarrow \text{const}$  with increasing  $R_-$ , in analogy to hydrodynamics, and the asymptote is a measure of the total energy transfer flux. The model was compared to DNS data for data sets that differ in their initial values of magnetic and cross helicities. At moderate to high  $R_-$ , we found good agreement with data with the model only using terms up to first order in  $1/R_-$ . However, at low  $R_-$  terms of second order in  $R_-$  cannot be neglected, in fact these terms improve the fit specifically at low  $R_-$ . This is expected from adding another term in the expansion and thus provides further justification of the validity of Eq. (20).

As predicted, the values of the respective asymptotes from the data sets differ, suggesting a dependence of  $C_\varepsilon$  on different values of the helicities, and thus a connection to the question of universality in MHD turbulence. This presents an interesting point for further research concerning the influence of other vector field correlations on the dissipation rate. Other questions concern the generalization of this approach to more general MHD systems such as flows with magnetic Prandtl numbers  $Pm \neq 1$ , compressive fluctuations, and to the presence of a background magnetic field, as well as to turbulent systems where the flow carries other quantities such as temperature or pollutants, and also the application to decaying hydrodynamic turbulence [20]. In the most general case in plasmas there will be a mean magnetic field, which leads to spectral anisotropy and the breakdown of the conservation of magnetic helicity [35] and thus might introduce several difficulties to be overcome when generalizing this method, as the spectral flux will then depend on the direction of the mean field [3,36] and a

more generalized description and role for the magnetic helicity would be needed.

Our model shows that different degrees of correlation in a turbulent plasma control the amount of energy that can effectively be transferred into the smallest scales. It could have several possible applications, e.g., for heating rates in the solar wind, especially as high values of the cross helicity inhibit such transfer to some extent. For situations where one is interested in sustaining a magnetic field over long times, thus trying to minimize dissipative effects, one could estimate from Eqs. (16)–(17) what type of correlations produce not only a low asymptotic value of the dissipation rate but also a fast approach to this asymptote. This would have relevance to cosmological and astrophysical [37] magnetic fields as well as terrestrial plasmas, such as in a tokamak reactor. Our results suggest that in cosmology, where a topical problem is the origin of large-scale magnetic fields, it is not only a nonzero value of magnetic helicity, but perhaps also the parameter range of other correlations such as the cross and kinetic helicities, that facilitate the presence of long-time magnetic fields. Moreover, this raises questions about the possible generation mechanisms for cosmological magnetic fields leading to different correlations between the vector fields such that they can sustain long evolution times.

The authors thank Samuel Yoffe for providing his hydrodynamic pseudospectral code for further development and the referees and the editor for insightful comments. This work has made use of the resources provided by ARCHER (<http://www.archer.ac.uk>), made available through the Edinburgh Compute and Data Facility (ECDF, <http://www.ecdf.ed.ac.uk>). A. B. is supported by STFC, M. F. L. and M. E. M. are funded by the UK Engineering and Physical Sciences Research Council (EP/K503034/1 and EP/M506515/1).

\* m.linkmann@ed.ac.uk

† ab@ph.ed.ac.uk

- [1] V. Dallas and A. Alexakis, *Phys. Fluids* **25**, 105106 (2013).
- [2] V. Dallas and A. Alexakis, *Phys. Rev. E* **88**, 063017 (2013).
- [3] M. Wan, S. Oughton, S. Servidio, and W. H. Matthaeus, *J. Fluid Mech.* **697**, 296 (2012).
- [4] A. A. Schekochihin, S. C. Cowley, and T. A. Yousef, in *IUTAM Symposium on Computational Physics and New Perspectives in Turbulence*, edited by Y. Kaneda (Springer, Berlin, 2008), p. 347.
- [5] P. D. Mininni, *Annu. Rev. Fluid Mech.* **43**, 377 (2011).
- [6] R. Grappin, A. Pouquet, and J. Léorat, *Astron. Astrophys.* **126**, 51 (1983).
- [7] A. Pouquet, P. Mininni, D. Montgomery, and A. Alexakis, in *IUTAM Symposium on Computational Physics and New Perspectives in Turbulence*, edited by Y. Kaneda (Springer, New York, 2008), p. 312.
- [8] A. Beresnyak, *Phys. Rev. Lett.* **106**, 075001 (2011).
- [9] S. Boldyrev, J. C. Perez, J. E. Borovsky, and J. J. Podesta, *Astrophys. J.* **741**, L19 (2011).
- [10] R. Grappin and W.-C. Müller, *Phys. Rev. E* **82**, 026406 (2010).
- [11] E. Lee, M. E. Brachet, A. Pouquet, P. D. Mininni, and D. Rosenberg, *Phys. Rev. E* **81**, 016318 (2010).
- [12] S. Servidio, W. H. Matthaeus, and P. Dmitruk, *Phys. Rev. Lett.* **100**, 095005 (2008).
- [13] A. Pouquet and G. S. Patterson, *J. Fluid Mech.* **85**, 305 (1978).
- [14] D. Biskamp, *Nonlinear Magnetohydrodynamics.*, 1st ed. (Cambridge University Press, Cambridge, England, 1993).
- [15] G. K. Batchelor, *The Theory of Homogeneous Turbulence*, 1st ed. (Cambridge University Press, Cambridge, England, 1953).
- [16] K. R. Sreenivasan, *Phys. Fluids* **27**, 1048 (1984).
- [17] K. R. Sreenivasan, *Phys. Fluids* **10**, 528 (1998).
- [18] W. D. McComb, *Homogeneous, Isotropic Turbulence: Phenomenology, Renormalization and Statistical Closures* (Oxford University Press, New York, 2014).
- [19] W. D. McComb, A. Berera, S. R. Yoffe, and M. F. Linkmann, *Phys. Rev. E* **91**, 043013 (2015).
- [20] J. C. Vassilicos, *Annu. Rev. Fluid Mech.* **47**, 95 (2015).
- [21] P. D. Mininni and A. G. Pouquet, *Phys. Rev. E* **80**, 025401 (2009).
- [22] V. Dallas and A. Alexakis, *Astrophys. J.* **788**, L36 (2014).
- [23] H. Politano and A. Pouquet, *Phys. Rev. E* **57**, R21 (1998).
- [24] S. Chandrasekhar, *Proc. R. Soc. London, Ser. A* **204**, 435 (1951).
- [25] J. J. Podesta, *J. Fluid Mech.* **609**, 171 (2008).
- [26] The scaling is ill defined for the (measure zero) cases  $\mathbf{u} = \pm \mathbf{b}$ , which correspond to exact solutions to the MHD equations where the nonlinear terms vanish. Thus no turbulent transfer is possible, and these cases are not amenable to an analysis which assumes nonzero energy transfer [23].
- [27] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.114.235001> for simulation details, definitions of the structure functions and details on some of the derivations.
- [28] W. Wasow, *Asymptotic Expansions for Ordinary Differential Equations* (John Wiley and Sons, New York, 1965).
- [29] T. S. Lundgren, *Phys. Fluids* **14**, 638 (2002).
- [30] Reverse transfer of magnetic energy has recently been discovered in nonhelical 3D MHD turbulence [31,32].
- [31] A. Brandenburg, T. Kahniashvili, and A. G. Tevzadze, *Phys. Rev. Lett.* **114**, 075001 (2015).
- [32] J. Zrake, *Astrophys. J.* **794**, L26 (2014).
- [33] S. R. Yoffe, Ph.D. thesis, University of Edinburgh, 2012, <http://arxiv.org/pdf/1306.3408v1.pdf>.
- [34] A. Berera and M. F. Linkmann, *Phys. Rev. E* **90**, 041003(R) (2014).
- [35] W. H. Matthaeus and M. L. Goldstein, *J. Geophys. Res.* **87**, 6011 (1982).
- [36] M. Wan, S. Servidio, S. Oughton, and W. H. Matthaeus, *Phys. Plasmas* **16**, 090703 (2009).
- [37] L. Sorriso-Valvo, R. Marino, V. Carbone, A. Noullez, F. Lepreti, P. Veltri, R. Bruno, B. Bavassano, and E. Pietropaolo, *Phys. Rev. Lett.* **99**, 115001 (2007).