



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

# What can MaxEnt reveal about high-density recordings and what can high-density recordings reveal about MaxEnt?

### Citation for published version:

Panas, D, Maccione, A, Berdondini, L & Hennig, M 2011, 'What can MaxEnt reveal about high-density recordings and what can high-density recordings reveal about MaxEnt?', *BMC Neuroscience*, vol. 12, no. Supplement 1, P146. <https://doi.org/10.1186/1471-2202-12-S1-P146>

### Digital Object Identifier (DOI):

[10.1186/1471-2202-12-S1-P146](https://doi.org/10.1186/1471-2202-12-S1-P146)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Publisher's PDF, also known as Version of record

### Published In:

BMC Neuroscience

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



POSTER PRESENTATION

Open Access


# What can MaxEnt reveal about high-density recordings and what can high-density recordings reveal about MaxEnt?

Dagmara Panas<sup>1\*</sup>, Alessandro Maccione<sup>2</sup>, Luca Berdondini<sup>2</sup>, Matthias H Hennig<sup>1</sup>

From Twentieth Annual Computational Neuroscience Meeting: CNS\*2011  
Stockholm, Sweden. 23-28 July 2011

Recent advances in neural recording techniques open exciting possibilities of better understanding whole populations of neurons. Devices such as APS MEA (Active Pixel Sensor Microelectrode Array) [1,2] allow for simultaneous recordings from 4096 channels (64x64

grid) at near-cellular resolution (electrode size: 21 $\mu$ m, electrode spacing: 42 $\mu$ m) and constitute a potentially very rich and detailed source of information on the dynamics of neural systems. Such volumes of data are however difficult to analyse: simple measures such as



**Figure 1** A Fraction of multiinformation captured by the MaxEnt model versus multiinformation (bits per bin, 5ms bins used). B Histograms of the values of fitted interaction parameters.

\* Correspondence: D.Panas@sms.ed.ac.uk

<sup>1</sup>Institute for Adaptive and Neural Computation, School of Informatics,  
University of Edinburgh, EH8 9AB, UK

Full list of author information is available at the end of the article

mean firing rates and correlations are often insufficient to capture interesting phenomena, while more sophisticated approaches can be computationally intensive and hard to interpret. Here we examine the applicability of pairwise maximum entropy (MaxEnt) [3-5] modelling to describe APS MEA data.

Pairwise maximum entropy model (equivalent to Ising model in physics), when fit to the data, yields a minimally structured probability distribution of network states that respects first and second order interactions. It is a convex, parsimonious and readily interpretable model that has been shown to characterize spiking patterns surprisingly robustly in many cases [3,4]. Additionally, it can provide a sensitive tool in detecting higher-order interactions. As reported in [5], the significant failure of the Ising model in close range (<300  $\mu\text{m}$ ) uncovers a high-order processing mode in local clusters of neurons, a mode of processing absent on larger scale (>600  $\mu\text{m}$ ) and undetectable with correlations.

In present work we examine the results and performance of the MaxEnt model fitting in different preparation types and parameter regimes; owing to high resolution recordings we can specifically focus on varying spatial scales. As can be seen in Fig.1, indeed even in cultured tissue data there are indicators of certain discrepancies between local populations and populations further apart. Firstly (panel A), it is in local populations where the advantage of Ising model over the independent model is most prominent. Secondly (panel B), the interactions within local populations reveal a different structure than those among groups of neurons spread further apart (Kolmogorov-Smirnov test,  $p < 0.05$ ); and, importantly, this is not a feature that can be shown by correlation analysis.

#### Acknowledgements

This work was funded by EPSRC and BBSRC as part of Doctoral Training Centre programme at School of Informatics, University of Edinburgh(DP); MRC Fellowship G0900425 (MHH); and IDEA, FP6-NEST Grant 516432 (AM and LB).

#### Author details

<sup>1</sup>Institute for Adaptive and Neural Computation, School of Informatics, University of Edinburgh, EH8 9AB, UK. <sup>2</sup>Department of Neuroscience and Brain Technologies, Italian Institute of Technology, 16163 Genova, Italy.

Published: 18 July 2011

#### References

1. Imfeld K, Neukom S, Maccione A, Bornat Y, Martinoa S, Farine PA, Koudelka-Hep M, Berdondini L: **Large-scale, high-resolution data acquisition system for extracellular recording of electrophysiological activity.** *IEEE Trans Biomed Eng* 2008, **55**:2064-2073.
2. Berdondini L, Imfeld K, Maccione A, Tedesco M, Neukom S, Koudelka-Hep M, Martinoa S: **Active pixel sensor array for high spatio-temporal resolution electrophysiological recordings from single cell to large scale neuronal networks.** *Lab Chip* 2009, **9**:2644-2651.
3. Schneidman E, Bialek W: **Ising models for networks of real neurons.** *Nature* 2006, **440**:1007-1012.
4. Tang A, Jackson D: **A Maximum Entropy Model Applied to Spatial and Temporal Correlations from Cortical Networks In Vitro.** *J Neurosci* 2008, **28**(2):505-518.
5. Ohiorhenuan IE, Mechler F, Purpura KP, Schmid AM, Hu Q, Victor JD: **Sparse coding and high-order correlations in fine-scale cortical networks.** *Nature* 2010, **466**:617-621.

doi:10.1186/1471-2202-12-S1-P146

**Cite this article as:** Panas *et al.*: What can MaxEnt reveal about high-density recordings and what can high-density recordings reveal about MaxEnt? *BMC Neuroscience* 2011 **12**(Suppl 1):P146.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

